



Contents lists available at [ScienceDirect](http://www.sciencedirect.com)

# International Journal for Parasitology: Parasites and Wildlife

journal homepage: [www.elsevier.com/locate/ijppaw](http://www.elsevier.com/locate/ijppaw)



Invited review

## Do managed bees drive parasite spread and emergence in wild bees?



Peter Graystock <sup>a, \*</sup>, Edward J. Blane <sup>b</sup>, Quinn S. McFrederick <sup>a</sup>, Dave Goulson <sup>c</sup>,  
William O.H. Hughes <sup>c</sup>

<sup>a</sup> Department of Entomology, University of California, Riverside, CA 92507, USA

<sup>b</sup> Natural England, Mail Hub Block B, Whittington Road, Worcester, WR5 2LQ, UK

<sup>c</sup> School of Life Sciences, University of Sussex, Brighton, BN1 9QG, UK

### ARTICLE INFO

**Article history:**  
Received 27 July 2015  
Received in revised form  
27 September 2015  
Accepted 1 October 2015

**Keywords:**  
Domesticated  
Apis  
Commercial  
Bumble bee  
Honey bee  
Bombus  
Farming  
Intensification  
Pollination service  
Disease

### ABSTRACT

Bees have been managed and utilised for honey production for centuries and, more recently, pollination services. Since the mid 20th Century, the use and production of managed bees has intensified with hundreds of thousands of hives being moved across countries and around the globe on an annual basis. However, the introduction of unnaturally high densities of bees to areas could have adverse effects. Importation and deployment of managed honey bee and bumblebees may be responsible for parasite introductions or a change in the dynamics of native parasites that ultimately increases disease prevalence in wild bees. Here we review the domestication and deployment of managed bees and explain the evidence for the role of managed bees in causing adverse effects on the health of wild bees. Correlations with the use of managed bees and decreases in wild bee health from territories across the globe are discussed along with suggestions to mitigate further health reductions in wild bees.

© 2015 The Authors. Published by Elsevier Ltd on behalf of Australian Society for Parasitology. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

### Contents

1. Introduction .....	65
2. Parasite facilitation .....	65
3. Parasite spillover .....	65
4. Parasite spillback .....	65
5. Is spillover/spillback between wild and managed bees common? .....	65
6. Success and domestication of pollinators .....	66
7. Honey bee domestication .....	66
8. Bumblebee domestication .....	66
9. Reports of possible spillover from managed bumblebee or honey bee populations .....	67
9.1. Case study 1: transmission between managed and wild honey bees globally .....	67
9.2. Case study 2: transmission between managed and wild bumblebees in Japan .....	68
9.3. Case study 3: transmission between managed and wild bumblebees in North America .....	68
9.4. Case study 4: transmission between managed and wild bumblebees in the United Kingdom and Ireland .....	69
9.5. Case study 5: transmission between managed honey bees and wild bumblebees in Europe .....	70
10. Conclusion .....	70
10.1. Mitigating the harmful effects of managed bees on wild bees .....	71
Acknowledgements .....	72
References .....	72

\* Corresponding author.  
E-mail address: [peter@graystock.info](mailto:peter@graystock.info) (P. Graystock).

## 1. Introduction

Growing demand for food and agricultural intensification is resulting in a growing demand for pollination services and, ironically, growing pressures on wild bees. Currently both managed and wild pollinators are suffering alarming declines in many parts of the world (Potts et al., 2010). In the last century, many wild bee populations have become reduced and fragmented, and with reduced genetic diversity are now facing increased vulnerability to infectious diseases and other stressors such as pesticides and reduced resources (McCallum and Dobson, 2002; Tarpy, 2003; Whitehorn et al., 2011). Over 25% of wild bees in Belgium and 47% of native bumblebee species in Hungary have exhibited recent declines (Sárospataki et al., 2005). Similar declines in wild pollinators are found in Britain, Ireland, Spain, France, Morocco, Corsica, Tibet, Brazil and Madagascar (Fitzpatrick et al., 2007; Kosior et al., 2007; Martins and Melo, 2010; Rasmont et al., 2005; Williams, 1982; Xie et al., 2008). In the North American mid-west, half of the native bumblebee species have declined during the mid-twentieth century (Grixti et al., 2009), while bumblebees have also declined in diversity, evenness and abundance in eastern North America between 1971 and 2006 (Colla and Packer, 2008). Today 11% of all bumblebee species worldwide are listed as threatened in the IUCN Red list (Williams and Osborne, 2009).

Emergent infectious diseases (EIDs) are ranked as one of the top five causes of species extinction worldwide (Daszak et al., 2000). They are defined broadly as diseases that have recently increased in either incidence, demographic or host range, or that have recently evolved or been discovered (Daszak et al., 2001, 2000; Lederberg et al., 1992; Morse, 1993). Many instances of EIDs in wild animals are the result of interactions with domesticated/managed species (Fig. 1). One of the main conservation concerns regarding the use, and particularly the importation, of managed bees, is that of exotic parasite (or parasite strain) introduction and subsequent spillover to wild populations (Blitzer et al., 2012; Goulson et al., 2008). By allowing managed and/or imported bees to mix with wild pollinators, there is the potential for disease emergence via direct transmission and facilitated by changes in host susceptibility.




Fig. 1. The key factors that may drive disease emergence within and between populations of managed and wild bees. Adapted from Daszak et al. (2000).

## 2. Parasite facilitation

As with other animals, bee susceptibility and subsequent pathology is often increased when individuals are stressed (Freestone et al., 2008; Jokela et al., 2005; MacKinnon, 1998; Møller et al., 1998). Dense populations of managed bees can cause stress to wild populations and have been linked to developmental retardation in wild bees (Elbgami et al., 2014; Goulson and Sparrow, 2008; Thomson, 2004). It's also likely that, as in other organisms, such competitive and nutritional stresses can influence their susceptibility to infection (Brown et al., 2003; Graystock et al., 2014; Hedtke et al., 2011; MacKinnon, 1998). This stress would therefore facilitate an increase in prevalence or intensity of natural parasite infections in wild bees, independent of any transmission between the two groups of bees.

## 3. Parasite spillover

Parasite spillover occurs when the reservoir population (usually managed stock) transmit disease to the sink (in this case, wild) population (Daszak et al., 2000; Fig. 2). Emergence of the parasite in wild populations may be self-supporting or may require continued mixing with the source/reservoir population. When intra-specific and/or inter-specific transmission rates are low, infection of susceptible individuals may occur in many instances without the parasite establishing in the wild population (Hatcher and Dunn, 2011). Closely related, sympatric hosts have a greater potential to transmit pathogens between them, though the virulence of the parasite may differ (Graystock et al., 2013a; Perlman and Jaenike, 2003; Schmid-Hempel, 2011).

## 4. Parasite spillback

In parasite spillback, a natural parasite infection in the wild host population transmits to the managed host population, increases in prevalence within it, and then spills back into the wild population, resulting in an increase in prevalence in wild populations (Hatcher and Dunn, 2011; Kelly et al., 2009; Fig. 2). Here, the newly infected animals may achieve unnaturally high parasite prevalence due to their high density (for example, honey bee colonies in apiaries are much denser than in wild honey bee populations). This unnaturally high parasite prevalence then spills back, driving parasite levels in the wild population higher than would naturally occur (Hatcher and Dunn, 2011; Kelly et al., 2009). Another factor in spillback from managed populations is that well managed, healthy animals may be able to tolerate pathogenic effects better than wild hosts. Such tolerance may result in a population of asymptomatic, yet infectious, managed bees.

## 5. Is spillover/spillback between wild and managed bees common?

Over the past century there have been huge advancements in agricultural intensification, combined with a rapid global demand for food (Aizen and Harder, 2009; Aizen et al., 2009; Gallai et al., 2009; Velthuis, 2002). One of the advances farmers increasingly utilise is the purchase of pollination services via managed honey bees or managed bumblebee colonies (VanEngelsdorp and Meixner, 2010; Velthuis and Van Doorn, 2006). The practices and regulations in place for managed bee use are not uniform across the globe and there is growing concern that managed bees may pose a disease risk to other pollinators (Goulson and Hughes, 2015; Graystock et al., 2014; Meeus et al., 2011). Here we review the domestication of bumblebees and honey bees before looking at specific examples from across the world where parasite




Fig. 2. Highlighting the three main mechanisms that influence parasite infections between managed and wild bee populations. Arrows represent direction of potential parasite spread as a result of the mechanism.

transmission is reported to have occurred following the use of managed bees for crop pollination. Finally, we consider the parasite threat that managed honey bees or managed bumblebees may pose to wild bees, highlighting ways to reduce this threat and possible avenues for further research.

## 6. Success and domestication of pollinators

In Europe, the yields of 84% of 264 crop species rely on, or are improved by insect pollination (Klein et al., 2007; Williams, 1994). Modern farming is increasingly intensive, creating large areas with limited proximity to pollinator-rich areas such as wild meadows or hedgerows. When flowering crops are used in such fields, they can suffer from suboptimal crop production due to the fields and surrounding area providing few or no habitat for pollinators such as bumblebees to nest (Free and Williams, 2009; Kremen et al., 2002). Additionally, areas can be planted in such densities that wild pollinator populations are unable to service all crops. To increase crop yields, it is therefore beneficial to boost pollinators numbers by utilizing managed bees (Lye et al., 2011a). Honey bees (*Apis*) and bumblebees (*Bombus*) are the two groups of bees which have been most successfully domesticated and used in crop pollination (VanEngelsdorp and Meixner, 2010; Velthuis, 2002).

## 7. Honey bee domestication

Out of about 20,000 known bee species only 7 of these are honey bees (Arias and Sheppard, 2005; Engel, 1999; Koeniger and Koeniger, 2000). All but one honey bee species occur only in Asia, while the other, *Apis mellifera*, has a natural range from central Asia into Europe and Africa (Ruttner, 1988; Seeley, 1985; Sheppard and Meixner, 2003). Managed for their honey since at least 2600 BC, the economic value of honeybees has ensured their translocation along with every large-scale human migration (Crane, 1999, 1975;

Ransome, 2004; VanEngelsdorp and Meixner, 2010). This movement across the world has caused some conservation concerns; the introduction of managed honey bees has been found to cause changes in seed set of natural floral and reduce the fitness of native pollinators (Goulson and Derwent, 2004; Goulson and Sparrow, 2008; Goulson, 2003; Hurn, 1997). Despite these concerns, *A. mellifera* is now the most commonly managed pollinator and is found in nearly all habitable regions (Engel, 1999; Ransome, 2004; VanEngelsdorp and Meixner, 2010). Fifty-two of the top 115 global food commodities depend on honey bee pollination for either fruit or seed set (Klein et al., 2007), with honey bees being capable of increasing the yields of 96% of animal-pollinated crops, making them the most important pollinator for most crops worldwide (Delaplane et al., 2000; Klein et al., 2007; McGregor, 1976). Recently it has been estimated that projected honey bee populations will not be able to satisfy agricultural pollination demands in the future (Aizen and Harder, 2009).

## 8. Bumblebee domestication

For many wildflowers and valuable crops such as raspberry and tomatoes, honey bees are inefficient pollinators (Batra, 1995; Cane, 2005; Greenleaf and Kremen, 2006; Velthuis and Van Doorn, 2006). Bumblebees have several adaptations such as the ability to buzz pollinate (sonication) and their insulated bodies making them ideal pollinators for these plant and crop species (De Luca et al., 2013; Goulson, 2010; Heinrich, 1993; Velthuis and Van Doorn, 2006). Unlike honey bees, bumblebees do not produce honey or hive material suitable for mass market sales. Their commercial use is purely based on demand for their pollination services where honey bees are not the most efficient pollinator (VanEngelsdorp and Meixner, 2010).

Prior to bumblebee domestication, crops requiring sonication such as tomatoes were mechanically vibrated to achieve

pollination. The cost of mechanically vibrating the plants was over €10,000 per ha per year in 1988 (Velthuis and Van Doorn, 2006). Like honey bees, bumblebees have more recently been translocated by humans to aid crop production, even before their true 'domestication'. Between 1882 and 1906, hundreds of bumblebee queens were translocated from the UK to New Zealand to establish wild colonies and improve the seed set of red clover, with four UK species (*Bombus hortorum*, *Bombus ruderatus*, *Bombus subterraneus* and *Bombus terrestris*) all becoming naturalised in New Zealand as a result (Hopkins, 1914; Lye et al., 2011b). Following this agricultural success in New Zealand, *B. ruderatus* queens from the naturalised New Zealand stock were translocated to Chile for their red clover pollination in 1982 and 1983 (Arretz and Macfarlane, 1986).

By the 1960's there was a clear demand to utilise bumblebees as pollinators on an intensive scale, which motivated their domestication (Free, 1970; Holm, 1966; Velthuis and Van Doorn, 2006; Velthuis, 2002). Unlike honey bees, bumblebees require diapause initiation after mating and then need to be stimulated to lay eggs following emergence (Goulson, 2010). Various protocols were developed to achieve this, such as the inclusion of honey bee workers to stimulate egg laying by bumblebee queens (Heemert et al., 1990; Ptacek, 1985; Van der Eijnde, 1990). By the late 1970's enough progress had been made to fully domesticate the most common European bumblebee species, *B. terrestris* (Röseler, 1977; Velthuis and Van Doorn, 2006). With the joining of both demand and technology, commercial production of bumblebee colonies began in 1987 and by 1989 three companies based in Belgium and the Netherlands had begun production (Velthuis and Van Doorn, 2006). This growing industry was estimated to be worth in excess of €55 million pa in 2006 and is served by over 30 factories (Velthuis and Van Doorn, 2006). Globally, over two million bumblebee colonies are reared and transported to various countries for farm placement. Placement of hives in fields are recommended to be at near natural densities of 1 per km<sup>2</sup> but farmers frequently buy more and site them in vertical 'stacks' creating zones of incredibly high density (6 colonies in 1 m<sup>2</sup>). Although the most utilised species in commercial colonies is *B. terrestris*, several others have been domesticated/managed including *Bombus lucorum*, *Bombus occidentalis*, *Bombus ignitus*, *Bombus canariensis* and *Bombus impatiens* (Asada and Ono, 1997; Velthuis and Van Doorn, 2006; Wu, 2008). This diversity is an attempt to mitigate environmental damage from the escape of exotic pollinators by the production of native stock for various countries (Dafni et al., 2010; Velthuis and Van Doorn, 2006), although non-native species or subspecies are still used in many instances (Velthuis and Van Doorn, 2006). In order to produce enough bumblebee colonies commercially to meet worldwide demand, it is estimated that around 500 tonnes of pollen is required to feed the colonies in the rearing facilities (Goulson, 2013). This pollen is harvested by honey bees and collected from their pollen baskets using 'pollen traps'. The bumblebee industry is therefore reliant on the honeybee market with Spain and China being key suppliers of pollen (Centre for the Promotion of Imports from developing countries, 2009).

## 9. Reports of possible spillover from managed bumblebee or honey bee populations

While this review is focused on the parasite threats from managed bees, other concerns have also been raised over the use of managed bees including non-native naturalisation, competition and changes in floral seed set (Goulson, 2003; Goulson et al., 2002; Ings et al., 2006; Kremen et al., 2002). Many of these other threats can be exacerbated by parasite emergence (Daszak and Cunningham, 1999; Jones et al., 2008). When considering the parasite risks that managed bees pose to wild bees, it is important

to note that spillover can act in either direction since both managed and native populations can act as a parasite reservoir. Transmission between different managed species could then allow unexpected parasites to spillover to wild bees (Graystock et al., 2013a; McMahon et al., 2015). In cases where parasites from wild bees transmit to managed bees, it is important to also look for signs of possible spillback (Kelly et al., 2009). To identify spillover and spillback, studies must account for: 1) whether pathogens have emerged in tandem with managed bee introduction, 2) if the dynamics of pathogen spread indicate that the health of wild bees near managed bees is different from areas without managed bees, and 3) ideally trace pathogen origin with genetic markers (Daszak et al., 2000; Kelly et al., 2009; Power and Mitchell, 2004; Wood et al., 2012). Frequently all three are lacking so identification of the type of parasite dynamics occurring cannot be fully reconciled. Despite this, the balance of evidence on a case-by-case basis can allow for a reasonable verdict. On this premise, evidence for managed bees causing harm to wild bees is presented below as a series of case studies.

### 9.1. Case study 1: transmission between managed and wild honey bees globally

The transmission of parasites to the European honey bee from their original host, the Asian honey bee, led to some of the most dramatic honey bee declines in recorded history. Two parasites are particularly noteworthy: the ectoparasitic *Varroa* mite, and the microsporidian *Nosema ceranae* (Chauzat et al., 2010; Neumann and Carreck, 2010; vanEngelsdorp et al., 2011).

*Varroa*: The ectoparasitic mite *Varroa jacobsoni* is native to Eastern Asia where it parasitizes Asian honey bees such as *Apis ceranae* (Oldroyd, 1999; Oudemans, 1904). Some time in the mid 20th century, *Varroa* spilled over from its natural host to managed colonies of *A. mellifera* that had been transported into Asia by apiculturists for honey production and pollination services (Rosenkranz et al., 2010). This host jump, determined by genetic changes in the *Varroa* population, resulted in a new species, *Varroa destructor* (Anderson and Trueman, 2000; Dietemann et al., 2012; Solignac et al., 2005). This occurred on a large scale at least twice, forming the Japanese and Korean strains (Anderson and Trueman, 2000). The first wave of infection started by *Varroa* jumping from *Apis cerana* to *A. mellifera* in Japan in 1957 then spreading on to Paraguay in 1971, Brazil in 1972 and then within the decade on to North America (Jong et al., 1982; Sakai and Okada, 1974; Solignac et al., 2005). The second wave of infection occurred when *Varroa* jumped from *A. cerana* to *A. mellifera* in the region of Primorsky Krai, Russia, then spread through the *A. mellifera* populations in Europe before being recognised as a globally emerging parasite (Crane, 1978; Solignac et al., 2005). This rapid spread was made possible by the transport of managed bees across countries and between continents, allowing *Varroa* mites from infested hives to spillover to new hives in new areas (Boecking and Genersch, 2008; Bowen-Walker et al., 1999; Solignac et al., 2005).

Following their spillover to *A. mellifera*, dramatic honey bee declines were reported in both managed and wild colonies with a host of symptoms resulting from infestation (Carreck et al., 2002; Le Conte et al., 2010; Martin et al., 2012). Honey bees from *Varroa* infected colonies are smaller and become cognitively impaired (Kralj and Fuchs, 2006; Kralj et al., 2007; Rosenkranz et al., 2010), their hives produce less honey (Emsen et al., 2013), and only survive 2–3 years (Fries et al., 2006a; Korpela et al., 1993). *Varroa* mites feed off the hemolymph of host bees, and are therefore excellent vectors of virus parasites. This direct injection of virus into the haemolymph of the host results in more intense infections with pathogenic effects, known as Varroosis (Boecking and Genersch,

2008; Dainat et al., 2012). While good bee husbandry can help reduce colony losses, the *Varroa* mite is now considered to be a significant contributor to *A. mellifera* losses worldwide, particularly of feral and wild honey bee populations which in the USA and Europe are thought to have been significantly reduced following the emergence of *Varroa* (Martin et al., 2012; Pohorecka et al., 2011). In the UK mainland and parts of Europe, it is believed that *Varroa* aided declines have resulted in the loss of truly wild populations of *A. mellifera*, with only feral colonies left (Jaffé et al., 2010; Moritz et al., 2007; Thompson et al., 2014).

*Nosema ceranae*: It is believed that the microsporidian *Nosema ceranae* was first detected in the Asian honey bee, *Apis ceranae*, in 1975, although earlier misidentifications may have occurred (Fries et al., 2006b; Singh, 1975). It was only in 1996 that it was formally described as *N. ceranae* (initially misidentified as *Nosema apis*), from samples of managed *A. cerana* in China (Fries, 2010; Fries et al., 1996). Cross-infectivity experiments showed *N. ceranae* to be infective to the European honey bee *A. mellifera* as well as its natural host (Fries, 1997). Symptoms of *N. ceranae* infections have been termed Nosemosis type C. They are stronger in younger bees, and include suppressed immune function, increased hunger and increased mortality, causing *N. ceranae* to be considered a contributing factor to colony losses in some, but not all, areas (Antúñez et al., 2009; Higes et al., 2010; Mayack and Naug, 2009; Roberts and Hughes, 2014).

In 2005, samples of *A. mellifera* from an apiary in Taiwan that also housed *A. cerana* honey bees were found to be infected with *N. ceranae*, suggesting spillover from its natural host could be occurring (Huang et al., 2007). This was also confirmed in Europe when *N. ceranae* was found in an *A. mellifera* apiary in Spain (Higes et al., 2006). Historical bee samples have shown *N. ceranae* started to spillover to *A. mellifera* earlier than this but an exact date is not yet resolved (Fries, 2010). It has been detected in *A. mellifera* collected before 1990 in Uruguay (Invernizzi et al., 2009), in North America from 1995 (Chen et al., 2008), and in Spain from 2000 (Botías et al., 2012). Temporal analysis shows that *N. ceranae* introduction, spread and prevalence has increased over time (Klee et al., 2007; Paxton et al., 2007). Today, *N. ceranae* is present on all continents where *A. mellifera* is present and has been found at similar prevalence in managed and nearby wild honey bees (Giersch et al., 2009; Klee et al., 2007; Thompson et al., 2014).

### 9.2. Case study 2: transmission between managed and wild bumblebees in Japan

The first reported instance of pathogen transmission between commercially reared bumblebees and wild bumblebees was in Japan. Commercial *B. terrestris* have been imported into Japan primarily for tomato production since 1991 (Goka et al., 2001). Within a decade, over 40,000 colonies of non-native *B. terrestris* were being imported annually into Japan from Europe (Velthuis and Van Doorn, 2006). On average, 20% of the commercially produced colonies of *B. terrestris* examined were found to be infected with the European strain of the tracheal mite *Locustacarus buchneri* (Goka et al., 2001, 2000). In a separate study, a viable microsporidian parasite (resembling *N. bombi*) was also detected in a commercially produced *B. terrestris* colony imported into Japan (Niwa et al., 2004). At the same time, a number of bumblebee species native to Japan, including *B. ignitus*, were taken back to Europe in an attempt to commercialise them. This process carried the risk of European parasites, such as *L. buchneri*, infecting the Japanese bumblebees in the production facilities and then being carried back with them into Japan.

High mite incidence in managed bumblebee colonies is a particular concern as the mite feeds, reproduces, and lives inside

the abdominal air sacks (trachea) of female bumblebees (Yoneda et al., 2008). Heavily infected bumblebees suffer from diarrhoea and lethargy, stop foraging, and have a reduced lifespan, thus retarding colony growth (Husband and Sinha, 1970; Otterstatter and Whidden, 2004). This reduces the ability of the commercially produced bumblebees to pollinate the intended crops efficiently and it could negatively impact wild bumblebees if spillover was to occur. Due to this concern, between 1997 and 1999 three native bumblebee species, *Bombus hypocrita sapporoensis*, *B. hypocrita hypocrita* and *B. ignitus*, were screened across Japan with 8%, 1% and 0% respectively being found to be infected with the mite (Goka et al., 2001). Molecular analysis suggested that spillover of *L. buchneri* from managed bumblebees to wild bumblebees had not at that time occurred. However, the fact that 20% and 17% of commercially produced *B. terrestris* and *B. ignitus* respectively were infected with the mite, together with the escape and naturalisation of commercially produced *B. terrestris* in Japan, and the finding that the European haplotype of *L. buchneri* was able to parasitize the native *B. ignitus*, suggested that the potential for spillover from bumblebees imported into Japan was high (Goka et al., 2001; Kondo et al., 2009). A similar study, run between 2000 and 2001 found wild *B. hypocrita* bumblebees infected with European mites, while and commercially produced bumblebees were infected Japanese mites, showing that spillover had occurred in both directions (Goka et al., 2006). By 2001, commercially produced bumblebees were no longer found to have *L. buchneri* infections and there are no subsequent reports of *L. buchneri* in commercially produced bumblebees in Japan. Studies in Poland also determined commercial bumblebees to be free from mites now, however a recent study did find them in south America (Reade et al., 2014; Rozej et al., 2012; Sachman-Ruiz et al., 2015). The more rigorous, and apparently successful, control of the mite in stocks exported to Japan was however, too late to prevent the spillover of the parasite into Japan (Goka et al., 2006). It is not known whether there are still mites of European descent circulating in wild bumblebees in Japan, or if the spillover was transient and dependent on continued introduction from managed bees.

### 9.3. Case study 3: transmission between managed and wild bumblebees in North America

In 1992, the Animal and Plant Health Inspection Service (APHIS) in North America granted a permit for the importation of native bumblebees *B. occidentalis* and *B. impatiens* from Europe (Wehling and Flanders, 2005). These North American bees had been reared in European facilities along with European *B. terrestris* and likely fed pollen collected from European honey bees (Velthuis and Van Doorn, 2006; Winter et al., 2006). While their importation back into North America was under condition of parasite screening, the methods used had limited sensitivity (Ghosh and Weiss, 2009; Klee et al., 2006; Ndao, 2009; Procop, 2007; Rinder et al., 1998; Yang and Rothman, 2004). Following concerns from conservationists, importation was stopped and the use of *B. occidentalis* was restricted to its native range in Western North America, with the use of *B. impatiens* being restricted to its range in Eastern North America (Wehling and Flanders, 2005). As a result, breeding lines of both species were maintained together in new facilities in Canada and the United States.

In 1997, *B. occidentalis* bumblebees in production factories in California started suffering from high levels of the microsporidian parasite *Nosema bombi* and stocks plummeted (Wehling and Flanders, 2005). This coincided with unusually high incidences of *N. bombi* in some wild bumblebee populations in North America and rapid declines in some species (including *Bombus affinis*, *B. occidentalis*, *B. pensylvanicus* and *Bombus terricola*; Flanders et al.,

2003; Thorp, 2005; Thorp and Shepherd, 2005; Winter et al., 2006). Bumblebees infected by *Nosema bombi* can become sluggish, have increased mortality, reduced egg production, and infected colonies tend to be smaller and produce fewer reproductives (Fantham and Porter, 1914; Otti and Schmid-Hempel, 2008, 2007; Rutrecht and Brown, 2008; Schmid-Hempel and Loosli, 1998; Van der Steen, 2008; Whittington and Winston, 2003). Molecular analysis on a small number of samples suggested that *N. bombi* found in declining wild bumblebees may have been from a recent introduction, suggesting its emergence via importation with commercially reared bumblebees (Cameron et al., 2011; Winter et al., 2006). The concurrent timing of the declines of the wild and managed bumblebee populations in North America is suggestive of the same factor being involved, with *N. bombi* then being the best candidate.

This evidence, however, is only correlational. *N. bombi* is known to have been present in North American bumblebees prior to the introduction of commercially produced bumblebees, and recent molecular analysis suggests isolates of *N. bombi* with both American and European origin may exist in North American bumblebee species (Cordes et al., 2012). Declining populations can be more prone to parasite infections, so cause and effect cannot always be clearly disentangled on the basis of correlational data alone (Brown, 2011). Alternatives to the spillover of European *N. bombi* into North American bumblebees are for American *N. bombi* to have spilled over into commercially produced bumblebees and then spilled back into wild bumblebees, or for the use of commercially produced bumblebees to have facilitated an increase in *N. bombi* levels in wild bumblebees due to increased competitive stress or parasite vectoring, or for levels in managed and wild bumblebees to have increased independently due to some other causal factor(s). Two later studies found that the prevalence of infections with the *N. bombi* and *Crithidia bombi* parasites was higher in wild bumblebees caught near sites using commercially produced bumblebees than in wild bees caught 2 km or further from such sites (Colla et al., 2006; Otterstatter and Thomson, 2008), suggesting that spillover, spillback or facilitation was occurring. A further meta-analysis of all possible causes of bumblebee declines in North America found that the declines of *B. terricola* and *Bombus pennsylvanicus*, but not *B. affinis*, were associated with the use of commercially produced bumblebees, making them the most probable cause for the declines of these species (Szabo et al., 2012).

In addition, in recent years there are several reports of managed bumblebees delivered to North America, being found with parasites including *L. buchneri*, *N. bombi*, *Apicystis bombi* and the viral pathogens deformed wing virus (DWV), black queen cell virus (BQCV), Israeli acute paralysis virus (IAPV), Chronic bee paralysis virus, and Kashmir bee virus (Sachman-Ruiz et al., 2015). The presence of these parasites upon delivery carries with it the potential for disease emergence and spillover with any interacting wild bees in the area.

#### 9.4. Case study 4: transmission between managed and wild bumblebees in the United Kingdom and Ireland

Managed *B. terrestris dalmatinus/terrestris* started being imported into the British Isles from Europe in 1989 (Ings et al., 2006; Velthuis and Van Doorn, 2006). In 2008, 74% of 68 (as yet unopened) commercially produced *B. terrestris* colonies examined (imported from two different suppliers) were found to be infected with either *C. bombi*, *N. bombi*, or *A. bombi* (Murray et al., 2013). In addition, bumblebees captured at six sites in Ireland with a history of using managed bumblebees were found to have elevated levels of parasite infection (Murray et al., 2013). The presence of parasites in hives upon delivery plus the increased prevalence of those parasites in conspecifics suggests parasite spillover taking place in Eastern Ireland (Murray et al., 2013). In 2010, following a

clarification of section 14 of the Wildlife and Countryside Act 1981, the release of non-native bumblebees subspecies required a release licence that was only issued if the bees were 'disease-free'. This screening of the bees was self-regulated by the producers in accordance with the Natural England licence conditions (Central Office of Information, 2009; Natural England, 2009). These conditions initially required stocks in the rearing facilities to adhere to a minimum of microscopy screening for *A. bombi*, *C. bombi*, and *N. bombi* of 2 workers from each 200 colonies every 10 weeks, with a minimum sample size of over 2000 per year. On the same sample size, bees were visually inspected for wing deformities and newly collected queens were to be visually screened for *Sphaerularia bombi*. In addition, at least routine visual inspection was required for *L. buchneri* (following dissection under binocular microscope), *Melittobia acasta* and small hive beetle (Natural England licencing conditions NNR/2007/30–33). Although not the most sensitive screening methods, the breadth of parasites screened for, particularly those specific to bumblebees was far superior to the honey bee-centric conditions of the European directive 92/65/EEC on imports.

By 2010, the screening requirements of non-native subspecies imports were somewhat undermined when commercial bumblebee breeders began importing the native subspecies *B. t. audax* into the UK from rearing facilities in continental Europe. This had been done to meet concerns from Natural England about the release of non-native bees. However, The regulating body for non-native animal releases in England, Natural England, can only enforce licence requirements for use of non-native bees and so the release of *B. t. audax* is largely unregulated. In 2010, no difference in the prevalence of *C. bombi* was found between wild bumblebees collected near farms using or not using commercially produced bumblebees, although the parasite did increase in prevalence during the summer only at sites using commercially produced bumblebees which could suggest spillback (Whitehorn et al., 2013). In 2011, wild bumblebees from around farms in England using commercially produced bumblebees were more likely to be infected with the parasites *A. bombi*, *C. bombi*, and *N. ceranae* than wild bumblebees near farms not using managed bumblebees (Graystock et al., 2014). *Nosema ceranae* was first detected in bumblebees in Argentina is believed to have spilled over from honey bees (Plischuk et al., 2009). Its effects on bumblebee fitness are still unclear. Initial descriptions detected damaged epithelial cells suggesting spore production in the gut (Plischuk et al., 2009). In one study, controlled inoculations of mixed-aged *B. terrestris* workers with *N. ceranae* confirmed infection and spore production, and found reduced host survival over a 14 day period (Graystock et al., 2013a). In another study, inoculations of 2 day old *B. terrestris* workers produced persistent infections over a 21 day period, but survival was not reduced (Fürst et al., 2014). In 2011, 2012, screening on arrival of 48 commercially produced *B. terrestris* colonies from 3 different suppliers found that 77% of colonies contained bees with at least one parasite species known to harm bumblebees or honey bees, including *A. bombi*, *C. bombi*, *N. ceranae*, DWV, *Nosema apis* and *N. bombi*, with several of these being confirmed experimentally as being infectious (Graystock et al., 2013b). The following year, Natural England tightened up the release regulations on non-native bumblebees while a thorough review of the threats was conducted (Natural England, 2012, 2009). During this time, suppliers were requested to use molecular methods to screen for the bumblebee parasites *N. bombi*, *C. bombi*, *A. bombi*, plus the traditionally termed honey bee parasites *N. apis*, *N. ceranae*, and DWV. An updated risk assessment on *B. terrestris* was carried out on behalf of the UK Non-Native Species Secretariat, which highlighted the risks of establishment and hybridisation with native *B. t. audax*. As a result, Natural England heavily

restricted the release of non-native *B. terrestris*, with the 2015 licence only permitting this if commercial stocks of *B. t. audax* were unattainable. Native stocks of *B. t. audax* are still permitted for release to the UK despite them having previously been found to be equally as likely to be infected as *B. t. dalmatinus/terrestris* (Graystock et al., 2013b). This is of huge concern since the subspecies now being released into the UK has no legal requirement for disease screening or escape proofing whatsoever (apart from the honey bee centric restrictions relating to importation) (Graystock et al., 2013b). This is likely to remain the case unless local licencing is changed to allow the native sub species to be regulated (via its addition to schedule 9 and 14ZA of the Wildlife and Countryside Act), or European licencing is changed to encompass the breadth (and sometimes asymptomatic) nature of harmful parasites found in bumblebees.

#### 9.5. Case study 5: transmission between managed honey bees and wild bumblebees in Europe

In 2004 10% of commercially reared bumblebee queens inspected had the symptoms of deformed wing virus (DWV), and *Bombus pascuorum* workers collected from a wild colony in Germany the same year had symptomatic DWV, which was the first evidence for this parasite in bumblebees (Genersch et al., 2006). Deformed wing virus is highly prevalent and usually asymptomatic in honey bees, but high intensity infections (normally related to vectoring by ectoparasitic *Varroa* mites) can cause physical deformities, inability to fly and possible colony collapse (Highfield et al., 2009). In bumblebees, controlled inoculation experiments have shown that DWV can reduce survival (Fürst et al., 2014; Graystock et al., in press). In 2007 and 2008, sampling of wild pollinators across England for a range of parasites detected DWV in wild *B. terrestris* and *B. pascuorum* bumblebees, *Vespa vulgaris* wasps and honey bees (Evison et al., 2012). Wild bumblebees across Great Britain and the Isle of Mann have DWV and *N. ceranae* at varying levels which mirror the prevalence of these parasites in local honey bees, with higher infection intensities in honey bees suggesting they may be the infection reservoir and that the parasites spillover to bumblebees (Fürst et al., 2014). Further analysis of these samples also identified BQCV, ABPV, SBPV and SBV in several sympatric bumblebee species and these too were associated with their prevalence in managed *A. mellifera* populations, further indicating spillover, spillback and/or parasite facilitation between populations of bumblebees and honey bees (McMahon et al., 2015). However, infection intensities of ABPV and SBV were greater in *Bombus lapidarius* and *B. hortorum* than in honey bees suggesting either spillover from bumblebees to honey bees, bidirectional spillover and spillback, or that intensity is influenced by host tolerance and life–history parameters too much to be an adequate

predictor of transmission direction (McMahon et al., 2015; Runckel et al., 2011). It is in general now unclear whether viruses such as DWV which were first reported as honey bee pathogens are ancestrally honey bee pathogens that have subsequently jumped host into other bees, or whether the viruses ancestrally had a much greater host range than was previously realised, and have been traditionally considered as honey bee pathogens somewhat erroneously.

## 10. Conclusion

The balance of evidence suggests clearly that managed bees can, and have, had negative effects on wild bees, through parasite spillover, spillback and facilitation. Bumblebee colonies, which were commercially produced and imported into multiple countries over the last three decades, carried multiple infectious parasites with them, and the occurrence of parasites in managed colonies of honey bees is of course well known. The striking associations between the use of managed bees and local declines and extinctions of wild bees suggest strongly that multiple instances of parasite spillover, spillback and facilitation have occurred between managed and wild bees (Figs. 3–5). While the production of managed bees can be artificially increased to compensate for this, wild bee populations have to naturally bounce back, which may not be possible for many (Goulson and Hughes, 2015; Whitehorn et al., 2011).

Parasites have been found unequivocally in commercial bumblebees of several different species or subspecies, in three different continents, and from multiple suppliers, so disease appears to be a universal problem to commercial bumblebee breeders and their bees. Releasing managed bees therefore places wild bees at risk. Even when bees are reared in the same country of release, they could still introduce parasites to wild bees from the rearing facility. If it was possible to maintain disease free facilities, food and bees, the artificial introduction of large densities of disease free bees still may result in parasite spillback and parasite facilitation, and reduce the health of wild bees. Increased pollinator density has been suggested as a cause behind *B. terrestris* suffering increased disease prevalence in urban verse rural areas (Goulson et al., 2012).

Regulations regarding bumblebee screening are either absent in some territories or inadequate, such as European law Council Directive 92/65/EEC, which ensures bumblebees imported from 1992 must not show signs of American foulbrood (AFB) disease and must have no suspicion of *Tropilaelaps* mite or small hive beetle infestation. All three of these organisms are known predominantly for their harm to honey bees and are not considered a risk to bumblebees. The diversity of harmful parasites in bumblebees is far greater than previously considered yet most regulations regarding bumblebee screening have yet to acknowledge this.



Fig. 3. Overview of parasite detection in managed bees in Japan and likely instances of parasite transmission between managed and wild bumblebees.




Fig. 4. Overview of parasite detection in managed bees in North America and likely instances of parasite transmission between managed and wild bumblebees.




Fig. 5. Overview of parasite detection in managed bees in the British Isles and likely instances of parasite transmission between managed and wild bumblebees.

For some wild bees, prevention of parasite spillover, spillback, and facilitation is too late. There are lessons to be learned, however, about the dangers associated with intensive management of pollinators. As bumblebees and other managed pollinators such as *Osmia spp.* are increasingly cultivated, these lessons will become ever more important.

#### 10.1. Mitigating the harmful effects of managed bees on wild bees

There are a number of ways in which the harm managed bees cause to wild bees can be reduced. Firstly, parasites within rearing facilities must be prevented. A likely route by which managed bumblebees come into contact with parasites is via the pollen (collected by managed honey bees) that the bumblebees are reared on (Goulson, 2013). Many known bee pathogens are transmittable

via ingestion (Schmid-Hempel, 1998) and pollinator parasites including *C. bombi*, *N. bombi*, *N. apis*, *N. ceranae*, *A. bombi*, DWV, BQCV, SBV, *Acosphaera* (chalk brood) and *Paenibacillus larvae* (American Foul brood) have all been found in honey bee pollen with several shown to be infective (Chen et al., 2006; Flores et al., 2005; Graystock et al., 2013b; Higes et al., 2008; Singh et al., 2010). Gamma irradiation of pollen has been shown to reduce the pathogenicity of the honey bee virus IAPV and two commercial bumblebee producers (Biobest, Westerlo, Belgium; Koppert, B.V., Berkel en Rodenrijs, The Netherlands) are known to have started treating their pollen this way in Europe (Meeus et al., 2014). Feeding bumblebees without introducing known (or unknown) parasites is an area in need of further research. Also, some bumblebee rearing facilities still mix honey bee workers in with bumblebee queens to encourage egg laying. Forcing these 2 species



to live together provides an environment, which promote spillover to queens, which will later sire colonies that interact with wild bees. Further research into the production of colonies that doesn't rely on such practices are encouraged.

Exotic parasite introduction via the presence of exotic host species poses a particular risk; so controlling such exotic host introductions is a second mitigation measure that can be taken. In Japan, the non-native *B. terrestris* was added to a list of invasive alien species, prohibiting importation unless permits had been obtained by the farmer and they can prove no escapes are possible (Goka, 2010; Reade et al., 2014; Yoneda et al., 2007). In North America, importation of the non-native *B. terrestris* is banned outright (Winter et al., 2006), and in the UK the importation of non-native subspecies of *B. terrestris* is heavily restricted. Molecular screening is essential to identify the origin of many parasite strains and until done more thoroughly across various sites, the true identification of parasite spillover or spillback cannot be discerned. In cases when parasites are found upon delivery of managed bees, it seems likely to be spillover to wild bees, but a clearer understanding would be beneficial.

One of the best ways to avoid the potentially devastating effects of spillover, spillback, and facilitation in wild bees is to prevent their mixing with managed bees (Daszak et al., 2000; Foufopoulos et al., 2002). Farmers in many countries are now encouraged to use mesh on glasshouse windows or queen excluders on the front of hives to reduce the escape of commercially produced bumblebees. Despite this encouragement, studies have identified wild bees foraging inside glasshouses (Kraus et al., 2010; Lye et al., 2011a), and managed bumblebees foraging outside their glasshouses (Morandin et al., 2001; Murray et al., 2013). This population mixing results in the shared use of flowers by managed and wild bees, which are potential sites for parasite transmission (Durrer and Schmid-Hempel, 1994; Graystock et al., 2015; Singh et al., 2010). This highlights the ease with which wild and managed populations can mix and transmit parasites amongst themselves.

Based on this, we encourage the following to mitigate future and continued harm to wild bees:

- Strict, unified international regulations are needed to ensure bee movements do not introduce exotic parasites to new countries.
- Improved screening for a wider range of parasites, particularly cryptic viral infections.
- Improved husbandry to eliminate parasites within rearing facilities.
- Improved barrier control on farm glasshouses and poly tunnels to avoid mixing between managed and wild bees.
- Increase hedgerows and wildflower field margins to increase wild pollination services and reduce reliance on managed bees.
- Increased conservation effort to small/fragmented wild bee populations that are under increased stress.

While this will undoubtedly impose short-term monetary costs, it will help to protect wild bees and the valuable pollination services they provide in the long term. For honey bees, barrier control is however impossible so practices will instead rely heavily on regular screening and mitigative actions upon disease discovery, including the prevention of further movement by diseased hives. In the case of bumblebees used in glasshouses, complete barrier protection is possible and is a success story in Japan where regular parasite screening is also performed as a second line of defence (Reade et al., 2014).

## Acknowledgements

Thanks to Professor RCA (Andrew) Thompson for inviting PG to

lead this review and to two anonymous reviewers. Funded by Natural Environment Research Council (NERC) 2015 Early Career Impact Award to PG.

## References

- Aizen, M.A., Garibaldi, L.A., Cunningham, S.A., Klein, A.M., 2009. How much does agriculture depend on pollinators? Lessons from long-term trends in crop production. *Ann. Bot.* 103, 1579–1588. <http://dx.doi.org/10.1093/aob/mcp076>.
- Aizen, M.A., Harder, L.D., 2009. The global stock of domesticated honey bees is growing slower than agricultural demand for pollination. *Curr. Biol.* 19, 915–918. <http://dx.doi.org/10.1016/j.cub.2009.03.071>.
- Anderson, D.L., Trueman, J.W.H., 2000. *Varroa jacobsoni* (Acari: Varroidae) is more than one species. *Exp. Appl. Acarol.* 24, 165–189. <http://dx.doi.org/10.1023/A:1006456720416>.
- Antúñez, K., Martín-Hernández, R., Prieto, L., Meana, A., Zunino, P., Higes, M., 2009. Immune suppression in the honey bee (*Apis mellifera*) following infection by *Nosema ceranae* (Microsporidia). *Environ. Microbiol.* 11, 2284–2290. <http://dx.doi.org/10.1111/j.1462-2920.2009.01953.x>.
- Arias, M.C., Sheppard, W.S., 2005. Phylogenetic relationships of honey bees (Hymenoptera: Apinae: Apini) inferred from nuclear and mitochondrial DNA sequence data. *Mol. Phylogenet. Evol.* 37, 25–35. <http://dx.doi.org/10.1016/j.ympev.2005.02.017>.
- Arretz, P.V., Macfarlane, R.P., 1986. The introduction of *Bombus ruderatus* to Chile for red clover pollination. *Bee World* 67, 15–22.
- Asada, S., Ono, M., 1997. Tomato pollination with Japanese native bumblebees (*Bombus* spp.). VII Int. Symp. Pollinat. 437, 289–292.
- Batra, S.W.T., 1995. Bees and pollination in our changing environment. *Apidologie* 26, 361–370. <http://dx.doi.org/10.1051/apido:19950501>.
- Blitzer, E.J., Dormann, C.F., Holzschuh, A., Klein, A.M., Rand, T. a., Tschamtk, T., 2012. Spillover of functionally important organisms between managed and natural habitats. *Agric. Ecosyst. Environ.* 146, 34–43. <http://dx.doi.org/10.1016/j.agee.2011.09.005>.
- Boecking, O., Genersch, E., 2008. Varroosis – the ongoing crisis in bee keeping. *J. Verbrauch. Leb.* 3, 221–228. <http://dx.doi.org/10.1007/s00003-008-0331-y>.
- Botías, C., Martín-Hernández, R., Garrido-Bailón, E., González-Porto, A., Martínez-Salvador, A., De la Rúa, P., Meana, A., Higes, M., 2012. The growing prevalence of *Nosema ceranae* in honey bees in Spain, an emerging problem for the last decade. *Res. Vet. Sci.* 93, 150–155. <http://dx.doi.org/10.1016/j.rvsc.2011.08.002>.
- Bowen-Walker, P., Martin, S., Gunn, A., 1999. The transmission of deformed wing virus between honeybees (*Apis mellifera* L.) by the ectoparasitic mite *Varroa jacobsoni* Oud. *J. Invertebr. Pathol.* 73, 101–106. <http://dx.doi.org/10.1006/jipa.1998.4807>.
- Brown, M.J.F., 2011. Conservation: the trouble with bumblebees. *Nature* 469, 169–170. <http://dx.doi.org/10.1038/469169a>.
- Brown, M.J.F., Schmid-Hempel, R., Schmid-Hempel, P., 2003. Strong context-dependent virulence in a host-parasite system: reconciling genetic evidence with theory. *J. Anim. Ecol.* 72, 994–1002.
- Cameron, S.A., Lozier, J.D., Strange, J.P., Koch, J.B., Cordes, N., Solter, L.F., Griswold, T.L., 2011. Patterns of widespread decline in North American bumble bees. *Proc. Natl. Acad. Sci.* 108, 662–667. <http://dx.doi.org/10.1073/pnas.1014743108>.
- Cane, J.H., 2005. Pollination potential of the bee *Osmia aglaia* for cultivated red raspberries and blackberries (Rubus: Rosaceae). *Hortscience* 40, 1705–1708.
- Carreck, N.L., Ball, B.V., Wilson, J.K., 2002. Virus succession in honeybee colonies infested with *Varroa destructor*. *Apiacta* 37, 33–38.
- Central Office of Information, 2009. *Wildlife Management and Invasive Non-native Species Report of Research Findings Among the General Public, vol. 1. Anglers and the Horticultural Retail Trade.*
- Centre for the Promotion of Imports from Developing Countries, 2009. *The Honey and Other Bee Products Market in the EU.*
- Chauzat, M.-P., Carpentier, P., Madec, F., Bougeard, S., Cougoule, N., Drajnudel, P., Clément, M.-C., Aubert, M.A., Faucon, J.-P., 2010. The role of infectious agents and parasites in the health of honey bee colonies in France. *J. Apic. Res.* 49, 31–39. <http://dx.doi.org/10.3896/IBRA.149.1.05>.
- Chen, Y., Evans, J.D., Feldlaufer, M., 2006. Horizontal and vertical transmission of viruses in the honey bee, *Apis mellifera*. *J. Invertebr. Pathol.* 92, 152–159. <http://dx.doi.org/10.1016/j.jip.2006.03.010>.
- Chen, Y., Evans, J.D., Smith, I.B., Pettis, J.S., 2008. *Nosema ceranae* is a long-present and wide-spread microsporidian infection of the European honey bee (*Apis mellifera*) in the United States. *J. Invertebr. Pathol.* 97, 186–188. <http://dx.doi.org/10.1016/j.jip.2007.07.010>.
- Colla, S.R., Otterstatter, M.C., Gegear, R.J., Thomson, J.D., 2006. Plight of the bumble bee: pathogen spillover from commercial to wild populations. *Biol. Conserv.* 129, 461–467.
- Colla, S.R., Packer, L., 2008. Evidence for decline in eastern North American bumblebees (Hymenoptera: Apidae), with special focus on *Bombus affinis* Cresson. *Biodivers. Conserv.* 17, 1379–1391.
- Cordes, N., Huang, W.-F., Strange, J.P., Cameron, S.A., Griswold, T.L., Lozier, J.D., Solter, L.F., 2012. Interspecific geographic distribution and variation of the pathogens *Nosema bombi* and *Crithidia* species in United States bumble bee populations. *J. Invertebr. Pathol.* 109, 209–216. <http://dx.doi.org/10.1016/j.jip.2011.11.005>.

- Crane, E., 1999. *The World History of Beekeeping and Honey Hunting*. Routledge, New York.
- Crane, E., 1978. The *Varroa* mite. *Bee world* 59, 164.
- Crane, E., 1975. *Honey: a Comprehensive Survey*. Bee Research Association.
- Dafni, A., Kevan, P.G., Gross, C.L., Goka, K., 2010. *Bombus terrestris*, pollinator, invasive and pest: an assessment of problems associated with its widespread introductions for commercial purposes. *Appl. Entomol. Zool.* 45, 101–113. <http://dx.doi.org/10.1303/aez.2010.101>.
- Dainat, B., Evans, J.D., Chen, Y.P., Gauthier, L., Neumann, P., 2012. Dead or alive: deformed wing virus and *varroa destructor* reduce the life span of winter honeybees. *Appl. Environ. Microbiol.* 78, 981–987. <http://dx.doi.org/10.1128/AEM.06537-11>.
- Daszak, P., Cunningham, A.A., 1999. Extinction by infection. *Trends Ecol. Evol.* 14, 279. [http://dx.doi.org/10.1016/S0169-5347\(99\)01665-1](http://dx.doi.org/10.1016/S0169-5347(99)01665-1).
- Daszak, P., Cunningham, A.A., Hyatt, A.D., 2001. Anthropogenic environmental change and the emergence of infectious diseases in wildlife. *Acta Trop.* 78, 103–116.
- Daszak, P., Cunningham, A.A., Hyatt, A.D., 2000. Emerging infectious diseases of wildlife—threats to biodiversity and human health. *Science* 287, 443–449.
- De Luca, P. a., Bussi ere, L.F., Souto-Vilaros, D., Goulson, D., Mason, A.C., Vallejo-Mar n, M., 2013. Variability in bumblebee pollination buzzes affects the quantity of pollen released from flowers. *Oecologia* 172, 805–816. <http://dx.doi.org/10.1007/s00442-012-2535-1>.
- Delaplane, K.K.S., Mayer, D.R., Mayer, D.F., 2000. *Crop Pollination by Bees*. CABI Publishing.
- Dietemann, V., Pflugfelder, J., Anderson, D., Charri re, J.-D., Chejanovsky, N., Dainat, B., de Miranda, J., Delaplane, K., Dillier, F.-X., Fuch, S., Gallmann, P., Gauthier, L., Imdorf, A., Koeniger, N., Kralj, J., Meikle, W., Pettis, J., Rosenkranz, P., Sammartaro, D., Smith, D., Ya nez, O., Neumann, P., 2012. *Varroa destructor*: research avenues towards sustainable control. *J. Apic. Res.* 51, 125–132. <http://dx.doi.org/10.3896/IBRA.1.51.1.15>.
- Durrer, S., Schmid-Hempel, P., 1994. Shared use of flowers leads to horizontal pathogen transmission. *Proc. R. Soc. B Biol. Sci.* 258, 299–302. <http://dx.doi.org/10.1098/rspb.1994.0176>.
- Elbgami, T., Kunin, W.E., Hughes, W.O.H., Biesmeijer, J.C., 2014. The effect of proximity to a honeybee apiary on bumblebee colony fitness, development, and performance. *Apidologie*. <http://dx.doi.org/10.1007/s13592-013-0265-y>.
- Emsen, B., Guzman-Novoa, E., Kelly, P.G., 2013. Honey production of honey bee (*Hymenoptera: Apidae*) colonies with high and low *Varroa destructor* (*Acari: Varroidae*) infestation rates in eastern Canada. *Can. Entomol.* 146, 236–240. <http://dx.doi.org/10.4039/tce.2013.68>.
- Engel, M.M.S., 1999. The taxonomy of recent and fossil honey bees (*Hymenoptera Apidae Apis*). *J. Hymenopt. Res.* 8, 165–196.
- Evison, S.E.F., Roberts, K.E., Laurenson, L., Pietravalle, S., Hui, J., Biesmeijer, J.C., Smith, J.E., Budge, G., Hughes, W.O.H., 2012. Pervasiveness of parasites in pollinators. *PLoS One* 7, e30641. <http://dx.doi.org/10.1371/journal.pone.0030641>.
- Fantham, H.B., Porter, A., 1914. The morphology, biology and economic importance of *Nosema bombi*, *N. sp.*, parasitic in various bumble bees (*Bombus* spp.). *Annu. Trop. Med. Parasitol.* 8, 623–638.
- Fitzpatrick, U., Murray, T.E., Paxton, R.J., Breen, J., Cotton, D., Santorum, V., Brown, M.J.F., 2007. Rarity and decline in bumblebees – a test of causes and correlates in the Irish fauna. *Biol. Conserv.* 136, 185–194.
- Flanders, R.K., Wehling, W., Craghead, L., 2003. Chapter 6 Laws and regulations on the import, movement, and release of Bees in the United States. In: *For Non-native Crops, Whence Pollinators of the Future*, pp. 99–111.
- Flores, J.M., Guti rrez, I., Espejo, R., 2005. The role of pollen in chalkbrood disease in *Apis mellifera*: transmission and predisposing conditions. *Mycologia* 97, 1171–1176.
- Foufopoulos, J., Altizer, S., Dobson, A., 2002. Interactions between wildlife and domestic livestock in the tropics. In: *Tropical Agroecosystems*, pp. 219–244.
- Free, J.B., 1970. *Insect Pollination of Crops*. Academic Press Inc, London.
- Free, J.B., Williams, I.H., 2009. Pollination as a factor limiting the yield of field beans (*Vicia faba* L.). *J. Agric. Sci.* 87, 395. <http://dx.doi.org/10.1017/S0021859600027714>.
- Freestone, P.P.E., Sandrini, S.M., Haigh, R.D., Lyte, M., 2008. Microbial endocrinology: how stress influences susceptibility to infection. *Trends Microbiol.* 16, 55–64. <http://dx.doi.org/10.1016/j.tim.2007.11.005>.
- Fries, I., 2010. *Nosema ceranae* in European honey bees (*Apis mellifera*). *J. Invertebr. Pathol.* 103 (Suppl. 1), S73–S79. <http://dx.doi.org/10.1016/j.jip.2009.06.017>.
- Fries, I., 1997. Protozoa. In: Morse, R.A., Flottum, K. (Eds.), *Honey Bee Pests, Predators and Diseases*. A.I. Root Company, Medina, Ohio, USA, pp. 59–76.
- Fries, I., Feng, F., DaSilva, A., Slemenda, S.B., Pieniazek, N.J., 1996. *Nosema ceranae* n sp (*Microspora, Nosematidae*), morphological and molecular characterization of a microsporidian parasite of the Asian honey bee *Apis cerana* (*Hymenoptera, Apidae*). *Eur. J. Protistol.* 32, 356–365.
- Fries, I., Imdorf, A., Rosenkranz, P., 2006a. Survival of mite infested (*Varroa destructor*) honey bee (*Apis mellifera*) colonies in a Nordic climate. *Apidologie* 37, 564–570. <http://dx.doi.org/10.1051/apido>.
- Fries, I., Mart n, R., Meana, A., Garc a-Palencia, P., Higes, M., 2006b. Natural infections of *Nosema ceranae* in European honey bees. *J. Apic. Res.* 47, 230–233. <http://dx.doi.org/10.3896/IBRA.1.45.4.13>.
- F rst, M.A., McMahon, D.P., Osborne, J.L., Paxton, R.J., Brown, M.J.F., 2014. Disease associations between honeybees and bumblebees as a threat to wild pollinators. *Nature* 506, 364–366. <http://dx.doi.org/10.1038/nature12977>.
- Gallai, N., Salles, J.M., Settele, J., Vaissiere, B.E., 2009. Economic valuation of the vulnerability of world agriculture confronted with pollinator decline. *Ecol. Econ.* 68, 810–821. <http://dx.doi.org/10.1016/j.ecolecon.2008.06.014>.
- Genersch, E., Yue, C., Fries, I., Miranda, J.R. De, 2006. Detection of deformed wing virus, a honey bee viral pathogen, in bumble bees (*Bombus terrestris* and *Bombus pascuorum*) with wing deformities. *J. Invertebr. Pathol.* 91, 61–63. <http://dx.doi.org/10.1016/j.jip.2005.10.002>.
- Ghosh, K., Weiss, L.M., 2009. Molecular diagnostic tests for microsporidia. *Interdiscip. Perspect. Infect. Dis.* 2009, 926521. <http://dx.doi.org/10.1155/2009/926521>.
- Giersch, T., Berg, T., Galea, F., Hornitzky, M., 2009. *Nosema ceranae* infects honey bees (*Apis mellifera*) and contaminates honey in Australia. *Apidologie* 40, 117–123. <http://dx.doi.org/10.1051/apido/2008065>.
- Goka, K., 2010. Introduction to the Special Feature for Ecological Risk Assessment of Introduced Bumblebees: status of the European bumblebee, *Bombus terrestris*, in Japan as a beneficial pollinator and an invasive alien species. *Appl. Entomol. Zool.* 45, 1–6. <http://dx.doi.org/10.1303/aez.2010.1>.
- Goka, K., Okabe, K., Niwa, S., Yoneda, M., 2000. Parasitic mite infestation in introduced colonies of European bumblebees, *Bombus terrestris*. *Jpn. J. Appl. Entomol. Zool.* 44, 47–50. <http://dx.doi.org/10.1303/jjaez.2000.47>.
- Goka, K., Okabe, K., Yoneda, M., 2006. Worldwide migration of parasitic mites as a result of bumblebee commercialization. *Popul. Ecol.* 48, 285–291. <http://dx.doi.org/10.1007/s10144-006-0010-8>.
- Goka, K., Okabe, K., Yoneda, M., Niwa, S., 2001. Bumblebee commercialization will cause worldwide migration of parasitic mites. *Mol. Ecol.* 10, 2095–2099.
- Goulson, D., 2013. *A Sting in the Tale*. Jonathan Cape, London.
- Goulson, D., 2010. *Bumblebees: Behaviour, Ecology, and Conservation, Bumblebees Their Behaviour and Ecology*. Oxford University Press.
- Goulson, D., 2003. Effects of introduced bees on native ecosystems. *Annu. Rev. Ecol. Evol. Syst.* 34, 1–26. <http://dx.doi.org/10.1146/annurev.ecolsys.34.011802.132355>.
- Goulson, D., Derwent, L.C., 2004. Synergistic interactions between an exotic honeybee and an exotic weed: pollination of *Lantana camara* in Australia. *Weed Res.* 44, 195–202. <http://dx.doi.org/10.1111/j.1365-3180.2004.00391.x>.
- Goulson, D., Hughes, W.O.H., 2015. Mitigating the anthropogenic spread of bee parasites to protect wild pollinators. *Biol. Conserv.* 191, 10–19. <http://dx.doi.org/10.1016/j.biocon.2015.06.023>.
- Goulson, D., Lye, G.C., Darvill, B., 2008. Decline and conservation of bumble bees. *Annu. Rev. Entomol.* 53, 191–208. <http://dx.doi.org/10.1146/annurev.ento.53.103106.093454>.
- Goulson, D., Sparrow, K.R., 2008. Evidence for competition between honeybees and bumblebees; effects on bumblebee worker size. *J. Insect Conserv.* 13, 177–181. <http://dx.doi.org/10.1007/s10841-008-9140-y>.
- Goulson, D., Stout, J., Kells, A., 2002. Do exotic bumblebees and honeybees compete with native flower-visiting insects in Tasmania? *J. Insect Conserv.* 6, 179–189.
- Goulson, D., Whitehorn, P.R., Fowley, M., 2012. Influence of urbanisation on the prevalence of protozoan parasites of bumblebees. *Ecol. Entomol.* 37, 83–89. <http://dx.doi.org/10.1111/j.1365-2311.2011.01334.x>.
- Graystock, P., Goulson, D., Hughes, W.O.H., 2015. Parasites in bloom: flowers aid dispersal and transmission of pollinator parasites within and between bee species. *Proc. R. Soc. B Biol. Sci.* 282. <http://dx.doi.org/10.1098/rspb.2015.1371>.
- Graystock, P., Goulson, D., Hughes, W.O.H., 2014. The relationship between managed bees and the prevalence of parasites in bumblebees. *PeerJ* 2, e522. <http://dx.doi.org/10.7717/peerj.522>.
- Graystock, P., Meeus, I., Smagghe, G., Goulson, D., Hughes, W.O.H., (in press) The effects of single and mixed infections of *Apicystis bombi* and deformed wing virus on the bumblebee *Bombus terrestris*. *Parasitology*.
- Graystock, P., Yates, K., Darvill, B., Goulson, D., Hughes, W.O.H., 2013a. Emerging dangers: deadly effects of an emergent parasite in a new pollinator host. *J. Invertebr. Pathol.* 114, 114–119. <http://dx.doi.org/10.1016/j.jip.2013.06.005>.
- Graystock, P., Yates, K., Evison, S.E.F., Darvill, B., Goulson, D., Hughes, W.O.H., 2013b. The Trojan hives: pollinator pathogens, imported and distributed in bumblebee colonies. *J. Appl. Ecol.* 50, 1207–1215. <http://dx.doi.org/10.1111/1365-2664.12134>.
- Greenleaf, S.S., Kremen, C., 2006. Wild bee species increase tomato production and respond differently to surrounding land use in Northern California. *Biol. Conserv.* 133, 81–87. <http://dx.doi.org/10.1016/j.biocon.2006.05.025>.
- Grixti, J.C., Wong, L.T., Cameron, S.A., Favret, C., 2009. Decline of bumble bees (*Bombus*) in the North American Midwest. *Biol. Conserv.* 142, 75–84. <http://dx.doi.org/10.1016/j.biocon.2008.09.027>.
- Hatcher, M., Dunn, A., 2011. *Parasites in Ecological Communities*. Cambridge University Press.
- Hedtke, K., Jensen, P.M., Jensen, A.B., Genersch, E., 2011. Evidence for emerging parasites and pathogens influencing outbreaks of stress-related diseases like chalkbrood. *J. Invertebr. Pathol.* 108, 167–173. <http://dx.doi.org/10.1016/j.jip.2011.08.006>.
- Heemert, C., Ruijter, A., Eijnde, J., Steen, J., 1990. Year-round production of bumble bee colonies for crop pollination. *Bee World* 71, 54–56.
- Heinrich, B., 1993. *The Hot-blooded Insects: Strategies and Mechanisms of Thermoregulation*. Harvard University Press.
- Higes, M., Mart n, R., Meana, A., Mart n, R., 2006. *Nosema ceranae*, a new microsporidian parasite in honeybees in Europe. *J. Invertebr. Pathol.* 92, 93–95. <http://dx.doi.org/10.1016/j.jip.2006.02.005>.
- Higes, M., Mart n-Hern ndez, R., Garrido-Bail n, E., Garc a-Palencia, P., Meana, A., 2008. Detection of infective *Nosema ceranae* (*Microsporida*) spores in corbicular pollen of forager honeybees. *J. Invertebr. Pathol.* 97, 76. <http://dx.doi.org/10.1016/j.jip.2008.02.005>.

- 10.1016/j.jip.2007.06.002.
- Higes, M., Martín-Hernández, R., Meana, A., 2010. *Nosema ceranae* in Europe: an emergent type C nosemosis. *Apidologie* 41, 375–392. <http://dx.doi.org/10.1051/apido/2010019>.
- Highfield, A.C., El Nagar, A., Mackinder, L.C.M., Noel, L., Hall, M.J., Martin, S.J., Schroeder, D.C., 2009. Deformed wing virus implicated in overwintering honeybee colony losses. *Appl. Environ. Microbiol.* 75, 7212–7220. <http://dx.doi.org/10.1128/aem.02227-09>.
- Holm, S.N., 1966. The utilization and management of bumble bees for Red Clover and Alfalfa seed production. *Annu. Rev. Entomol.* 11, 155–182. <http://dx.doi.org/10.1146/annurev.en.11.010166.001103>.
- Hopkins, I., 1914. *History of the Humble-bee in New Zealand: its Introduction and Results*.
- Huang, W.F., Jiang, J.H., Chen, Y.W., Wang, C.-H., 2007. A *Nosema ceranae* isolate from the honeybee *Apis mellifera*. *Apidologie* 38, 30–37. <http://dx.doi.org/10.1051/apido>.
- Huryan, V.M., 1997. Ecological impacts of introduced honey bees. *Q. Rev. Biol.* 72.
- Husband, R.W., Sinha, R.N., 1970. A revision of genus *Locustacarus* with a key to genera of Family Podapolipidae (Acarina). *Entomol. Soc. Am.* 63, 1152.
- Ings, T.C., Ward, N.L., Chittka, L., 2006. Can commercially imported bumble bees outcompete their native conspecifics? *J. Appl. Ecol.* 43, 940–948. <http://dx.doi.org/10.1111/j>.
- Invernizzi, C., Abud, C., Tomasco, I.H., Harriet, J., Ramallo, G., Campá, J., Katz, H., Gardiol, G., Mendoza, Y., 2009. Presence of *Nosema ceranae* in honeybees (*Apis mellifera*) in Uruguay. *J. Invertebr. Pathol.* 101, 155–153. <http://dx.doi.org/10.1016/j.jip.2009.03.006>.
- Jaffé, R., Dietemann, V., Allsopp, M.H., Costa, C., Crewe, R.M., Dall'olio, R., De La Rúa, P., El-Niweiri, M.A.A., Fries, I., Kezic, N., Meusel, M.S., Paxton, R.J., Shaibi, T., Stolle, E., Moritz, R.F.A., 2010. Estimating the density of honeybee colonies across their natural range to fill the gap in pollinator decline censuses. *Conserv. Biol.* 24, 583–593. <http://dx.doi.org/10.1111/j.1523-1739.2009.01331.x>.
- Jokela, J., Taskinen, J., Mutikainen, P., Kopp, K., 2005. Virulence of parasites in hosts under environmental stress: experiments with anoxia and starvation. *Oikos* 108, 156–164. <http://dx.doi.org/10.1111/j.0030-1299.2005.13185.x>.
- Jones, K.E., Patel, N.G., Levy, M.A., Storeygard, A., Balk, D., Gittleman, J.L., Daszak, P., 2008. Global trends in emerging infectious diseases. *Nature* 451, 990–993. <http://dx.doi.org/10.1038/nature06536>.
- Jong, D. De, Morse, R. a, Eickwort, G.C., 1982. Mite pests of honey bees. *Annu. Rev. Entomol.* 27, 229–252.
- Kelly, D., Paterson, R., Townsend, C., 2009. Parasite spillback: a neglected concept in invasion ecology? *Ecology* 90, 247–2056.
- Klee, J., Besana, A.M., Genersch, E., Gisder, S., Nanetti, A., Tam, D.Q., Chinh, T.X., Puerta, F., Ruz, J.M., Kryger, P., Message, D., Hatjina, F., Korpela, S., Fries, I., Paxton, R.J., 2007. Widespread dispersal of the microsporidian *Nosema ceranae*, an emergent pathogen of the western honey bee, *Apis mellifera*. *J. Invertebr. Pathol.* 96, 1–10. <http://dx.doi.org/10.1016/j.jip.2007.02.014>.
- Klee, J., Tek Tay, W., Paxton, R.J., 2006. Specific and sensitive detection of *Nosema bombi* (Microsporidia: Nosematidae) in bumble bees (*Bombus* spp.; Hymenoptera: Apidae) by PCR of partial rRNA gene sequences. *J. Invertebr. Pathol.* 91, 98–104. <http://dx.doi.org/10.1016/j.jip.2005.10.012>.
- Klein, A.-M., Vaissière, B.E., Cane, J.H., Steffan-Dewenter, I., Cunningham, S. a, Kremen, C., Tscharntke, T., 2007. Importance of pollinators in changing landscapes for world crops. *Proc. R. Soc. B Biol. Sci.* 274, 303–313. <http://dx.doi.org/10.1098/rspb.2006.3721>.
- Koeniger, N., Koeniger, G., 2000. Reproductive isolation among species of the genus *Apis*. *Apidologie* 31, 313–339. <http://dx.doi.org/10.1051/apido:2000125>.
- Kondo, N.I., Yamanaka, D., Kanbe, Y., Kunitake, Y.K., Yoneda, M., Tsuchida, K., Goka, K., 2009. Reproductive disturbance of Japanese bumblebees by the introduced European bumblebee *Bombus terrestris*. *Naturwissenschaften* 96, 467–475. <http://dx.doi.org/10.1007/s00114-008-0495-4>.
- Korpela, S., Aarhus, A., Fries, I., Hansen, H., 1993. *Varroa jacobsoni* Oud. in cold climates: population growth, winter mortality and influence on survival of honey bee colonies. *J. Apic. Res.* 31, 157–164.
- Kosior, A., Celary, W., Olejniczak, P., Fijał, J., Król, W., Solarz, W., Plonka, P., 2007. The decline of the bumble bees and cuckoo bees (Hymenoptera: Apidae: Bombini) of Western and Central Europe. *Oryx* 41, 79–88. <http://dx.doi.org/10.1017/S0030605307001597>.
- Kralj, J., Brockmann, A., Fuchs, S., Tautz, J., 2007. The parasitic mite *Varroa destructor* affects non-associative learning in honey bee foragers. *Apis mellifera* L. *J. Comp. Physiol. A Neuroethol. Sens. Neural, Behav. Physiol.* 193, 363–370. <http://dx.doi.org/10.1007/s00359-006-0192-8>.
- Kralj, J., Fuchs, S., 2006. Parasitic *Varroa destructor* mites influence flight duration and homing ability of infested *Apis mellifera* foragers. *Apidologie* 37, 577–587.
- Kraus, F.B., Szentgyörgyi, H., Rozej, E., Rhode, M., Morón, D., Woyciechowski, M., Moritz, R.F.A., 2010. Greenhouse bumblebees (*Bombus terrestris*) spread their genes into the wild. *Conserv. Genet.* 12, 187–192. <http://dx.doi.org/10.1007/s10592-010-0131-7>.
- Kremen, C., Williams, N.M., Thorp, R.W., 2002. Crop pollination from native bees at risk from agricultural intensification. *Proc. Natl. Acad. Sci.* 99, 16812–16816. <http://dx.doi.org/10.1073/pnas.262413599>.
- Le Conte, Y., Ellis, M., Ritter, W., 2010. *Varroa* mites and honey bee health: can *Varroa* explain part of the colony losses? *Apidologie* 41, 353–363. <http://dx.doi.org/10.1051/apido/2010017>.
- Lederberg, J., Shope, R.E., Oakes, S.C., 1992. *Emerging Infections: Microbial Threats to Health in the United States*. Institute of Medicine. National Academy Press, Washington D.C.
- Lye, G.C., Jennings, S.N., Osborne, J.L., Goulson, D., 2011a. Impacts of the use of nonnative commercial bumble bees for pollinator supplementation in raspberry. *J. Econ. Entomol.* 104, 107–114. <http://dx.doi.org/10.1603/EC10092>.
- Lye, G.C., Lepais, O., Goulson, D., 2011b. Reconstructing demographic events from population genetic data: the introduction of bumblebees to New Zealand. *Mol. Ecol.* 20, 2888–2900. <http://dx.doi.org/10.1111/j.1365-294X.2011.05139.x>.
- MacKinnon, B., 1998. Host factors important in sea lice infections. *ICES J. Mar. Sci.* 55, 188–192. <http://dx.doi.org/10.1006/jmsc.1997.0361>.
- Martin, S.J., Highfield, A.C., Brettell, L., Villalobos, E.M., Budge, G.E., Powell, M., Nikaido, S., Schroeder, D.C., 2012. Global honey bee viral landscape altered by a parasitic mite. *Science* 336, 1304–1306. <http://dx.doi.org/10.1126/science.1220941>.
- Martins, A., Melo, G., 2010. Has the bumblebee *Bombus bellicosus* gone extinct in the northern portion of its distribution range in Brazil? *J. Insect Conserv.* 14, 207–210. <http://dx.doi.org/10.1007/s10841-009-9237-y>.
- Mayack, C., Naug, D., 2009. Energetic stress in the honeybee *Apis mellifera* from *Nosema ceranae* infection. *J. Invertebr. Pathol.* 100, 185–188. <http://dx.doi.org/10.1016/j.jip.2008.12.001>.
- McCallum, H., Dobson, A., 2002. Disease, habitat fragmentation and conservation. *Proc. R. Soc. B Biol. Sci.* 269, 2041–2049. <http://dx.doi.org/10.1098/rspb.2002.2079>.
- McGregor, S.E., 1976. *Insect Pollination of Cultivated Crop Plants*. Agricultural Research Service, US Department of Agriculture.
- McMahon, D.P., Fürst, M. a., Caspar, J., Theodorou, P., Brown, M.J.F., Paxton, R.J., 2015. A sting in the spit: widespread cross-infection of multiple RNA viruses across wild and managed bees. *J. Anim. Ecol.* 84, 615–624. <http://dx.doi.org/10.1111/1365-2656.12345>.
- Meeus, I., Brown, M.J.F., de Graaf, D.C., Smagghe, G., 2011. Effects of invasive parasites on bumble bee declines. *Conserv. Biol.* 25, 662–671. <http://dx.doi.org/10.1111/j.1523-1739.2011.01707.x>.
- Meeus, I., Mosallanejad, H., Niu, J., de Graaf, D.C., Wäckers, F., Smagghe, G., 2014. Gamma irradiation of pollen and eradication of Israeli acute paralysis virus. *J. Invertebr. Pathol.* 121, 10–13. <http://dx.doi.org/10.1016/j.jip.2014.06.012>.
- Møller, A.P., Milinski, M., Slater, P.J.B., 1998. Stress and behavior. In: *Advances in the Study of Behavior*. Academic Press, p. 552.
- Morandini, L.A., Laverty, T.M., Kevan, P.G., Khlosa, S., Shipp, L., 2001. Bumble bee (Hymenoptera: Apidae) activity and loss in commercial tomato greenhouses. *Can. Entomol.* 133, 883–893.
- Moritz, R.F.A., Kraus, F.B., Kryger, P., Crewe, R.M., 2007. The size of wild honeybee populations (*Apis mellifera*) and its implications for the conservation of honeybees. *J. Insect Conserv.* 11, 391–397.
- Morse, S.S., 1993. Examining the origins of emerging viruses. In: *Emerging Viruses*. Oxford University Press, New York, pp. 10–28.
- Murray, T.E., Coffey, M.F., Kehoe, E., Horgan, F.G., 2013. Pathogen prevalence in commercially reared bumble bees and evidence of spillover in conspecific populations. *Biol. Conserv.* 159, 269–276. <http://dx.doi.org/10.1016/j.biocon.2012.10.021>.
- Natural England, 2012. Licence (Class) to Permit the Use of Non Native Subspecies of the Bumblebee (*Bombus terrestris*) in Commercial Glass-houses or Poly-tunnels for Crop Pollination WML-CL22.
- Natural England, 2009. Use of Non-native *Bombus terrestris* Subspecies for Crop Pollination in England - Licence Disease Screening Requirements NNR/2007/30-33, Licence (Class) to Permit the Use of Non Native Subspecies of the Bumblebee (*Bombus terrestris*) in Commercial Glass-houses or Poly-tunnels for Crop Pollination WML-CL22.
- Ndao, M., 2009. Diagnosis of parasitic diseases: old and new approaches. *Interdiscip. Perspect. Infect. Dis.* 2009, 278246. <http://dx.doi.org/10.1155/2009/278246>.
- Neumann, P., Carreck, N.L., 2010. Honey bee colony losses. *J. Apic. Res.* 49, 1–6. <http://dx.doi.org/10.3896/ibra.149.1.01>.
- Niwa, S., Iwano, H., Asada, S.I., Matsuura, M., Goka, K., 2004. A microsporidian pathogen isolated from a colony of the European bumblebee, *Bombus terrestris*, and infectivity on Japanese bumblebee. *Jpn. J. Appl. Entomol. Zool.* 48, 60–64. <http://dx.doi.org/10.1303/jjaez.2004.60>.
- Oldroyd, B.P., 1999. Coevolution while you wait: *Varroa jacobsoni*, a new parasite of western honeybees. *Trends Ecol. Evol.* 14, 312–315. [http://dx.doi.org/10.1016/S0169-5347\(99\)01613-4](http://dx.doi.org/10.1016/S0169-5347(99)01613-4).
- Otterstatter, M.C., Thomson, J.D., 2008. Does pathogen spillover from commercially reared bumble bees threaten wild pollinators? *PLoS One* 3, e2771. <http://dx.doi.org/10.1371/journal.pone.0002771>.
- Otterstatter, M.C., Whidden, T.L., 2004. Patterns of parasitism by tracheal mites (*Locustacarus buchneri*) in natural bumble bee populations. *Apidologie* 35, 351–357. <http://dx.doi.org/10.1051/apido:2004024>.
- Otti, O., Schmid-Hempel, P., 2008. A field experiment on the effect of *Nosema bombi* in colonies of the bumblebee *Bombus terrestris*. *Ecol. Entomol.* 33, 577–582. <http://dx.doi.org/10.1111/j.1365-2311.2008.00998.x>.
- Otti, O., Schmid-Hempel, P., 2007. *Nosema bombi*: a pollinator parasite with detrimental fitness effects. *J. Invertebr. Pathol.* 96, 118–124. <http://dx.doi.org/10.1016/j.jip.2007.03.016>.
- Oudemans, A.C., 1904. On a new genus and species of parasitic acari. *Notes Leyden Mus.* 24, 216–222. <http://dx.doi.org/10.1017/S0031182000012919>.
- Paxton, R.J., Klee, J., Korpela, S., Fries, I., 2007. *Nosema ceranae* has infected *Apis mellifera* in Europe since at least 1998 and may be more virulent than *Nosema apis*. *Apidologie* 38, 558.

- Perlman, S.J., Jaenike, J., 2003. Infection success in novel hosts: an experimental and phylogenetic study of *Drosophila*-parasitic nematodes. *Evolution* (N. Y.) 57, 544–557.
- Plischuk, S., Martín-Hernández, R., Prieto, L., Lucía, M., Botías, C., Meana, A., Abrahamovich, A.H., Lange, C., Higes, M., 2009. South American native bumblebees (Hymenoptera: Apidae) infected by *Nosema ceranae* (Microsporidia), an emerging pathogen of honeybees (*Apis mellifera*). *Environ. Microbiol. Rep.* 1, 131–135.
- Pohorecka, K., Bober, A., Skubida, M., Zdanska, D., 2011. Epizootic status of apiaries with massive losses of bee colonies (2008–2009). *J. Apic. Sci.* 55, 137–151.
- Potts, S.G., Biesmeijer, J.C., Kremen, C., Neumann, P., Schweiger, O., Kunin, W.E., 2010. Global pollinator declines: trends, impacts and drivers. *Trends Ecol. Evol.* 25, 345–353. <http://dx.doi.org/10.1016/j.tree.2010.01.007>.
- Power, A.G., Mitchell, C.E., 2004. Pathogen spillover in disease epidemics. *Am. Nat.* 164 (Suppl. 1), S79–S89. <http://dx.doi.org/10.1086/424610>.
- Procop, G.W., 2007. Molecular diagnostics for the detection and characterization of microbial pathogens. *Clin. Infect. Dis.* 45 (Suppl. 2), S99–S111. <http://dx.doi.org/10.1086/519259>.
- Ptacek, V., 1985. Testing three methods of bumble bee rearing. *Sb. Ved. Pr. Vyzk. a selchitelský Ust. picnicnarský V. Troubsko u Brna* 9, 59–67.
- Ransome, H.M., 2004. *The sacred bee. In: Ancient Times and Folklore.* Dover Publications, New York.
- Rasmont, P., Pauly, A., Terzo, M., 2005. The Survey of Wild Bees (Hymenoptera, Apoidea) in Belgium and France. Food and Agriculture Organisation, Rome. FAO report [online]. <http://www.fao.org/ag/AGP/C-AGPS/C-CAB/Castudies/pdf/1-010.pdf>.
- Reade, C., Goka, K.K., Thorp, R., Mitsuhashi, M., Wasbauer, M., 2014. CSR, biodiversity and Japan's Stakeholder Approach to the global bumble Bee trade. *J. Corp. Citizsh.* 53–66. <http://dx.doi.org/10.9774/GLEAF.4700.2014.de.00006>.
- Rinder, H., Janitschke, K., Aspöck, H., Da Silva, A.J., Deplazes, P., Fedorko, D.P., Franzen, C., Futh, U., Hüniger, F., Lehmacher, a, Meyer, C.G., Molina, J.M., Sandfort, J., Weber, R., Löscher, T., 1998. Blinded, externally controlled multicenter evaluation of light microscopy and PCR for detection of microsporidia in stool specimens. The Diagnostic Multicenter Study Group on Microsporidia. *J. Clin. Microbiol.* 36, 1814–1818.
- Roberts, K.E., Hughes, W.O.H., 2014. Immunosenescence and resistance to parasite infection in the honey bee, *Apis mellifera*. *J. Invertebr. Pathol.* 121, 1–6. <http://dx.doi.org/10.1016/j.jip.2014.06.004>.
- Röseler, P., 1977. Rearing bumblebee colonies. In: *Proceedings of the 8th International Congress of the IJSSI, Wageningen.*
- Rosenkranz, P., Aumeier, P., Ziegelmann, B., 2010. Biology and control of Varroa destructor. *J. Invertebr. Pathol.* 103, S96–S119. <http://dx.doi.org/10.1016/j.jip.2009.07.016>.
- Rozej, E., Witaliński, W., Szentgyörgyi, H., Wantuch, M., Morón, D., Woyciechowski, M., 2012. Mite species inhabiting commercial bumblebee (*Bombus terrestris*) nests in Polish greenhouses. *Exp. Appl. Acarol.* 56, 271–282. <http://dx.doi.org/10.1007/s10493-012-9510-8>.
- Runkel, C., Flenniken, M.L., Engel, J.C., Ruby, J.G., Ganem, D., Andino, R., DeRisi, J.L., 2011. Temporal analysis of the honey bee microbiome reveals four novel viruses and seasonal prevalence of known viruses, *Nosema*, and *Crithidia*. *PLoS One* 6, e20656. <http://dx.doi.org/10.1371/journal.pone.0020656>.
- Rutrecht, S.T., Brown, M.J.F., 2008. Within colony dynamics of *Nosema bombi* infections: disease establishment, epidemiology and potential vertical transmission. *Apidologie* 39, 504–514.
- Ruttner, F., 1988. *Biogeography and taxonomy of honeybees.* Springer-Verlag Berlin and Heidelberg GmbH & Co. K.
- Sachman-Ruiz, B., Narváez-Padilla, V., Reynaud, E., 2015. Commercial *Bombus impatiens* as reservoirs of emerging infectious diseases in central Mexico. *Biol. Invasions* 17, 2043–2053. <http://dx.doi.org/10.1007/s10530-015-0859-6>.
- Sakai, T., Okada, I., 1974. The present beekeeping in Japan. *Glean. Bee Cult.* 356–357.
- Sárospataki, M., Novák, J., Molnár, V., 2005. Assessing the threatened status of bumble bee species (Hymenoptera: Apidae) in Hungary. *Central Europe. Biodivers. Conserv.* 14, 2437–2446. <http://dx.doi.org/10.1007/s10531-004-0152-y>.
- Schmid-Hempel, P., 2011. *Evolutionary Parasitology. The Integrated Study of Infections, Immunology, Ecology and Genetics.* Oxford University Press.
- Schmid-Hempel, P., 1998. *Parasites in Social Insects.* Princeton University Press, Princeton University Press.
- Schmid-Hempel, P., Loosli, R., 1998. A contribution to the knowledge of *Nosema* infections in bumble bees, *Bombus* spp. *Apidologie* 29, 525–535.
- Seeley, T.D., 1985. *Honey Bee Ecology.* Princeton University Press.
- Sheppard, W.S., Meixner, M.D., 2003. *Apis mellifera* pomonella, a new honey bee subspecies from Central Asia. *Apidologie* 34, 367–375. <http://dx.doi.org/10.1051/apido:2003037>.
- Singh, R., Levitt, A.L., Rajotte, E.G., Holmes, E.C., Ostiguy, N., VanEngelsdorp, D., Lipkin, W.I., Depamphilis, C.W., Toth, A.L., Cox-Foster, D., 2010. RNA viruses in hymenopteran pollinators: evidence of inter-taxa virus transmission via pollen and potential impact on non-*Apis* hymenopteran species. *PLoS One* 5, e14357. <http://dx.doi.org/10.1371/journal.pone.0014357>.
- Singh, Y., 1975. *Nosema* in Indian honey bee (*Apis cerana indica*). *Am. Bee J.* 115, 59.
- Solignac, M., Cornuet, J.-M., Vautrin, D., Le Conte, Y., Anderson, D., Evans, J., Cros-Arteil, S., Navajas, M., 2005. The invasive Korea and Japan types of *Varroa destructor*, ectoparasitic mites of the Western honeybee (*Apis mellifera*), are two partly isolated clones. *Proc. Biol. Sci.* 272, 411–419. <http://dx.doi.org/10.1098/rspb.2004.2853>.
- Szabo, N.D., Colla, S.R., Wagner, D.L., Gall, L.F., Kerr, J.T., 2012. Do pathogen spillover, pesticide use, or habitat loss explain recent North American bumblebee declines? *Conserv. Lett.* 5, 232–239. <http://dx.doi.org/10.1111/j.1755-263X.2012.00234.x>.
- Tarpy, D.R., 2003. Genetic diversity within honeybee colonies prevents severe infections and promotes colony growth. *Proc. R. Soc. B Biol. Sci.* 270, 99–103. <http://dx.doi.org/10.1098/rspb.2002.2199>.
- Thompson, C.E., Biesmeijer, J.C., Allnutt, T.R., Pietravalle, S., Budge, G.E., 2014. Parasite pressures on feral honey Bees (*Apis mellifera* sp.). *PLoS One* 9, e105164. <http://dx.doi.org/10.1371/journal.pone.0105164>.
- Thomson, D., 2004. Competitive interactions between the invasive European honey bee and native bumble bees. *Ecology* 85, 458–470. <http://dx.doi.org/10.1890/02-0626>.
- Thorp, R., 2005. *Bombus franklini* Frison, 1921. Franklin's bumble Bee (Hymenoptera: Apidae: Apinae: Bombini). In: *Red List Pollinator Insects North Am. CO-*, vol. 1, pp. 1–8.
- Thorp, R.W., Shepherd, M.D., 2005. Profile: Subgenus *Bombus*. Latreille 1802 (Apidae: Apinae: Bombini). In: Shepherd MD, Vaughan DM Red List of Pollinator Insects of North America. The Xerxes Society for Invertebrate Conservation, Portland, OR.
- Van der Eijnde, J., 1990. Method for continuous rearing of *Bombus terrestris* and the production of bumblebee colonies for pollination purposes. *Apidologie* 21, 330–332.
- Van der Steen, J.J.M., 2008. Infection and transmission of *Nosema bombi* in *Bombus terrestris* colonies and its effect on hibernation, mating and colony founding. *Apidologie* 39, 273–282. <http://dx.doi.org/10.1051/apido:2008006>.
- vanEngelsdorp, D., Hayes Jr., J., Underwood, R.M., Caron, D., Pettis, J., 2011. A survey of managed honey bee colony losses in the USA, fall 2009 to winter 2010. *J. Apic. Res.* 50, 1–10. <http://dx.doi.org/10.3896/IBRA.1.50.1.01>.
- VanEngelsdorp, D., Meixner, M.D., 2010. A historical review of managed honey bee populations in Europe and the United States and the factors that may affect them. *J. Invertebr. Pathol.* 103 (Suppl. 1), S80–S95.
- Velthuis, H.H.W., 2002. The historical background of the domestication of the bumble-bee, *Bombus terrestris*, and its introduction in agriculture. In: Kevan, P., Imperatriz Fonseca, V.L. (Eds.), *Pollinating Bees. The Conservation Link Between Agriculture and Nature.* Ministry of Environment, Brasilia, Brazil, pp. 177–184.
- Velthuis, H.H.W., Van Doorn, A., 2006. A century of advances in bumblebee domestication and the economic and environmental aspects of its commercialization for pollination. *Apidologie* 37, 421–451. <http://dx.doi.org/10.1051/apido>.
- Wehling, W.F., Flanders, R.V., 2005. *The Risks of International Trade in Bees: Laws and Regulations on the Import, Movement and Release of Bees in the United States.* BRAVE 301.
- Whitehorn, P.R., Tinsley, M.C., Brown, M.J.F., Darvill, B., Goulson, D., 2011. Genetic diversity, parasite prevalence and immunity in wild bumblebees. *Proc. R. Soc. B Biol. Sci.* 278, 1195–1202. <http://dx.doi.org/10.1098/rspb.2010.1550>.
- Whitehorn, P.R., Tinsley, M.C., Brown, M.J.F., Goulson, D., 2013. Investigating the impact of deploying commercial *Bombus terrestris* for crop pollination on pathogen dynamics in wild bumble bees. *J. Apic. Res.* 52, 149–157. <http://dx.doi.org/10.3896/IBRA.1.52.3.06>.
- Whittington, R., Winston, M.L., 2003. Effects of *Nosema bombi* and its treatment fumagillin on bumble bee (*Bombus occidentalis*) colonies. *J. Invertebr. Pathol.* 84, 54–58. [http://dx.doi.org/10.1016/S0022-2011\(03\)00123-X](http://dx.doi.org/10.1016/S0022-2011(03)00123-X).
- Williams, I.H., 1994. The dependences of crop production within the European Union on pollination by honey bees. *Agric. Sci. Rev.* 6, 229–257.
- Williams, P.H., 1982. The distribution and decline of British bumble bees (*Bombus* Latr.). *J. Apic. Res.* 21, 236–245.
- Williams, P.H., Osborne, J.L., 2009. Bumblebee vulnerability and conservation world-wide. *Apidologie* 40, 367–387. <http://dx.doi.org/10.1051/apido/2009025>.
- Winter, K., Adams, L., Thorp, R., Inouye, D., Day, L., Ascher, J., Buchmann, S.L., 2006. Importation of Non-native Bumble Bees into North America: Potential Consequences of Using *Bombus terrestris* and Other Non-native Bumble Bees for Greenhouse Crop Pollination in Canada, Mexico, and the United States. San Francisco.
- Wood, J.L.N., Leach, M., Waldman, L., Macgregor, H., Fooks, A.R., Jones, K.E., Restif, O., Dechmann, D., Hayman, D.T.S., Baker, K.S., Peel, A.J., Kamins, A.O., Fahr, J., Ntiamao-Baidou, Y., Suu-Ire, R., Breiman, R.F., Epstein, J.H., Field, H.E., Cunningham, A.A., 2012. A framework for the study of zoonotic disease emergence and its drivers: spillover of bat pathogens as a case study. *Philos. Trans. R. Soc. B Biol. Sci.* 367, 2881–2892. <http://dx.doi.org/10.1098/rstb.2012.0228>.
- Wu, J., 2008. Comparison of the colony development of two native bumblebee species *Bombus ignitus* and *Bombus lucorum* as candidates for commercial pollination in China. *J. Apic. Res.* 22–26. <http://dx.doi.org/10.3896/IBRA.1.47.1.04>.
- Xie, Z., Williams, P.H., Tang, Y., 2008. The effect of grazing on bumblebees in the high rangelands of the eastern Tibetan Plateau of Sichuan. *J. Insect Conserv.* 12, 695–703. <http://dx.doi.org/10.1007/s10841-008-9180-3>.
- Yang, S., Rothman, R.E., 2004. PCR-based diagnostics for infectious diseases: uses, limitations, and future applications in acute-care settings. *Lancet Infect. Dis.* 4, 337–348. [http://dx.doi.org/10.1016/S1473-3099\(04\)01044-8](http://dx.doi.org/10.1016/S1473-3099(04)01044-8).
- Yoneda, M., Furuta, H., Kanbe, Y., Tsuchida, K., Okabe, K., Goka, K., 2008. Reproduction and transmission within a colony of bumblebee tracheal mite *Locustacarus buchneri* (Acari: Podapolipidae) in *Bombus terrestris* (Hymenoptera: Apidae). *Appl. Entomol. Zool.* 43, 391–395. <http://dx.doi.org/10.1303/aez.2008.391>.
- Yoneda, M., Yokoyama, J., Tsuchida, K., Osaki, T., Itoya, S., Goka, K., 2007. Preventing *Bombus terrestris* from escaping with a net covering over a tomato greenhouse in Hokkaido. *Jpn. J. Appl. Entomol. Zool.* 51, 39–44. <http://dx.doi.org/10.1303/jjaez.2007.39>.