

# Prognostic value of expression levels of miR-148a, miR-152 and HLA-G in colon cancer

LEVENT EMIRZEOGLU<sup>1</sup>, OZGUR OLMEZ<sup>1</sup>, FATMA NIHAN AKKOC MUSTAFAYEV<sup>1</sup>, UFUK BERBER<sup>2</sup>, ISMAIL YILMAZ<sup>2</sup>, SERKAN CELIK<sup>3</sup>, BALA BASAK OVEN<sup>3</sup> and MEHMET ALPASLAN OZGUN<sup>1</sup>

Departments of <sup>1</sup>Medical Oncology and <sup>2</sup>Pathology, University of Health Sciences, Sultan II. Abdulhamid Khan Educational and Research Hospital, 34660 Istanbul;

<sup>3</sup>Department of Medical Oncology, Yeditepe University Koşuyolu Hospital, 34718 Istanbul, Turkey

Received January 31, 2022; Accepted April 26, 2022

DOI: 10.3892/ol.2022.13347

**Abstract.** Aberrant expression of human leukocyte antigen G (HLA-G), an immunosuppressive molecule, has been observed in various cancer types. The exact mechanism of HLA-G expression is unclear. HLA-G expression is associated with low expression levels of microRNA (miR)-148a and miR-152. To the best of our knowledge, the prognostic value of the expression levels of miR-148a, miR-152 and HLA-G has not been investigated in colon cancer. The aim of the present study was to investigate the relationship between the presence of HLA-G molecules and the expression levels of miR-148 and miR-152 in colon cancer. In addition, the association of both HLA-G and miR expression with survival results and clinicopathological data was determined. An immunohistochemical method was used for detection of HLA-G expression in the tumor tissues and adjacent healthy tissues of 108 patients with colon cancer. Reverse transcription-quantitative PCR was used for miR analysis. HLA-G was expressed in 82 (75.9%) of the cancer samples, and there was no staining in normal tissues. HLA-G expression >20%, which was found in 41.7% (45/108) of the patients with colon cancer, was significantly positively associated with patient survival (P=0.039). Additionally, miR-148a levels were lower in the tumor tissues compared with the adjacent healthy tissues. No differences were found for miR-152 expression. In addition, miR-148a levels were found to be significantly lower (P=0.001) in HLA-G-positive tumor tissues than HLA-G-negative tumor tissues, but no such relationship was found for miR-152 levels. The presence of HLA-G expression was associated with poor survival outcomes. HLA-G is one of the prognostic factors in colon cancer, and decreased

miR-148a expression in colon cancer tissues may be associated with HLA-G-mediated carcinogenesis.

## Introduction

Colon cancer is one of the most common cancers in the world. Many mechanisms responsible for colon cancer have been identified in detail, including activation of oncogenic signaling pathways resulting from genetic mutations, inactivation of tumor suppressor genes, and epigenetic changes (1). Important features of tumor cells during colon cancer development are their ability to suppress of immune system cells and their avoidance of immune destruction. In this process, tumor cells secrete cytokines such as transforming growth factor beta, which reorganize the microenvironment and inhibit the infiltration of T cells. As a result of major histocompatibility complex class 1 mutations, tumor neoantigens are lost, and immune system's ability to recognize cancer cells decreases. Cytotoxic activity is suppressed by the overexpression of HLA-G on the cell surface as a consequence of the production of miR (2).

HLA-G is a non-classical major histocompatibility class 1b molecule, the presence of which was first demonstrated in placental extravillous trophoblast cells (3). HLA-G creates an immunotolerant environment and prevents fetal rejection by maternal immune system cells (4,5). While its physiological presence is limited to pancreatic island cells, thymic epithelial cells, activated monocytes, corneal keratocytes, and erythroblasts, its pathological expression has been demonstrated on the surface of various cancer cells (6). HLA-G can inhibit the activation and proliferation of T and B cells, the cytotoxic function of T and NK cells, and neutrophil functions. HLA-G shows an immunosuppressive feature by binding to various receptors on the cell surface, including immunoglobulin-like transcript receptor 2 on lymphoid and myelomonocytic cells (ILT-2) on lymphoid and myelomonocytic cells, ILT-4 on dendritic cells (4,5). While its physiological presence is limited to pancreatic island cells, thymic epithelial cells, activated monocytes, corneal keratocytes and erythroblasts, its pathological expression has been demonstrated on the surface of various cancer cells (6). HLA-G can inhibit the activation and proliferation of T and B cells, the cytotoxic function of

---

*Correspondence to:* Dr Levent Emirzeoglu, Department of Medical Oncology, University of Health Sciences, Sultan II. Abdulhamid Khan Educational and Research Hospital, Tibbiye Street, 34660 Istanbul, Turkey  
E-mail: emirzeoglulevent@hotmail.com

**Key words:** colon cancer, human leukocyte antigen G, microRNA-148a, microRNA-152, survival

T and NK cells, and neutrophil functions. HLA-G shows an immunosuppressive feature by binding to various receptors on the cell surface some of which are immunoglobulin-like transcript (ILT) receptor 2 on lymphoid and myelomonocytic cells, ILT-4 expressed by dendritic cells, and killer cell immunoglobulin-like receptor on natural killer cells (7-9).

An miR is a 19 to 23-nucleotide long, non-protein-coding, single-stranded RNA molecule. The miR acts as a post-transcriptional regulator by binding to the 3'UTR region of protein-coding genes. Thus, miRs control various cellular mechanisms, such as metabolism, proliferation, differentiation, apoptosis, tumor development, and metastasis (10). Recently, there has been an increase in studies on miRs in various types of cancer, including colon cancer. It has been reported that various miRs interact with genes such as *KRAS*, *NRAS*, and *BRAF*, which act as a kinase in intracellular signaling pathways and show pro- or anti-tumorigenic effects (11).

The mechanisms controlling HLA-G expression are largely unknown. The 3'UTR region in the HLA-G structure has an important role in regulating the expression of the molecule and functions as the target of miRs. Previous studies have shown that miR148-a and miR-152 bind to the 3'UTR terminal of the HLA-G molecule, and downregulate it. HLA-G overexpression is accompanied by low miR-148a and/or low miR-152 levels in these studies (12-14). Based on previous studies, we hypothesized that there may be a similar relationship of HLA-G expression with miR-148a and miR-152 levels in colon cancer.

## Materials and methods

**Patients and tissue samples.** Pairs of colon tumor and adjacent nontumorous tissues of 108 patients with colon cancer were collected. All patients were treated with surgery between 2013 and 2016 at the GATA Haydarpasa Training Hospital (GHTH). We included patients who were operated from the beginning of January 2013 to the end of December 2016 in our study. The last patient we included in the study was followed up for 5 years. Previous patients had longer follow-up. Our study protocol complied with the principles of the Declaration of Helsinki. Informed consent forms were obtained from all patients. Patients who did not agree to give informed consent for participating in the study were excluded. Formalin-fixed, paraffin-embedded tissue blocks were retrieved from the pathology archives. Follow-up was conducted every 3 months for the first 2 years, then every 6 months for 3 years, and then yearly. During follow-up visits, whole and routine blood examinations were performed, serum carcinoembryonic antigen levels were tested, and a complete physical examination was conducted. Patients also received abdominal ultrasound and chest X-rays or a chest, abdominal, and pelvic computerized tomography scan once yearly. Colonoscopy was performed 1 year after surgical resection and repeated in Years 3 and 5. The patients were followed for at least 5 years post-operation. Clinicopathological data were retrieved from the patients' medical files and pathology reports. In addition, patients' survival times were confirmed via telephone inquiry until September 2021. Time of recurrence or death was noted. Our study was approved by the ethics committee of GHTH with approval number 37/24.02.2016.

**Immunohistochemical staining.** We determined the expression level of HLA-G using the immunohistochemistry method. Blocks greater than 0.5 cm in diameter and containing at least 50% tumor were chosen for evaluation. We cut 4- $\mu$ m-thick sections from paraffin-embedded tissue blocks and mounted them on poly-lysine-coated slides. The slides were deparaffinized in xylene and rehydrated in 70% ethanol. Endogenous peroxidase activity was blocked using a 0.3% hydrogen peroxide solution containing methanol at room temperature for 30 min. Antigen retrieval was performed at 120°C (autoclave) for 5 min in a 10-nmol/l sodium citrate buffer (pH 6.0), and then anti-HLA-G mouse monoclonal antibody (4H84:sc-21799; Santa Cruz Biotechnology, Inc.; dilution range: 1/100) was incubated for 2 h. Afterwards, the samples were washed in a 0.01-M phosphate-buffered saline solution. Subsequently, the binding sites of the primary antibody were visualized using the ultraView Universal DAB Detection Kit (Ventana Company), and the Ventana BenchMark automated system; Ventana Company. Finally, sections were counterstained with hematoxylin and mounted with glycerol gelatin. The entire tumor area in the samples was examined by two experienced pathologists under a light microscope at 4x magnification. In cancer cells, HLA-G stained cytoplasmically. The percentage of staining in the entire area was recorded. The presence of HLA-G expression in colon cancer tissue was found in different proportions, from negative to 95% positivity. Overall, 75.9% (n=82) of colon cancer samples were HLA-G positive. Kaplan-Meier survival analysis was performed to determine the minimum proportion of HLA-G expression which reaches statistic significance to patient survival. The data showed that the cut-off value for HLA-G was 20%, which was significantly associated with patient survival (P=0.039). Therefore, 41.7% (n=45) of the patients were considered HLA-G positive.

**RNA extraction and polyadenylation.** Total RNA was isolated using the RecoverAll Total Nucleic Acid Isolation kit (Life Technologies; Thermo Fisher Scientific, Inc.) according to manufacturer's instructions. The quantity of the isolated RNA was assessed using a NanoDrop 1000 spectrophotometer; Thermo Fisher Scientific, Inc. Polyadenylation of total RNA was performed with *Escherichia coli* poly (A) polymerase (Life Technologies; Thermo Fisher Scientific, Inc.) at 37°C for 30 min according to the manufacturer's instructions for the poly(A) tailing kit. RNAs were stored at 70°C.

**Reverse transcriptase reaction.** Single-stranded cDNA was synthesized using the miScript II Reverse Transcription Kit (Qiagen Sample and Assay Technologies). A reverse transcriptase mixture was prepared by adding 4  $\mu$ l of a HiSpec buffer, 2  $\mu$ l of a miScript nucleic mix, 2  $\mu$ l of Script RT, 2  $\mu$ l of template RNA, and 10  $\mu$ l RNase-free water. This mixture was kept at 37°C for 60 min and then incubated at 95°C for 5 min to deactivate the reverse transcriptase. For reverse transcription-quantitative PCR (RT-qPCR) assays of miRNAs, cDNA was generated from equal amounts of total RNA per sample. The cDNAs were stored undiluted at 20°C.

**qPCR.** qPCR was done using a miScript SYBR-Green Kit (Qiagen Sample and Assay Technologies). The 25- $\mu$ l PCR mixture contained 10  $\mu$ l of Quntitect SYBR-Green PCR

master mix, 1  $\mu$ l of miScript universal forward and reverse primer, 5  $\mu$ l of template cDNA, and 8  $\mu$ l of RNase-free water. The nucleotide sequences for miR-148a were forward 5'-TCA GTGACTACAGA AACTTTGT-3' and reverse 5'-GCTGTC AACGATACGCTACGT-3' and for miR-152 were forward 5'-TCAGTGCATGACAGA AACTTGGAA-3' and reverse 5'-GCTGTCAACGATACGCTACGT-3'. All amplifications were conducted with a Rotor-Gene Q 5Plex real-time PCR system (Qiagen Sample and Assay Technologies). The cycles were completed with qPCR at 94°C for 5 sec, 58°C for 20 sec, and 72°C for 30 sec, with a total of 45 amplification cycles. In order to evaluate the specificity of the PCR products formed, melting curve analyses were performed between 55 and 95°C, with a sample temperature increase rate of 0.1°C/sec. RNU6B small nuclear RNA with serial dilutions was used as a reference for normalizing the expression levels of miR-148a and miR-152. It has been shown that the RNU6B mRNA level remains relatively stable in most systems in the human body and is not affected by various conditions such as cancer and infection (15). The relative expression levels of miRNAs were calculated using the  $2^{-\Delta\Delta CT}$  method (16). The RNU6B amplification curves were compared with the threshold values (CT:cycle threshold) of the amplification curve of each miRNA, and the CT of the target miRNAs were determined. For each miRNA,  $\Delta CT$  (Delta Cycle Threshold) values were calculated with the  $\Delta CT = CT \text{ miRNA} - CT \text{ U6RNA}$  formula in both tumor tissue and non-tumor tissue. The Rotor-Gene Q Series software v. 1.7 (Qiagen Sample and Assay Technologies) was used for this calculation. The relative expression ratio of miR-148a and miR-152 was presented as the fold change. A relative expression ratio of <1.0 was considered low expression in cancer cells relative to the nontumorous control.

**Statistical analysis.** Statistical analyses were performed using SPSS v. 25.0 (IBM Corp.). Disease-free survival (DFS) was calculated from the date of surgery to the date of disease relapse or death from any cause. Overall survival (OS) was calculated from the date of diagnosis to the date of death or last follow-up. Survival analyses were performed using the Kaplan-Meier analysis method. The effects of various prognostic factors related to tumor and patient characteristics on DFS and OS were investigated with the log-rank test. The effects of multiple prognostic factors on DFS and OS were investigated using the multivariate Cox regression test. Comparisons between groups for quantitative data were made with the Mann-Whitney U test and Kruskal-Wallis H test. Categorical comparisons between groups were calculated with Pearson's Chi-squared test, continuity correction, and Fisher's exact test. Spearman's correlation test was used to determine the level of correlation between miR-148a and miR-152 levels. Wilcoxon's signed-rank test was used to determine whether there was a difference between the levels. The results were evaluated at the 95% confidence interval and a significance level of  $P < 0.05$ .

## Results

**Patient characteristics.** Analyses of HLA-G and miRNA expression were performed on 108 patients with colon cancer stages I-III. The mean age at the time of diagnosis was

69.31 $\pm$ 10.35 years. Of the 108 samples analyzed, 60 were from male and 48 from female subjects. Patients' clinicopathological features are shown in Table I.

**HLA-G expression in colon cancer.** In the immunohistochemical staining, HLA-G was detected as a brown stained product in tumor tissue but not stained in normal colon tissue (Fig. 1). Comparing the HLA-G-positive and negative patient groups, the former group showed significantly higher rates of lymph node positivity (53 vs. 29%,  $P = 0.003$ ) and presence of perineural invasion (49 vs. 21%,  $P = 0.004$ ) and significantly lower pathologically early-stage disease rates (7 vs. 22%,  $P = 0.038$ ). In addition, miR-148a levels were significantly lower ( $P = 0.001$ ) in the HLA-G-positive tumor tissues than in negative ones (Table I). For miR-148a expression, levels were downregulated 2.5 fold or more in tumor tissue compared with normal colon tissue, but no such relationship was found for miR-152 expression (Table II).

**Survival analyses.** The mean follow-up period was 87.46 $\pm$ 26.29 months, and the median follow-up period was 90 (14-127) months. A total of 32 (29.6%) patients received adjuvant fluoropyrimidine-based chemotherapy. The rate of adjuvant therapy was significantly higher (46.6 vs. 17.4%,  $P = 0.041$ ) in the HLA-G positive group. The mean disease-free survival (DFS) time was 80.41 $\pm$ 34.13 months, and the median DFS time was 90 (7-127) months. During the follow-up period, 26 patients (24.1%) relapsed, 23 patients (21.3%) died due to disease, and 3 patients (2.8%) died due to other causes. According to the Kaplan-Meier survival analysis results, the 10-year DFS, and OS (overall survival) probabilities of the patients were 75.9, 72.8 and 65.2%, respectively. Patients with positive HLA-G expression had a lower 10-year DFS rate (92.1 vs. 53.3%,  $P < 0.001$ ). According to the results of the univariate Cox regression analysis, miR-148a levels were found to be associated with DFS ( $P = 0.014$ ) (Table III). According to the multivariate Cox regression analysis, the presence of HLA-G expression, grade 3 tumor, and N2 status were shown to be independent factors that negatively affected survival (Table IV). Survival curves according to HLA-G status are shown in Figs. 2 and 3.

## Discussion

In our study, we found that the level of miR-148a in cancer tissue was significantly lower than in healthy adjacent tissue, but the difference between miR-152 values was not significant. Specifically, among the 108 patients with colon cancer, miR-148a was weakly expressed in 82 cases (76%;  $P < 0.001$ ), and miR-152 was weakly expressed in 48 cases (44%;  $P = 0.123$ ). However, no significant differences in miR-152 expression was found between the tumor tissue and non-tumor tissue. The mean fold changes for miR-148a and miR-152 were 0.68 and 0.88, respectively. In a study conducted with 101 gastric cancer patients, it was shown that 70% of cases had low expression of miR-148a and 58% of cases had low expression of miRNA-152 when compared to nontumor tissue. The median fold change was 0.28 and 0.50, respectively (17). In a similar study, although miR-152 levels were found to be low in ovarian cancer tissue, no differences were found for miR-148a levels (18). Among

Table I. HLA-G expression and clinicopathological characteristics of patients.

Features	HLA-G expression		$\chi^2/Z$	P-value
	Negative (n=63)	Positive (n=45)		
Mean age (SD)	68.98 (10.59)	69.76 (10.10)	-0.069 <sup>a</sup>	0.945
Age group, n (%)				
<70 years	26 (41.3)	24 (53.3)	1.090 <sup>b</sup>	0.297
≥70 years	37 (58.7)	21 (46.7)		
Sex, n (%)				
Male	37 (58.7)	23 (51.1)	0.347 <sup>b</sup>	0.556
Female	26 (41.3)	22 (48.9)		
Tumor localization, n (%)				
Cecum	9 (14.3)	3 (6.7)	6.043 <sup>b</sup>	0.418
Ascending colon	8 (12.7)	9 (20.0)		
Hepatic flexura	5 (7.9)	5 (11.1)		
Transvers colon	4 (6.3)	2 (4.4)		
Splenic flexura	4 (6.3)	7 (15.6)		
Descending colon	12 (19.0)	9 (20.0)		
Sigmoid colon	21 (33.3)	10 (22.2)		
Tumor side, n (%)				
Right colon	26 (41.3)	19 (42.2)	0.000 <sup>b</sup>	>0.999
Left colon	37 (58.7)	26 (57.8)		
Histopathology, n (%)				
Adenocarcinoma	55 (87.3)	37 (82.2)	0.210 <sup>b</sup>	0.647
Mucinous carcinoma	8 (12.7)	8 (17.8)		
Tumor invasion, n (%)				
pT1/2	15 (23.8)	4 (8.9)	- <sup>c</sup>	0.071
pT3/4	48 (76.2)	41 (91.1)		
Nodal status, n (%)				
pN0	45 (71.4)	21 (46.7)	11.788 <sup>b</sup>	0.003 <sup>d</sup>
pN1	15 (23.8)	12 (26.7)		
pN2	3 (4.8)	12 (26.7)		
Stage, n (%)				
I	14 (22.2)	3 (6.7)	6.518 <sup>b</sup>	0.038 <sup>d</sup>
II	30 (47.6)	20 (44.4)		
III	19 (30.2)	22 (48.9)		
Grade, n (%)				
I/II	58 (92.1)	36 (80.0)	- <sup>c</sup>	0.084
III	5 (7.9)	9 (20.0)		
Lymphovascular invasion, n (%)				
Negative	42 (66.7)	22 (48.9)	2.739 <sup>b</sup>	0.098
Positive	21 (33.3)	23 (51.1)		
Perineural invasion, n (%)				
Negative	50 (79.4)	23 (51.1)	8.320 <sup>b</sup>	0.004 <sup>d</sup>
Positive	13 (20.6)	22 (48.9)		
Adjuvant chemotherapy, n (%)				
Yes	11 (17.4)	21 (46.6)		0.041 <sup>d</sup>
No	52 (82.6)	24 (53.4)		
Mean miR-148a expression in tumor tissues (SD)	0.83 (0.44)	0.43 (0.31)	-5.038 <sup>a</sup>	<0.001 <sup>d</sup>
Mean miR-152 expression in tumor tissues (SD)	0.86 (0.40)	0.82 (0.42)	-0.545 <sup>a</sup>	0.586

<sup>a</sup>Z, Mann-Whitney U test; <sup>b</sup> $\chi^2$ ,  $\chi^2$  test; <sup>c</sup> $\chi^2$ , Fisher's exact test. <sup>d</sup>P<0.05. HLA-G, human leukocyte antigen G; miR, microRNA.



Table II. miRNA levels in non-tumor and tumor tissues.

Variable	$\Delta$ CT mean (SD)	$\Delta$ CT median (minimum-maximum)	Z <sup>a</sup>	P-value
miR-148a				
Non-tumor	1.33 (0.57)	0.92 (0.02-3.02)	-5.238	<0.001 <sup>b</sup>
Tumor	0.66 (0.44)	0.68 (0.01-2.87)		
miR-152				
Non-tumor	0.84 (0.41)	0.85 (0.01-2.62)	-1.279	0.123
Tumor	0.93 (0.41)	0.88 (0.01-2.74)		

Z<sup>a</sup>=Wilcoxon Signed Ranks test  $\Delta$ CT=CT miRNA-CT U6RNA. <sup>b</sup>P<0.05.  $\Delta$ CT, delta cycle threshold; miRNA/miR, microRNA.




Figure 1. Representative image of immunohistochemical HLA-G expression. HLA-G showed cytoplasmic staining in tumor tissues. Scale bar, 500  $\mu$ m (A, C and E) or 100  $\mu$ m (B, D and F). (A) Negative HLA-G staining in cancer tissue (magnification, x4). (B) Negative HLA-G staining in cancer tissue (magnification, x20). (C) HLA-G staining <20% in cancer tissue (magnification, x4). (D) HLA-G staining <20% in cancer tissue (magnification, x20). (E) Positive HLA-G staining in cancer tissue (magnification, x4). (F) Positive HLA-G staining in cancer tissue (magnification, x20). HLA-G, human leukocyte antigen G.

Table III. Five-year survival rates of patients.

Category	No.	5-year DFS, %	P-value	5-year DSS, %	P-value	5-year OS, %	P-value
Age group							
<70 years	50	72.8	0.292	72.8	0.445	70.1	0.558
≥70 years	58	79.3		78.1		60.2	
Sex							
Male	60	71.7	0.266	70.1	0.174	67.8	0.163
Female	48	81.2		84.5		73.9	
Tumor localization							
Cecum	12	66.7	0.625	64.3	0.353	64.3	0.143
Ascending colon	17	88.2		80.2		80.1	
Hepatic flexura	10	70.2		70.5		70.4	
Transvers colon	6	83.3		83.3		83.3	
Splenic flexura	11	63.6		63.6		63.6	
Descending colon	21	81.3		86.6		86.6	
Sigmoid colon	31	74.2		80.1		70.4	
Tumor side							
Right colon	45	77.8	0.673	81.7	0.702	62.3	0.889
Left colon	63	74.6		75.9		68.2	
Histopathology							
Adenocarcinoma	92	75.8	0.634	78.1	0.835	66.4	0.851
Mucinous carcinoma	16	81.3		81.3		81.3	
Invasion							
pT1/2	19	100	0.011 <sup>a</sup>	100	0.021 <sup>a</sup>	100	0.014 <sup>a</sup>
pT3/4	89	70.8		71.4		59.1	
Nodal status							
pN0	66	84.8	<0.001 <sup>a</sup>	86.8	<0.001 <sup>a</sup>	82.7	<0.001 <sup>a</sup>
pN1	27	81.5		81.5		76.4	
pN2	15	26.7		37.3		37.3	
Stage							
I	17	100	0.009 <sup>a</sup>	100	0.023 <sup>a</sup>	100	0.003 <sup>a</sup>
II	50	78.2		80.5		75.5	
III	41	63.4		67.2		55.1	
Grade							
I/II	94	79.8	0.002 <sup>a</sup>	82.4	0.006 <sup>a</sup>	71.4	0.001 <sup>a</sup>
III	14	50.3		57.1		57.1	
LVI							
Negative	64	84.4	0.007 <sup>a</sup>	84.4	0.001 <sup>a</sup>	82.6	<0.001 <sup>a</sup>
Positive	44	63.6		64.2		53.5	
PNI							
Negative	73	83.6	0.003 <sup>a</sup>	84.0	0.001 <sup>a</sup>	76.3	<0.001 <sup>a</sup>
Positive	35	60.0		61.0		56.6	
HLA-G expression							
Negative	63	92.1	<0.001 <sup>a</sup>	93.1	<0.001 <sup>a</sup>	76.8	<0.001 <sup>a</sup>
Positive	45	53.3		58.1		48.4	
miR-148a tumor							
≤0.7	62	69.4	0.005 <sup>a</sup>	79.0	0.028 <sup>a</sup>	69.7	0.070
>0.7	46	89.1		91.1		80.1	
miR-148a non-tumor							
<1	62	88.7	0.006 <sup>a</sup>	93.5	0.005 <sup>a</sup>	89.6	0.021 <sup>a</sup>
≥1	46	65.2		76.1		64.6	
miR-152 tumor							
<1	74	78.4	0.542	83.4	0.914	77.0	0.745
≥1	34	82.4		85.3		78.6	

Table III. Continued.

Category	No.	5-year DFS, %	P-value	5-year DSS, %	P-value	5-year OS, %	P-value
miR-152 non-tumor							
<1	80	77.5	0.347	79.3	0.200	70.1	0.385
≥1	28	82.1		84		77.0	

<sup>a</sup>P<0.05. DFS, disease-free survival; DSS, disease-specific survival; HLA-G, human leukocyte antigen G; HR, hazard ratio; LVI, lymphovascular invasion; miR, microRNA; OS, overall survival; PNI, perineural invasion.

Table IV. Results of multivariate Cox regression analysis.

Factors	Disease-free survival		Disease-specific survival		Overall survival	
	HR (95% CI)	P-value	HR (95% CI)	P-value	HR (95% CI)	P-value
Nodal status						
pN0	1 (Reference)		1 (Reference)		1 (Reference)	
pN1	1.10 (0.33-3.69)	0.876	1.25 (0.32-4.81)	0.749	1.53 (0.45-5.26)	0.500
pN2	5.94 (1.92-18.42)	0.002 <sup>a</sup>	2.09 (0.57-7.68)	0.269	2.65 (0.75-9.34)	0.129
HLA-G expression						
Negative	1 (Reference)		1 (Reference)		1 (Reference)	
Positive	6.07 (1.75-20.99)	0.004 <sup>a</sup>	5.19 (1.49-18.11)	0.010 <sup>a</sup>	3.37 (1.19-9.56)	0.022 <sup>a</sup>
Tumor grade						
I/II	1 (Reference)		1 (Reference)		1 (Reference)	
III	3.56 (1.20-10.58)	0.022 <sup>a</sup>	1.55 (0.50-4.75)	0.445	1.84 (0.67-5.10)	0.239
LVI						
Negative	1 (Reference)		1 (Reference)		1 (Reference)	
Positive	2.14 (0.41-11.16)	0.366	3.79 (0.67-21.65)	0.134	2.12 (0.44-10.32)	0.352
PNI						
Negative	1 (Reference)		1 (Reference)		1 (Reference)	
Positive	2.15 (0.40-11.49)	0.373	1.60 (0.27-9.43)	0.605	1.05 (0.24-4.65)	0.952
miR-148a tumor						
≤0.7	1 (Reference)		1 (Reference)		1 (Reference)	
>0.7	0.62 (0.20-1.93)	0.407	0.76 (0.25-2.36)	0.636	0.78 (0.30-2.08)	0.624
miR-148a non-tumor						
<1	1 (Reference)		1 (Reference)		1 (Reference)	
≥1	1.26 (0.52-3.09)	0.612	1.86 (0.72-4.75)	0.198	1.47 (0.62-3.49)	0.348

<sup>a</sup>P<0.05; Cox Regression Analysis (Method=Enter). 95% CI, 95% confidence interval; HLA-G, human leukocyte antigen G; HR, hazard ratio; LVI, lymphovascular invasion; miR, microRNA; PNI, perineural invasion.

the 78 patients with ovarian cancer, miR-152 was weakly expressed in 52 cases, and miR-148a was weakly expressed in 49 cases. However, no significant differences in miR-148a expression were found between the ovarian cancer and normal epithelium tissues. The mean fold-change was 0.70 and 0.37, respectively. In addition to these studies, suppressed miR-148a expression has been demonstrated in gastrointestinal cancers such as gastric cancer (19), colorectal cancer (20,21), pancreatic cancer (22), and esophageal cancer (23). Similarly, miRNA levels were similar in all other studies. Thus, our results were in agreement with previous studies of gastrointestinal cancers

in terms of miRNA levels. In addition, we found that miR-148a expression was lower in HLA-G-positive tumor tissue compared to HLA-G-negative tumor tissue. In a study on the role of miRNAs on HLA-G expression in kidney cancer, an inverse correlation was observed for miR-148a and miR133a with HLA-G protein expression *in situ* and *in vitro* (24). In a similar study in esophageal cancer, HLA-G miRNA and HLA-G protein levels were decreased in EC9706 cells transfected with miR-148a (25).

In this study, we showed that the HLA-G protein was expressed at varying rates in 75.9% of colon cancer samples.






Figure 2. Association between HLA-G status and DFS. DFS was shorter ( $P < 0.001$ ) in the HLA-G-positive group compared with the HLA-G-negative group. Cum, cumulative; DFS, disease-free survival; HLA-G, human leukocyte antigen G.




Figure 3. Association between HLA-G status and OS. OS was shorter ( $P < 0.001$ ) in the HLA-G-positive group compared with the HLA-G-negative group. Cum, cumulative; HLA-G, human leukocyte antigen G; OS, overall survival.

HLA-G expression above 20%, which was found in 41.7% ( $n=45$ ) of patients, was significantly correlated with patient survival ( $P=0.039$ ). Previous studies have found that HLA-G expression varied between 50 and 80% in colon cancer and other gastrointestinal system cancers (26,27). Those studies also found no HLA-G expression in normal colon tissue, similar to our study.

Numerous studies have demonstrated that high HLA-G expression was associated with poor prognosis in gastrointestinal cancer patients (28-32). However, there are few studies showing that there is no significant relationship between HLA-G expression and the prognosis of gastrointestinal cancers (33-35). These differences between studies may be related to the autoantibody used. Lin *et al* (36) demonstrated that HLA-G expression was related to a poor prognosis when detected with antibody 4H84, but the results were inconsistent when using the 5A6G7 antibody. We used 4H84 antibodies

for HLA-G detection in our study. The 4H84 antibody has been confirmed by international conferences as a reference tool for evaluating HLA-G expression in paraffin-embedded specimens (37). Limited studies of other antibodies could not justify the lack of prognostic value of HLA-G in colon cancer. Moreover, the invasive nature of the disease and the tumor microenvironment are different in gastrointestinal cancers, and the expression of HLA-G differs between different cancer types, which may explain the differing results between studies (38,39). The number of studies showing that HLA-G expression is a poor prognostic factor in colon cancer is increasing. Ye *et al* (40) studied HLA-G expression in colorectal carcinoma using immunohistochemistry and observed positive HLA-G expression in 64.6% of colorectal patients. Their multivariate analysis showed that HLA-G may serve as an independent prognostic factor for colorectal cancer patients. Guo *et al* (41) showed that HLA-G expression is significantly correlated with OS and can serve as an independent factor in OS. Zhang *et al* (39) showed that different rates of HLA-G expression in colorectal cancer patients affect patient survival and that a combination of HLA-G expression status with traditional clinical risk factors can improve the prediction of specific clinical outcomes in subpopulations of colorectal cancer patients. The results of current studies suggest that HLA-G expression is most likely associated with the prognosis of colorectal cancers. Similarly, our study showed that HLA-G-positive patients had worse outcomes in terms of both OS and DFS than HLA-G-negative patients.

In conclusion, we found that HLA-G is expressed in many colorectal carcinomas, and miR-148a levels in HLA-G-positive colon cancer tissues are lower than in HLA-G-negative tissues. This is the first study to investigate the relationship between miR-148a, miR-152 and HLA-G in colon cancer. Using the Kaplan-Meier analysis, we found a significant correlation between HLA-G expression and OS in colorectal cancer patients. Furthermore, our multivariate analysis results showed that the expression HLA-G can serve as an independent factor for OS. According to the results of the univariate Cox regression analysis, miR-148a levels were found to be a factor associated with DFS. Low expression of miR-148a in tumor tissue is associated with shorter DFS. Our results suggest that HLA-G and miR-148a may be new prognostic markers in colon cancer. Further investigation is needed to confirm these results.

#### Acknowledgements

Not applicable.

#### Funding

The present study was funded by GATA Haydarpasa Training Hospital (grant no. DS2016TO002).

#### Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.



### Authors' contributions

LE and OO designed the study and prepared the manuscript. LE and OO confirm the authenticity of all the raw data. FNAM reviewed medical documents and collected patients' clinical characteristics and was involved in the analysis of data and validation. UB, IY, BBO, SC and MAO contributed to study design, data analysis and manuscript editing. All authors have read and approved the final manuscript.

### Ethics approval and consent to participate

All the procedures implemented in studies involving human participants were consistent with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki Declaration and its later amendments or comparable ethical standards. All patients provided written informed consent for participation and patient anonymity was preserved. This retrospective study was approved by the ethics committee of GATA Haydarpaşa Training Hospital (approval number 2016/37; Istanbul, Turkey).

### Patient consent for publication

Written informed consent was obtained from the patients for publication of the data.

### Competing interests

The authors declare that they have no competing interests.

### References

- Raskov H, Pommergaard HC, Burcharth J and Rosenberg J: Colorectal carcinogenesis-update and perspectives. *World J Gastroenterol* 20: 18151-18164, 2014.
- Yi M, Xu L, Jiao Y, Luo S, Li A and Wu K: The role of cancer-derived microRNAs in cancer immune escape. *J Hematol Oncol* 13: 1-14, 2020.
- Kovats S, Main EK, Librach C, Stubblebine M, Fisher SJ and DeMars R: A class I antigen, HLA-G, expressed in human trophoblasts. *Science* 248: 220-223, 1990.
- Carosella ED, Moreau P, Le Maoult J, Le Discorde M, Dausset J and Rouas-Freiss N: HLA-G molecules: From maternal-fetal tolerance to tissue acceptance. *Ad Immunol* 81: 199-252, 2003.
- Tilburgs T, Evans JH, Crespo AC and Strominger JL: The HLA-G cycle provides for both NK tolerance and immunity at the maternal-fetal interface. *Proc Natl Acad Sci USA* 112: 13312-13317, 2015.
- Loustau M, Anna F, Dréan R, Lecomte M, Langlade-Demoyen P and Caumartin J: HLA-G neo-expression on tumors. *Front Immunol* 11: 1685, 2020.
- Carosella ED, Moreau P, LeMaoult J and Rouas-Freiss N: HLA-G: From biology to clinical benefits. *Trends Immunol* 29: 125-132, 2008.
- Favier B, LeMaoult J, Lesport E and Carosella ED: ILT2/HLA-G interaction impairs NK-cell functions through the inhibition of the late but not the early events of the NK-cell activating synapse. *FASEB J* 24: 689-699, 2010.
- Morandi F, Ferretti E, Bocca P, Prigione I, Raffaghello L and Pistoia V: A novel mechanism of soluble HLA-G mediated immune modulation: Downregulation of T cell chemokine receptor expression and impairment of chemotaxis. *PLoS One* 5: e11763, 2010.
- Calin GA and Croce CM: MicroRNA-cancer connection: The beginning of a new tale. *Cancer Res* 66: 7390-7394, 2006.
- Yaghoubi N, Avval FZ, Khazaei M and Aghaee-Bakhtiari SH: MicroRNAs as potential investigative and predictive biomarkers in colorectal cancer. *Cell Signal* 80: 109910, 2021.
- Manaster I, Goldman-Wohl D, Greenfield C, Nachmani D, Tsukerman P, Hamani Y, Yagel S and Mandelboim O: MiRNA-mediated control of HLA-G expression and function. *PLoS One* 7: e33395, 2012.
- Tan Z, Randall G, Fan J, Camoretti-Mercado B, Brockman-Schneider R, Pan L, Solway J, Gern JE, Lemanske RF, Nicolae D and Ober C: Allele-specific targeting of microRNAs to HLA-G and risk of asthma. *Am J Hum Genet* 81: 829-834, 2007.
- Zhu XM, Han T, Wang XH, Li YH, Yang HG, Luo YN, Yin GW and Yao YQ: Overexpression of miR-152 leads to reduced expression of human leukocyte antigen-G and increased natural killer cell mediated cytotoxicity in JEG-3 cells. *Am J Obstet Gynecol* 202: 592-e1-7, 2010.
- Peskoe SB, Barber JR, Zheng Q, Meeker AK, De Marzo AM, Platz EA and Lupold SE: Differential long-term stability of microRNAs and RNU6B snRNA in 12-20 year old archived formalin-fixed paraffin-embedded specimens. *BMC Cancer* 17: 1-7, 2017.
- Livak KJ and Schmittgen TD: Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) method. *Methods* 25: 402-408, 2001.
- Chen Y, Song Y, Wang Z, Yue Z, Xu H, Xing C and Liu Z: Altered expression of MiR-148a and MiR-152 in gastrointestinal cancers and its clinical significance. *J Gastrointest Surg* 14: 1170-1179, 2010.
- Zhou X, Zhao F, Wang ZN, Song YX, Chang H, Chiang Y and Xu HM: Altered expression of miR-152 and miR-148a in ovarian cancer is related to cell proliferation. *Oncol Rep* 27: 447-454, 2012.
- Zheng G, Xiong Y, Xu W, Wang Y, Chen F, Wang Z and Yan Z: A two-microRNA signature as a potential biomarker for early gastric cancer. *Oncol Lett* 7: 679-684, 2014.
- Takahashi M, Cuatrecasas M, Balaguer F, Hur K, Toiyama Y, Castells A, Boland CR and Goel A: The clinical significance of MiR-148a as a predictive biomarker in patients with advanced colorectal cancer. *PLoS One* 7: e46684, 2012.
- Tsai HL, Yang IP, Huang CW, Ma CJ, Kuo CH, Lu CY, Juo SH and Wang JY: Clinical significance of microRNA-148a in patients with early relapse of stage II stage and III colorectal cancer after curative resection. *Transl Res* 162: 258-268, 2013.
- Hanoun N, Delpu Y, Suriawinata AA, Bournet B, Bureau C, Selves J, Tsongalis GJ, Dufresne M, Buscail L, Cordelier P and Torrisani J: The silencing of microRNA 148a production by DNA hypermethylation is an early event in pancreatic carcinogenesis. *Clin Chem* 56: 1107-1118, 2010.
- Hummel R, Hussey DJ, Michael MZ, Haier J, Bruewer M, Senninger N and Watson DI: MiRNAs and their association with locoregional staging and survival following surgery for esophageal carcinoma. *Ann Surg Oncol* 18: 253-260, 2011.
- Jasinski-Bergner S, Stoehr C, Bukur J, Massa C, Braun J, Hüttelmaier S, Spath V, Wartenberg R, Legal W, Taubert H, *et al*: Clinical relevance of miR-mediated HLA-G regulation and the associated immune cell infiltration in renal cell carcinoma. *Oncoimmunology* 4: e1008805, 2015.
- Chen Q, Luo G and Zhang X: MiR-148a modulates HLA-G expression and influences tumor apoptosis in esophageal squamous cell carcinoma. *Exp Ther Med* 14: 4448-4452, 2017.
- Hansel DE, Rahman A, Wilentz RE, Shih IM, McMaster MT, Yeo CJ and Maitra A: HLA-G upregulation in pre-malignant and malignant lesions of the gastrointestinal tract. *Internat J Gastrointest Cancer* 35: 15-23, 2005.
- Swets M, König MH, Zaalberg A, Dekker-Ensink NG, Gelderblom H, van de Velde CJ, van den Elsen PJ and Kuppen PJK: HLA-G and classical HLA class I expression in primary colorectal cancer and associated liver metastases. *Hum Immunol* 77: 773-779, 2016.
- Murdaca G, Calamaro P, Lantieri F, Pigozzi S, Mastracci L, Grillo F, Magnani O, Ceppà P, Puppo F and Fiocca R: HLA-G expression in gastric carcinoma: Clinicopathological correlations and prognostic impact. *Virchows Arch* 473: 425-433, 2018.
- Shahraki PK, Alavian-Mehr A and Farjadian S: HLA-G: Facts and fictions. *Asian Pac J Cancer Biol* 3: 37-45, 2018.
- Tuncel T, Karagoz B, Haholu A, Ozgun A, Emirzeoglu L, Bilgi O and Kandemir EG: Immunoregulatory function of HLA-G in gastric cancer. *Asian Pac J Cancer Prev* 14: 7681-7684, 2013.
- Wang Y, Ye Z, Meng XQ and Zheng SS: Expression of HLA-G in patients with hepatocellular carcinoma. *Hepatobiliary Pancreat Dis Int* 10: 158-163, 2011.

32. Peng Y, Xiao J, Li W, Li S, Xie B, He J and Liu C: Prognostic and clinicopathological value of human leukocyte antigen G in gastrointestinal cancers: A meta-analysis. *Front Oncol* 11: 642902, 2021.
33. Ishigami S, Natsugoe S, Miyazono F, Nakajo A, Tokuda K, Matsumoto M, Okumura H, Douchi T, Hokita S and Aikou T: HLA-G expression in gastric cancer. *Anticancer Res* 26: 2467-2472, 2006.
34. Leelawat K, Engprasert S, Pongchai-rerk U, Suthipintawong C and Leardkamolkarn V: No expression of human leukocyte antigen G (HLA-G) in colorectal cancer cells. *J Med Assoc Thai* 87: 816-818, 2004.
35. Reimers MS, Engels CC, Putter H, Morreau H, Liefers GJ, van de Velde CJ and Kuppen PJK: Prognostic value of HLA class I, HLA-E, HLA-G and Tregs in rectal cancer: A retrospective cohort study. *BMC Cancer* 14: 1-12, 2014.
36. Lin A, Chen HX, Zhu CC, Zhang X, Xu HH, Zhang JG, Wang Q, Zhou WJ and Yan WH: Aberrant human leukocyte antigen-G expression and its clinical relevance in hepatocellular carcinoma. *J Cell Mol Med* 14: 2162-2171, 2010.
37. Lin A, Zhang X, Zhou WJ, Ruan YY, Xu DP, Wang Q and Yan WH: Human leukocyte antigen-G expression is associated with a poor prognosis in patients with esophageal squamous cell carcinoma. *Int J Cancer* 129: 1382-1390, 2011.
38. Farjadian S, Tabebordbar M, Mokhtari M, Safaei A, Malekzadeh M and Ghaderi A: HLA-G expression in tumor tissues and soluble HLA-G plasma levels in patients with gastrointestinal cancer. *Asian Pac J Cancer Prev* 19: 2731-2735, 2018.
39. Zhang RL, Zhang X, Dong SS, Hu B, Han QY, Zhang JG, Zhou WJ, Lin A and Yan WH: Predictive value of different proportion of lesion HLA-G expression in colorectal cancer. *Oncotarget* 8: 107441-107451, 2017.
40. Ye SR, Yang H, Li K, Dong DD, Lin XM and Yie SM: Human leukocyte antigen G expression: As a significant prognostic indicator for patients with colorectal cancer. *Mod Pathol* 20: 375-383, 2017.
41. Guo ZY, Lv YG, Wang L, Shi SJ, Yang F, Zheng GX, Wen WH and Yang AG: Predictive value of HLA-G and HLA-E in the prognosis of colorectal cancer patients. *Cell Immunol* 293: 10-16, 2015.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.