

RESEARCH ARTICLE

Open Access

Enterovirus genotypes causing hand foot and mouth disease in Shanghai, China: a molecular epidemiological analysis

Menghua Xu, Liyun Su, Lingfeng Cao, Huaqing Zhong, Niuniu Dong and Jin Xu*

Abstract

Background: A rapid expansion of hand, foot, and mouth disease (HFMD) outbreaks has occurred and caused deaths in China in recent years, but little is known about the other etiologic agents except enterovirus 71 (EV71) and coxsackievirus A 16 (CA16). The objective of this study is to determine the genotype compositions of enterovirus causing HFMD in Shanghai and identify any associations between enterovirus types and clinical manifestations.

Methods: Stool specimens were collected from patients hospitalized for treatment of HFMD, from May 2010 to April 2011. Enterovirus was detected by reverse transcription PCR and directly genotyped by sequencing the PCR products. Phylogenetic analysis was based on the VP1 partial gene.

Results: Of 290 specimens, 277 (95.5%) tested positive for enterovirus. The major genotypes were EV71 (63.8%), CA10 (9.0%), CA6 (8.3%), CA16 (6.9%), CA12 (2.4%), and CA4 (1.4%). The EV71 strains belonged to the C4a subtype and CA16 belonged to the B subtype. CA6 was closely related to strains detected in Japan, Taiwan and China, and CA10, CA12 and CA4 were phylogenetically similar to other strains circulating in China. Mean hospital stays and the prevalence of complications in patients with EV71 infection were higher than those in patients in CA6, CA10 or CA16 infection ($P < 0.05$ for all comparisons). Children with CA12 infection were the youngest, and most likely have the highest risk of complications when compared to the other non-EV71 infection groups.

Conclusions: This study demonstrated a diversified pathogen compositions attributing to HFMD and clinical symptoms differing in enterovirus genotypes. It deserves our attention as early identification of enterovirus genotypes is important for diagnosis and treatment of HFMD patients.

Keywords: Hand foot and mouth disease (HFMD), Human enterovirus (HEV), Genotype

Background

Hand, foot, and mouth disease (HFMD), first reported in New Zealand in 1957, is a global and common infectious disease in young children, particularly in those less than 5 years old. Vesicular exanthema develops on the hands, feet, mouth and buttocks. The disease usually resolves spontaneously, but severe complications, including death, can occur [1-4]. Large-scale outbreaks have occurred in China and caused 126 and 353 deaths in the year of 2008 and

2009, respectively (http://www.chinacdc.cn/tjsj/fdcrrbbg/201002/t20100224_25293.htm) [1-7].

Hand, foot, and mouth disease is caused by human enterovirus, a genus of the *Picornavirus* family, and is characterized by a single 7.4kb of positive-strand genomic RNA. The genomes contain only a single long open reading frame that encodes four structural proteins (VP1 through VP4) and seven non-structural proteins (2A, 2B, 2C, 3AB, 3C and RNA-dependent RNA polymerase 3D), which are responsible for viral replication and protein processing. The VP1 is the most external and immunodominant *picornavirus* capsid protein and contains the major neutralization epitopes that has been used for virus serotype identification and evolutionary studies [8-10].

* Correspondence: jin030101@aliyun.com
Laboratory Medicine Center, Pediatric Institute, Children's Hospital of Fudan University, Shanghai 201102, China

Enterovirus genotypes are classified into four species, A, B, C and D, on the basis of genome organization, sequence similarity and biological properties [11]. Enterovirus 71 (EV71) and coxsackievirus A16 (CA16) belonging to species A are the most common causes of HFMD worldwide [1]. Severe EV71-associated clinical syndromes, such as acute flaccid paralysis, brainstem encephalitis, rapid fatal pulmonary edema, and hemorrhage, have been observed in many outbreaks and is considered an important cause of severe HFMD [2,3]. CA16 associated HFMD has a milder outcome and a much lower incidence of severe complications [12].

Though HFMD has become a public issue in China, little is known about the other etiologic agents except EV71 and CA16. Sporadic reports demonstrated that various enterovirus genotypes, such as CA2, CA4, CA5, CA6, CA8, CA10 and CA12 from species A, CB2 and CB4 from species B and some echoviruses can cause HFMD [13-17]. However, studies on the relationship of these genotypes with clinical symptoms are lacking.

In this study, we aimed to investigate the genotypes of enterovirus contributing to HFMD and to identify any associations between enterovirus types and clinical manifestations in Shanghai, China.

Methods

Sample collection

From 1 May 2010 to 30 April 2011, stool specimens were collected from patients hospitalized for HFMD at children's Hospital of Fudan University, Shanghai, China. All the patients were diagnosed by the Ministry of Health diagnostic criteria (<http://www.nhfpc.gov.cn/yzygj/s3593g/201306/6d935c0f43cd4a1fb46f8f71acf8e245.shtml>). The complications involve diseases in neurologic, respiratory or circulatory system that caused by HFMD, such as aseptic meningitis, encephalitis, acute flaccid paralysis, pulmonary oedema or cardiorespiratory failure, which were defined in "A Guide to Clinical Management and Public Health Response for Hand, Foot and Mouth Disease (HFMD)" (<http://www.wpro.who.int/publications/docs/GuidancefortheclinicalmanagementofHFMD.pdf>).

Patients' demographic data, clinical symptoms, and major complications were collected retrospectively from medical history. This study was approved by the Ethics Committee in the Children's Hospital of Fudan University. Because the specimens were collected in the normal course of patient care, no informed consent was required according to the Ethics Committee.

Human enterovirus testing

From supernatants of 10% (V/V) stool specimens, RNA was extracted by Trizol (Invitrogen, CA, USA) according to the manufacturer's instructions. The RNA was dissolved in 20 μ L DEPC (Diethylpyrocarbonate) water. cDNA was synthesized using 4 μ L of extracted RNA, 100 μ mol of random

primers, and 2.5 U of reverse transcriptase (PrimeScript™ RT kit, Takara, Dalian, China). The reverse transcriptase reaction was carried out at 37°C for 30 min and 85°C for 5 s.

Human enterovirus was preliminarily detected with highly conserved 5'UTR primers. The first PCR step was performed in a reaction volume of 25 μ L, including 2 μ L cDNA, 0.5 μ mol each of outer primers (EV1F: 5'-CGGCCCTGAATGCGGC-3', EV1R: 5'-CACCGGATGGCCAATCCA-3'), 50 μ mol dNTP, and 0.75U of ExTaq DNA polymerase (Takara, Dalian, China). The PCR protocol was to use an initial temperature of 94°C for 1 min, followed by 35 cycles of 94°C for 30 s, 55°C for 30s, 72°C for 30 s, and 72°C for 7 min. Then 1 μ L of DNA from the first round of PCR was used as the template in the second round of PCR with the inner primers (EV2F: 5'-CCCCTGAATGCGGCTAAT-3', EV2R: 5'-ATTGTCACATAAGCAGCCA-3') under the same reaction system and cycling conditions. The PCR products were subjected to electrophoresis in a 2% agarose gel to identify positive samples with a predicted size of 146 bp.

Enterovirus genotyping

For enterovirus genotyping, samples tested positive by the 5'UTR were amplified by nested PCR using primers from the VP1 junction region of enterovirus as described [18]. For the first round of PCR, 2 μ L of cDNA was used in a volume of 25 μ L containing 0.5 μ mol each of outer primers, 50 μ mol of dNTP, and 0.75U of ExTaq DNA polymerase in the following cycling conditions: 35 cycles of 94°C for 30 s, 54°C for 45 s, 72°C for 30 s, and a final incubation of 72°C for 7 min. A volume of 2 μ L of the product was used for the second round of PCR with 0.5 μ mol each of inner primers, 50 μ mol of dNTP, and 0.75U of ExTaq DNA polymerase in a volume of 25 μ L undergoing the same PCR conditions as described above. The final product of the different groups of enterovirus was visualized by agarose gel electrophoresis. DNA fragments were purified and the nucleotide sequence of each PCR product was bi-directional sequenced on a 3730 sequencer (Pekin-Elmer Applied Biosystems, Foster City, CA). The PCR products used for the sequencing were about 683 bp for the group A, 619 bp for the group B and 497 bp for the 5'UTR.

Phylogenetic analysis and sequence submission

Sequence analysis of the PCR product of each strain was analyzed with SeqScanner software (Applied Biosystems, USA), and genetic identity was determined by comparing the sequence with standard strains in Genbank (US National Center for Biotechnology Information, NCBI). A multiple-sequence alignment was constructed using ClustalW and phylogenetic trees were performed applying the neighbor-joining method with the 1000-bootstrap re-sampling implemented in the Molecular

Evolutionary Genetics Analysis, version 5.0 program. All the VP1 gene sequences were submitted to GenBank sequence database (NCBI, Bethesda, MD, USA) with given accession numbers KC834832-KC834854 for CA6, KC834855-KC834877 for CA10, KC834878-KC834892 for CA16 and KC834893-KC835055 for EV71. The phylogenetic trees have been deposited in TreeBASE (# 14766).

Statistical methods

The means of the continuous variables and the proportions of the categorical variables were compared using

Student's *t* test, χ^2 tests, and Fisher's exact test, as appropriate. All data met the assumptions of the tests used to analyze them. Alpha was set at 0.05. Data analysis was performed using SPSS version 17.0 for windows.

Results

Enterovirus genotypes

Of 290 specimen, 277 (95.5%) tested positive for enterovirus. Except for 8 specimens that could not be typed, nine different genotypes were identified with different




Figure 1 Phylogenetic analysis with the neighbor-joining method based on the alignment of the 468 nucleotide VP1 gene of 163 Shanghai EV71 strains. The 59 reference strains were obtained from the US National Center for Biotechnology Information's Genbank.

detection rates: EV71 (63.8%, 185/290), CA10 (9.0%, 26/290), CA6 (8.3%, 24/290), CA16 (6.9%, 20/290), CA12 (2.4%, 7/290), CA4 (1.4%, 4/290), CA14 (0.3%, 1/290), Echo6 (0.3%, 1/290), and HEV-C (0.3%, 1/290). No mixed infections were found.

Phylogenetic analysis of enterovirus genotypes

The partial VP1 sequence of the 163 EV71 strains from each of the patients was used for phylogenetic analysis.

The nucleotide homologies in all the EV71 strains ranged from 90.7% to 100%. The molecular epidemiology of the Shanghai EV71 strains was determined with a phylogenetic dendrogram, with the reference EV71 strains from Genbank representing all the known sub-genotypes (A, B0-B5, C1-C5). The dendrogram showed that all the EV71 strains clustered in the same lineages as a sub-genotype C4. The sequences in sub-genotype C4 could be further divided into C4a and C4b clusters.





Figure 2 Phylogenetic analysis based on the alignment of the 488 nucleotide VP1 gene of 15 Shanghai CA16 strains. The 25 reference strains were obtained from the US National Center for Biotechnology Information's Genbank.

The Shanghai EV71 strains clustered in C4a, closely related to strains like Fuyang 22, DTID/ZJU-62 which were detected in China in 2008 (Figure 1).

The nucleotide homologies within all the CA16 strains ranged from 88.4% to 99.1%. Phylogenetic analysis of the partial VP1 coding sequences showed all the CA16

strains belonged to the B genotype. Six sequences that were closely related to strains circulating in China (such as HK08-7 and QH0269T) belonged to a B1a cluster, and nine sequences belonged to a B1b cluster, which are closely related to strains identified in Vietnam, Australia, Malaysia and China (Figure 2).



Of the 23 CA6 partial VP1 gene sequences used for phylogenetic analysis, the nucleotide homologies ranged from 89.8% to 99.0%. The CA6 strains appeared in two branches, 15 of which were closely related to the 09s81 and 09s82 strains detected in Japan in 2009, and another 8 strains were closely related to strains circulating in China and Taiwan (Figure 3).

The nucleotide homologies of CA10 strains ranged from 95.8% to 99.5%. These strains were closely related to the CA10 strains found in China in 2009 and distantly related to the CA10 strain reported in China in 2004 and 2006 and to strains in France and India (Figure 4).

Both the CA12 and CA4 strains were most closely related to the other CA12 and CA4 strains found previously in Jiangsu and Yunnan provinces of China.

Clinical features of infected patients

All the patients ranged from 6 to 153 months ($31.6 \pm 19.2M$), and 63.1% (183/290) were boys. Almost half the patients had complications such as neurologic, respiratory or circulatory system symptoms.

Compared to patients without EV71 infection, patients with EV71 infection were older ($33.8 \pm 20.3M$ vs $27.8 \pm 16.3M$, $P < 0.05$), had longer hospital stays ($4.1 \pm 2.4d$ vs




Figure 4 Phylogenetic analysis based on the alignment of the 355 nucleotide VP1 gene of 23 Shanghai CA10 strains. The 25 reference strains were obtained from the US National Center for Biotechnology Information's Genbank.

3.2 ± 1.1d, $P < 0.05$), and higher complication rates (57.8% vs 33.3%, $P < 0.001$).

When we divided patients into 5 groups infected with different enterovirus genotypes, patients with EV71 infection had significantly longer mean hospital stays and a higher incidence of complications than those patients in the CA6, CA10, and CA16 infection groups ($P < 0.05$), but not than those in the CA12 infection group. Patients infected with CA12 were the youngest of the groups and had a higher risk of complications than that of the other non-EV71 infection groups, but the differences were not statistically significant (Table 1).

Discussion

In this study, we identified a variety of enterovirus genotypes, including EV71, CA10, CA6, CA16, CA12, CA4, CA14, Echo6, and EV-C, from stool samples of patients hospitalized for HFMD in Shanghai, China. Our results show a diversified pathogen compositions which are similar with reports from other areas of China, Korea and Singapore [13-15].

The most common genotypes in our study all belonged to species A, in which EV71 occupied the predominant position with the highest detection rate of 63.8%. Since the late 1990s, EV71-related HFMD outbreaks have often been reported in the Asia-Pacific region, including Malaysia, Taiwan, Perth, Japan and China [7,19-25]. Thus, EV71 has been identified as the most prevalent genotype causing HFMD in recent years.

EV71 is classified into three genotypes (A, B, and C) on the basis of their phylogenetic relationship of the VP1 structural sequences. At present, the B genotype contains 6 subgenotypes (B0 through B5), whereas the C genotype contains 5 subgenotypes (C1 through C5) [26-32]. The frequency of these subgenotypes differs over time and area. In Malaysia, the prevalent subgenotypes were C1, C2, B3, and B4 in 1997, but C1 and

C3 in 2000 and B5 and C1 in 2003 and 2005, respectively [33]. The epidemic genotypes in Taiwan were C2 in 1998 but B4 from 2000 to 2003, C5 in 2006, and B5 in 2008 [34,35].

The predominant subgenotype of EV71 in Shanghai during 2010 and 2011 was subgenotype C4, cluster C4a, which is most closely related to the Fuyang22 and DTID/ZJU-62 strains isolated in China. During the past 10 years, the endemic circulation of subgenotype C4 has changed. From 1998 to 2004, EV71 belonged to cluster C4b, but after 2004, cluster C4a replaced C4b and became the predominant virus circulating in China [6,13,36,37]. Our results further confirmed the absolutely dominant position of the EV71 C4a subgenotype in China.

Another major pathogen, CA16, was thought to be associated with mild HFMD without severe neurological complications. The rate of evolution of CA16 strains is far below that of EV71 strains, but sequence information is limited [12]. CA16 has been divided into genogroups A and B, while genogroup B could be further divided into subgenotype B1 and B2 [38]. In our study, B1 was the only prevalent subgenotype, as it was in the CA16 strains in China, Taiwan, Malaysia, Thailand, Vietnam and other areas isolated during 1999 to 2008 [31,39-42].

The higher detection rate of CA6 and CA10, when compared to that of CA16, indicates the importance of these two pathogens in HFMD. Occasional CA6- or CA10-related HFMD outbreaks have been reported. For example, in Singapore, both CA6 and CA10 had a detection rate of 35.3% in a HFMD outbreak in 2008 [14]. The prevalence of CA6 and CA10 was as high as 71% and 28%, respectively, in Finland in 2008, and 28% and 39.9%, respectively, in France in 2010 [43,44]. A CA6-related HFMD outbreak was also reported in Taiwan in 2010 [45]. However, phylogenetic trees analysis was rarely conducted for these two viruses.

Table 1 Characteristics and symptoms of HFMD patients according to different enterovirus genotypes

Characteristics	EV71 (n = 185)	CA6 (n = 24)	CA10 (n = 26)	CA12 (n = 7)	CA16 (n = 20)	Total (n = 290)
Age (M)	33.8 ± 20.3	26.4 ± 14.8	25.6 ± 12.2*	16.6 ± 7.8 ^{*/#}	33.8 ± 8.5	31.6 ± 19.2
Male, n(%)	119 (64.3)	16 (66.7)	17 (65.4)	6 (85.7)	9 (45)	183 (63.1)
Hospital stay (d)	4.1 ± 2.4	2.9 ± 0.8*	2.9 ± 1.0*	3.4 ± 1.4	3.3 ± 1.4*	3.7 ± 1.5
Fever (>37°C), n(%)	175 (94.6)	20 (83.3)*	26 (100)	7 (100)	20 (100)	275 (94.8)
Vomiting, n(%)	72 (39.0)	6 (25)	0 (0)*	0 (0)*	5 (25)	98 (33.8)
Limb shaking, n(%)	41 (22.2)	1 (4.2)*	3 (11.5)	0 (0)	0 (0)*	47 (16.2)
Babinski's sign, n(%) and/or Brudzinski's sign, n(%)	5 (2.7)	0 (0)	0 (0)	0 (0)	0 (0)	7 (2.4)
Complication, n(%)	107 (57.8)	4 (16.7)*	7 (26.9)*	3 (42.9)	5 (25)*	142 (49.0)

*: Compared with EV71 infection group, $P < 0.05$.

#: CA12 infection group compared with CA16 infection group, $P < 0.05$.

In this study, we compared the sequences from our samples to those in Genbank and found that the Shanghai CA6 strain was closely related to strains isolated from Japan, Taiwan, and China. The Shanghai CA6 strain belonged to two evolutionary clusters with a high nucleotide homology, whereas all the CA10 strains were most closely related to the strains found in the Shandong and Yunnan provinces of China.

Although not prevalent, CA4 and CA12 were identified as pathogens attributing to HFMD in Shanghai. CA4 caused a high infection rate in preschool children in Taiwan from 2006 to 2008, whereas only one case of CA4 infection were reported in Korea in 2009 [14,41]. Sporadic HFMD cases with CA12 infection were seen in other areas of China between 2008 and 2009 [13,46,47].

Most enterovirus infections are self-limited and do not require hospitalization, but EV71 infection in young children frequently cause complications and can progress quickly [3,48,49]. However the clinical characteristics of enterovirus genotypes, other than EV71 and CA16, have not been well studied. We found that HFMD cases infected with CA6, CA10 and CA16 caused less complications compared to those infected with EV71. But worth of note, the patients infected with CA12 had the youngest age of onset and most likely have the highest incidence of complications than any of the other non-EV71 infection groups. Since there were only 7 cases in this group, more study and data are needed for accurately identifying the pathogenetic characteristic of CA12 infections in China. In addition, we did not observe the clinical symptoms, such as skin ulceration and an obvious onychomadesis, that were associated with CA6-caused HFMD in Finland, Taiwan and Spain [45,50,51].

One limitation of this study is that different samples were not taken in the patients i.e. throat swab, vesicle and CSF which may influence the integrality of the data and bring bias to the conclusion. Due to the retrospective nature of the study, stools were the only specimen sent routinely to virology laboratory for enterovirus testing and collection of more types of samples could not be performed. However, it is widely recognized that HFMD is a common disease of children mostly associated with the human enterovirus species A. All the stools were from hospitalized patients clinically diagnosed with HFMD, and all the specimens were collected during the acute phase of the illness. Thus we think that the enterovirus identified in the stool specimen could represent the pathogen causing HFMD and had a great relationship with the clinical symptoms. In the future, to elucidate further the epidemiology of the pathogens for HFMD, a prospective study will be developed and multiple clinical specimens from the same patient need to be taken and evaluated.

Conclusions

In summary, this study provides useful epidemiological data on the features of the pathogen compositions of HFMD as well as clinical characteristics differing in enterovirus genotypes in Shanghai, China. It deserves our attention that early identification of enterovirus genotypes is important for diagnosis and treatment of HFMD patients.

Abbreviations

HFMD: Hand foot and mouth disease; EV: Enterovirus; EV71: Enterovirus 71; CA16: Coxsackievirus A16; NCBI: US national center for biotechnology information; DEPC: Diethylpyrocarbonate.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

JX designed the study and contributed to manuscript writing. MX performed the experiment, analyzed the data and wrote the manuscript. LS, LC, ND and HZ contributed to specimen collection, reagents preparation, setting up the assay and data analysis. All authors have read and approved the final manuscript.

Acknowledgements

This work was supported by the Program for New Century Excellent Talents in University (NECT-09-0320) and supported by the Program for Excellent Academic Leaders in Public Health in Shanghai (GWDTR201221). Thank Tom Lang who is a scientific English editor in U.S.A for revising this manuscript on English language.

Received: 29 September 2012 Accepted: 17 October 2013

Published: 22 October 2013

References

1. Huang CC, Liu CC, Chang YC, Chen CY: **Neurologic complications in children with enterovirus 71 infection.** *N Engl J Med* 1999, **341**:936–942.
2. Ho M, Chen ER, Hsu KH, Twu SJ, Chen KT, Tsai SF, Wang JR, Shih SR: **An epidemic of enterovirus 71 infection in Taiwan. Taiwan enterovirus epidemic working group.** *N Engl J Med* 1999, **341**:929–935.
3. Ishimaru Y, Nakano S, Yamaoka K, Takami S: **Outbreaks of hand, foot, and mouth disease by enterovirus 71. High incidence of complication disorders of central nervous system.** *Arch Dis Child* 1980, **55**:583–588.
4. Blomberg J, Lycke E, Ahlfors K, Johnsson T, Wolontis S, von Zeipel G: **New enterovirus type associated with epidemic of aseptic meningitis and/or hand, foot, and mouth disease.** *Lancet* 1974, **2**:112.
5. Schmidt NJ, Lennette EH, Ho HH: **An apparently new enterovirus isolated from patients with disease of the central nervous system.** *J Infect Dis* 1974, **129**:304–309.
6. Zhang Y, Tan XJ, Wang HY, Yan DM, Zhu SL, Wang DY, Ji F, Wang XJ, Gao YJ, Chen L, An HQ, Li DX, Wang SW, Xu AQ, Wang ZJ, Xu WB: **An outbreak of hand, foot, and mouth disease associated with subgenotype C4 of human enterovirus 71 in Shandong, China.** *J Clin Virol* 2009, **44**:262–267.
7. Zhang Y, Zhu Z, Yang W, Ren J, Tan X, Wang Y, Mao N, Xu S, Zhu S, Cui A, Zhang Y, Yan D, Li Q, Dong X, Zhang J, Zhao Y, Wan J, Feng Z, Sun J, Wang S, Li D, Xu W: **An emerging recombinant human enterovirus 71 responsible for the 2008 outbreak of hand foot and mouth disease in Fuyang city of China.** *Virology* 2010, **7**:94–102.
8. Oberste MS, Maher K, Kilpatrick DR, Pallansch MA: **Molecular evolution of the human enteroviruses: correlation of serotype with VP1 sequence and application to picornavirus classification.** *J Virol* 1999, **73**:1941–1948.
9. Oberste MS, Nix WA, Maher K, Pallansch MA: **Improved molecular identification of enteroviruses by RT-PCR and amplicon sequencing.** *J Clin Virol* 2003, **26**:375–377.
10. Nasri D, Bouzlama L, Pillet S, Bourlet T, Aouni M, Pozzetto B: **Basic rationale, current methods and future directions for molecular typing of human enterovirus.** *Expert Rev Mol Diagn* 2007, **7**:419–434.

11. King AMQ, Adams MJ, Carstens EB, Lefkowitz EJ: *Virus taxonomy: classification and nomenclature of viruses: ninth report of the international committee on taxonomy of viruses*. San Diego: Elsevier; 2012:855–880.
12. Li L, He Y, Yang H, Zhu J, Xu X, Dong J, Zhu Y, Jin Q: **Genetic characteristics of human enterovirus 71 and coxsackievirus A16 circulating from 1999 to 2004 in Shenzhen, People's Republic of China.** *J Clin Microbiol* 2005, **43**:3835–3839.
13. Yang F, Zhang T, Hu Y, Wang X, Du J, Li Y, Sun S, Sun X, Li Z, Jin Q: **Survey of enterovirus infections from hand, foot and mouth disease outbreak in China, 2009.** *Virology* 2011, **8**:508–511.
14. Park SH, Choi SS, Oh SA, Kim CK, Cho SJ, Lee JH, Ryu SH, Pak SH, Jung SK, Lee JI, Park GY, Choi SM, Chae YZ, Kang BH, Cheon DS, Kim HS: **Detection and characterization of enterovirus associated with herpangina and hand, foot, and mouth disease in Seoul, Korea.** *Clin Lab* 2011, **57**:959–967.
15. Wu Y, Yeo A, Phoon MC, Tan EL, Poh CL, Quak SH, Chow VT: **The largest outbreak of hand; foot and mouth disease in Singapore in 2008: the role of enterovirus 71 and coxsackievirus A strains.** *Int J Infect Dis* 2010, **14**:e1076–1081.
16. De W, Changwen K, Wei L, Monagin C, Jin Y, Cong M, Hanri Z, Jun S: **A large outbreak of hand, foot, and mouth disease caused by EV71 and CA16 in Guangdong, China, 2009.** *Arch Virol* 2011, **156**:945–953.
17. Shah VA, Chong CY, Chan KP, Ng W, Ling AE: **Clinical characteristics of an outbreak of hand, foot and mouth disease in Singapore.** *Ann Acad Med Singapore* 2003, **32**:381–387.
18. Leitch EC, Harvala H, Robertson I, Ubillos I, Templeton K, Simmonds P: **Direct identification of human enterovirus serotypes in cerebrospinal fluid by amplification and sequencing of the VP1 region.** *J Clin Virol* 2009, **44**:119–124.
19. Lum LC, Wong KT, Lam SK, Chua KB, Goh AY, Lim WL, Ong BB, Paul G, AbuBakar S, Lambert M: **Fatal enterovirus 71 encephalomyelitis.** *J Pediatr* 1998, **133**:795–798.
20. Podin Y, Gias EL, Ong F, Leong YW, Yee SF, Yusof MA, Perera D, Teo B, Wee TY, Yao SC, Yao SK, Kiyu A, Arif MT, Cardoso MJ: **Sentinel surveillance for human enterovirus 71 in Sarawak, Malaysia: lessons from the first 7 years.** *BMC Public Health* 2006, **6**:180–189.
21. Chang LY, Huang YC, Lin TY: **Fulminant neurogenic pulmonary oedema with hand, foot, and mouth disease.** *Lancet* 1998, **352**:367–368.
22. Wang JR, Tuan YC, Tsai HP, Yan JJ, Liu CC, Su IJ: **Change of major genotype of enterovirus 71 in outbreaks of hand-foot-and-mouth disease in Taiwan between 1998 and 2000.** *J Clin Microbiol* 2002, **40**:10–15.
23. McMinn P, Stratov I, Nagarajan L, Davis S: **Neurological manifestations of enterovirus 71 infection in children during an outbreak of hand, foot, and mouth disease in Western Australia.** *Clin Infect Dis* 2001, **32**:236–242.
24. **Infectious Disease Surveillance Centre, National Institute of Infectious Diseases, Japan: hand, foot and mouth disease, 2000–2003, Japan.** *Infect Agents Surv Rep* 2004, **25**:224–225.
25. AbuBakar S, Sam IC, Yusof J, Lim MK, Misbah S, MatRahim N, Hooi PS: **Enterovirus 71 outbreak, Brunei.** *Emerg Infect Dis* 2009, **15**:79–82.
26. Brown BA, Oberste MS, Alexander JP, Kennett ML, Pallansch MA: **Molecular epidemiology and evolution of enterovirus 71 strains isolated from 1970 to 1998.** *J Virol* 1999, **73**:9969–9975.
27. McMinn P, Lindsay K, Perera D, Chan HM, Chan KP, Cardoso MJ: **Phylogenetic analysis of enterovirus 71 strains isolated during linked epidemics in Malaysia, Singapore, and Western Australia.** *J Virol* 2001, **75**:7732–7738.
28. Cardoso MJ, Perera D, Brown BA, Cheon HM, Chan KP, Cho H, Cho H, McMinn P: **Molecular epidemiology of human enterovirus 71 strains and recent outbreaks in the Asia-Pacific region: comparative analysis of the VP1 and VP4 genes.** *Emerg Infect Dis* 2003, **9**:461–468.
29. Shimizu H, Utama A, Onnimala N, Li C, Li-Bi Z, Yu-Jie M, Cho H, McMinn P: **Molecular epidemiology of enterovirus 71 infection in the Western Pacific Region.** *Pediatr Int* 2004, **46**:231–235.
30. Ooi MH, Wong SC, Podin Y, Akin W, del Sel S, Mohan A, Chieng CH, Perera D, Clear D, Wong D, Blake E, Cardoso J, Solomon T: **Human enterovirus 71 disease in Sarawak, Malaysia: a prospective clinical, virological, and molecular epidemiological study.** *Clin Infect Dis* 2007, **44**:646–656.
31. Tu PV, Thao NT, Perera D, Huu TK, Tien NT, Thuong TC, How OM, Cardoso MJ, McMinn PC: **Epidemiologic and virologic investigation of hand, foot, and mouth disease, southern Vietnam, 2005.** *Emerg Infect Dis* 2007, **13**:1733–1741.
32. van der Sanden S, Koopmans M, Uslu G, van der Avoort H, Dutch Working Group for Clinical Virology: **Epidemiology of enterovirus 71 in the Netherlands, 1963 to 2008.** *J Clin Microbiol* 2009, **47**:2826–2833.
33. Chua KB, Kasri AR: **Hand foot and mouth disease due to enterovirus 71 in Malaysia.** *Virology* 2011, **26**:221–228.
34. Huang YP, Lin TL, Kuo CY, Lin MW, Yao CY, Liao HW, Hsu LC, Yang CF, Yang JY, Chen PJ, Wu HS: **The circulation of subgenogroups B5 and C5 of enterovirus 71 in Taiwan from 2006 to 2007.** *Virus Res* 2008, **137**:206–212.
35. Lee MS, Lin TY, Chiang PS, Li WC, Luo ST, Tsao KC, Liou GY, Huang ML, Hsia SH, Huang YC, Chang SC: **An investigation of epidemic enterovirus 71 infection in Taiwan, 2008: clinical, virologic, and serologic features.** *Pediatr Infect Dis J* 2010, **29**:1030–1034.
36. Yu H, Chen W, Chang H, Tang R, Zhao J, Gan L, Liu B, Chen J, Wang M: **Genetic analysis of the VP1 region of enterovirus 71 reveals the emergence of genotype A in central China in 2008.** *Virus Genes* 2010, **41**:1–4.
37. Zhu Q, Hao Y, Ma J, Yu S, Wang Y: **Surveillance of hand, foot, and mouth disease in mainland China (2008–2009).** *Biomed Environ Sci* 2011, **24**:349–356.
38. Perera D, Yusof MA, Podin Y, Ooi MH, Thao NT, Wong KK, Zaki A, Chua KB, Malik YA, Tu PV, Tien NT, Puthavathana P, McMinn PC, Cardoso MJ: **Molecular phylogeny of modern coxsackievirus A16.** *Arch Virol* 2007, **152**:1201–1208.
39. Iwai M, Masaki A, Hasegawa S, Obara M, Horimoto E, Nakamura K, Tanaka Y, Endo K, Tanaka K, Ueda J, Shiraki K, Kurata T, Takizawa T: **Genetic changes of coxsackievirus A16 and enterovirus 71 isolated from hand, foot, and mouth disease patients in Toyama, Japan between 1981 and 2007.** *Jpn J Infect Dis* 2009, **62**:254–259.
40. Zhang Y, Wang D, Yan D, Zhu S, Liu J, Wang H, Zhao S, Yu D, Nan L, An J, Chen L, An H, Xu A, Xu W: **Molecular evidence of persistent epidemic and evolution of subgenotype B1 coxsackievirus A16-associated hand, foot, and mouth disease in China.** *J Clin Microbiol* 2010, **48**:619–622.
41. Wu PC, Huang LM, Kao CL, Fan TY, Cheng AL, Chang LY: **An outbreak of coxsackievirus A16 infection: comparison with other enteroviruses in a preschool in Taipei.** *J Microbiol Immunol Infect* 2010, **43**:271–277.
42. Puenpa J, Theamboonlers A, Korkong S, Linsuwanon P, Thongmee C, Chatproedprai S, Poovorawan Y: **Molecular characterization and complete genome analysis of human enterovirus 71 and coxsackievirus A16 from children with hand, foot and mouth disease in Thailand during 2008–2011.** *Arch Virol* 2011, **156**:2007–2013.
43. Mirand A, Henquell C, Archimbaud C, Ughetto S, Antona D, Bailly JL, Peigue-Lafeuille H: **Outbreak of hand, foot and mouth disease/herpangina associated with coxsackievirus A6 and A10 infections in 2010, France: a large citywide, prospective observational study.** *Clin Microbiol Infect* 2012, **18**:E110–118.
44. Blomqvist S, Klemola P, Kajjalainen S, Paananen A, Simonen ML, Vuorinen T, Roivainen M: **Co-circulation of coxsackieviruses A6 and A10 in hand, foot and mouth disease outbreak in Finland.** *J Clin Virol* 2010, **48**:49–54.
45. Wei SH, Huang YP, Liu MC, Tsou TP, Lin HC, Lin TL, Tsai CY, Chao YN, Chang LY, Hsu CM: **An outbreak of coxsackievirus A6 hand, foot, and mouth disease associated with onychomadesis in Taiwan, 2010.** *BMC Infect Dis* 2011, **11**:346–351.
46. Zhu B, Zhong JY, Xia HM, Gong ST, Xiao MS, Xie JH, Zhang YY, Hua L, Lian GW: **Etiology of hand, foot and mouth disease in Guangzhou in 2008.** *Zhonghua Er Ke Za Zhi* 2010, **48**:127–130.
47. Wang ZG, Liu XL, Yang TT, Yi Y: **Etiology of hand, foot and mouth disease in Qingdao during 2008–2009.** *Bing Du Xue Bao* 2011, **27**:438–441.
48. Chong CY, Chan KP, Shah VA, Ng WY, Lau G, Teo TE, Lai SH, Ling AE: **Hand, foot and mouth disease in Singapore: a comparison of fatal and non-fatal cases.** *Acta Paediatr* 2003, **92**:1163–1169.
49. Xu W, Liu CF, Yan L, Li JJ, Wang LJ, Qi Y, Cheng RB, Xiong XY: **Distribution of enteroviruses in hospitalized children with hand, foot and mouth disease and relationship between pathogens and nervous system complications.** *Virology* 2012, **9**:8.
50. Osterback R, Vuorinen T, Linna M, Susi P, Hyypiä T, Waris M: **Coxsackievirus A6 and hand, foot, and mouth disease, Finland.** *Emerg Infect Dis* 2009, **15**:1485–1488.
51. Clementz GC, Mancini AJ: **Nail matrix arrest following hand-foot-mouth disease: a report of five children.** *Pediatr Dermatol* 2000, **17**:7–11.

doi:10.1186/1471-2334-13-489

Cite this article as: Xu et al.: Enterovirus genotypes causing hand foot and mouth disease in Shanghai, China: a molecular epidemiological analysis. *BMC Infectious Diseases* 2013 **13**:489.