

CORRECTION

Correction: Predicting Response Trajectories during Cognitive-Behavioural Therapy for Panic Disorder: No Association with the BDNF Gene or Childhood Maltreatment

Martí Santacana, Bárbara Arias, Marina Mitjans, Albert Bonillo, María Montoro, Sílvia Rosado, Roser Guillamat, Vicenç Vallès, Víctor Pérez, Carlos G. Forero, Miquel A. Fullana

The [S1 Fig](#) legend is incorrect. The MET-CARRIERS and VAL-VAL groups are swapped. The authors have provided a corrected version here.

Supporting Information

S1 Fig. Mean scores of the PDSS-SR throughout the treatment and follow-up for Met-carriers and Val-Val.

(JPG)

Reference

1. Santacana M, Arias B, Mitjans M, Bonillo A, Montoro M, Rosado S, et al. (2016) Predicting Response Trajectories during Cognitive-Behavioural Therapy for Panic Disorder: No Association with the *BDNF* Gene or Childhood Maltreatment. PLoS ONE 11(6): e0158224. doi: [10.1371/journal.pone.0158224](https://doi.org/10.1371/journal.pone.0158224) PMID: [27355213](https://pubmed.ncbi.nlm.nih.gov/27355213/)



OPEN ACCESS

Citation: Santacana M, Arias B, Mitjans M, Bonillo A, Montoro M, Rosado S, et al. (2016) Correction: Predicting Response Trajectories during Cognitive-Behavioural Therapy for Panic Disorder: No Association with the BDNF Gene or Childhood Maltreatment. PLoS ONE 11(12): e0167833. doi:10.1371/journal.pone.0167833

Published: December 1, 2016

Copyright: © 2016 Santacana et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.