

## Research Article

# Sulfur Dioxide Enhances Endogenous Hydrogen Sulfide Accumulation and Alleviates Oxidative Stress Induced by Aluminum Stress in Germinating Wheat Seeds

Dong-Bo Zhu,<sup>1</sup> Kang-Di Hu,<sup>1</sup> Xi-Kai Guo,<sup>1</sup> Yong Liu,<sup>1</sup> Lan-Ying Hu,<sup>1</sup>  
Yan-Hong Li,<sup>1</sup> Song-Hua Wang,<sup>2</sup> and Hua Zhang<sup>1</sup>

<sup>1</sup>School of Biotechnology and Food Engineering, Hefei University of Technology, Hefei 230009, China

<sup>2</sup>Life Science College, Anhui Science and Technology University, Bengbu 233100, China

Correspondence should be addressed to Hua Zhang; hzhanglab@gmail.com

Received 28 October 2014; Accepted 21 November 2014

Academic Editor: Guangdong Yang

Copyright © 2015 Dong-Bo Zhu et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Aluminum ions are especially toxic to plants in acidic soils. Here we present evidences that SO<sub>2</sub> protects germinating wheat grains against aluminum stress. SO<sub>2</sub> donor (NaHSO<sub>3</sub>/Na<sub>2</sub>SO<sub>3</sub>) pretreatment at 1.2 mM reduced the accumulation of superoxide anion, hydrogen peroxide, and malondialdehyde, enhanced the activities of guaiacol peroxidase, catalase, and ascorbate peroxidase, and decreased the activity of lipoxygenase in germinating wheat grains exposed to Al stress. We also observed higher accumulation of hydrogen sulfide (H<sub>2</sub>S) in SO<sub>2</sub>-pretreated grain, suggesting the tight relation between sulfite and sulfide. Wheat grains germinated in water for 36 h were pretreated with or without 1 mM SO<sub>2</sub> donor for 12 h prior to exposure to Al stress for 48 h and the ameliorating effects of SO<sub>2</sub> on wheat radicles were studied. SO<sub>2</sub> donor pretreatment reduced the content of reactive oxygen species, protected membrane integrity, and reduced Al accumulation in wheat radicles. Gene expression analysis showed that SO<sub>2</sub> donor pretreatment decreased the expression of Al-responsive genes TaWali1, TaWali2, TaWali3, TaWali5, TaWali6, and TaALMT1 in radicles exposed to Al stress. These results suggested that SO<sub>2</sub> could increase endogenous H<sub>2</sub>S accumulation and the antioxidant capability and decrease endogenous Al content in wheat grains to alleviate Al stress.

## 1. Introduction

Aluminum ions (Al<sup>3+</sup>) together with silicon and iron are the three most abundant mineral elements in soil. Whereas silicon and iron are required for plant growth, Al is toxic. Many different mechanisms have been advanced to explain Al toxicity in plants [1, 2]. One of the primary causes of Al toxicity is oxidative stress due to accumulation of reactive oxygen species (ROS), such as the superoxide anion (O<sub>2</sub><sup>•-</sup>) and hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>), bringing about lipid peroxidation in plant cells [3–5]. Plants have developed several strategies to counteract oxidative stress caused by Al, such as activation of antioxidants, and exudation of organic acids as a mechanism for Al exclusion [6]. Recently, a range of signaling molecules, such as inositol 1,4,5-triphosphate (IP<sub>3</sub>), salicylic acid, hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) and nitric oxide

(NO), carbon monoxide (CO), and hydrogen sulfide (H<sub>2</sub>S), were found to participate in plant's resistance to Al-induced oxidative stress [4, 7–10].

Sulfur dioxide (SO<sub>2</sub>) is a colorless, nonflammable gas with a penetrating odor. Low concentrations of SO<sub>2</sub> have been found to play a physiological role *in vivo* in animal models, participating in various biological processes [11]. The physiological processes regulated by SO<sub>2</sub> in animals include cardiac function [11], inhibition of L-calcium channels in cardiomyocytes [12], and improvement in pulmonary vascular structural remodeling [13]. In plants, the toxic effects of SO<sub>2</sub> on growth and development have been extensively studied [14, 15]. Exposure to high concentrations of SO<sub>2</sub> can cause visible foliar damage, a decline in photosynthesis, an inhibition of plant growth, and structural disorganization and cell death [16–19]. On the other hand, many reports show

that low levels of atmospheric SO<sub>2</sub> might be beneficial to plants [20]. SO<sub>2</sub> can be metabolized and used as a sulfur source for plant growth, especially when the sulfur supply in soil is insufficient for normal growth [20]. Recently, low concentrations of SO<sub>2</sub> were found to induce transcriptome reprogramming associated with oxidative signaling and biotic defence responses in plants, suggesting a physiological role of SO<sub>2</sub> in plant [21].

In plants, sulfate is taken up from soil by high-affinity transporters. Sulfate is largely transported to shoots where it can be activated by ATP via ATP sulfurylase in the leaves. The product is reduced by 5'-adenylylsulfate (APS) reductase to sulfite which can be reduced to H<sub>2</sub>S by sulfite reductase [22]. SO<sub>2</sub> can also be produced endogenously from sulfur-containing amino acids [23]. The endogenous production of SO<sub>2</sub> also suggests that it has a physiological role in plants.

In order to establish the role of SO<sub>2</sub> in alleviating Al stress, we investigated the effects of SO<sub>2</sub> pretreatment on H<sub>2</sub>S and ROS accumulation and the antioxidant system in whole wheat grains and in wheat radicles. We also analyzed endogenous H<sub>2</sub>S and Al content as a means of understanding the mechanism of the role of SO<sub>2</sub>. We speculated that SO<sub>2</sub> might act as an antioxidant molecule to alleviate Al toxicity during wheat grain germination.

## 2. Materials and Methods

**2.1. Materials and Treatments.** Wheat (*Triticum aestivum* L.) grains were supplied by the Anhui Aidi Agricultural Technology Co., Ltd., Anhui Province, China. Sodium bisulfite (NaHSO<sub>3</sub>) and anhydrous sodium sulfite (Na<sub>2</sub>SO<sub>3</sub>) were used as sulfur dioxide (SO<sub>2</sub>) donors according to Laisk et al. [24]. Wheat grains were sterilized by 0.1% HgCl<sub>2</sub> for 3 min and washed extensively with H<sub>2</sub>O and then dried with filter papers. Wheat grains of similar size were selected and allocated randomly in Petri dish (9 cm diameter × 1.2 cm depth, 50 grains per dish). Wheat grains were germinated in H<sub>2</sub>O or aqueous solutions of AlCl<sub>3</sub> at 5, 10, 15, 20, 25, 30, 60, and 90 mM for 48 h at 25°C and the length of coleoptiles and radicles and radicle number were recorded. To test the protective role of SO<sub>2</sub> on germination and seedling growth of wheat grains under Al stress, grains were pretreated with H<sub>2</sub>O or 0.4, 0.8, 1.2, 1.6, or 2.0 mM SO<sub>2</sub> donor for 12 h and subsequently subjected to a semi-inhibitory AlCl<sub>3</sub> concentration (15 mM). AlCl<sub>3</sub> solutions were renewed every 12 h and germinating grains were sampled every 12 h for further analysis.

**2.2. Determination of MDA, O<sub>2</sub><sup>•-</sup>, and H<sub>2</sub>O<sub>2</sub>.** The contents of MDA, O<sub>2</sub><sup>•-</sup>, and H<sub>2</sub>O<sub>2</sub> were determined by the method of Zhang et al. [25].

**2.3. Assays of LOX, CAT, APX, and POD Activities.** Activity of lipoxygenase (LOX, EC 1.13.11.12) was determined following the description by Surrey [26] and those of catalase (CAT, EC 1.11.1.6), ascorbate peroxidase (APX, EC 1.11.1.11), and guaiacol peroxidase (POD, EC 1.11.1.7) were assayed according to Hu et al. [27]. Wheat grains were homogenized in ice-cold

50 mM phosphate buffer (pH 7.8) containing 1.0 mM EDTA. The homogenate was centrifuged at 15,000 g at 4°C for 10 min. The supernatant was used for activity determination.

**2.4. Assays of Reducing Sugars and Soluble Protein.** Wheat grains (0.5 ± 0.05 g) were ground in 5 mL of phosphate buffer (pH 7.0, 200 mM), the homogenate was centrifuged at 10,000 g for 30 min, and the supernatant was used for detection of reducing sugars and soluble protein content. Reducing sugar content was measured according to Miller [28].

For detection of soluble protein, 0.1 mL supernatant was mixed with 0.9 mL H<sub>2</sub>O and 5 mL Coomassie brilliant blue for 5 min and the absorbance recorded at 595 nm using the method described by Bradford [29].

**2.5. Preparation of Wheat Radicles.** Wheat grains were germinated in H<sub>2</sub>O for 36 h in the dark at 25°C and the average of radicle length was approximately 1.0 cm. The germinated wheat grains were pretreated with or without 1 mM SO<sub>2</sub> donor for 12 h and then exposed to 0 or 400 μM AlCl<sub>3</sub> for 48 h.

**2.6. Detection of Plasma Membrane Integrity, Al Accumulation, and ROS Production in Radicles.** Plasma membrane integrity of wheat radicles was detected following the method of Yamamoto et al. [30]. Radicles were stained with Evans blue solution (0.025% [w/v] Evans blue in 100 μM CaCl<sub>2</sub>, pH 5.6) for 10 min, then washed three times with 100 μM CaCl<sub>2</sub> solutions, and photographed. Staining intensity of Evans blue is positively correlated with more damaged plasma membrane.

Al content in radicles was visualized by staining tissues with hematoxylin. Hematoxylin stain was prepared as described by Polle et al. [31]. Wheat radicles were washed with H<sub>2</sub>O for 30 min and then stained with solution of 0.2% hematoxylin and 0.02% NaIO<sub>3</sub> for 30 min at room temperature. Radicles were then immersed in H<sub>2</sub>O for 30 min to remove excess stain and photographed. Staining intensity of hematoxylin is positively correlated with Al uptake.

ROS distribution in radicle tips was detected by 2',7'-dichlorofluorescein diacetate (DCFH-DA) following the method of LeBel et al. [32]. Radicle tips were incubated in a solution containing 100 μM CaCl<sub>2</sub> and 10 μM DCFH-DA for 20 min and then washed three times with H<sub>2</sub>O. The fluorescence was detected with a Nikon 80i microscope (excitation at 488 nm and emission at 525 nm). For each treatment, ten individual roots from ten seedlings were examined and similar results were obtained.

**2.7. Real-Time Quantitative RT-PCR Analysis in Wheat Radicles.** Radicle tips were prepared for RNA extraction according to Li et al. [33]. Total RNA was isolated by grinding with liquid nitrogen according to the manufacturer's instructions (CWBIO, Beijing, China). cDNA was generated from total RNA with a reverse transcription kit (Prime Script RT Master Mix, Takara, Kyoto, Japan). Quantitative PCR was performed using a StepOnePlus Real-Time PCR

TABLE 1: Inhibitory effect of Al stress on the germination of wheat grains. Wheat grains were exposed to 0, 5, 10, 15, 20, 25, 30, 60, or 90 mM  $\text{AlCl}_3$  for 48 h.

$\text{Al}^{3+}$ concentration (mM)	Germination percentage (%)	Radicle length (cm)	Coleoptile length (cm)	Radicle number (50 grains)
0	64 ± 1.2 <sup>a</sup>	3.1 ± 0.8 <sup>a</sup>	1.5 ± 0.3 <sup>ab</sup>	178 ± 7.8 <sup>a</sup>
5	66 ± 1.1 <sup>a</sup>	2.7 ± 0.5 <sup>ab</sup>	1.6 ± 0.2 <sup>a</sup>	168 ± 8.9 <sup>a</sup>
10	51 ± 2.3 <sup>b</sup>	1.9 ± 0.3 <sup>b</sup>	1.3 ± 0.3 <sup>ab</sup>	162 ± 7.6 <sup>a</sup>
15	35 ± 3.8 <sup>c</sup>	1.1 ± 0.2 <sup>c</sup>	1.1 ± 0.2 <sup>bc</sup>	142 ± 6.3 <sup>b</sup>
20	28 ± 4.2 <sup>cd</sup>	0.7 ± 0.3 <sup>cd</sup>	0.8 ± 0.2 <sup>cd</sup>	80 ± 5.6 <sup>c</sup>
25	21 ± 5.1 <sup>de</sup>	0.4 ± 0.3 <sup>d</sup>	0.6 ± 0.2 <sup>de</sup>	68 ± 6.1 <sup>c</sup>
30	15 ± 4.7 <sup>ef</sup>	0.2 ± 0.2 <sup>d</sup>	0.3 ± 0.3 <sup>e</sup>	45 ± 5.3 <sup>d</sup>
60	8 ± 5.2 <sup>f</sup>	0.1 ± 0.1 <sup>d</sup>	0.3 ± 0.2 <sup>e</sup>	22 ± 3.5 <sup>e</sup>
90	7 ± 6.3 <sup>f</sup>	0 ± 0 <sup>e</sup>	0.2 ± 0.2 <sup>e</sup>	0 ± 0 <sup>f</sup>

Values are the means ± SD ( $n = 6$ ). Values are the means ± SD ( $n = 6$ ). Different letters mean significance of difference between different treatments ( $P < 0.05$ ).

System (Applied Biosystems, Foster City, CA, USA) with SYBR Premix Ex Taq (TaKaRa Bio Inc., China) according to the manufacturer's instructions. cDNA was amplified by PCR using the following primers:  $\text{Ta}\beta$ -actin forward (5'-CTATCCTTCGTTTGGACCTT-3') and reverse (5'-AGC-GAGCTTCTCCTTTATGT-3');  $\text{TaWali1}$  forward (5'-CTG-ATGGAGTCGAGCAAGG-3') and reverse (5'-CCGAAG-TAGCGATTTAGGAGT-3');  $\text{TaWali2}$  forward (5'-AGC-CTACTGCTCCGCCTTGT-3') and reverse (5'-CGTTTC-GTCGGCATCTCC-3');  $\text{TaWali3}$  forward (5'-GACGAG-CCCTAAGAAGACG-3') and reverse (5'-CACGGAGCA-ATGACAACAG-3');  $\text{TaWali5}$  forward (5'-TGGACCCTG-CAAGAAGTAC-3') and reverse (5'-GCTGAACAACAA-GCAACACC-3');  $\text{TaWali6}$  forward (5'-TACGGAATAGAC-AGGACAAGG-3') and reverse (5'-CAGCATTTTCGGG-AACTCG-3');  $\text{TaALMT1}$  forward (5'-TGCCACGCTGAG-TAAAGG-3') and reverse (5'-CGCTGACGCTACGAA-GAA-3'). Relative gene expression was presented as values relative to the corresponding gene expression in control, after normalization to the control  $\text{Ta}\beta$ -actin transcript levels.

**2.8. Statistical Analysis.** Statistical significance was tested by one-way ANOVA, and the results are expressed as the mean values ± SD (standard deviation) of three independent experiments. Each experiment was repeated three times.

### 3. Results

**3.1. Inhibitory Effect of Al on Wheat Grain Germination.** The effect of Al stress on wheat seedling growth and development was examined following incubation of grain in  $\text{AlCl}_3$  with concentrations ranging from 5 mM to 90 mM (Table 1). At concentrations of 5 mM or below, germination percentage, coleoptile length, and radicle number are almost unaffected, but radicle length was reduced by 13%, suggesting that the radicle is the primary target of Al toxicity. At 15 mM Al, germination percentage was almost halved compared with that of control and this concentration was selected for further experiments. At 90 mM Al, radicle growth was completely inhibited, but very stunted coleoptile growth was still observed.

**3.2.  $\text{SO}_2$  Donor Ameliorates Al Stress in Germinating Wheat Grain.** To establish whether the  $\text{SO}_2$  donor  $\text{Na}_2\text{SO}_3/\text{NaHSO}_3$  had a toxic effect on wheat grain germination, grains were germinated in different  $\text{SO}_2$  donor concentrations ranging from 0.4 to 2.0 mM for 36 h (see Table S1 in Supplementary Material available online at <http://dx.doi.org/10.1155/2015/612363>). Table S1 shows there was no significant change in germination percentage, coleoptile length, radicle length, or radicle number between water control and  $\text{SO}_2$  donor treatment, establishing that the concentrations of  $\text{SO}_2$  donor used in this work exhibited no visible toxic effects. To test the ability of  $\text{SO}_2$  donor to alleviate Al stress, wheat grains were pretreated with  $\text{SO}_2$  donor concentrations ranging from 0.4 to 2.0 mM for 12 h prior to incubation with 15 mM Al (Table 2 and Figure 1). At all  $\text{SO}_2$  donor concentrations used,  $\text{SO}_2$  pretreatment was effective in alleviating the toxic effects of Al in a dose dependent manner. The optimal  $\text{SO}_2$  donor concentration for alleviating Al stress was 1.2 mM, a concentration where the germination percentage was increased by 51%, radicle and coleoptile length by 28% and 26%, respectively, compared with those exposed to Al. This result clearly shows that  $\text{SO}_2$  alleviates Al-induced inhibition of wheat grain germination and seedling growth.

**3.3. Effect of  $\text{SO}_2$  Donor on the Contents of Reducing Sugars and Soluble Protein in Al-Stressed Wheat Grain.** Figure 2(a) shows the changes in reducing sugars in germinating wheat grains preincubated in  $\text{SO}_2$  donor or  $\text{H}_2\text{O}$  for 12 h followed by incubation in Al for 48 h. Within 12 h pretreatment in  $\text{H}_2\text{O}$  and 24 h of Al treatment, the content of reducing sugar decreased gradually, whereas reducing sugar in the  $\text{SO}_2$  donor pretreatment remained stable and slightly increased at 24 h. Thereafter reducing sugar content increased steadily in both treatments followed by a slight decrease at 48 h. The content of reducing sugars in  $\text{SO}_2$  donor pretreated grain was always significantly higher than the counterpart of only Al treatment.

The content of soluble protein increased gradually and peaked on 24 h of Al stress followed by a slight decrease (Figure 2(b)). Though the mean values of soluble protein in

TABLE 2: Effects of SO<sub>2</sub> donor pretreatment on wheat grain germination under 15 mM Al<sup>3+</sup> stress. Wheat grains were pretreated with 0, 0.4, 0.8, 1.2, 1.6, and 2.0 mM SO<sub>2</sub> for 12 h and subsequently subjected to 15 mM AlCl<sub>3</sub> for further 48 h, and then germination was investigated.

SO <sub>2</sub> donor concentration (mM)	0.0	0.4	0.8	1.2	1.6	2.0
Germination percentage (%)	37 ± 3.3 <sup>a</sup>	42 ± 2.7 <sup>a</sup>	44 ± 3.5 <sup>a</sup>	56 ± 3.8 <sup>a</sup>	48 ± 2.7 <sup>a</sup>	47 ± 3.1 <sup>a</sup>
Length of radicle (cm)	1.42 ± 0.6 <sup>a</sup>	1.72 ± 0.4 <sup>a</sup>	1.80 ± 0.7 <sup>a</sup>	1.82 ± 0.7 <sup>a</sup>	1.78 ± 0.8 <sup>a</sup>	1.62 ± 0.4 <sup>a</sup>
Length of coleoptile (cm)	4.64 ± 0.4 <sup>a</sup>	4.70 ± 0.6 <sup>a</sup>	5.20 ± 0.5 <sup>a</sup>	5.83 ± 0.8 <sup>a</sup>	5.40 ± 0.8	5.23 ± 0.6 <sup>a</sup>
Radicle number (50 grains)	127 ± 7.3 <sup>a</sup>	135 ± 8.1 <sup>a</sup>	139 ± 8.1 <sup>a</sup>	148 ± 7.9 <sup>a</sup>	130 ± 6.7 <sup>a</sup>	119 ± 7.1 <sup>a</sup>

Values are the means ± SD ( $n = 6$ ). Different letters mean significance of difference between different treatments ( $P < 0.05$ ).




FIGURE 1: Effects of SO<sub>2</sub> pretreatment on wheat grain germination under 15 mM Al stress. Wheat grains were pretreated with 0, 0.4, 0.8, 1.2, 1.6, and 2.0 mM SO<sub>2</sub> for 12 h, subsequently subjected to 15 mM Al for further 48 h, and then photographed.

SO<sub>2</sub> donor pretreatment were higher than those pretreated in H<sub>2</sub>O, they are not significantly different.

**3.4. Effect of SO<sub>2</sub> Donor Pretreatment on Contents of Endogenous H<sub>2</sub>S, O<sub>2</sub><sup>•-</sup>, H<sub>2</sub>O<sub>2</sub>, and MDA.** H<sub>2</sub>S, which can be produced from sulfite, is involved in plant growth regulation including various abiotic stresses [8, 22]. To investigate whether exogenous SO<sub>2</sub> application can induce endogenous H<sub>2</sub>S production, we measured the concentration of H<sub>2</sub>S in Al-stressed wheat grain. Generally, H<sub>2</sub>S accumulated during wheat grain germination following pretreatment with water or SO<sub>2</sub>, but SO<sub>2</sub> donor pretreatment significantly enhanced H<sub>2</sub>S concentration at 12 h of pretreatment and 12 h, 36 h of Al stress (Figure 3(a)).

To study the protective role of SO<sub>2</sub> in the Al-stressed wheat grain, reactive oxygen species O<sub>2</sub><sup>•-</sup>, H<sub>2</sub>O<sub>2</sub>, and malondialdehyde (MDA) were determined with time. As shown in Figure 3(b), a rapid accumulation of O<sub>2</sub><sup>•-</sup> was observed when H<sub>2</sub>O-pretreated grains were exposed to Al. During the first 12 h of Al exposure, the increase in O<sub>2</sub><sup>•-</sup> content was very rapid, but this was followed by a slow decrease. In contrast, the content of O<sub>2</sub><sup>•-</sup> in SO<sub>2</sub> pretreatment increased slowly till 36 h of Al stress followed by a decrease. SO<sub>2</sub> pretreatment maintained significantly lower level of O<sub>2</sub><sup>•-</sup> in Al-stressed wheat grains compared with grains incubated in H<sub>2</sub>O and exposed to Al.

H<sub>2</sub>O<sub>2</sub> in both treatments increased gradually during pretreatment time and 36 h of Al stress and decreased at 48 h (Figure 3(c)). However, H<sub>2</sub>O<sub>2</sub> content in SO<sub>2</sub> pretreatment was significantly lower than that in water pretreatment when exposed to Al stress.

During the 12 h pretreatment time, no significant difference was observed in MDA content in wheat grains whether pretreated with SO<sub>2</sub> donor or H<sub>2</sub>O (Figure 3(d)). After exposure to Al, the content of MDA in water pretreated grains increased rapidly till 48 h of Al stress. An increase of MDA content was also observed in SO<sub>2</sub> pretreatment at 12 h of Al stress, but thereafter MDA content remained stable until 36 h. SO<sub>2</sub> pretreatment dramatically reduced the amount of MDA from 24 h to 48 h of Al stress in comparison with grains pretreated in water.

**3.5. Effects of SO<sub>2</sub> Donor Pretreatment on POD, CAT, APX, and LOX Activities.** Activities of POD, CAT, APX, and LOX were determined with time in SO<sub>2</sub> donor and H<sub>2</sub>O-pretreated grains exposed to Al (Figure 4). Figure 4(a) shows the time course of POD activity following pretreatment in SO<sub>2</sub> donor or H<sub>2</sub>O for 12 h when POD activity showed almost a twofold increase. During Al stress, POD activity exhibited a gradual increase in both treatments, but SO<sub>2</sub> pretreatment maintained significantly higher level of POD activity during Al stress.

The activity of CAT increased almost twofold during 12 h pretreatment with H<sub>2</sub>O or SO<sub>2</sub> donor (Figure 4(b)). After exposure to Al, CAT activity in water pretreatment decreased gradually till 48 h of Al stress, suggesting that CAT activity is very sensitive to Al stress. In contrast, CAT activity in SO<sub>2</sub> pretreatment increased steadily and decreased only slightly at 48 h of Al stress.

As shown in Figure 4(c), SO<sub>2</sub> pretreatment enhanced APX activity in Al-stressed wheat grain. A rapid increase in APX activity occurred during the pretreatment time in H<sub>2</sub>O and SO<sub>2</sub>. Within the first 12 h of Al stress, APX activity in H<sub>2</sub>O-pretreated grains decreased sharply, whereas SO<sub>2</sub> donor pretreatment enhanced APX activity slightly. Thereafter APX activity increased steadily in water pretreated grain, whereas its activity in SO<sub>2</sub> donor pretreatment fluctuated slightly. The APX activity in SO<sub>2</sub> donor pretreated grains was always significantly higher than the counterpart of water pretreatment.

An increase in LOX activity was observed during the first 24 h of Al stress in SO<sub>2</sub> and H<sub>2</sub>O-pretreated grains (Figure 4(d)). However, the increase of LOX activity in water pretreatment was more rapid than after SO<sub>2</sub> pretreatment. Thereafter LOX activity in water pretreatment showed a sharp decrease at 36 h of Al stress, while its activity in SO<sub>2</sub> pretreatment decreased at 48 h. At 12 and 24 h of Al stress, SO<sub>2</sub> pretreatment maintained significantly lower level of




FIGURE 2: Effect of SO<sub>2</sub> pretreatment on the contents of reducing sugar and soluble protein in Al-treated grain as shown in (a) and (b), respectively. Wheat grains were pretreated with water (Al) or 1.2 mM SO<sub>2</sub> donor (S + Al) for 12 h (shown from 0' to 12'/0 h of pretreatment time) and then exposed to 15 mM Al for further 48 h (shown as 12'/0, 12, 24, 36, and 48 h). The symbols \* and \*\* in this figure and following ones stand for significant difference between Al-treated grains with and without SO<sub>2</sub> pretreatment at  $P < 0.05$  and  $P < 0.01$ , respectively.

LOX, while at 36 h LOX activity in SO<sub>2</sub> pretreatment was higher than that of water pretreatment.

**3.6. Effects of SO<sub>2</sub> Donor Pretreatment on Localization of Al, Lipid Peroxidation, and ROS Production.** To detect ROS production in the radicle tips, we used DCFH-DA fluorescence to indicate ROS accumulation. As shown in Figure 5(a), Al treatment induced higher level of ROS in radicle as intense DCFH-DA fluorescence, while SO<sub>2</sub> donor pretreatment for 12 h followed by Al stress significantly reduced fluorescence. Figure 5(b) shows DCFH-DA fluorescence in maturation zone in radicles. Similarly, intense fluorescence in SO<sub>2</sub> donor pretreatment followed by Al stress was much weaker than that in water pretreated plus Al-stressed radicles, suggesting that SO<sub>2</sub> donor was effective in alleviating oxidative stress in radicles. SO<sub>2</sub> donor treatment alone showed comparable fluorescence intensity as observed in water control.

The radicles were stained with Evans blue to show membrane integrity. The radicles treated with Al alone were stained extensively with Evans blue, while Al-stressed radicles pretreated with SO<sub>2</sub> donor for 12 h were less stained (Figure 5(c)), suggesting SO<sub>2</sub> donor serves to protect cell membrane from Al-induced damage. SO<sub>2</sub> donor treatment alone showed similar Evans blue staining to water control, suggesting no visible damaging effect of SO<sub>2</sub> on radicles.

The hematoxylin staining was used to detect Al accumulation in radicles. As shown in Figure 5(d), the radicles of water control and SO<sub>2</sub> treatment incubated with hematoxylin showed no dark staining but wheat radicles treated with Al alone were stained intensively. In contrast, radicles pretreated with SO<sub>2</sub> donor for 12 h and then exposed to Al for 48 h

showed much weaker staining compared with Al stress, especially in the elongation zone.

**3.7. Effect of SO<sub>2</sub> Donor Pretreatment on the Relative Expressions of Aluminum Stress Related Genes.** We determined the changes in gene expression of aluminum stress related genes in wheat radicles. Radicles were pretreated with or without 1 mM SO<sub>2</sub> donors for 12 h and then exposed to Al for 48 h. As shown in Figure 6, Al stress induced higher expression of TaWali1, TaWali2, TaWali3, TaWali5, and TaWali6 (wheat aluminum induced) in radicles, while pretreatment with SO<sub>2</sub> donor for 12 h followed by Al stress alleviated such expression increase. Besides, the gene expression of TaALMT1 (Al-activated malate transporter) was also attenuated by SO<sub>2</sub> pretreatment.

## 4. Discussion

In solution, SO<sub>2</sub> is dissociated from its sulfite derivatives (NaHSO<sub>3</sub>/Na<sub>2</sub>SO<sub>3</sub> 1:3 M/M) [34]. Thus NaHSO<sub>3</sub>/Na<sub>2</sub>SO<sub>3</sub> (1:3 M/M) was chosen as an SO<sub>2</sub> donor in our study. Similar to the observation that H<sub>2</sub>S could promote wheat grain germination and alleviate oxidative damage against Al stress [8], our results show that SO<sub>2</sub> donor pretreatment alleviates Al stress in germinating wheat seedlings. Wheat grains pretreated for 12 h with the SO<sub>2</sub> donor show an increase in germination percentage, coleoptile length, radicle length, and radicle numbers of wheat. The increase in the contents of reducing sugars and soluble protein suggests that nutrients in wheat grains pretreated with SO<sub>2</sub> donor are rapidly mobilized to provide energy to grain germination. SO<sub>2</sub> donor maintained lower level of H<sub>2</sub>O<sub>2</sub>, O<sub>2</sub><sup>•-</sup>, and MDA probably




FIGURE 3: Effects of  $SO_2$  pretreatment on the accumulation of endogenous  $H_2S$  (a), superoxide anion ( $O_2^{\cdot-}$ ) (b), hydrogen peroxide ( $H_2O_2$ ) (c), and malondialdehyde (MDA) (d) in germinating wheat grains under Al stress. The numbers (0', 12'/0, 12, 24, 36, and 48) or letters (CK or  $SO_2$ ) presented are the same as mentioned in Figure 2. Al: Al stress without  $SO_2$  pretreatment; S + Al: Al stress with  $SO_2$  pretreatment.

by activation of the antioxidant system. These results suggest that  $SO_2$  acts as an antioxidant and may function in a way that is similar to what the effects of  $H_2S$ , CO, and NO do in plants exposed to heavy metal stress [10, 35].

Sulfite can be reduced by sulfite reductase to  $H_2S$ , which is incorporated into O-acetylserine via O-acetyl(thiol)lyase to form cysteine [22]. In RNA interfered mutant of sulfite reductase (SiR), sulfide synthesis in younger leaves was decreased by the impaired SiR activity [36]. In the present study, exogenous  $SO_2$  application can induce endogenous  $H_2S$  production in Al-stressed wheat grains (Figure 3(a)), suggesting the interplay between sulfite and the formation of  $H_2S$ .

Consistent with previous observations [7], our results show that Al stress caused overproduction of ROS in wheat.

To mitigate and repair oxidative damage, plants have evolved an efficient antioxidant system that includes enzymes such as SOD, CAT, and APX that function to scavenge ROS [37]. SOD catalyzes the dismutation of the superoxide radical  $O_2^{\cdot-}$  and  $H^+$  into  $H_2O_2$ . CAT, APX, and POD are responsible for the elimination of  $H_2O_2$  generated by SOD. Al stress brings about a dramatic increase in  $H_2O_2$  and  $O_2^{\cdot-}$ . The elevated levels of  $H_2O_2$  and  $O_2^{\cdot-}$  suggest that antioxidant enzymes in Al-stressed wheat do not efficiently scavenge the overproduction of ROS, and this can result in lipid peroxidation or plasma membrane inhibiting grain germination and seedling growth [8]. Our data show that pretreatment of wheat with  $SO_2$  donor activates antioxidant enzymes including POD, CAT, and APX.




FIGURE 4: Effect of SO<sub>2</sub> donor pretreatment on the activities of POD (a), CAT (b), APX (c), and LOX (d) in germinating wheat grains under 15 mM Al stress. Grains were treated and the number or letters presented are the same as mentioned in Figure 2. Al: Al stress without SO<sub>2</sub> pretreatment; S + Al: Al stress with SO<sub>2</sub> pretreatment.

LOX, which catalyzes oxygenation of polyunsaturated fatty acids into lipid hydroperoxides, is considered an indicator of oxidative stress during responses to various environmental stresses [9]. Pretreatment with SO<sub>2</sub> donor lowers LOX activity in Al-stressed wheat radicles compared to seedlings pretreated with H<sub>2</sub>O and exposed to Al. The lowering of LOX by SO<sub>2</sub> pretreatment also helps to explain the lower MDA content of Al-stressed grain. Taken together, these data suggest that SO<sub>2</sub> donor reduced oxidative stress by modulation of the antioxidant system.

Our data indicate that the radicle is the primary target for Al toxicity. DCFH-DA fluorescence assay shows that Al incubation induces higher accumulation of ROS in radicle

tips and maturation zone. SO<sub>2</sub> donor pretreatment effectively reduces ROS content in subsequent Al stress, suggesting the role of SO<sub>2</sub> in alleviating oxidative stress. Correspondently, Al stress causes membrane injury to radicles, while SO<sub>2</sub> donor effectively alleviates such injury. To understand whether SO<sub>2</sub> donor helps to reduce Al accumulation in radicles, hematoxylin staining was used to indicate Al and the results show that SO<sub>2</sub> donor obviously reduces Al content in radicles, implying a potential role of SO<sub>2</sub> donor treatment as a strategy to reduce Al uptake.

In response to Al stress, many gene expressions are activated, for instance, TaWali (wheat aluminum induced), aluminum-activated malate transporter (TaALMT1) [38–41].




FIGURE 5: ROS staining ((a) on radicle tips; (b) on maturation zone; bar: 200  $\mu\text{m}$ ), Evans blue staining (c), and hematoxylin staining (d) in wheat radicles. Initially, wheat grains were germinated in water for 36 h. Then four treatment groups were done as follows, control, 60 h in  $\text{H}_2\text{O}$ ; Pre-S, pretreatment with 1 mM  $\text{SO}_2$  donor for 12 h, and then exposed to  $\text{H}_2\text{O}$  for 48 h; Al, 12 h in  $\text{H}_2\text{O}$  prior to exposure to 400  $\mu\text{M}$   $\text{AlCl}_3$  for 48 h; Pre-S + Al, 12 h in 1 mM  $\text{SO}_2$  donor pretreatment followed by 400  $\mu\text{M}$   $\text{AlCl}_3$  for 48 h.






FIGURE 6: Effect of SO<sub>2</sub> donor pretreatment on relative gene expression of TaWali1, TaWali2, TaWali3, TaWali5, TaWali6, and TaALMT1 in wheat radicals exposed to Al stress. Initially, wheat grains were germinated in water for 36 h. Then four treatment groups were done as follows, control, 60 h in H<sub>2</sub>O; Pre-S, pretreatment with 1 mM SO<sub>2</sub> donor for 12 h, and then exposed to H<sub>2</sub>O for 48 h; Al, 12 h in H<sub>2</sub>O prior to exposure to 400 μM AlCl<sub>3</sub> for 48 h; Pre-S + Al, 12 h in 1 mM SO<sub>2</sub> donor pretreatment followed by 400 μM AlCl<sub>3</sub> for 48 h.

Relative gene expression analysis shows that Al treatment induces higher expression of TaWali, while these gene expression levels are reduced by SO<sub>2</sub> donor pretreatment, suggesting the response to Al stress is attenuated in SO<sub>2</sub> donor pretreatment.

## 5. Conclusion

In the present study, SO<sub>2</sub> acts as an antioxidant signal to reduce ROS damage in wheat grains and radicles caused by Al stress. Besides, SO<sub>2</sub> also decreases Al uptake. The induced higher level of H<sub>2</sub>S suggests an intricate interplay of SO<sub>2</sub>

and H<sub>2</sub>S in plants. Exogenous application of SO<sub>2</sub> may be reduced to H<sub>2</sub>S by sulfite reductase, thus contributing to H<sub>2</sub>S production. H<sub>2</sub>S in itself acts as an antioxidant signaling molecule in plants' response to abiotic stress. Thus the nature of SO<sub>2</sub>/sulfite functions in alleviating Al stress still needs further research.

## Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

## Authors' Contribution

Dong-Bo Zhu, Kang-Di Hu, and Xi-Kai Guo contributed equally to this work.

## Acknowledgments

This work was supported by the Natural Science Foundation of China (31271803, 31301820, 31300133, and 31470013), the Scientific Research Foundation for Returned Overseas Chinese Scholars (SRF for ROCS, MOE), the Natural Science Foundations of Anhui Province (H040606M85), and the Anhui Provincial Education Department (2012AJZR0028, ZD200910).

## References

- [1] P. R. Ryan, S. D. Tyerman, T. Sasaki et al., "The identification of aluminium-resistance genes provides opportunities for enhancing crop production on acid soils," *Journal of Experimental Botany*, vol. 62, no. 1, pp. 9–20, 2011.
- [2] Z. Q. Wang, X. Y. Xu, Q. Q. Gong et al., "Root proteome of rice studied by iTRAQ provides integrated insight into aluminum stress tolerance mechanisms in plants," *Journal of Proteomics*, vol. 98, no. 26, pp. 189–205, 2014.
- [3] W. Huang, X. Yang, S. Yao et al., "Reactive oxygen species burst induced by aluminum stress triggers mitochondria-dependent programmed cell death in peanut root tip cells," *Plant Physiology and Biochemistry*, vol. 82, no. 12, pp. 76–84, 2014.
- [4] Y.-S. Wang and Z.-M. Yang, "Nitric oxide reduces aluminum toxicity by preventing oxidative stress in the roots of *Cassia tora* L.," *Plant and Cell Physiology*, vol. 46, no. 12, pp. 1915–1923, 2005.
- [5] K. Tahara, T. Yamanoshita, M. Norisada et al., "Aluminum distribution and reactive oxygen species accumulation in root tips of two *Melaleuca* trees differing in aluminum resistance," *Plant and Soil*, vol. 307, no. 1-2, pp. 167–178, 2008.
- [6] Z. J. Ding, J. Y. Yan, X. Y. Xu, G. X. Li, and S. J. Zheng, "WRKY46 functions as a transcriptional repressor of *ALMT1*, regulating aluminum-induced malate secretion in Arabidopsis," *The Plant Journal*, vol. 76, no. 5, pp. 825–835, 2013.
- [7] S. J. Zheng and J. L. Yang, "Target sites of aluminum phytotoxicity," *Biologia Plantarum*, vol. 49, no. 3, pp. 321–331, 2005.
- [8] H. Zhang, Y. H. Li, L. Y. Hu, S. H. Wang, F. Q. Zhang, and K. D. Hu, "Effects of exogenous nitric oxide donor on antioxidant metabolism in wheat leaves under aluminum stress," *Russian Journal of Plant Physiology*, vol. 55, no. 4, pp. 469–474, 2008.
- [9] H. Zhang, Z.-Q. Tan, L.-Y. Hu, S.-H. Wang, J.-P. Luo, and R. L. Jones, "Hydrogen sulfide alleviates aluminum toxicity in germinating wheat seedlings," *Journal of Integrative Plant Biology*, vol. 52, no. 6, pp. 556–567, 2010.
- [10] J. Xu, W. Xuan, B. Huang et al., "Carbon monoxide-induced adventitious rooting of hypocotyl cuttings from mung bean seedling," *Chinese Science Bulletin*, vol. 51, no. 6, pp. 668–674, 2006.
- [11] S. Q. Zhang, J. B. Du, H. F. Jin et al., "Endogenous sulfur dioxide aggravates myocardial injury in isolated rat heart with ischemia and reperfusion," *Transplantation*, vol. 87, no. 4, pp. 517–524, 2009.
- [12] R. Y. Zhang, J. B. Du, Y. Sun et al., "Sulfur dioxide derivatives depress L-type calcium channel in rat cardiomyocyte," *Clinical and Experimental Pharmacology and Physiology*, vol. 38, no. 7, pp. 416–422, 2011.
- [13] H.-F. Jin, S.-X. Du, X. Zhao et al., "Effects of endogenous sulfur dioxide on monocrotaline-induced pulmonary hypertension in rats," *Acta Pharmacologica Sinica*, vol. 29, no. 10, pp. 1157–1166, 2008.
- [14] I. Ziegler, "The effect of SO<sub>2</sub> pollution on plant metabolism," *Residue Reviews*, vol. 56, no. 1, pp. 79–105, 1975.
- [15] R. Sandhu, Y. Li, and G. Gupta, "Sulphur dioxide and carbon dioxide induced changes in soybean physiology," *Plant Science*, vol. 83, no. 1, pp. 31–34, 1992.
- [16] D. Yarmolinsky, G. Brychkova, R. Fluhr, and M. Sagi, "Sulfite reductase protects plants against sulfite toxicity," *Plant Physiology*, vol. 161, no. 2, pp. 725–743, 2013.
- [17] M. Noji, M. Saito, M. Nakamura, M. Aono, H. Saji, and K. Saito, "Cysteine synthase overexpression in tobacco confers tolerance to sulfur-containing environmental pollutants," *Plant Physiology*, vol. 126, no. 3, pp. 973–980, 2001.
- [18] R. Rakwal, G. K. Agrawal, A. Kubo et al., "Defense/stress responses elicited in rice seedlings exposed to the gaseous air pollutant sulfur dioxide," *Environmental and Experimental Botany*, vol. 49, no. 3, pp. 223–235, 2003.
- [19] J. J. Yin, X. Liu, H. L. Yi, and M. L. Yang, "Sulfur dioxide induces guard cell death in *Vicia faba*," *Acta Scientiae Circumstantiae*, vol. 30, no. 12, pp. 2512–2517, 2010.
- [20] H. Rennenberg, "The fate of excess sulfur in higher plants," *Annual Review of Plant Physiology*, vol. 35, no. 4, pp. 121–153, 1984.
- [21] E. Giraud, A. Ivanova, C. S. Gordon, J. Whelan, and M. J. Considine, "Sulphur dioxide evokes a large scale reprogramming of the grape berry transcriptome associated with oxidative signalling and biotic defence responses," *Plant, Cell and Environment*, vol. 35, no. 2, pp. 405–417, 2012.
- [22] T. Rausch and A. Wachter, "Sulfur metabolism: a versatile platform for launching defence operations," *Trends in Plant Science*, vol. 10, no. 10, pp. 503–509, 2005.
- [23] M. H. Stipanuk, J. E. Dominy Jr., J.-I. Lee, and R. M. Coloso, "Mammalian cysteine metabolism: new insights into regulation of cysteine metabolism," *Journal of Nutrition*, vol. 136, no. 6, pp. 1652S–1659S, 2006.
- [24] A. Laisk, H. Pfan, and U. Heber, "Sulfur-dioxide fluxes into different cellular compartments of leaves photosynthesizing in a polluted atmosphere: II. Consequences of SO<sub>2</sub> uptake as revealed by computer analysis," *Planta*, vol. 173, no. 2, pp. 241–252, 1988.
- [25] H. Zhang, S.-L. Hu, Z.-J. Zhang et al., "Hydrogen sulfide acts as a regulator of flower senescence in plants," *Postharvest Biology and Technology*, vol. 60, no. 3, pp. 251–257, 2011.
- [26] K. Surrey, "Spectrophotometric method for determination of lipoxidase activity," *Plant Physiology*, vol. 39, no. 1, pp. 65–70, 1964.
- [27] L.-Y. Hu, S.-L. Hu, J. Wu et al., "Hydrogen sulfide prolongs postharvest shelf life of strawberry and plays an antioxidative role in fruits," *Journal of Agricultural and Food Chemistry*, vol. 60, no. 35, pp. 8684–8693, 2012.
- [28] G. L. Miller, "Use of dinitrosalicylic acid reagent for determination of reducing sugar," *Analytical Chemistry*, vol. 31, no. 3, pp. 426–428, 1959.
- [29] M. M. Bradford, "A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein dye binding," *Analytical Biochemistry*, vol. 72, no. 1-2, pp. 248–254, 1976.

- [30] Y. Yamamoto, Y. Kobayashi, S. R. Devi, S. Rikiishi, and H. Matsumoto, "Aluminum toxicity is associated with mitochondrial dysfunction and the production of reactive oxygen species in plant cells," *Plant Physiology*, vol. 128, no. 1, pp. 63–72, 2002.
- [31] E. Polle, C. F. Konzak, and J. A. Kittrick, "Visual detection of aluminum tolerance levels in wheat by hematoxylin staining of seeding roots," *Crop Science*, vol. 18, no. 5, pp. 823–827, 1978.
- [32] C. P. LeBel, H. Ischiropoulos, and S. C. Bondy, "Evaluation of the probe 2',7'-dichlorofluorescein as an indicator of reactive oxygen species formation and oxidative stress," *Chemical Research in Toxicology*, vol. 5, no. 2, pp. 227–231, 1992.
- [33] S.-P. Li, K.-D. Hu, L.-Y. Hu et al., "Hydrogen sulfide alleviates postharvest senescence of broccoli by modulating antioxidant defense and senescence-related gene expression," *Journal of Agricultural and Food Chemistry*, vol. 62, no. 5, pp. 1119–1129, 2014.
- [34] R. Shapiro, "Genetic effects of bisulfite (sulfur dioxide)," *Mutation Research*, vol. 39, no. 2, pp. 149–175, 1977.
- [35] K.-D. Hu, L.-Y. Hu, Y.-H. Li, F.-Q. Zhang, and H. Zhang, "Protective roles of nitric oxide on germination and antioxidant metabolism in wheat seeds under copper stress," *Plant Growth Regulation*, vol. 53, no. 3, pp. 173–183, 2007.
- [36] D. Yarmolinsky, G. Brychkova, A. Kurmanbayeva et al., "Impairment in sulfite reductase leads to early leaf senescence in tomato plants," *Plant Physiology*, vol. 165, no. 4, pp. 1505–1520, 2014.
- [37] P. Sharma and R. S. Dubey, "Involvement of oxidative stress and role of antioxidative defense system in growing rice seedlings exposed to toxic concentrations of aluminum," *Plant Cell Reports*, vol. 26, no. 11, pp. 2027–2038, 2007.
- [38] K. C. Snowden and R. C. Gardner, "Five genes induced by aluminum in wheat (*Triticum aestivum* L.) roots," *Plant Physiology*, vol. 103, no. 3, pp. 855–861, 1993.
- [39] A. Ligaba, I. Dreyer, A. Margaryan, D. J. Schneider, L. Kochian, and M. Piñeros, "Functional, structural and phylogenetic analysis of domains underlying the Al sensitivity of the aluminum-activated malate/anion transporter, TaALMT1," *The Plant Journal*, vol. 76, no. 5, pp. 766–780, 2013.
- [40] N. Yoshimoto, H. Takahashi, F. W. Smith, T. Yamaya, and K. Saito, "Two distinct high-affinity sulfate transporters with different inducibilities mediate uptake of sulfate in *Arabidopsis* roots," *The Plant Journal*, vol. 29, no. 4, pp. 465–473, 2002.
- [41] T. Kataoka, A. Watanabe-Takahashi, N. Hayashi et al., "Vacuolar sulfate transporters are essential determinants controlling internal distribution of sulfate in *Arabidopsis*," *The Plant Cell*, vol. 16, no. 10, pp. 2693–2704, 2004.