

Research article

Open Access

Cloning of the cytochrome p450 reductase (*crtR*) gene and its involvement in the astaxanthin biosynthesis of *Xanthophyllomyces dendrorhous*

Jennifer Alcaíno, Salvador Barahona, Marisela Carmona, Carla Lozano, Andrés Marcoleta, Mauricio Niklitschek, Dionisia Sepúlveda, Marcelo Baeza and Víctor Cifuentes*

Address: Departamento de Ciencias Ecológicas, Facultad de Ciencias, Universidad de Chile, Santiago, Chile

Email: Jennifer Alcaíno - jennyalcaino@vtr.net; Salvador Barahona - barahona.salvador@gmail.com; Marisela Carmona - marisela.carmona@gmail.com; Carla Lozano - carlalozan@gmail.com; Andrés Marcoleta - amarcoleta@gmail.com; Mauricio Niklitschek - maurobiotec@gmail.com; Dionisia Sepúlveda - dony@uchile.cl; Marcelo Baeza - mbaeza@uchile.cl; Víctor Cifuentes* - vcifuentes@uchile.cl

* Corresponding author

Published: 6 October 2008

Received: 1 July 2008

BMC Microbiology 2008, 8:169 doi:10.1186/1471-2180-8-169

Accepted: 6 October 2008

This article is available from: <http://www.biomedcentral.com/1471-2180/8/169>

© 2008 Alcaíno et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: The yeast *Xanthophyllomyces dendrorhous* synthesizes astaxanthin, a carotenoid with high commercial interest. The proposed biosynthetic route in this organism is isopentenyl-pyrophosphate (IPP) → geranylerythryl pyrophosphate (GGPP) → phytoene → lycopene → β-carotene → astaxanthin. Recently, it has been published that the conversion of β-carotene into astaxanthin requires only one enzyme, astaxanthin synthase or CrtS, encoded by *crtS* gene. This enzyme belongs to the cytochrome P450 protein family.

Results: In this work, a *crtR* gene was isolated from *X. dendrorhous* yeast, which encodes a cytochrome P450 reductase (CPR) that provides CrtS with the necessary electrons for substrate oxygenation. We determined the structural organization of the *crtR* gene and its location in the yeast electrophoretic karyotype. Two transformants, CBSTr and T13, were obtained by deleting the *crtR* gene and inserting a hygromycin B resistance cassette. The carotenoid composition of the transformants was altered in relation to the wild type strain. CBSTr forms yellow colonies because it is unable to produce astaxanthin, hence accumulating β-carotene. T13 forms pale colonies because its astaxanthin content is reduced and its β-carotene content is increased.

Conclusion: In addition to the *crtS* gene, *X. dendrorhous* requires a novel gene, *crtR*, for the conversion of β-carotene to astaxanthin.

Background

Carotenoids are natural pigments of yellow, orange or red color. More than 600 different chemical structures have been described to date [1]. They are terpenoids with the

isopentenyl- pyrophosphate (IPP) molecule as the basic unit. Astaxanthin is a carotenoid with a high commercial interest due to its use as a food additive for trout and salmon flesh pigmentation in aquaculture [2]. Its biosyn-

thesis is limited to a few microorganisms such as the microalgae *Haematococcus pluvialis* and the basidiomycete yeast *Xanthophyllomyces dendrorhous* [3,4].

The first step in the biosynthesis of astaxanthin in *X. dendrorhous* is the isomerization of IPP into dimethylallyl pyrophosphate (DMAPP) by IPP isomerase, which is encoded by the *idi* gene [5]. Three IPP molecules are sequentially added to one DMAPP molecule, where the GGPP synthase (*crtE* gene) is involved, forming geranylgeranyl pyrophosphate (GGPP) [6]. Then, the condensation of two GGPP molecules produces phytoene, which corresponds to the first carotenoid synthesized in the pathway. This step is catalyzed by a bi-functional enzyme called phytoene- β -carotene synthase (PBS), which also has lycopene cyclase activity and is encoded by the *crtYB* gene [7]. Subsequently, phytoene goes through four successive desaturation reactions, catalyzed by the phytoene desaturase enzyme (*crtI* gene), producing lycopene [8]. Then, both lycopene ends are cyclized by the lycopene cyclase activity of the bi-functional PBS enzyme, forming β -carotene.

In the last step for the formation of astaxanthin from β -carotene, two enzymatic activities are required, ketolase, to incorporate a keto group in positions 4 and 4' of β -carotene, and hydroxylase, to introduce a hydroxyl group at positions 3 and 3' of β -carotene [9,10]. Unlike other organisms, where two independent genes are needed for this step, in *X. dendrorhous* a single *crtS* gene has been isolated that encodes an astaxanthin synthase which can probably perform both activities [11-13]. However, it has been suggested that CrtS may require an auxiliary enzyme for the formation of astaxanthin from β -carotene [14]. The deduced CrtS amino acid sequence strongly suggests that the *X. dendrorhous* enzyme belongs to the cytochrome P450 protein family [12,14].

Cytochrome P450 enzymes require an electron donor for the addition of oxygen-bearing functional groups to a substrate. In eukaryotes, this electron donor is cyto-

chrome P450 reductase (CPR), encoded by the *cpr* gene [15]. Although several genes for different cytochrome P450 enzymes can exist in an organism, in most species only one *cpr* gene exists. Some exceptions have been observed in plants and zygomycetes that contain two or three *cpr* genes [16].

In another study, an *Escherichia coli* strain was transformed with three compatible plasmids. The first plasmid contained the necessary genes for the synthesis of β -carotene, the second contained the *X. dendrorhous crtS* gene, and the third contained the *Saccharomyces cerevisiae cpr* gene [12]. Through this system, it was possible to obtain oxygenated β -carotene derivatives, but not astaxanthin, suggesting the need for an appropriate CPR to complement the astaxanthin biosynthetic pathway. Prior to this study, a *X. dendrorhous* gene encoding for a CPR for astaxanthin biosynthesis had not been isolated. This paper describes for the first time the sequence of the *X. dendrorhous crtR* gene and its encoded polypeptide. Moreover, our results reveal that the *crtR* gene product is essential for astaxanthin biosynthesis in *X. dendrorhous*.

Methods

Strains, plasmids, media and enzymes

The strains and plasmids used in this study are listed in Tables 1 and 2. The wild type *X. dendrorhous* UCD 67-385 strain was used for genomic and cDNA library construction. *E. coli* DH-5 α strain was used as a host for plasmid propagation and genomic libraries. Two genomic libraries were used in this study. The first library consisted of genomic DNA partially digested with *Bam*HI and cloned into the *Bam*HI site of the Yip5 vector [6]. The second library consisted of approximately 5 to 6 kb genomic DNA fragments digested with *Sal*I and cloned into the *Sal*I site of the pBluescript SK- vector. The cDNA library was constructed with the pBluescript II XR cDNA library construction kit from Stratagene [6].

crtR gene deletion bearing plasmids (pBsiWI Δ *crtR*::*hph* and pNdeI Δ *crtR*::*hph*) were built from pCPR1.3 (Table 2),

Table 1: *E. coli* and *X. dendrorhous* strains used in this work

| Strains | Genotype or relevant features | Reference |
|-----------------------|--|-----------|
| <i>E. coli</i> | | |
| DH-5 α | F- ϕ 80d <i>lacZ</i> Δ M15 Δ (<i>lacZY-argF</i>)U169 <i>deoR recA1 endA1 hsdR17</i> (<i>r_k-m_k⁺</i>) <i>phoA supE44I-thi-1 gyrA96 relA1</i> | [18] |
| DS1B | β -carotene producer strain with a yellow phenotype which corresponds to BL21-Gold strain carrying pDS1B | [6] |
| <i>X. dendrorhous</i> | | |
| UCD 67-385 | ATCC 24230, wild-type | ATCC |
| CBS-6938 | ATCC 96594, wild-type | ATCC |
| T13 | (<i>crtR</i> Δ BsiWI:: <i>hph/crtR⁺). Heterozygote transformant from UCD 67-385 parental wild-type strain, containing an allele of the <i>crtR</i> gene with a BsiWI deletion and a hygromycin B resistance cassette.</i> | This work |
| CBSTr | (<i>crtR</i> Δ NdeI:: <i>hph</i>). Transformant from CBS-6938 parental wild-type strain. β -carotene accumulating mutant. | This work |

Table 2: Plasmids used in this work

| Plasmid | Genotype or relevant features | Source of reference |
|--|--|-----------------------------|
| pBluescript SK- pBAD33 | ColEI ori; Amp ^R ; cloning vector with blue-white selection p15 ori; Cam ^R ; cloning vector | Stratagene Beckwith Lab. |
| Ylp5 | <i>S. cerevisiae</i> /E. coli cloning vector. pMB1 ori; Amp ^R ; TET ^R ; URA3. | ATCC 37061 |
| pET101/D-TOPO | ColEI ori; Amp ^R ; expression vector | Invitrogene |
| pDSIB | pBAD33 vector carrying the carotenogenesis genes <i>crtE</i> , <i>crtB</i> , <i>crtI</i> and <i>crtY</i> and <i>crtX</i> from <i>Erwinia uredovora</i> [40]. | This work |
| pPR52.P.44 | Ylp5 bearing an insert of 12.8 kb that contains the <i>crtR</i> gene from base n° 88 of exon I. | This work |
| pCPR1.3 | pBluescript SK- bearing the DNA fragment from site <i>Bam</i> HI 4,529 to <i>Bam</i> HI 9,895 (Figure 1). pCPR1.3 was the <i>crtR</i> deletion platform. | This work |
| pcXDA.44.29 | pBluescript SK- bearing the <i>crtR</i> gene cDNA. The cDNA measures 2,680 bp. | This work |
| pPR32.AA.51 | Ylp5 bearing an insert of 4.5 kb that contains the <i>crtR</i> gene up to base n° 87 of exon I. | This work |
| pCPRB6 | pBluescript SK- bearing an insert of 5.9 kb that contains the complete genomic version of <i>crtR</i> gene. Isolated from a <i>SalI</i> partial genomic library. | This work |
| p <i>Bsi</i> WIΔ <i>crtR</i> :: <i>hph</i> | pCPR1.3 <i>Bsi</i> WI deletion and <i>hph</i> cassette insertion. | This work |
| p <i>Nde</i> IΔ <i>crtR</i> :: <i>hph</i> | pCPR1.3 <i>Nde</i> I deletion and <i>hph</i> cassette insertion. | This work |

which has the *X. dendrorhous crtR* gene. The deletions were created by digesting pCPR1.3 with *Bsi*WI or *Nde*I and then filling the ends with Klenow polymerase to enable the ligation of the blunt ended hygromycin B resistance cassette (*hph*) [6] for the selection of *X. dendrorhous* transformants.

X. dendrorhous strains were grown at 22°C with constant agitation in YM medium (1% glucose, 0.3% yeast extract, 0.3% malt extract and 0.5% peptone) or MM_v + 2% glucose medium [17]. UCD 67–385 strain fermentation was performed in a 12 l jar fermentor (New Brunswick) containing 8.8 l of MM_v + 2% glucose medium, shaking at 300 rpm and 13 l/min sterile air injection. Antifoam agent (1520 US, Dow Corning) was automatically added when required. *E. coli* strains were grown with constant agitation at 37°C in Luria-Bertani (LB) medium and supplemented with 100 µg/ml ampicillin for plasmid selection and 40 µl of a 2% solution of X-gal (5-bromo-4chloro-3-indolyl-β-D-galactopyranoside) for recombinant clone selection [18]. For *E. coli* complementation experiments, the LB growth medium was supplemented with 10 µM heme and 0.5 mM δ-aminolevulinic acid, which supports heme prosthetic group formation in *E. coli* [12,19]. The expression was induced with 1 mM IPTG. Enzymes were purchased from Promega (*Taq*DNA pol, restriction enzymes, Klenow polymerase, M-MLV reverse transcriptase).

DNA amplification

Primers were purchased from Alpha DNA (Montreal, Canada) and are listed in Table 3. PCR reactions were performed in a final volume of 25 µl containing 2 U of *Taq*DNA pol, 2.5 µl of 10× *Taq* buffer, 0.5 µl of 10 mM dNTPs, 1 µl of 50 mM MgCl₂, 1 µl of 25 µM of each primer and 10–20 ng DNA. Amplification was done in a DNA

thermal cycler 2400 (Perkin-Elmer) as follows: initial denaturation at 95°C for 3 min; 35 cycles of denaturation at 94°C for 30 s, annealing at 55°C for 30 s, synthesis at 72°C for 3 min and a final extension step at 72°C for 10 min. Samples were kept at 4°C until checked. The amplicons were separated by 0.8% agarose gel electrophoresis in TAE buffer containing 0.5 µg/ml ethidium bromide [18] followed by DNA purification using glassmilk [20] for sequencing.

DNA sequence and bioinformatic tools

Nucleotide sequences were obtained from an ABI 3100 Avant genetic analyzer, using a DYEnamic ET terminator Kit (Amersham Bioscience). DNA sequences were analyzed with Vector NTI Suite 10 (Informax), CLUSTAL W 1.8 and BLAST programs. Protein analysis was performed with the online programs InterProScan <http://www.ebi.ac.uk/InterProScan/> and TMPRED http://www.ch.embnet.org/software/TMPRED_form.html. Phylogenetic analysis was carried out using BioEdit 7.0.0 and Treecon 1.3b.

Pulse field gel electrophoresis

Chromosomal DNA was separated by contour-clamped homogeneous electric fields (CHEF) in a CHEF II BioRad system in 0.9% agarose gels in TBE 0.5× buffer (45 mM Tris-borate, 1 mM EDTA pH 8.0) at 14°C [21]. The pulses used were 90 s for 24 h, followed by 120 s for 24 h at 6 V/cm.

Southern blot hybridization

Southern blot hybridization was performed according to Sambrook and Russell [18]. *crtR* probe was obtained by PCR amplification with primers CPR3 and CPR7 of the *crtR* gene. *hph* probe was acquired from PCR amplification

Table 3: Primers used in this work

| Primer | Direction | Sequence 5' to 3' | Target |
|------------|-----------|----------------------------|------------------------|
| CPR3 | F | CARACTGGKACDGGCHGARGATT | <i>crtR</i> gene |
| CPR4 | F | CAAACCTGGTACGGCTGAAGATT | <i>crtR</i> gene |
| CPR5 | R | WGGDCCRATCATGAYRACTGG | <i>crtR</i> gene |
| CPR6 | R | AGGTCCAATCATGACGACTGG | <i>crtR</i> gene |
| CPR7 | R | CRGTACCWGGDCCRATCATGA | <i>crtR</i> gene |
| CPR8 | R | CCAGTACCAGGTCCAATCATGA | <i>crtR</i> gene |
| CPR9 | F | GGATCCGCGACATCGAAGAGTATGAC | <i>crtR</i> gene |
| CPR12 | R | GGATCCCTTCCAAGCGAGGTAGTCTT | <i>crtR</i> gene |
| CPR13 | R | AGAAGACTGTGCGATCGTGT | <i>crtR</i> gene |
| CPR31 | R | TCAAGCAATTGGTGTGGTC | <i>crtR</i> gene |
| CPR56 | R | TCGTTGTGTTGTTGCTATTCCG | <i>crtR</i> gene |
| CPR3A | F | TACAACGTCGTCGGTAGACA | <i>crtR</i> gene |
| CPR7A | R | TATTCACACCGTGGTGTTC | <i>crtR</i> gene |
| CPR3D | F | GGTCTCACTTCTCCAGAGAA | <i>crtR</i> gene |
| CPRfwdC | F | GCACAGGAAGTTGGTTGGAT | <i>crtR</i> gene |
| CPRlexC | F | GCCCACTCTCCGATCTTGT | <i>crtR</i> gene |
| CPREx1fwdB | F | TCAGGACCCTGTACAGTCAGC | <i>crtR</i> gene |
| CPREx1rev | R | TCCCAACAGTCGATCCTTGT | <i>crtR</i> gene |
| Pef | F | GATATCGGCTCATCAGCCGAC | EF-1 α Promoter |
| Gpd | R | ATGAGAGATGACGGAGATG | GPDH Terminator |
| HR | R | CTATTCCTTTGCCCTCGGAC | <i>hph</i> gene |
| HF | F | ATGAAAAAGCCTGAACCTACC | <i>hph</i> gene |
| HygSecR | R | GTATTGACCGATTCTTGGCG | <i>hph</i> gene |
| HygSecF | F | TCGCCAACATCTTCTTCTGG | <i>hph</i> gene |
| M13R | R | GGAAACAGCTATGACCATG | pBluescript SK- |
| M13F | F | TGTAACACGACGGCCAGT | pBluescript SK- |
| ACT3 | F | ACTCCTACGTTGGTGACGAG | Actin gene |
| ACT4 | R | TCAAGTCTCGACCGCCAAG | Actin gene |
| AST1 | F | GCCACCTACTTTCTCCATATGT | <i>crtS</i> gene |
| AST2 | R | GAGCCATGACGTCCAGAGTA | <i>crtS</i> gene |

F: forward, R: reverse

of the *hph* cassette using primers HF and HR. Probes were labeled with [³²P] α -dCTP using the Promega prime-a-gene labeling system.

RNA extraction and cDNA synthesis

Total RNA extraction was performed according to a modified protocol of Chomczynski and Sacchi [22,23]. The determination of the relative levels of *crtS* and *crtR* mRNAs was performed using a semi-quantitative RT-PCR method. The intensities of the *crtS* and *crtR* amplification bands were normalized with the intensity of the actin [24] amplification product [23,25]. The amplicons were quantified with ImageJ 1.40 using a 100 bp DNA ladder (Fermentas) as the standard.

X. dendrorhous transformation

X. dendrorhous transformation was performed by electroporation according to [26] and [27]. Electrocompetent cells were prepared from an exponential culture with DO_{600 nm} = 1.2 grown in YM medium [6] and electroporated using a BioRad gene pulser \times cell with PC and CE modules under the following conditions: 125 mF, 600 Ω , 0.45 kV. Transformant selection was performed in YM

1.5% agar plates supplemented with 10 μ g/ml hygromycin B. The transformants were identified as *X. dendrorhous* by ITS1, 5.8 rRNA gene and ITS2 DNA sequences [28].

Pigment extraction and RP-HPLC

Carotenoid extraction was carried out from cellular pellets according to the acetone extraction method [29]. Carotenoids were quantified by absorbance at 465 nm using an absorption coefficient of A_{1%} = 2,100. The analyses were performed in triplicate, and pigments were normalized relative to the dry weight of the yeast. Carotenoids were separated by RP-HPLC using a reverse phase RP-18 Lichrocart 125-4 (Merck) column with acetonitrile: methanol: isopropyl (85:10:5 v/v) as the mobile phase with a 1 ml/min flux under isocratic conditions. The elution spectra were recuperated using a diode array detector.

Results and discussion

Cloning of *crtR* gene

Three cytochrome P450 reductase genes were used for primer design: from the zygomycete *Cunninghamella echinulata* [GenBank:AF195660], the ascomycete *Saccharomyces cerevisiae* [GenBank:D13788] and the basidiomyc-

ete *Rhodotorula minuta* [GenBank:AB055119]. Based on the conserved DNA segments, degenerate primers were designed (Table 3). PCR reactions were performed using genomic DNA from UCD 67–385 *X. dendrorhous* as a template and different combinations of forward and reverse primers. An amplicon of 1.6 kb was obtained when PCR reactions were performed with primers CPR3 and CPR7. This amplicon was completely sequenced and was found to be homologous to CPR genes by BLAST analysis. We named this gene *crtR* and designed specific *crtR* primers.

The Ylp5 genomic library was screened for the *X. dendrorhous crtR* gene by PCR [6], and one recombinant plasmid carrying the *crtR* gene (pPR52.P.44) was isolated. This plasmid contained a DNA insert of approximately 12.8 kb, carrying six *Bam*HI fragments of 5.4 kb, 4.4 kb, 0.9 kb (doublet), 0.8 kb and 0.4 kb. Each fragment was sub-cloned into the pBluescript SK- vector for sequencing. The sequence analysis showed that the *crtR* gene was in one end of the 5.4 kb fragment (pCPR1.3), but it was incomplete. The promoter region and part of the first exon were missing due to the presence of a *Bam*HI site in the *crtR* gene.

A cDNA library was screened for the *X. dendrorhous crtR* cDNA, and a recombinant plasmid, pcXDA.44.29, was isolated, which contained a 2,680 bp insert corresponding to the cDNA of the *crtR* gene [GenBank:EU884134]. We used this cDNA sequence for specific primer design (primers CPREx1fwd and CPREx1rev, Table 3) from the remaining *crtR* gene sequence, and this allowed the isolation of the pPR32.AA.51 plasmid from the Ylp5 genomic library. pPR32.AA.51 contains the promoter and 5' region of the first exon of the *crtR* gene. Sequence analysis indicated that the 5' region of the *crtR* gene is at one end of the 4.5 kb *Bam*HI DNA fragment.

The *in silico* DNA sequence analysis of the inserts from pPR32.AA.51 and pPR52.P.44 indicated that the *crtR* gene is entirely contained in a 5.9 kb *Sall* DNA fragment. This result was corroborated by Southern blot hybridization (data not shown). The recombinant plasmid pCPRB.6, which contains an insert of 5,896 bp that harbors the *crtR* gene, was isolated from a partial *Sall* genomic library. The complete *crtR* gene sequence [GenBank:EU884133] is displayed in Figure 1.




Figure 1
Graphical representation and sequence of the *crtR* gene from *X. dendrorhous*. The three *crtR* exons are highlighted in red and white letters. The restriction enzymes used in this work and their recognition sites are in parentheses. Underlined and bold black letters represent the donor and acceptor splice sites. The sequence is numbered in accordance with its graphical representation. The complete sequence of the DNA fragment that contains the *crtR* gene is in GenBank [GenBank:EU884133]. pCPR1.3 corresponds to the vector pBluescript SK- bearing the DNA fragment from site *Bam*HI 4529 to *Bam*HI 9895. *crtR* deletions were constructed from pCPR1.3.

Localization of the *crtR* gene in the yeast electrophoretic karyotype

To study the *crtR* gene's genomic organization, its localization in the electrophoretic karyotype of the UCD 67–385 strain was performed by Southern blot hybridization. The results showed that the *crtR* gene is in the first two chromosomal bands (Figure 2), where each band corresponds to a triplet [21]. As it was recently demonstrated that this strain is diploid [30], it is possible to conclude that the *crtR* gene is in at least two chromosomes and could be homologous polymorphic. In different *X. dendrorhous* strains, a high chromosomal polymorphism has been revealed [31], so it is possible that these homologous chromosomes have experienced genetic rearrangement, thus explaining their different sizes.

Sequence analysis of *crtR* gene

The *crtR* gene consists of three exons of 222, 317 and 1702 bp and two introns of 240 and 138 bp. This gene encodes a 746 amino acid protein (Figure 3) with a predicted molecular weight of 81.9 kDa. CrtR has 22 hydrophobic amino acids in its amino terminal region, a typical trans-membrane segment of known CPRs that anchors the protein to the endoplasmic reticulum membrane in the cell [32]. CrtR also has recognizable CPR conserved domains such as a FMN binding domain, a FAD binding domain between amino acids 275 and 512, and a NAD(P)H binding domain between amino acids 582 and 709. The flavodoxin signature, which is only found in certain bacteria and algae, was identified between amino acids 80 and

222. However, the FMN binding domain of CPRs is homologous to bacterial flavodoxins [33]. Two putative P450-binding regions (P450-1 and P450-2) have been suggested [32], and one (P450-2) was studied by site-directed mutagenesis in rat CPR where the importance of acidic amino acid residues in this region was demonstrated [34]. The corresponding regions in CrtR were characterized by the presence of acidic amino acids (aspartic acid at positions: 112, 117, 122, 127, 210, 211, 218; glutamic acid at positions: 114, 115, 119, 126, 216, 217). The aspartic acid residues at position 210, 211 and 218 are extremely conserved in all known CPRs [32]. The P450 binding regions are located in the FMN binding domain. This is consistent with the electron flow from NAD(P)H to FAD to FMN and finally to the cytochrome P450 monooxygenase enzyme.

A phylogenetic comparison was performed using available fungal CPR protein sequences in the database by the Neighbor Joining method after 1000 replications of bootstrap analysis (Figure 4). The phylogenetic tree shows the evolutionary relationship of CrtR and allows the differentiation of three fungal clusters: Basidiomycota, Ascomycota and Zygomycota. CrtR from *X. dendrorhous* is most closely related to the basidiomycetes *C. versicolor* and to *P. chrysosporium*, with sequence identities of 60.2% and 58.3%, respectively.

Expression of *crtR* gene

The relationship between the *crtS* and *crtR* gene expression along the growth curve at the mRNA level was studied in the wild type UCD 67–385 *X. dendrorhous* strain, grown in MM_v supplemented with glucose as a fermentable carbon source. It was observed that *crtS* mRNA level reaches a maximum at 66 h of cultivation, equivalent to approximately half of the exponential growth phase. However, the levels of *crtR* mRNA remained constant along the growth curve (Figure 5). Despite the fact that both proteins are involved in the same stage of astaxanthin biosynthesis, there is no relationship, at the mRNA level, in their gene expression. It is possible to conclude that the two genes are not regulated in the same way at the transcriptional level.

According to this last result, in other work done in our laboratory, a dramatic decrease of transcripts levels from the *crtYB*, *crtI* and *crtS* genes (carotenogenic genes) was observed during the stationary phase [35]. This was not the case for the *idi* and *crtE* genes, where the level of the *idi* transcript showed a slight decline and the *crtE* transcript remained constant during the stationary phase. This suggests the presence of higher levels of the enzymes IPP isomerase and GGPP synthase at this stage. This situation would allow for the flow of the isoprenoid precursors towards carotenoid biosynthesis, as well as other meta-




Figure 2
***X. dendrorhous* electrophoretic karyotype.** (A) CHEF of *X. dendrorhous* UCD 67–385 strain. (B) Hybridization with *crtR* probe.





Figure 3
Deduced amino acid sequence from *X. dendrorhous* CrtR. (A) Graphical representation of CPR conserved domains in *X. dendrorhous* CrtR. (B) Deduced CrtR protein sequence. Conserved domains are shown in white text. TM: in italics and lower case. Highlighted in yellow: FMN BD, green: FAD BD and dark green: NAD(P)H BD. The two acidic regions involved in the binding to cytochrome P450 are also shown. TM: transmembrane region, BD: binding domain.

bolic pathways such as ergosterol biosynthesis [35]. Similarly, the absence of a relationship between the pattern of *crtR* gene expression and the rest of the carotenogenic genes suggests the existence of a different regulatory mechanism for *crtR* gene expression due to its involvement in other metabolic pathways of the yeast.

An example of *cpr* gene regulation has been described for *Aspergillus niger*. In this case, it was observed that the regulation of *cpr* gene expression was particularly complex in the P450-CPR system involved in benzoate hydroxylation [36]. In addition to benzoate induction at the transcriptional level, there were other regulatory mechanisms such as differential promoter use and post-translational regulatory mechanisms [36]. The decoupling of a P450-CPR system would imply the release of electrons to the cytoplasm that could generate active oxygen molecules, so a strict regulation of CPR is of vital importance for the cell. Similarly, it is not surprising that the *crtS* and *crtR* genes are regulated differently because, as mentioned earlier, several genes for different cytochrome P450 enzymes can exist in an organism, but generally there is a single gene for a unique cytochrome P450 reductase. Therefore, complex regulation for CrtR protein expression is necessary to adjust the levels of cytochrome P450 reductase activity in a cell with different levels of diverse cytochrome P450 proteins.

Functional study of the *crtR* gene

The participation of the *crtR* gene product in the formation of astaxanthin from β -carotene was studied in a bacterial heterologous system. For this, the DS1B *E. coli* strain (Table 1) was transformed with a pET101/D-TOPO expression vector carrying the cDNA of *crtS* and *crtR* genes from *X. dendrorhous*, and the carotenoid production was analyzed by RP-HPLC. This system was not successful, as it was not possible to produce any xanthophylls in the bacteria. A similar situation was observed in another study, where only canthaxanthin (β -carotene-4, 4'-dione, which is an intermediate in the conversion of β -carotene into astaxanthin) in very small quantities was produced when the *S. cerevisiae* *cpr* gene was used [12]. These authors concluded that the *X. dendrorhous* astaxanthin synthase has very low activity in *E. coli* [12]. It should be noted that the P450 systems are associated with membranes; therefore, *E. coli* may not provide a proper environment for its functionality. Recently, it was reported that several attempts to produce carotenoids in *E. coli* using the *X. dendrorhous* carotenogenic genes resulted in poor enzyme expression and carotenoid production [37]. Hence, a eukaryotic system would be more appropriate to attempt the expression of *X. dendrorhous* carotenogenic genes.

**Figure 4**

Phylogenetic tree for cytochrome P450 reductases. Distance analysis and tree construction was done using the Neighbor Joining method. Numbers at nodes indicate the percent support for specific nodes after 1000 replications of bootstrap analysis. The scale bar indicates 10% change estimated sequence difference. The percentage of sequence identity with *X. dendrorhous* CrtR is in parenthesis. Accession numbers: *Neurospora crassa*: [GenBank:XP_964443]; *Gibberella fujikuroi*: [GenBank:AJ576025]; *Cochliobolus lunatus*: [GenBank:ABW86977]; *Coccidioides immitis*: [GenBank:XP_001246449]; *Saccharomyces cerevisiae*: [GenBank:D13788]; *Candida tropicalis*: [GenBank:M35199]; *Rhodotorula minuta*: [GenBank:AB055119]; *Xanthophyllomyces dendrorhous*: [GenBank:EU884134] and [GenBank:EU884133]; *Coriolus versicolor*: [GenBank:AB065368]; *Phanerochaete chrysosporium*: [GenBank:AF193060]; *Cunninghamella elegans*: [GenBank:AF195659]; *Rhizopus stolonifer*: [GenBank:AF290425]; *Arabidopsis thaliana*: [GenBank:A75959].

As the *E. coli* expression system was not successful, in order to demonstrate the importance of the *crtR* gene in the astaxanthin biosynthetic pathway, *X. dendrorhous crtR* deletion mutant strains were generated. The wild type strains UCD 67-385 and CBS-6938 were transformed with the linearized plasmid pBsiWIAcrtR::hph and circular plasmid pNdeIAcrtR::hph, respectively. Through homologous recombination events, the wild type *crtR* gene in the wild type strains was exchanged by the DNA fragment containing the deletion and resistance marker (Figure 6).

Through this experiment, a yellow (CBSTr, which derives from CBS-6938) and a pale red-orange (T13, which derives from UCD 67-385) hygromycin B resistant trans-

formant strains were obtained. The color phenotype of these transformants is an indicator of an alteration in astaxanthin biosynthesis because astaxanthin has a strong red-orange color. The strain UCD 67-385 is diploid [30]; therefore, we suggest that T13 is heterozygote for the *crtR* gene, with a mutant and a wild type allele. Moreover, by a gene-dose effect, T13 produces less astaxanthin and accumulates more β -carotene than its parental wild type strain. On the other hand, the ploidy level of CBS-6938 is unknown. However, based on random mutagenesis results with physical and chemical mutagens performed in our laboratory (data not shown) and transformation with carotenogenic genes [38], it was concluded that this strain could be haploid. The mutagenesis results suggest




Figure 5

Relative expression of *crtR* and *crtS* genes along the *X. dendrorhous* growth curve. Wild type UCD 67–385 strain was grown at 22°C in a fermentor with MM_V + 2% glucose medium. (Δ): *crtS* relative expression, (□): *crtR* relative expression and (▲): n° cells per ml.

that, in the case of CBSTr, the only *crtR* gene copy is mutated and, therefore, it is not capable of producing astaxanthin and accumulates β-carotene.

In order to confirm the T13 (*crtR*^{ΔBsiWI::hph}/*crtR*⁺) and CBSTr (*crtR*^{ΔNdel::hph}) transformant genotype modifications, PCR reactions were performed using specific primers for the *crtR* and *hph* genes (Table 2), and genomic DNA samples from wild type and transformant strains were used as templates. The amplicon sequences validated the representation shown in Figure 6. The results indicated that the CBS-6938 strain has one *crtR* gene copy and its deletion is not lethal. In the case of T13, it was shown that it is heterozygote for the *crtR* gene, having a wild type and a mutant allele, which supports the gene-dose effect hypothesis in the production of astaxanthin. These results were also corroborated by Southern blot hybridization (data not shown).

Several attempts were performed to obtain a β-carotene accumulating transformant derived from UCD 67–385. T13 was retransformed with a DNA fragment harboring a *crtR* gene deletion and a G418 resistance cassette marker. No G418 resistant transformants were obtained. In order to produce a homozygous *crtR* mutation, T13 was subjected to the Double Recombinant Method [6], but it was not possible to obtain a double mutant strain for the *crtR* gene.

The *cpr* gene deletion in *S. cerevisiae* was not lethal, suggesting the existence of an alternative electron donor such as a cytochrome b5 (*cytb5*) [39]. When a *cytb5* deletion in the wild type was done, no new phenotype was generated. However, it was lethal when *cytb5* and *cpr* were simultaneously disrupted. This result demonstrated that, in mutants with single disruptions of *cpr* or *cytb5*, both enzymes can complement each other [39]. Likewise, the disruption of the *cpr* gene from the fungus *Gibberella fujikuroi* was not




Figure 6

Graphical representation of transformation events. Wild type strains UCD 67–385 and CBS-6938 were transformed with the linearized plasmid *pBsiWIΔcrtR::hph* and circular plasmid *pNdelΔcrtR::hph*, respectively. I, II, III: represent *crtR* exons I, 2 and 3; P: EF-1α promoter; Term: GPDH transcription terminator; *hph*: hygromycin B resistance gene; pBS: pBluescript; *BsiWI*** and *Ndel***::: modified restriction sites after hygromycin B cassette insertion and arrows represent primer binding sites.

Table 4: Carotenoid composition of wild-type and transformant *X. dendrorhous* strains in ppm

| Carotenoid | UCD 67–385 wt-strain | T13 (<i>crtR</i> ^{Δ<i>BsiWI</i>::<i>hph</i>} / <i>crtR</i> ⁺) | CBS-6938 wt-strain | CBSTr (<i>crtR</i> ^{Δ<i>Ndel</i>::<i>hph</i>}) |
|-------------------------|----------------------|---|--------------------|---|
| astaxanthin | 102.0 ± (4.8) | 24.0 ± (3.6) | 152.0 ± (8.5) | ND |
| phoenicoxanthin | 7.0 ± (0.4) | 8.0 ± (0.4) | 8.0 ± (0.5) | ND |
| canthaxanthin | 1.0 ± (0.1) | 3.00 ± (0.06) | 1.0 ± (0.1) | ND |
| hydroxy-keto-γ-carotene | 6.0 ± (0.2) | 1.00 ± (0.06) | ND | ND |
| HO-keto-torulene | ND | ND | 6.0 ± (1.1) | ND |
| keto-γ-carotene | 9.0 ± (0.7) | 11.0 ± (0.7) | 8.0 ± (1.2) | 1.0 ± (0.3) |
| HO-echinenone | ND | ND | 2.0 ± (0.4) | 1.0 ± (0.4) |
| echinenone | 5.0 ± (0.6) | 16.0 ± (0.5) | 5.0 ± (0.7) | 8.0 ± (1.2) |
| lycopene | ND | 1.0 ± (0.01) | ND | ND |
| neurosporene | ND | 1.00 ± (0.02) | ND | ND |
| γ-carotene | 1.0 ± (0.1) | 2.00 ± (0.05) | 1.0 ± (0.1) | 3.0 ± (0.2) |
| β-carotene | 9.0 ± (1.7) | 50.0 ± (3.7) | 9.0 ± (1.5) | 114.5 ± (6.6) |
| phytoene | 0.4 ± (0.3) | 4.2 ± (1.9) | 3.3 ± (1.2) | 1.0 ± (0.4) |
| NI (25 min)* | ND | 1.0 ± (0.1) | ND | 1.0 ± (0.1) |
| NI (34 min)* | ND | 2.00 ± (0.04) | ND | 4.0 ± (0.9) |
| Total carotenoids | 139.0 ± (3.2) | 122.8 ± (8.0) | 195.3 ± (8.0) | 131.9 ± (7.5) |

ND: Not detected, NI: Not identified carotenoid. Table values are the average results of three independent experiments, SD in parenthesis. (*)*: NI retention time.




Figure 7
RP-HPLC analysis of carotenoids from wild type (UCD 67–385, CBS-6983) and transformant (T13 ($crtR^{\Delta BsiWI::hph}/crtR^+$), CBSTr ($crtR^{\Delta Ndel::hph}$)) strains. Numbers indicate different carotenoids as follows: 1: astaxanthin; 2: phoeni-coxanthin; 3: canthaxanthin; 4: hydroxy-keto- γ -carotene; 5: keto- γ -carotene; 6: hydroxy-echinenone; 7: echinenone; 8: lycopene; 9: neurosporene; 10: γ -carotene and 11: β -carotene. In the upper right of the chromatogram, colony phenotype is shown.

lethal, proving the existence of another electron donor [16]. In light of these results and evidence for the existence of a high level of polymorphisms in different *X. dendrorhous* strains [21,30,31], we suggest that both wild type strains may have a different genetic background. Moreover, the results obtained in this work suggest that, in the CBS-6938 wild type strain, an alternative electron donor such as a cytochrome b5 may exist because the *crtR* gene disruption was not lethal. On the other hand, this is not the case for the UCD 67–385 strain due to the impossibility to obtain a homozygous mutant for the *crtR* gene. However, the possible alternative electron donor in the CBS-6938 strain cannot support astaxanthin synthesis, as the *crtR* gene mutant (CBSTr) accumulates β -carotene.

To check the carotenoid composition, total pigments were extracted from both wild type and transformant strains, and the composition was analyzed by RP-HPLC (Figure 7 and Table 4). The RP-HPLC analysis showed that the carotenoid composition is completely different between transformants and their parental strain. CBSTr accumulates β -carotene and is unable to synthesize astaxanthin. T13 has a reduced astaxanthin production at the expense of increased β -carotene production, which is the astaxanthin precursor.

Conclusion

A new *crtR* gene has been isolated and characterized from *X. dendrorhous*, which plays an auxiliary role to the *crtS* gene and is also essential for astaxanthin biosynthesis. This gene consists of three exons, and is localized in at least two chromosomes of the UCD 67–385 *X. dendrorhous* wild type strain. *crtR* encodes a 746 amino acid protein with characteristic domains of cytochrome P450 reductases (FMN, FAD and NAD(P)H binding domains) and a small amino terminal transmembrane region. Although both *crtS* and *crtR* proteins participate in the same stage of yeast carotenogenesis, the mRNA level pattern from these genes along the growth curve is different. Thus, both genes are regulated in a different manner at the transcriptional level.

Authors' contributions

JA participated in the design of this study and carried out the *crtR* gene isolation, sequence analysis, *crtR* gene expression study, *X. dendrorhous* transformation, pigment extraction, RP-HPLC and drafted the manuscript. SB constructed the *X. dendrorhous* YIp5 genomic library. MC participated in the isolation of the *crtS* gene genomic and cDNA versions. CL participated in the UCD 67–385 strain fermentation and RT-PCR analysis. AM constructed the *X. dendrorhous* cDNA library. MN constructed the hygromycin B resistance cassette. DS participated in DNA sequencing. MB participated in the study design. VC conceived the

study and participated in its design and coordination. All authors read and approved the final manuscript.

Acknowledgements

This work was supported by U. de Chile ENL 08/06 and: MECESUP UCH 0106 by graduate scholarships to JA and MN, Fundación María Ghilardi by a graduate scholarship to CL, CONICYT by a graduate scholarship to AM.

References

1. Takaichi S, Sandmann G, Schnurr G, Satomi Y, Suzuki A, Misawa N: **The carotenoid 7,8-dihydro- ψ end group can be cyclized by the lycopene cyclases from the bacterium *Erwinia uredovora* and the higher plant *Capsicum annuum*.** *Eur J Biochem* 1996, **241**:291-296.
2. Johnson E, Conklin D, Lewis M: **The yeast *Phaffia rhodozyma* as a dietary pigment source for salmonids and crustaceans.** *J Fish Res Board Can* 1997, **34**:2417-2421.
3. Andrewes AG, Phaff HJ, Starr MP: **Carotenoids of *Phaffia rhodozyma*. A red pigmented fermenting yeast.** *Phytochem* 1976, **15**:1003-1007.
4. Fraser P, Miura Y, Misawa N: **In vitro characterization of astaxanthin biosynthetic enzymes.** *J Biol Chem* 1997, **272**:6128-6135.
5. Kajiwara S, Fraser PD, Kondo K, Misawa N: **Expression of an exogenous isopentenyl diphosphate isomerase gene enhances isoprenoid biosynthesis in *Escherichia coli*.** *Biochem J* 1997, **324**:421-426.
6. Niklitschek M, Alcaíno J, Barahona S, Sepúlveda D, Carmona M, Wozniak A, Marcoleta A, Lodato P, Baeza M, Cifuentes V: **Genomic organization of the structural gene controlling the astaxanthin biosynthesis pathway of *Xanthophyllomyces dendrorhous*.** *Biol Res* 2008, **41**:93-108.
7. Verdoes JC, Krubasik P, Sandmann G, van Ooyen AJJ: **Isolation and functional characterisation of a novel type of carotenoid biosynthetic gene from *Xanthophyllomyces dendrorhous*.** *Mol Gen Genet* 1999, **262**:453-461.
8. Verdoes JC, Misawa N, van Ooyen AJJ: **Cloning and characterization of the astaxanthin biosynthetic gene encoding phytoene desaturase of *Xanthophyllomyces dendrorhous*.** *Biotechnol Bioeng* 1999, **63**:750-755.
9. Cunningham FX, Gantt E: **Genes and enzymes of carotenoid biosynthesis in plants.** *Annu Rev Plant Physiol Plant Mol Biol* 1998, **49**:557-583.
10. Linden H: **Carotenoid hydroxylase from *Haematococcus pluvialis*: cDNA sequence, regulation and functional complementation.** *Biochim Biophys Acta* 1999, **1446**:203-212.
11. Hoshino T, Ojima K, Setoguchi Y: **Astaxanthin synthase.** *Hoffman-LaRoche, European patent no. 1035206-A3* 2000.
12. Ojima K, Breitenbach J, Visser H, Setoguchi Y, Tabata K, Hoshino T, Berg J van den, Sandmann G: **Cloning of the astaxanthin synthase gene from *Xanthophyllomyces dendrorhous* (*Phaffia rhodozyma*) and its assignment as a beta-carotene 3-hydroxylase/4-ketolase.** *Mol Genet Genomics* 2006, **275**:148-158.
13. Alvarez V, Rodriguez-Saiz M, de la Fuente JL, Gudina EJ, Godio RP, Martín JF, Barredo JL: **The *crtS* gene of *Xanthophyllomyces dendrorhous* encodes a novel cytochrome-P450 hydroxylase involved in the conversion of beta-carotene into astaxanthin and other xanthophylls.** *Fungal Genet Biol* 2006, **43**:261-272.
14. Martín JF, Gudiña E, Barredo JL: **Conversion of beta-carotene into astaxanthin: Two separate enzymes or a bifunctional hydroxylase-ketolase protein?** *Microb Cell Fact* 2008, **20**:7-13.
15. van den Brink HM, van Gorcom RF, van den Hondel CA, Punt PJ: **Cytochrome P450 enzyme systems in fungi.** *Fungal Genet Biol* 1998, **23**:1-17.
16. Malonek S, Rojas MC, Hedden P, Gaskin P, Tudzynski B: **The NADPH-cytochrome P450 reductase gene from *Gibberella fujikuroi* is essential for gibberellin biosynthesis.** *J Biol Chem* 2004, **279**:25075-25084.
17. Retamales P, Hermosilla G, León R, Martínez C, Jiménez A, Cifuentes V: **Development of the sexual reproductive cycle of *Xanthophyllomyces dendrorhous*.** *J Microbiol Methods* 2002, **48**:87-93.
18. Sambrook J, Russell DW: *Molecular cloning. A laboratory manual* 3rd edition. Cold Spring Harbor NY: Cold Spring Harbor Laboratory Press; 2001.

19. Guengerich FP, Martin MV, Guo Z, Chun YJ: **Purification of functional recombinant P450s from bacteria.** *Methods Enzymol* 1996, **272**:35-44.
20. Boyle JS, Lew AM: **An inexpensive alternative to glassmilk for DNA purification.** *TIG* 1995, **11**:8.
21. Cifuentes V, Hermosilla G, Martínez C, León R, Pincheira G, Jiménez A: **Genetics and electrophoretic karyotyping of wild type and astaxantin mutants strains from *Phaffia rhodozyma*.** *Antonie Van Leeuwenhoek* 1997, **72**:111-117.
22. Chomczynski P, Sacchi YN: **Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction.** *Anal Biochem* 1987, **162**:156-159.
23. Lodato P, Alcaíno J, Barahona S, Retamales P, Jiménez A, Cifuentes V: **Study of the expression of carotenoid biosynthesis gene in wild type and deregulated strains of *Xanthophyllomyces dendrorhous* (Ex *Phaffia rhodozyma*).** *Biol Res* 2004, **37**:83-93.
24. Wery J, Dalderup MJ, Ter Linde J, Boekhout T, van Ooyen AJJ: **Structural and phylogenetic analysis of the actin gene from the yeast *Phaffia rhodozyma*.** *Yeast* 1996, **12**:641-651.
25. Ngiam C, Jeenes DJ, Punt PJ, Hondel CA van den, Archer DB: **Characterization of a foldase, protein disulfide isomerase A, in the protein secretory pathway of *Aspergillus niger*.** *Appl Environ Microbiol* 2000, **66**:775-782.
26. Adrio J, Veiga M: **Transformation of the astaxanthin-producing yeast *Phaffia rhodozyma*.** *Biotech Techniques* 1995, **9**:509-512.
27. Kim IG, Nam SK, Sohn JH, Rhee SK, An GH, Lee SH, Choi ES: **Cloning of the ribosomal protein L41 gene of *Phaffia rhodozyma* and its use as a drug resistance marker for transformation.** *Appl Environ Microbiol* 1998, **64**:1947-1949.
28. Reyes E, Barahona S, Fischman O, Niklitschek M, Baeza M, Cifuentes V: **Genetic polymorphism of clinical and environmental strains of *Pichia anomala*.** *Biol Res* 2004, **37**:747-57.
29. An GH, Schuman DB, Johnson EA: **Isolation of *Phaffia rhodozyma* mutants with increased astaxanthin content.** *Appl Environ Microbiol* 1989, **55**:116-124.
30. Hermosilla G, Martínez C, Retamales P, León R, Cifuentes V: **Genetic determination of ploidy level in *Xanthophyllomyces dendrorhous*.** *Antonie van Leeuwenhoek* 2003, **84**:279-287.
31. Nagy A, Garamszegib N, Vágvölgyib C, Ferenczyb L: **Electrophoretic karyotypes of *Phaffia rhodozyma* strains.** *FEMS Microbiol Lett* 1994, **123**:315-318.
32. Yadav JS, Loper JC: **Cytochrome P450 oxidoreductase gene and its differentially terminated cDNAs from the white rot fungus *Phanerochaete chrysosporium*.** *Curr Genet* 2000, **37**:65-73.
33. Porter TD, Kasper CB: **NADPH-cytochrome P-450 oxidoreductase: flavin mononucleotide and flavin adenine dinucleotide domains evolved from different flavoproteins.** *Biochem* 1986, **25**:1682-1687.
34. Shen AL, Kasper CB: **Role of acidic residues in the interaction of NADPH-cytochrome P450 oxidoreductase with cytochrome P450 and cytochrome c.** *J Biol Chem* 1995, **270**:27475-27480.
35. Lodato P, Alcaíno J, Barahona S, Niklitschek M, Carmona M, Wozniak A, Baeza M, Jiménez A, Cifuentes V: **Expression of the carotenoid biosynthesis gene in *Xanthophyllomyces dendrorhous*.** *Biol Res* 2007, **40**:73-84.
36. van den Brink JM, Punt PJ, van Gorcom RF, van den Hondel CA: **Regulation of expression of the *Aspergillus niger* benzoate parahydroxylase cytochrome P450 system.** *Mol Gen Genet* 2000, **263**:601-609.
37. Verwaal R, Wang J, Meijnen JP, Visser H, Sandmann G, Berg JA van den, van Ooyen AJJ: **High-level production of beta-carotene in *Saccharomyces cerevisiae* by successive transformation with carotenogenic genes from *Xanthophyllomyces dendrorhous*.** *Appl Environ Microbiol* 2007, **73**:4342-4350.
38. Verdoes JC, Sandmann G, Visser H, Diaz M, van Mossel M, van Ooyen AJJ: **Metabolic Engineering of the Carotenoid Biosynthetic Pathway in the Yeast *Xanthophyllomyces dendrorhous* (*Phaffia rhodozyma*).** *Appl Environ Microbiol* 2003, **69**(7):3728-38.
39. Truan G, Epinat JC, Rougeulle C, Cullin C, Pompon D: **Cloning and characterization of a yeast cytochrome b5-encoding gene which suppresses ketoconazole hypersensitivity in a NADPH-P-450 reductase-deficient strain.** *Gene* 1994, **142**:123-127.
40. Misawa N, Nakagawa M, Kobayashi K, Yamano S, Izawa Y, Nakamura K, Harashima K: **Elucidation of the *Erwinia uredovora* caroten-**

oid biosynthetic pathway by functional analysis of gene products expressed in *Escherichia coli*. *J Bacteriol* 1990, **172**:6704-6712.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

