




Review

Basic Substances, a Sustainable Tool to Complement and Eventually Replace Synthetic Pesticides in the Management of Pre and Postharvest Diseases: Reviewed Instructions for Users

Gianfranco Romanazzi ^{1,*}, Yann Orçonneau ², Marwa Moumni ¹, Yann Davillerd ²
and Patrice André Marchand ²

¹ Department of Agricultural, Food and Environmental Sciences, Marche Polytechnic University, Via Breccie Bianche, 60131 Ancona, Italy; m.moumni@staff.univpm.it

² Institut Technique de l'Agriculture et de l'Alimentation Biologiques (ITAB), 149 rue de Bercy, 75012 Paris, France; y.orco17@gmail.com (Y.O.); yann.davillerd@itab.asso.fr (Y.D.); patrice.marchand@itab.asso.fr (P.A.M.)

* Correspondence: g.romanazzi@univpm.it; Tel.: +39-071-220-4336

Abstract: Synthetic pesticides are widely used to protect crops from pathogens and pests, especially for fruits and vegetables, and this may lead to the presence of residues on fresh produce. Improving the sustainability of agriculture and, at the same time, reducing the adverse effects of synthetic pesticides on human health requires effective alternatives that improve the productivity while maintaining the food quality and safety. Moreover, retailers increasingly request fresh produce with the amounts of pesticides largely below the official maximum residue levels. Basic substances are relatively novel compounds that can be used in plant protection without neurotoxic or immune-toxic effects and are still poorly known by phytosanitary consultants (plant doctors), researchers, growers, consumers, and decision makers. The focus of this review is to provide updated information about 24 basic substances currently approved in the EU and to summarize in a single document their properties and instructions for users. Most of these substances have a fungicidal activity (calcium hydroxide, chitosan, chitosan hydrochloride, *Equisetum arvense* L., hydrogen peroxide, lecithins, cow milk, mustard seed powder, *Salix* spp., sunflower oil, sodium chloride, sodium hydrogen carbonate, *Urtica* spp., vinegar, and whey). Considering the increasing requests from consumers of fruits and vegetables for high quality with no or a reduced amount of pesticide residues, basic substances can complement and, at times, replace the application of synthetic pesticides with benefits for users and for consumers. Large-scale trials are important to design the best dosage and strategies for the application of basic substances against pathogens and pests in different growing environments and contexts.

Keywords: European Union; fungicide residues; plant protection; regulation EU 1107/2009



Citation: Romanazzi, G.; Orçonneau, Y.; Moumni, M.; Davillerd, Y.; Marchand, P.A. Basic Substances, a Sustainable Tool to Complement and Eventually Replace Synthetic Pesticides in the Management of Pre and Postharvest Diseases: Reviewed Instructions for Users. *Molecules* **2022**, *27*, 3484. <https://doi.org/10.3390/molecules27113484>

Academic Editor: James Barker

Received: 15 April 2022

Accepted: 24 May 2022

Published: 28 May 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

The world population continues to grow and will reach 9.7 billion by 2050 [1]. For this, increasing food production is the primary objective of all countries. According to the latest estimates of the Food and Agriculture Organization of the United Nations [2], up to 40% of food crops worldwide are lost every year due to pests and plant diseases. Crop losses caused by plant disease alone cost the global economy \$220 billion annually [3]. Crop protection is essential to reduce yield losses, improve food quality, and increase grower profitability. The application of plant protection products (PPPs) is the main way to protect crops against pathogens, pests, and weeds [4]. However, human, animal, and environmental risks associated with the use of chemical PPPs are a growing concern. All these concerns have encouraged the onset of research to develop alternative approaches to control plant diseases [5]. Reducing the use of pesticides being a major challenge in

developed countries, European Union Member States are required to implement National Action Plans that set quantitative objectives, timetables, and indicators related to reducing the impact of pesticide use (Directive 2009/128/CE) [6,7]. The use of basic substances is approved in the European Union under Article 23 of EC Regulation No 1107/2009 and which are listed in Part C of the Annex of the Regulation (EC) No 540/2011 [8]. In the EU, Integrated Pest Management (IPM) has been mandatory since January 2014, and among the rules of the IPM is the reduction of the application of synthetic pesticides whenever possible [9]. For sustainable and qualitative food production, respectful of the need to produce in sufficient quantities, biocontrol has grown tremendously through the last few years [10]. The PPP EU Regulation (EC) 1107/2009 was established to ensure a level of protection of humans, animals, and the environment and, at the same time, to unify for the entire EU the rules on the placing on the market of plant protection products [11,12]. Basic substances are sources of interest for research as alternative to synthetic pesticides, since they are used in human medicine or as a food ingredient, so they have no residue concerns and then no maximum residue limit (MRL) and, usually, no preharvest interval [13,14]. The lack of MRL contributes to a better prevention of contamination in plant protection, a better control of the residues and a reduction of analytical problems, of decommissioning, and of market withdrawal [14]. Another benefit of basic substances, and perhaps the most important, is their very low ecological impact. Basic substances are products that are used as 'foodstuffs', as defined in Article 2 of Regulation (EC) 178/2002 [15] cosmetic, and does not have an inherent capacity to cause endocrine-disrupting, neurotoxic or immunotoxic effects, but they are also plant protection means and not placed on the market as a plant protection product. Article 28 of Regulation (EC) No. 1107/2009 set the absence of marketing authorizations and usages allowance for basic substances. Regulation (EC) No. 1107/2009 introduced the new category of 'basic substances', which are defined by recital 18 as 'certain substances which are not predominantly used as plant protection products may be of value for plant protection, but the economic interest of applying for approval may be limited. Therefore, specific provisions should ensure that such substances, as far as their risks are acceptable, may also be approved for plant protection use'. The properties of basic substances are described in Article 23 of the EU Regulation (EC) No 1107/2009 [11]. In 2021, the Euphresco project 'BasicS' contributed to demonstrate the effectiveness toward pests and pathogens of basic substances, with potential benefits for the farmers, the consumer, and the environment [16,17]. The basic substances have a positive impact on crop health when applied preventively. Certain basic substances, such as chitosan, stimulate the defense system of crops against several classes of pathogens, including fungi, viruses, bacteria, and phytoplasma [18]. According to the EU pesticides database, 24 basic substances were approved for use, 7 were withdrawn, 18 applications were not approved and 8 are still pending [19,20]. This review includes currently approved basic substances that have a protective potential and are a valuable addition to the range of measures and protection methods intended for use. Detailed information about basic substances and updates on new available compounds can be found at the page <https://ec.europa.eu/food/plant/pesticides/eu-pesticides-database/active-substances> (accessed on 23 May 2022). The standard-folder for approval of a basic substance, called 'Basic Substance Application Template (BSAT)', is based on the structure of the European Union evaluation report of an active substance that can be used for plant protection purposes. BSAT refers to all areas of risk assessment in the regulation of phytopharmaceutical product uses and shall be considered as a structured model to build a file collating all available information and enabling to demonstrate that the evaluated substance meets the eligibility criteria of a basic substance (SANCO 10,363 rev.10, 2021). Therefore, nowadays, a full deposit under International Uniform Chemical Information Database (IUCLID) software is mandatory since March 2021. Basic substances are submitted individually (Annex I inclusion dossier) at the first stage; then, later, an automatic inclusion was adopted for food/foodstuff basic substance from plant or animal origin [21,22]. Recently, an automatic consideration procedure (without any Annex I inclusion dossier) by Expert Group for Technical advice on

Organic Production (EGTOP)/Directorate-General for the Agriculture and Rural Development (DGAgri) of positive ongoing basic substance approval (from Directorate-General Health and Food Safety—DGSanté to DGAgri) to generate an automatic EGTOP/DGAgri outcome for inclusion (or not). This provision bypasses the traditional route of substances in organic production in plant protection through dossiers submitted to Member States, but so far, no basic substance has been rejected by the Regulatory Committee of Organic Production (RCOP), and with the current procedure, are no longer studied than substances of mineral origin (or non-foods).

This review aimed to highlight the properties of approved basic substances, summarize, and provide this information for phytosanitary consultants, scientists, growers, stakeholders, companies, and consumers.

2. Results

Out of the 86 basic substance application submitted to the European Commission until now, less than one-third have been approved (24) (Tables 1 and 2), 19 have been refused, 6 have been withdrawn during their assessment (Table 3), 8 are currently being processed by the EC (Table 4 and Figure 1), and 2 already successfully submitted via IUCLID software (Ginger extract and *Capsicum frutescens*).

Currently, 24 basic substances are approved, of which 21 are also approved in organic production; for example, talc was validated in 2021 following EGTOP PPP VII and is being currently voted on at RCOP [23] and clayed charcoal was submitted. Recently, voted chitosan does not seem to be acceptable directly in organic production as the basic substance from its microorganism's origin, although in the context of food quality. Basic substances are approved by EU Regulations, so the application month, where reported in Table 1, is related to the Northern Hemisphere.




Figure 1. Total of the basic substance applications (BSA) and extensions presented by the results (%).

Table 1. Application of the basic substances approved.

Basic Substance	Approval Regulation and Applicant	Crops and/or Situation	Function in Plant Protection	Pests or Group of Pests Target	Application				Application Rates			Notes		
					Method	Growth Stage & Season	No. Min/Max	IBA ¹ (Days)	Min–Max	Water L/ha Min–Max	Total Rate		PHI ¹	
<i>Equisetum arvense</i> L.	Reg. (EU) No 462/2014 ITAB	Fruit trees Apple fruit (<i>Malus pumila</i> , <i>Malus domestica</i>) Peach-tree (<i>Prunus persica</i>)	Fungicide	Foliar fungi like scab disease (<i>Venturia inaequalis</i>), Powdery mildews (<i>Podosphaera leucotricha</i>) Peach leaf curl (<i>Taphrina deformans</i>)	Foliar application spraying	From green leaf tip (BBCH 53) to flowers fading (BBCH 67) Spring	2–6	7		500–1000	1000–2000 g/ha	Na ¹	Plant homogenate extracted with hot water and filtered to be used 24 h after preparation	
		Grapevine (<i>Vitis vinifera</i>)		Downy mildew (<i>Plasmopara viticola</i>), Powdery mildew (<i>Erysiphe necator</i>)		From 1st shoots (BBCH 10) to cluster tightening (BBCH 57) Spring to summer			200 g/hL	100–300	200–600 g/ha	Na		
		Cucumber (<i>Cucumis sativus</i>) roots		Powdery mildew (<i>Podosphaera fusca</i>) Root fungi like common root rot, seedling blight (<i>Pythium</i> spp.)	Root feeding application and foliar application spraying	From (9th leaf unfolded on main stem—BBCH 19) to 9 or more primary side shoots visible (BBCH 49)	2	3–4			300	600 g/ha		15
		Tomato (<i>Lycopersicon esculentum</i>)		Early blight (<i>Alternaria solani</i>), Septoria blight (<i>Septoria lycopersici</i>)	Foliar application spraying	First inflorescence visible (BBCH 51) to BBCH 59 summer		14						
		Strawberry (<i>Fragaria</i> × <i>Ananassa</i>) Raspberry (<i>Rubus idaeus</i>)		Gray mold (<i>Botrytis cinerea</i>), Powdery mildew (<i>Podosphaera aphanis</i>), red core (<i>Phytophthora fragariae</i>), other fungi like <i>Colletotrichum acutatum</i>	Foliar application spraying ²	Growth restart till end of fructification. Early spring till end of summer Stage BBCH 1 to BBCH 89	4–8	5–14	225 g/hL		300	675 g/ha		
		Potato (<i>Solanum tuberosum</i>)		Late blight (<i>Phytophthora infestans</i>), early blight (<i>Alternaria solani</i>), powdery mildew (<i>Erysiphe cichoracearum</i>)		Stage BBCH 1 until BBCH 9								Na
		Ornamental trees use of which <i>Prunus</i> spp. Roses <i>Rosa</i> spp.		Ornamental fungal diseases, rose black spot (<i>Marsonia</i> spp.), Rose rust (<i>Phragmidium mucronatum</i>), leaf curl diseases, monilioses, oidium and mildew	Included in mulch	Not relevant	1	Na	Na	Na	Na	9000 g/ha		

Table 1. Cont.

Basic Substance	Approval Regulation and Applicant	Crops and/or Situation	Function in Plant Protection	Pests or Group of Pests Target	Application				Application Rates			Notes				
					Method	Growth Stage & Season	No. Min/Max	IBA ¹ (Days)	Min–Max	Water L/ha Min–Max	Total Rate		PHI ¹			
Chitosan hydrochloride	Reg. (EU) 2021/1446 ChiPro	Fruits berries and small fruit	Elicitor, having a fungicide and bactericide effect via the stimulation of natural defence mechanisms	Plant elicitor, plant resistance against pathogenic fungi and bacteria	Low–Medium volume spraying	From 1 leaf development (main shoot) to 7 development of fruit	4–8	14	50–200 g/hL	100–800 g/ha		0				
		Vegetables							50–100 g/hL	100–400 g/ha						
		Cereals								200–400						
		Spices								100–400 g/ha						
		Crops for animal feed							Low volume spraying	Before sowing	1		Na	50–100 g/hL	Na	Na
		Cereals Seed treatment												Na	Na	
		Potatoes Seed treatment												Na	Na	
		Sugar beet Seed treatment							Low–Medium volume spraying	Germination (BBCH 00–01)	1–8		5–7	50–200 g/hL	Na	Na
		Ornamental bulbous plants												50–100 g/hL	200–800	100–800 g/ha
		Beet crops							Low–Medium volume spraying	Leaf development–senescence (BBCH 10–92)	50–200 g/hL		200–400			

Table 1. Cont.

Basic Substance	Approval Regulation and Applicant	Crops and/or Situation	Function in Plant Protection	Pests or Group of Pests Target	Application				Application Rates			Notes				
					Method	Growth Stage & Season	No. Min/Max	IBA ¹ (Days)	Min–Max	Water L/ha Min–Max	Total Rate		PHI ¹			
Sucrose	Reg. (EU) No 916/2014 ITAB IRBI	Apple trees / orchards (<i>Malus pumila</i> , <i>Malus domestica</i>)	Elicitor, having an insecticidal and fungicidal effect via the stimulation of natural defence mechanisms	Fruits borer like Codling moth (<i>Cydia pomonella</i>) ³	Foliar application spraying early in the morning before 9 AM (Solar time)	From spring BBCH stage 6 to summer BBCH stage 89	15	10 g/hL	200	60–100 g/ha	Na	Cold water solution prepared just before application				
		Sweet Maize (Sweet corn) (<i>Zea mays</i> L. convar. <i>saccharata</i> Koern)		Corn borer (<i>Ostrinia nubilalis</i> Hbn.) ³		From the BBCH stage 12 to 89							3–4			
		Maize (corn grain) (<i>Zea mays</i> subsp. <i>mays</i> (L.) and corn seed		Corn borer (<i>Ostrinia nubilalis</i> Hbn.) ³		From the BBCH stage 12 to 51							3–4			
		Grapevine (<i>Vitis vinifera</i>)		Vine leafhopper (<i>Scaphoideus titanus</i>) ³		From the BBCH stage 17 to 57							3			
		Grapevine (<i>Vitis vinifera</i>)		Downy mildew (<i>Plasmopara viticola</i>) ³		From 1st shoots to cluster tightening spring (BBCH 10–57)							up to 2			
Calcium hydroxide	Reg. (EU) 2015/762 IFOAM	Pome fruit	Fungicide	<i>Neonectria galligena</i>	Sprinkler application	Leaf drops end of October till end of December	2–7	5–14	500–1000 L/ha	104–208 L/ha ⁴ 1460 L/ha ⁵	5000– 10,000 L/ha	25–50 kg/ha 350 kg/ha ³	Na			
		Pome fruit and stone fruit		<i>Neonectria galligena</i> and other diseases						Brush application directly on pruning wounds and old cancers on stems ⁶	Winter to March	1–2		21	With products at 24% 63–104 L/ha ⁴ 728 L/ha ⁵ with products at 33.12% 45–76 L/ha ⁴ 532 L/h ⁵	15–25 kg/ha ⁴ 175 kg/ha ⁵
														With products at 24% 450 L/ha ³ 900 L/ha ⁴ with products at 33.12% 450 L/ha ⁴ 900 L/ha ⁵	No extra water ⁶	149.04 kg ⁴ 299.08 kg ⁵

Table 1. Cont.

Basic Substance	Approval Regulation and Applicant	Crops and/or Situation	Function in Plant Protection	Pests or Group of Pests Target	Application				Application Rates			Notes
					Method	Growth Stage & Season	No. Min/Max	IBA ¹ (Days)	Min–Max	Water L/ha Min–Max	Total Rate	
Vinegar	Reg. (EU) No 540/2011 Reg. (EU) 2015/1108 Reg. (EU) 2019/149 ITAB	Wheat seeds (<i>Triticum vulgare</i>), common wheat (<i>Triticum aestivum</i>), durum wheat (<i>Triticum durum</i>), spelt (<i>Triticum spelta</i>)	Fungicide, bactericide and herbicide	Common bunt (<i>Tilletia caries</i> , <i>Tilletia foetida</i>)	Seed treatment just before seeding	Autumn	1	Na	25–50 ⁷ per 100 kg of seed	Not applicable	24–100 ^{7,8}	Na
		Barley seeds (<i>Hordeum vulgare</i>)		Barley leaf stripe (<i>Pyrenophora graminea</i>)		Autumn to spring			Seeds are temporary soaked in the dilution then removed		Seeds are temporary soaked in the preparation then removed	
		Market vegetables Gardening like carrot (<i>Daucus carota</i>), tomato (<i>Solanum lycopersicum</i>), bell pepper (<i>Capsicum</i> spp.)		<i>Alternaria</i> spp.								
		Market vegetables gardening like tomato (<i>Solanum Lycopersicum</i>), bell pepper (<i>Capsicum</i> spp.), cabbage (<i>Brassica oleracea</i>)		<i>Clavibacter michiganensis</i> , <i>Clavibacter michiganensis</i> subsp. <i>michiganensis</i> , <i>Pseudomonas syringae</i> pv. <i>tomato</i> , <i>Xanthomonas campestris</i> pv. <i>vesicatoria</i> , <i>Botrytis aclada</i>								

Table 1. Cont.

Basic Substance	Approval Regulation and Applicant	Crops and/or Situation	Function in Plant Protection	Pests or Group of Pests Target	Application				Application Rates			Notes	
					Method	Growth Stage & Season	No. Min/Max	IBA ¹ (Days)	Min–Max	Water L/ha Min–Max	Total Rate		PHI ¹
	ITAB/ITEIPMAI	Medicinal aromatic and perfume crops		Weeds	Spray ¹⁰	Pre crop emergence	1	Na	10 kg/hL	100 L vinegar (no dilution)	10 kg/ha	>120	Phytotoxic to plant, may kill the young plants ¹¹
	Charbonneaux-Brabant	paths, borders, sidewalks and terraces		Weeds	Direct spray (spot application)	Vegetation Period of the weeds	1–2	7–21	6 kg/hL	100 L (diluted vinegar)	6–12 kg/ha	Na	Temp > 20 °C phytotoxic to plant, may kill the young plants ¹²
Salix spp. cortex	Reg. (EU) 2015/1107 ITAB	Fruit trees, Peach tree (<i>Prunus persica</i>)	Fungicide	Foliar fungi like <i>Taphrina deformans</i>	Foliar application spraying	From 1st shoots (BBCH 10) to cluster tightening (BBCH 57) spring	2–6	7	222.2 g/hL	500–1000 L/ha	1111.1–2222.2 g/ha	Na	Plant homogenate extracted with hot water (infusion), filtered and diluted by 3, to be used up to a maximum of 24 h after preparation. The product cannot be applied in case of hot temperature. It is used in case of rainy period
		Apple fruit (<i>Malus pumila</i> , <i>Malus domestica</i>)		Foliar fungi like scab disease (<i>Venturia inaequalis</i>), powdery mildew (<i>Podosphaera leucotricha</i>)		From green leaf tip (BBCH 53) to flowers fading (BBCH 67) spring							
		Grapevine (<i>Vitis vinifera</i>)		Downy mildew (<i>Plasmopara viticola</i>), Powdery mildew (<i>Erysiphe necator</i>)		From 1st shoots (BBCH 10) to cluster tightening (BBCH 57) spring to summer				100–300			

Table 1. Cont.

Basic Substance	Approval Regulation and Applicant	Crops and/or Situation	Function in Plant Protection	Pests or Group of Pests Target	Application				Application Rates		Notes			
					Method	Growth Stage & Season	No. Min/Max	IBA ¹ (Days)	Min–Max	Water L/ha Min–Max		Total Rate	PHI ¹	
Lecithins	Reg. (EU) No 540/2011 Reg. (EU) 2015/1116 ITAB DAE	Fruit trees Apple fruit (<i>Malus pumila</i>) Peach tree (<i>Prunus persica</i>)	Fungicide	Powdery mildew (<i>Podosphaera leucotricha</i>) Peach leaf curl (<i>Taphrina deformans</i>)	Spray application	BBCH 03 to BBCH 79	3–12	5	75 g/hL	500–1000	375–750 g/ha	5		
		Gooseberry Ribes uva-crispa		Powdery mildew (<i>Microsphaera grossulariae</i>)		BBCH 10 to BBCH 85	2–4		200 g/hL	1000–2000 g/ha				
		Market vegetables gardening like cucumber (<i>Cucumis sativus</i>)		Powdery mildew (<i>Podosphaera fusca</i>)			2–6							
		Lettuce (<i>Lactuca sativa</i>)		<i>Erysiphe cichoracearum</i>		BBCH 10 to BBCH 89	2		7					
		Mash (<i>Valerianella locusta</i>)		<i>Erysiphe polyphaga</i>			1		Na	150 g/hL	1000–1500		1500–2250 g/ha	
		Tomato (<i>Lycopersicon esculentum</i>)		Tomato late blight (<i>Phytophthora infestans</i>)			2 to 6		7					
		Endive (<i>Cichorium endivia</i> L.)		<i>Alternaria cichorii</i>										
		Ornamentals, especially roses		Powdery mildew and other fungal diseases										
		Grapevine (<i>Vitis vinifera</i>)		Downy mildew (<i>Plasmopara viticola</i>), Powdery mildew (<i>Erysiphe necator</i>)		BBCH 11 to BBCH 85				75 g/hL	100–300		75–225 g/ha	30
		Strawberry (<i>Fragaria × Ananassa</i>) Raspberrry (<i>Rubus idaeus</i>)		Powdery mildew and other fungal diseases, i.e., <i>Podosphaera aphanis</i> , Red core (<i>Phytophthora fragariae</i>)		Growth restart till end of fructification Early spring till end of Summer Stage BBCH 10 to BBCH 89 (2nd crop, other strawberries have reached them specific color)	3–12		5	200 g/hL	300–500		600–1000 g/ha	Na
Potato (<i>Solanum tuberosum</i>)	Late blight (<i>Phytophthora infestans</i>)	Stage BBCH 10 until BBCH 90	3–12			100–400	200–800 g/ha							
Carrot (<i>Daucus carota</i> subsp. <i>sativus</i>)	Powdery mildew (<i>Leveillula taurica</i>)	BBCH 19 to BBCH 90	4	14		1000	2000 g/ha							

Table 1. Cont.

Basic Substance	Approval Regulation and Applicant	Crops and/or Situation	Function in Plant Protection	Pests or Group of Pests Target	Application				Application Rates			Notes	
					Method	Growth Stage & Season	No. Min/Max	IBA ¹ (Days)	Min–Max	Water L/ha Min–Max	Total Rate		PHI ¹
Fructose	Reg. (EU) 2015/1392 ITAB IRBI	Apple fruit (<i>Malus pumila</i> , <i>Malus domestica</i>)	Elicitor, having an insecticidal and fungicidal effect via the stimulation of natural defence mechanisms	Fruits borer like Codling Moth (<i>Cydia pomonella</i>) ¹³	Foliar application spraying early in the morning before 9 AM (solar time)	From spring BBCH stage 6 to summer BBCH stage 65	5–7	21		600–1000	60–100 g/ha	Na	Cold water solution prepared just before application
		Maize (Corn grain) (<i>Zea mays</i> subsp. <i>mays</i> L.) Sweet Maize (Sweet corn) (<i>Zea mays</i> L. convar. <i>saccharata</i> Koern)		Symphylans (<i>Scutigerella immaculata</i>) ¹³	Treatment in seedling line before 9 AM (solar time)	-	1	Na		40	40 g/ha		
		<i>Zea mays</i> subsp. <i>mays</i> L.			Foliar application Spraying early in the morning before 9 AM (solar time)	1 application at 2–3 leaves (BBCH 12–13) + 1 application at 4 leaves (BBCH 14)	2	1–2		82	8.2 g/ha		
		Grapevine (<i>Vitis vinifera</i>)		Vine leafhopper (<i>Scaphoideus titanus</i>) ⁴	Foliar application spraying early in the morning before 9 AM (solar time)	From the BBCH stage 17 to 57	3	3		150	15 g/ha		
		Grapevine (<i>Vitis vinifera</i>)		Downy mildew (<i>Plasmopara viticola</i>) ⁴	Foliar application spraying early in the morning before 9 AM (solar time)	From 1st shoots to cluster tightening Spring (BBCH 10–57)	up to 12	>12		100–200	10–20 g/ha		

Table 1. Cont.

Basic Substance	Approval Regulation and Applicant	Crops and/or Situation	Function in Plant Protection	Pests or Group of Pests Target	Application				Application Rates			PHI ¹	Notes
					Method	Growth Stage & Season	No. Min/Max	IBA ¹ (Days)	Min–Max	Water L/ha Min–Max	Total Rate		
Sodium hydrogen carbonate	Reg. (EU) 2015/2069 Reg. (EU) 2015/2069 Danish Environmental Protection Agency	Vegetables Soft fruit Ornamentals	Fungicide and herbicide	Mildews (<i>Sphaerotheca</i> spp., <i>Oidium</i> spp.)	Broad cast using field spray or greenhouse spray	BBCH 12 to 89	1–8	10	333–1000 g/hL	300–600	2000–5000 g/ha or 0.33–1.0% Max 1% Dose adjusted depending on water volume	1	Different crops have different sensitivity. Check concentrations for phytotoxic effects before widely used
		Grapevine (<i>Vitis vinifera</i>)		Powdery mildew (<i>Erysiphe necator</i>)	Broadcast using air blast orchard sprayer	BBCH 12 to 89	1–8		420–2000 g/hL	200–600	2500–5000 g/ha or 0.42–2.0%		Volumes and doses will vary according to crop canopy size. Conc. higher than 1–2% can be phytotoxic
		Apple		Apple scab (<i>Venturia inaequalis</i>)	Broadcast using air blast orchard sprayer	BBCH 10 to 85	1–8		500–1000 g/hL	500–1000	2500–5000 g/ha or 0.5–1.0%		Dose rates between 1–4% has been tested
		Fruit of different types (oranges, cherries, apples, papaya)		Storage diseases like Blue mold (<i>Penicillium italicum</i>) Green mold (<i>Penicillium digitatum</i>)	Dipping or surface treatment	Harvested fruit	1–2		1000–4000 g in 100 L water	1–4%			
		Potted plants		Liverwort/Bryophyte (thallose, <i>Lumularia cruciata</i>) Green thallus of liverwort plus, fruiting bodies	Direct application of powder	Post emergence late summer or winter	1		Na	Na	Na		122 kg/ha

Table 1. Cont.

Basic Substance	Approval Regulation and Applicant	Crops and/or Situation	Function in Plant Protection	Pests or Group of Pests Target	Application				Application Rates		PHI ¹	Notes					
					Method	Growth Stage & Season	No. Min/Max	IBA ¹ (Days)	Min–Max	Water L/ha Min–Max			Total Rate				
Whey	Reg. (EU) 2016/560 ITAB	Cucumber (<i>Cucumis sativus</i>), zucchini squash (<i>Cucurbita pepo</i>)	Fungicide and virucide	<i>Podospaera fusca</i> , <i>Podospaera xanthii</i> , <i>Golovinomyces cichoracearum</i> , <i>Erysiphe orontii</i> , <i>Sphaerotheca fuliginea</i> , <i>Leveillula cucurbitacearum</i>	Foliar spray ¹²	From three weeks after sowing (9th leaf unfolded on main stem) to 9 or more primary side shoots visible (BBCH 19–49) ¹⁴	3–5	7	0.6–3 L (0.036–0.24 kg/hL)	1000–1500	6–30 L (0.36–2.4 kg/ha)	Na	Whey should be used rapidly after collection, not stored in metal vessel				
		Grapevine (<i>Vitis vinifera</i>)		Powdery mildew (<i>Erysiphe necator</i>)										From 1st shoots to cluster tightening Spring ¹⁵	7–10	6–30 L (0.36–2.4 kg/hL)	100–30,0 ¹⁵
		Vegetable Gardening, Tomato (<i>Lycopersicon esculentum</i>)		Tomato (Sinaloa) yellow leaf curl virus Begomovirus										First inflorescence visible Summer (BBCH 10–51) ¹⁵	3–4	0.6–3 L (0.036–0.24 kg/hL)	1000–1500
		Glove fingertips and mechanical cutting tools All crops		Viruses (Mechanically transferable) e.g., Tobacco mosaic virus (TMV), Tomato mosaic virus (ToMV), Pepper mild mottle virus (PMMV), Cucumber green mottle mosaic virus (CGMMV), Tomato brown rugose fruit virus (ToBRFV)	Dipping	On tools and glove fingertips	Before/after every plant contact ¹⁶	Na	Na	Na	Na			Dipping for 5 s for gloves and 5 min for mechanical cutting tools. For reasons of efficacy use whey protein powder with at least 80% protein content. Replace the whey solution regularly (e.g., after each crop row) to prevent cross contamination of the plant			

Table 1. Cont.

Basic Substance	Approval Regulation and Applicant	Crops and/or Situation	Function in Plant Protection	Pests or Group of Pests Target	Application				Application Rates			Notes	
					Method	Growth Stage & Season	No. Min/Max	IBA ¹ (Days)	Min–Max	Water L/ha Min–Max	Total Rate		PHI ¹
Diammonium phosphate	Reg. (EU) 2016/548 ITAB	Orchards including cherry tree (<i>Prunus</i> spp.)	Attractant	Mediterranean fruit fly (<i>Ceratitis capitata</i>), Cherry fly (<i>Rhagoletis cerasi</i>)	Placed in physical traps	Na	Mass trapping: 1 trap per tree up to 100 traps/ha	42–56 ¹⁷	max 4 kg/hL	Mass trapping: max 100	Mass trapping: max 4 kg/ha	Na	
		Olive trees (<i>Olea europaea</i>)		Olive fly (<i>Bactrocera oleae</i>)									
		Citrus spp. Other crops where <i>C. capitata</i> cause damage		Mediterranean fruit fly (<i>Ceratitis capitata</i>)									
Sunflower oil	Reg. (EU) 2016/1978 ITAB	Tomato (<i>Lycopersicon esculentum</i>)	Fungicide	Tomato powdery mildew (<i>Pseudoidium neolycopersici</i>)	Foliar application spraying	BBCH 32–37 then BBCH 61–71	2 to 4	8	0.092 kg/hL (0.1 L) –0.46 kg/hL (0.5 L)	500 to 1000	0.46 kg/hL (0.5 L)– 4.6 kg/hL (5 L)	2	Precautions must be taken to avoid overwatering and spilling of the dispersion. Treatment should be avoided during flowering time

Table 1. Cont.

Basic Substance	Approval Regulation and Applicant	Crops and/or Situation	Function in Plant Protection	Pests or Group of Pests Target	Application				Application Rates		PHI ¹	Notes	
					Method	Growth Stage & Season	No. Min/Max	IBA ¹ (Days)	Min–Max	Water L/ha Min–Max			Total Rate
<i>Urtica</i> spp.	Reg. (EU) 2017/419 ITAB	Fruit trees Apple tree (<i>Malus domestica</i>), Plum tree (<i>Prunus domestica</i>), Peach tree (<i>Prunus persica</i>), Red currant (<i>Ribes rubrum</i>), Walnut tree (<i>Juglans</i> sp.), Cherry tree (<i>Prunus</i> sp.)	Insecticide, fungicide, acaricide	Peach-potato Aphid (<i>Myzus persicae</i> , <i>Macrosiphum rosae</i>), woolly Apple aphid (<i>Eriosoma lanigerum</i>), Currant aphid (<i>Cryptomyzus ribis</i>), Walnut aphid (<i>Callaphis juglandis</i>), Black cherry aphid (<i>Myzus cerasi</i>)	Foliar spraying or Shoot spraying Directly on aphids	Spring summer until BBCH 87 (fruit ripe for picking)	1–5	7–15	1500 g/hL (dry matter) ¹⁸	300–900 L/ha	4500–13,500 g/ha ¹⁷	7	Preventive treatment is inefficient 24 h of maceration at 20 °C is enough
		Bean, for example French bean (<i>Phaseolus vulgaris</i>)		Black bean aphid (<i>Aphis fabae</i>)	Na	Spring Summer until BBCH 89 (fully ripe)	Na	Na	Na	4500–7500 g/ha ¹⁸	Na		
		Potato (<i>Solanum tuberosum</i>)		Peach-potato aphid (<i>Myzus persicae</i>)	Na	Spring Summer until BBCH 49 (end of tuber formation)	Na	Na	Na	300–500 L/ha ¹⁸	4500–10,000 g/ha ¹⁷	Na	
		Leaf Vegetables: Lettuce (<i>Lactuca sativa</i>), Cabbage (<i>Brassica oleracea</i>)		Aphids, for example: cabbage aphid (<i>Brevicoryne brassicae</i>), <i>Nazonoviaribis nigri</i>)	Foliar spraying or shoot spraying directly on aphids	Spring Summer until BBCH 19 (9 or more true leaves unfolded)	1–5	7–15	1500 g/hL (dry matter) ¹⁸	4500–7500 g/ha ¹⁸	7	Preventive treatment is inefficient 24 h of maceration at 20 °C is enough	
		Elder tree (<i>Sambucus racemosa</i>)		Elder aphid (<i>Aphis sambuci</i>)	Na	Spring Summer	Na	Na	Na	400–800	6000–12,000 g/ha ¹⁸	7	Preventive treatment is inefficient 24 h of maceration at 20 °C is enough
		Rose (<i>Rosa</i> sp.)		Rose aphid (<i>Macrosiphum rosae</i>)	Na	Spring Summer	Na	Na	Na	300–600	4500–9000 g/ha ¹⁸	7	Preventive treatment is inefficient 24 h of maceration at 20 °C is enough
		<i>Spiraea</i> sp.		<i>Aphis spiraeaphaga</i>	Na	Spring Summer	Na	Na	Na	300–600	4500–9000 g/ha ¹⁸	7	Preventive treatment is inefficient 24 h of maceration at 20 °C is enough

Table 1. Cont.

Basic Substance	Approval Regulation and Applicant	Crops and/or Situation	Function in Plant Protection	Pests or Group of Pests Target	Application				Application Rates			Notes	
					Method	Growth Stage & Season	No. Min/Max	IBA ¹ (Days)	Min–Max	Water L/ha Min–Max	Total Rate		PHI ¹
		Brassicaceae (cabbage— <i>Brassica oleracea</i> , rapeseed— <i>Brassica napus</i> , radish— <i>Raphanus sativus</i>)		Fleabeetle (<i>Phyllotreta nemorum</i>)	Foliar spraying	Spring Summer Until BBCH 19 (9 or more true leaves unfolded)	1–6			300–500	4500–10,000 g/ha ¹⁸		
			Diamondback moth (<i>Plutella xylostella</i>)	Spring Summer until BBCH 49 (Typical leaf mass reached)									
		Apple tree (<i>Malus domestica</i>), Pear tree (<i>Pyrus communis</i>)		Codling moth (<i>Cydia pomonella</i>)		2 treatments in April, 1 treatment in May	3	15		300–900	4500–13,500 g/ha ¹⁸		
		Bean, for example French bean (<i>Phaseolus vulgaris</i>)		Two-spotted spider mite (<i>Tetranychus urticae</i>)		Spring Summer Until BBCH 89 (fully ripe)	1–6 (commonly 3)	7–21		300–500	4500–7500 g/ha ¹⁸	7	24 h of maceration at 20 °C is enough
		Grapevine (<i>Vitis vinifera</i>)		Two-spotted spider mite (<i>Tetranychus urticae</i>), red spider mite (<i>Tetranychus telarius</i>)		Spring Summer Until BBCH 89 stage	1–6 (three before flowering, three after flowering)			300–600	4500–9000 g/ha ¹⁸		
		Brassicaceae (Mustard family, <i>Brassica</i> sp., <i>Sinapis</i> sp., radish— <i>Raphanus sativus</i>)		<i>Alternaria</i> sp.		Spring Summer until BBCH 49 (typical leaf mass reached)	1–6	7–15	1500 g/hL (Based on dry matter) ¹⁸	300–500	4500–7500 g/ha ¹⁸	7	
		Cucurbitaceae (Cucumber— <i>Cucumis sativus</i>)		Powdery mildew (<i>Erysiphe polygoni</i>), <i>Alternaria alternata</i> f. sp. <i>cucurbitae</i>		Until BBCH 89 (typical fully ripe colour)							

Table 1. Cont.

Basic Substance	Approval Regulation and Applicant	Crops and/or Situation	Function in Plant Protection	Pests or Group of Pests Target	Application				Application Rates			Notes	
					Method	Growth Stage & Season	No. Min/Max	IBA ¹ (Days)	Min–Max	Water L/ha Min–Max	Total Rate		PHI ¹
Sodium chloride	Reg. (EU) 2017/1529 Reg. (EU) 2021/556 ITAB AHDB	Grapevine (<i>Vitis vinifera</i>)	Fungicide, insecticide, herbicide	Fungal diseases Powdery mildews (<i>Erysiphe necator</i>)	Foliar application spraying	From 1st shoots (BBCH 10) to cluster tightening (BBCH 57) Spring to summer	1–2	Na	600–2000 g/hL	200	1200–4000	30	In case of 2 applications: one at 20 g/L + one at only 10 g/L. Maximum total rate of salt shall not exceed 6 kg/ha per year. Careful application should be controlled in terms of spray and target should be only the foliage. Low volumes are recommended in order to avoid spill. It is recommended not to spray every year, only in emergency cases. Maximum total rate of sodium chloride shall not exceed 6 kg/ha per year

Table 1. Cont.

Basic Substance	Approval Regulation and Applicant	Crops and/or Situation	Function in Plant Protection	Pests or Group of Pests Target	Application				Application Rates			Notes	
					Method	Growth Stage & Season	No. Min/Max	IBA ¹ (Days)	Min–Max	Water L/ha Min–Max	Total Rate		PHI ¹
		Mushrooms like <i>Agaricus bisporus</i>		Fungal diseases like cobweb disease (<i>Cladobotryum</i> strains—i.e., <i>Mycophilum</i>), dry bubble disease (<i>Lecanicillium fungicola</i>), wet bubble disease (<i>Mycogone pernicioso</i>)	Hand trowel cup scoop	On finding the pathogen. No earlier than 16 days into grow cycle	1	Na	0.03 g/kg	–Dry	80–100 g/ha	Na	Salt is used as a spot treatment to cover incidents of disease. On a well-managed farm, disease will be spotted early with specialist teams identifying and spot treating. This avoids harvesters accidentally spreading disease thorough contamination of personal protective equipment (PPE) and transfer to other areas. This in turn will keep on site disease levels low and avoid the use of large volumes of salt.

Table 1. Cont.

Basic Substance	Approval Regulation and Applicant	Crops and/or Situation	Function in Plant Protection	Pests or Group of Pests Target	Application				Application Rates			Notes	
					Method	Growth Stage & Season	No. Min/Max	IBA ¹ (Days)	Min–Max	Water L/ha Min–Max	Total Rate		PHI ¹
		Grapevine (<i>Vitis vinifera</i>)		European grapevine moth (<i>Lobesia botrana</i>)	Foliar application spraying	1st late April to May (BBCH 55–57) 2nd July (BBCH 75–77) 3rd September (BBCH 83–91)	1–3	Depen- ding on egg stage	600 g/ha	200	1200–3600 g/ha	30	Careful application should be controlled in terms of spray and target should be only the foliage. Low volumes are recommended to avoid spill. It is recommended not to spray every year, only in emergency cases
		Salt swamps and salt marshes		<i>Baccharis halimifolia</i>	Spot application on drilled tree stump or on soil in direct vicinity of tree stump	November–February	1	Na	Na	Na	10–100 g per tree stump ²⁰	Na	Treatment is allowed only in salt marshes and salt swamps zones as defined by national or local authorities. Treatment should be performed outside the rainy period
Beer	Reg. (EU) 2017/2090 ITAB	All edible and nonedible crops	Molluscicide	Pest slugs and snails	Specific traps for slugs	At the beginning of infestation	1–5	Na	Not applicable (because ready to use liquid)	Na	Na	Na	

Table 1. Cont.

Basic Substance	Approval Regulation and Applicant	Crops and/or Situation	Function in Plant Protection	Pests or Group of Pests Target	Application				Application Rates			PHI ¹	Notes				
					Method	Growth Stage & Season	No. Min/Max	IBA ¹ (Days)	Min–Max	Water L/ha Min–Max	Total Rate						
Mustard seed powder	Reg. (EU) 2017/2066 ITAB	Wheat seeds (<i>Triticum vulgare</i> , <i>Triticum aestivum</i>), Durum wheat (<i>Triticum durum</i>), Spelt (<i>Triticum spelta</i>)	Fungicide for seed treatment	Fungi like Common Bunt (<i>Tilletia caries</i> , <i>Tilletia foetida</i>)	Seed application before sowing	Summer to Autumn	1	Na	Na	Na	1.5 kg/100 kg seeds	Na	Mix 1.5 kg of mustard seeds powder with 4.5 L water. Treat 100 kg seeds with the slurry created				
Talc E553B	Reg. (EU) 2018/691 COMPO Expert France SAS	Fruit trees i.e., Apple fruit (<i>Malus Domestica</i>), Pear tree (<i>Pyrus sp.</i>), Olive tree (<i>Olea europea</i>), etc.	Insectifuge, fungifuge	Physical barrier, Insectifuge: Insects and mites like <i>Cacopsylla pyri</i> , <i>Cacopsylla fulguralis</i> , <i>Drosophila suzukii</i> , <i>Panonychus ulmi</i> , <i>Bactrocera oleae</i>	Foliar application spraying	From BBCH 41	2–5	21–28	1st application: 2.13 to 3.54 kg/hL succeeding applications: 1.7 to 2.83 kg/hL	600–1000	1st application: 21.25 kg/ha succeeding applications: 17 kg/ha	Na	Water solution prepared just before application and maintained stirred				
		Fruit trees i.e., Apple fruit (<i>Malus Domestica</i>), Pear tree (<i>Pyrus sp.</i>)		Physical barrier, Fungifuge: Foliar fungi like mildews (<i>Venturia inaequalis</i> , <i>Erysiphe necator</i>)										3–5	14–21	1.28–2.13 kg/hL	12.75 kg/ha
		Grapevine (<i>Vitis vinifera</i>)												From BBCH 20	2–5	21–28	4.25–8.5 kg/hL
Onion oil	Reg. (EU) 2018/1295 Bionext	Carrots, celery, parsnip, parsley root	Repellent, scent masking	Carrot root fly (<i>Psilla rosae</i>)	Masking the smell of the umbelliferous crop by onion oil evaporated from dispensers	Shortly after planting or crop emergence (around mid–April) until end of November (before harvest)	1	Na	Na	Pot dispensers 0.08–0.160 L/ha Granule Dispenser 17.6–35.2 g/ha	Na	Na	4–8 dispensers per ha professional use only				

Table 1. Cont.

Basic Substance	Approval Regulation and Applicant	Crops and/or Situation	Function in Plant Protection	Pests or Group of Pests Target	Application					Application Rates			Notes
					Method	Growth Stage & Season	No. Min/Max	IBA ¹ (Days)	Min–Max	Water L/ha Min–Max	Total Rate	PHI ¹	
L-cysteine	Reg. (EU) 2020/642 Soleo-EcoSolutions	All crops and forestry in tropical areas	Insecticide	Leaf cutting ants	Hand held spreader	Post swarming (July)	1–3	30	3–36 kg granules/ha	Na	Min 0.015 kg/ha Max 2.88 kg/ha ²¹	Na	Used as an insecticide against ants. Application is made by hand on nest of ants. The application can be renewed, if necessary, with a maximum of 3 applications. Minimum/ Maximum number of nests by hectare: 10–120
Cow milk	Reg. (EU) 2020/1004 Basic-Eco-Logique	Grapevine (<i>Vitis vinifera</i>)	Fungicide and virucide	Powdery mildew (<i>Erysiphe necator</i>)	Foliar application Spraying	From 1st shoots (BBCH 07) to inflorescences fully developed; flowers separating (BBCH 57) ²²	3–6	6–8	10–40 L/hL	100–300	10–120 L/ha	Na	No application in presence of fruits
		Vegetable Gardening pumpkin (<i>Cucurbita pepo</i>)		Pumpkins powdery mildew (<i>Podosphaera fusca</i>)		From leaf development (BBCH 01) until flowering (BBCH 06) ²³	3–4	7–12	50 L/hL	400	200 L/ha		
		Flower Gerbera (<i>Gerbera jamesonii</i>)		Powdery mildew (<i>Erysiphe cichoracearum</i>)		Before and during flowering (BBCH 51–69)	3–4		16 L/hL	500–1000	80–160 L/ha	8	
		Cucumber (<i>Cucumis sativus</i>), Zucchini squash (<i>Cucurbita pepo</i>)		Powdery Mildew (<i>Podosphaera fuliginea</i>)		From three weeks after sowing (9th leaf unfolded on main stem) to 9 or more primary side shoots visible (BBCH 19–49) ²⁴	3–4	7	5–10 L/hL	1000–1500	50–150 L/ha	Na	

Table 1. Cont.

Basic Substance	Approval Regulation and Applicant	Crops and/or Situation	Function in Plant Protection	Pests or Group of Pests Target	Application				Application Rates			Notes
					Method	Growth Stage & Season	No. Min/Max	IBA ¹ (Days)	Min–Max	Water L/ha Min–Max	Total Rate	
		Soybean (<i>Glycine max</i> (L.) Merr)		Soybean Powdery mildew (<i>Erysiphe diffusa</i>)		On leaves (BBCH 19–49)	3–4	7	18 L/hL	1000–1500	180–270 L/ha	
		Glove fingertips and mechanical cutting tools All crops		Viruses (mechanically transferable) e.g., Tobacco mosaic virus (TMV), Tomato mosaic virus (ToMV), Pepper mild mottle virus (PMMV), Cucumber green mottle mosaic virus (CGMMV)	Dipping	On tools	Before/after every plant contact	Before/after every plant contact	Before/after every plant contact	Na	Na	Dipping for 2 s. For reasons of efficacy use milk with at least 3,5% protein content. Replace the milk regularly (e.g., after each crop row) to prevent cross-contamination of the plants
<i>Allium cepa</i> bulb extract	Reg. (EU) 2021/81 ITAB	Potatoes (<i>Solanum tuberosum</i>)	Fungicide	Early blight (<i>Alternaria solani</i>)	Spray	BBCH 21–85	3–5	7	1 kg/hL	600–1000	6–10 L/ha (0.3–0.5 kg onion bulb/ha)	Na
		Vegetable Gardening Tomato (<i>Lycopersicon esculentum</i>)		Tomato late blight (<i>Phytophthora infestans</i>)		75 days after planting BBCH 21–75		3–4		1500	15 L/ha (0.75 kg onion bulb/ha)	
		Cucumber (<i>Cucumis sativus</i>)		Cucumber gray mold (<i>Botrytis cinerea</i>)				7				

Table 1. Cont.

Basic Substance	Approval Regulation and Applicant	Crops and/or Situation	Function in Plant Protection	Pests or Group of Pests Target	Application				Application Rates		PHI ¹	Notes	
					Method	Growth Stage & Season	No. Min/Max	IBA ¹ (Days)	Min–Max	Water L/ha Min–Max			Total Rate
Chitosan	Reg. (EU) 2022/456 KitoZyme	Horticulture	Fungicide	Plant elicitor, plant resistance against pathogenic fungi and bacteria	Spray	BBCH 09 to BBCH 89	4–8	2 weeks	50–100 g/hL	200–400	100–400	Na	Chitosan can be prepared for use following any of the two recipes provided in Appendix of Reg. (EU) 2022/456 (preparation for use).
		Low–Medium volume spraying			From 1st new leaf development BBCH 10 to development of fruit BBCH 71	800–3200				800–7800			
		olive trees											
		grapes								200–600	800–7800		
		grass				BBCH 09 to BBCH 89				200–400	800–3200		
		postharvest fruit treatment		Pathogenic fungi and bacteria	Immersion	Postharvest BBCH 89+	1	-	1	-	-		

¹ IBA: Interval between applications; PHI: minimum preharvest interval; Na: Data not available; ² The product cannot be applied in case of hot temperature. It is used in case of rainy period; ³ Indirect actions, no direct insecticide and fungicide properties; ⁴ maximum of rate per application; ⁵ maximum total rate per crop/season; ⁶ The aqueous solutions in this application are applied with few or without dilution. Here the case without dilution is calculated. Usually, not all trees are treated with brush application but only injured trees. In the calculation of maximum rate, it was assumed that 3000 trees per ha are treated with 0.15 L product per tree. This means that all trees of an orchard would be treated with several big wounds, which would be really the maximum rate and in reality, is very improbable; ⁷ Expressed as acetic acid. 1/1 dilution of vinegar/water L/L; ⁸ Considering 0.9 to 2 qt of seeds per ha; ⁹ Expressed as acetic acid. 50 mL/1 L dilution of vinegar/water for vinegar at 8% acetic acid; ¹⁰ Of main active substance acetic acid for vinegar at 10% acetic acid; ¹¹ Expressed as acetic acid in a preparation with 60% vinegar (diluted in water), for vinegar at 10% acetic acid; ¹² Treatments must be delayed 24–48 h or more after rain; ¹³ Spray when there is sun (preferably morning); ¹⁴ Do not apply when any plant is at a later growth stage than BBCH 49; ¹⁵ With a maximum of 10% concentration (30 L in 300 L); ¹⁶ Do not apply on treating fingertips right before or during harvest of edible commodities; ¹⁷ Depending upon environmental factors such as climate and topography; ¹⁸ The quantities of fresh nettle (or dry matter) written represents the quantities of nettle used in the recipe, but not the quantities that are effectively put in field—there is a filtration before; ¹⁹ Treatment, just before sowing; ²⁰ Assuming plant density of between 0.1/m² to 1/m²; ²¹ 300 g of granules per nest multiplied by 120 nest/ha = 36 kg product/ha. Considering a maximum of 8% L-cysteine in the product, the maximum application rate per treatment of L-cysteine is 2.88 kg/ha; ²² Do not apply when any plant is at a later growth stage than BBCH 57; ²³ Do not apply when any plant in the greenhouse is at a later growth stage than BBCH 06 and in presence of fruits; ²⁴ Do not apply when any plant in the greenhouse is at a later growth stage than BBCH 49.

Table 2. Typical uses of the basic substances.

Substance Name	Use(s)	Application	Recipe	Formulation Type
<i>Equisetum arvense</i>	Fungicide	Spraying on crops	2–2.25% water dilution 200 to 225 g/100 L water The product cannot be applied in case of hot temperature. It is used in case of rainy period	Dispersible concentrate
		Dry	9 kg/100 kg mulch	Mulch
Chitosan hydrochloride	Elicitor	Spraying on crops or seeds	0.05–0.2% water dilution 50 to 200 g/100 L water Must be applied within 24 h	Soluble powder, paste
Sodium hydrogen carbonate	Fungicide	Aerial parts spraying	0.33–2% water dilution 333 to 2000 g/100 L water	Soluble powder
		Postharvest dipping	1–4% water dilution 1 to 4 kg/100 L water	
	Herbicide	Direct dusting	10 g for a 50 cm Ø pot	Dry powder
Sunflower oil	Fungicide	Foliar spraying	0.1–0.5% water dilution 100 to 500 mL/100 L water	Oil dispersion
Hydrogen peroxide	Seed treatment	Seeds soaking	Ready-to-use solution (<5%)	Ready-to-use solution
<i>Urtica</i> spp.	Fungicide Insecticide	Spraying	3–4 days maceration in water at 20 °C Fresh leaves (75 g/L) or dried leaves (15 g/L) Water dilution by 6 of filtered maceration	Dispersible concentrate
		Mulch incorporation	Addition of dried aerial parts. 83 g/kg of mulch	Mulch
Clayed charcoal	Protectant	Soil burying	Buried. 500 kg/hectare maximum	Pellet
Sodium chloride	Fungicide Insecticide	Foliar spraying	0.6–2% water dilution 600 to 2000 g/100 L water	Soluble powder
		Substrate burying	Mix salt in the substrate. 30 g/kg substrate (3%)	Pellet
Beer	Molluscicide	Trap	Covered slug traps. 1 trap per m ² maximum	Pure product
Di Ammonium Phosphate	Attractant	Trap	Place in traps/bottle, 30 g/L.	Soluble powder
Onion oil	Odor mask	Oil dispenser	Fill the dispenser with onion oil only (20 mL) Fill the dispenser with oil then add the pellets (4.4 g oil per 30 g granule)	Oil or pellet
L-cysteine	Insecticide	Hand-held spreader	Mixture with matrix (flour, food grade) at a concentration of maximum 8%	Bait (ready for use)

Table 2. Cont.

Substance Name	Use(s)	Application	Recipe	Formulation Type
Cow milk	Fungicide	Foliar spraying	5–50% water dilution = 0.5 to 5 L of cow milk filled up with water to 10 L	Soluble concentrate
		Dipping	Dipping tools for 2 s in undiluted cow milk. For reasons of efficacy use milk with at least 3.5% protein content	
<i>Allium cepa</i> L. bulb extract	Fungicide	Spray application	Boil 500 g of chopped onions in 10 L of water for ten minutes then let infuse for a quarter of an hour and filter the mixture	Dispersible concentrate
Chitosan	Fungicide	Spray application & Immersion	Preparation 1: added to a half-filled water tank, making sure the powder is evenly distributed over the water surface to avoid aggregation. The mixture should be stirred vigorously while adding the remaining water. The mixture should be used as soon as possible. Preparation 2: dissolved in water with pH < 5. The pH of water should be regulated by adding 7 mL vinegar (8% of acetic acid) per 1 L of water).	Soluble powder
Vinegar	Fungicide	Seed treatment	Vinegar to be diluted in compliance with the rates of application reported in Appendix II.	Liquid for seed treatment
	Herbicide	Tools disinfection	Undiluted for uses as herbicide on medicinal aromatic and perfume crops.	Liquid
	pH modifier	Spray or spot application/	For the herbicidal use in spot applications on paths, borders, sidewalks and terraces, vinegar needs to be diluted to a concentration of 60% vinegar in water (60/40 vinegar/water).	Liquid

Some applications were not validated by DG Santé and Member States during discussion and votes. Some were withdrawn (Table 3) by applicants during evaluation or discussions with no regulatory trace, while some were processed up to the vote and finally non-approved with corresponding Implementing Regulations (Table 4).

Table 3. Basic substance applications retired during the evaluation process.

Basic Substances Removed/Withdrawn during Evaluation				
Substance Name	Intended Use(s)		EFSA Opinion	Reason(s)
<i>Castanea and Schinopsis sp. tannins</i>	Bactericide, fungicide and nematocide		EN 1363	Limited number of studies about toxicity and residues led to a doubt concerning exposure assessment. Non-dietary exposure considered as hazardous
Honey from rhododendron	Rodenticide		EN 1155	Lack of studies concerning substance composition and efficacy on rodents. Rodents in traps might suffer 'too long'
Extract from rhododendron	Rodenticide		EN 1596	Lack of studies concerning substance composition and efficacy on rodents. Rodents in traps might suffer 'too long'
<i>Quassiaamara</i> extract	Insecticide and repellent		EN 1382	Data gaps were identified for genotoxicity, residues, environmental risk and exposure assessment. Concerns were raised regarding reproductive and endocrine toxicity
<i>Valeriana officinalis</i>	Frost protection		None	Potential neurotoxicity, Valerian herbal tea makes it easier to fall asleep
<i>Citrus pulp</i>	-		None	-
<i>Potassium metabisulfite</i>	-		None	-
<i>Didecyl-dimethylammonium chloride (DDAC)</i>	-		RN-214	Toxic to aquatic organisms

Table 4. Basic substance applications refused (non-approval).

Substances Not Approved by the European Commission				
Substance Name	Intended Use(s)	Implementing Regulation	EFSA Opinion	Reason(s)
<i>Achillea millefolium</i> L.	Fungicide and insecticide	EU no. 2017/2057	EN 1093	Risk assessment for toxicology and ecotoxicology not comprehensive enough left doubts and substance is not considered as foodstuff
<i>Arctium lappa</i> L. aerial parts	Fungicide and insecticide	EU no. 2082/2015	EN 699	Risk assessment for toxicology and ecotoxicology not comprehensive enough left doubts and substance is not considered as foodstuff
<i>Artemisia absinthium</i> L.	Fungicide, nematocide and insecticide	EU no. 2015/2046	EN 665	Risk assessment for toxicology and ecotoxicology not comprehensive enough left doubts and Regulation (EC) 1334/2008 fixes limits for this substance
<i>Artemisia vulgaris</i> L.	Insecticide/repellent	EU no. 2015/1191	EN 644	Risk assessment for toxicology and ecotoxicology not comprehensive enough left doubts and Regulation (EC) 1334/2008 fixes limits for this substance

Table 4. Cont.

Substances Not Approved by the European Commission				
Substance Name	Intended Use(s)	Implementing Regulation	EFSA Opinion	Reason(s)
<i>Capsicum annuum</i> L. var. <i>annuum</i> , <i>longum</i> group, cayenne, extract (Oleoresin capsicum)	Repellent	EU no.2021/464	EN 1838	Risk assessment for toxicology show genotoxicity, causing serious eye damage, being harmful if swallowed and also as cause of skin irritation, although substance is considered as foodstuff
Caffeine	Molluscicide	EU no. 2022/xx	EN 6423	Proposal for non-approval under discussion
Carbon dioxide	Rodenticide	EU no. 2021/80	None	-
Comfrey steeping	Fungicide and insecticide	EU no. 2021/809	EN 1753	Risk assessment for toxicology and ecotoxicology not comprehensive enough left doubts and Regulation (EC) 1334/2008 fixes limits for this substance
Dimethyl Sulfide	Attractant	EU no. 2021/1451	EN 1911	Risk assessment for toxicology and ecotoxicology not provided for long-term toxicity and carcinogenicity concern
Grape (<i>Vitis vinifera</i>) cane tannins	Fungicide	EU no. 2020/29	EN 1414	Risk assessment for toxicology and ecotoxicology not comprehensive enough left doubts and substance is not considered as foodstuff
Landes pine tar	Protectant and repellent	EU no. 2018/1294	EN 1311	It may contain substances of concern, so there is a lack of data, so risk assessment is not comprehensive enough and left doubts
<i>Origanum vulgare</i> L. essential oil	Fungicide, bactericide and insecticide	EU no. 2017/241	EN 1054	Risk assessment for toxicology and ecotoxicology not comprehensive enough left doubts
Paprika extract E160c	Repellent	EU no. 2017/2067	EN 1096	Risk assessment for toxicology and ecotoxicology not comprehensive enough left doubts
Potassium sorbate	Fungicide	EU no. 2017/2058	EN 1232	Lack of data concerning residues lead to an impossibility concerning exposition assessment
Propolis (water soluble extract)	Fungicide and bactericide	EU no. 2020/640	EN-1494	Defined as a skin sensitizer, risk assessment for genotoxicity and endocrine disruption toxicity left doubts. No safe limit for the use. Substance is not considered as foodstuff
<i>Rheum officinale</i> roots extract	Fungicide	EU no. 2015/707	EN 617	Risk assessment for toxicology and ecotoxicology not comprehensive enough left doubts and substance is not considered as foodstuff

Table 4. Cont.

Substances Not Approved by the European Commission				
Substance Name	Intended Use(s)	Implementing Regulation	EFSA Opinion	Reason(s)
<i>Saponaria officinalis</i> L. roots	Acaricide and elicitor	EU no. 2020/643	EN 1263	Risk assessment for toxicology and ecotoxicology not comprehensive enough left doubts
<i>Satureja montana</i> L. essential oil	Fungicide and bactericide	EU no. 2017/240	EN 1051	Risk assessment for toxicology and ecotoxicology not comprehensive enough left doubts
<i>Tanacetum vulgare</i> L.	Repellent	EU no. 2015/2083	EN 666	Risk assessment for toxicology and ecotoxicology not comprehensive enough left doubts and substance is not considered as foodstuff
Willow bark and stem extract	Plant growth and defense elicitor	EU no.2022/	EN 1872	Previously proposed for non-approval since not sold for other uses, proposal under discussion, may be accepted.

The scientific literature dealing with basic substances is relatively limited but increasing in recent years (Figure 2), and there is poor information about the effectiveness in field trials of basic substances toward pests and pathogens.
















Figure 2. Number of documents available on Scopus through searches with keywords ‘basic substances’ in ‘Article title, Abstract, and Keywords’ (histograms) or in ‘All fields’ (linear) published over the last 10 years (Source: Scopus, <https://www.scopus.com>, accessed on 11 May 2022).

In the last decade, MRLs for pesticides with agricultural trade are becoming important. In the EU, there are increasing requirements from retailers to their suppliers to provide fruits and vegetables with an amount of pesticide residue below the MRLs (Table 5).

Table 5. Examples of requests from the retailer of the amount of the Maximum Residue Level (MRL) and Acute reference doses (ARfD).

Retailer		Max. %MRL/Active Substance	Max. Sum %MRL/Sample	Max. %ARfD/Active Substance	Max. Sum %ARfD/Sample	Max. Number of Active Substances/Samples
ALDI/ HOFER		70%	80%	70%	80%	3–5
ALBERT HEIJN		50%	-	50%	-	-
ASDA		80%	-	-	-	-
BILLA		100%	-	100%	-	-
DOHLA		-	70%	-	70%	3–5
EDEKA		70%	-	100%	-	5
EDEKA OWN BRANDS		50%	-	70%	-	5
GLOBUS		70%	-	70%	100%	5
LIDL		33.3%	80%	100%	-	5
KAUFLAND		33.3%	80%	50%	50%	5
NORMA		-	70%	-	70%	5
METRO		50%	80%	70%	100%	5
MIGROS		-	-	-	-	6
NETTO		70%	-	100%	-	5
REWE		50%	100%	70%	100%	5
REWE OWN BRANDS		50%	100%	50%	-	5
TEGUT		70%	-	70%	-	Max. 4 (>0.01 mg/kg)
TENGEL MANN		70%	150%	70%	100%	-

The substances tested during Casdar programs ‘4P’, ‘Carie’, ‘Sweet’, ‘HE’, Ecophyto ‘Usage’ and some from projects have already been described (Marchand, 2016) (Table 6). New projects are ongoing to develop extensions of use, describe better efficacy through better positioning during the season or to investigate compatibility/incompatibility with other biocontrol agents (i.e., reduce copper and macro-organisms). This is the ongoing work for Coperreplace, ABAPIC (ITAB), Vitinnova (UNIVPM), and Euphresco BasicS (Euphresco Network).

Table 6. Examples of the applications of the basic substances in research projects.

Substance Name	Use(s)	Program	Reference
Horsetail (<i>Equisetum arvense</i> L.)	Fungicide	Casdar ‘4P’ Coppereplace	[24–26]
White willow bark (<i>Salix cortex</i>)	Fungicide	Casdar ‘4P’	[24,25]
Vinegar	Seed treatment	Casdar ‘Carie’	[27]; http://itab.asso.fr/programmes/carie-ble.php , accessed on 23 May 2022.
Mustard seed powder	Seed treatment		
Sucrose	Elicitor	Ecophyto ‘Usage’ and Casdar ‘Sweet’, ABAPIC	[28]; https://ecophytopic.fr/cuivre-viticulture/protoger/micro-doses-de-sucres , accessed on 23 May 2022. [29]; https://ecophytopic.fr/sites/default/files/USAGE.pdf , accessed on 23 May 2022.
Fructose	Elicitor		
Lecithin	Fungicide	Casdar ‘HE’	[30]; https://ecophytopic.fr/recherche-innovation/protoger/projet-he , accessed on 23 May 2022.
Talc	Fungicide		[31]
Whey	Fungicide		[32]
Di-ammonium phosphate (DAP)	Attractant	out of program	[33]; https://ecophytopic.fr/pic/protoger/protoger-ses-oliviers-de-la-mouches-en-limitant-les-traitements , accessed on 23 May 2022.
Calcium hydroxide	Fungicide		[34]; https://www.researchgate.net/publication/279636728_The_post-infection_activity_of_hydrated_lime_against_conidia_of_Venturia_inaequalis , accessed on 23 May 2022.
Chitosan hydrochloride	Fungicide	Vitinnova	[35]; www.vitinnova.it/en , accessed on 23 May 2022.
		Euphresco BasicS	[16]; https://www.researchgate.net/project/EUPHRESCO-Basic-substances-as-an-environmentally-friendly-alternative-to-synthetic-pesticides-for-plant-protection-BasicS , accessed on 23 May 2022.
		PRIMA StopMedWaste	[36]; www.stopmedwaste.eu , accessed on 23 May 2022.
		ZeroSprechi	[37]; www.zerosprechi.info/en/zerosprechi , accessed on 23 May 2022.
		CleanSeed	[38]; https://www.cleansed.it/en/cleansed-2/ , accessed on 23 May 2022.

Each use of plant extracts and natural products, such as decoctions, herbal teas, or aqueous solutions, have been defined and tested in the field or identified from the literature then controlled or cross-referenced with producer surveys. Whenever water is mentioned in these tests, it is either natural spring water or rainwater. Each basic substance preparation is described in Section 2.5 of Basic substances applications in EU 2012. The evaluation process of the basic substance application is getting longer, and legal delays fixed by EC are not consistently respected. The evaluation process lasts an average of 19 months (Supplementary Table S1 and Figure S1), while the legal maximum delay is fixed at 18 months until basic substance application admissibility. Even not considering admissibility evaluation delays that are considered outside of the evaluation process, this process becomes longer from year to year, resulting in a delay in availability of additional basic substances.

3. Discussion

The use of pesticides, if not appropriate, may lead to problems like contamination of the water, potential damage to sensitive species (e.g., bees), contamination of final food products and water, with up to 90% of applied pesticides not reaching the target species, and, also, because of the development of resistant pathogens and pests [39]. A high number of PPPs were not reauthorized (or companies did not provide the dossier for the reregistration

of products out of patent, due to high costs and uncertain benefits) and leaves a gap for several uses. It is important that authorities provide a good number of options to growers to protect their crops, since farmers cannot stand without PPPs for certain crops and uses, and there is an increasing need, because a lot of substance prohibition dates are fixed without substitution mean. Just as an example, this occurred with the fungicide mancozeb in January 2022 and a risk to occur in 2025 with copper, that is fundamental for plant protection in organic agriculture and a good support to prevent the appearing of resistant isolates in IPM. In France, the use of neonicotinoids, known as dangerous insecticides, is extended when there is no other way to preserve crops and productivity. With Farm to Fork Strategy of the European Green Deal, the European Commission is committed to reduce the use of the most dangerous synthetic pesticides of 50% and achieve at least 25% of the EU agricultural land under organic farming by 2030, although the decrease of synthetic pesticides is already ongoing. These trends, together with the implementation of sustainable development goals—SDGs by the United Nations—are demanding for new alternatives, such as basic substances, to tackle some of these issues. To achieve these goals, more research is needed to advance the design of better farming systems and the development of alternatives to synthetic pesticides and to copper formulations.

Three decades ago, the concept of MRLs was poorly known, while, in recent years, MRLs for pesticides arguably have become the first action growers should consider in their pest management decisions [40]. Trying to interpret consumer demands, retailers are increasingly required to reduce pesticide residues even more than the allowed thresholds (MRLs), which are defined considering a wide security factor (e.g., $\times 100$) using the presence of pesticide residues as a factor of competition among companies. Requests from the retailers and consumer to reduce synthetic pesticide residues from fresh produce even more than the allowed threshold, such that the rules defined by the public administration have become more limiting for farmers in terms of the active ingredients allowed and MRLs [40,41]. The reduction of the presence of fungicide residues well beyond MRL may allow the pathogen to develop after harvest, resulting food loss and waste along the value chain. These developments have driven the search for alternative management strategies that are effective and not reliant just on conventional fungicide applications [5,42,43]. European regulation followed and carried this development with the introduction of new classes of phytosanitary products, in particular basic substances, but also new laws and simplification accompanied by the reduction of registration processes of low-risk substances, theoretically. Basic substances are approved for use in the EU and are products that are already sold for certain purposes, e.g., as a foodstuff or a cosmetic. Basic substances may be of major importance in biocontrol and several advantages can explain it. Basic substance regulatory application is simplified [44] and particularly reduced compared to other substances, therefore representing a lower cost to applicant (around 35–40 kEuro for approval of a basic substance and overall around 45 kEuro including approval for organic agriculture), thanks to the fact that these substances are already on the market for another purpose than plant protection, and safety is not an issue to be demonstrated. These substances are good alternatives available today and wide targets. Basic substances can be used in the crop protection as fungicide, bactericide, insecticide, etc., and most of them are allowed in organic production [18,45–47]. The basic substances are in order from 2014, when was the first approved application of *Equisetum arvense* L., chitosan hydrochloride, and sucrose until 2022, when a second chitosan formulation was approved. In some conditions basic substances were already at farm level, with a level of pest management not different than the standard. Just as example, chitosan hydrochloride was also applied in commercial conditions, in the field, and postharvest treatments, and several studies proved that it could have an effectiveness comparable to some commercial PPPs [42,48]. Basic substances, probably less efficient and practical to use than other active substances authorized as PPPs, are known and used by producers since decades as substitution means and have already demonstrated their effectiveness. Basic substances were the perfect tool to provide to producers as known, easy-to-use, less dangerous, and environmentally

more respectful. Today, there is a consensus among a wide range of stakeholders that synthetic pesticide used need to be gradually reduced to a level that is effectively required to ensure crop production and that risks of pesticide application should be reduced as far as possible. Basic substances are good alternatives available today in our hands. The use of these substances needs to be integrated in vocational education, training, and technical advice to farmers. Further research around the world on the efficacy of basic substances may prove in the future that these substances can replace pesticides without reducing yields or increasing production costs. To develop the uses and the field trials we listed here the main usages of basic substances. However, rates included in the approval schedule may not produce a significant containment of diseases and pests in specific pathosystems. Just as example, the advised application rate of chitosan hydrochloride is between 100 and 800 g/ha, equal to a concentration ranging among 0.05 and 0.2% with 200–400 L/ha, while trials in commercial vineyards found a good effectiveness delivering the chitosan hydrochloride, with a concentration of at least 0.5% and with a volume of at least 500 L/ha [34,49]. For this reasons, large-scale trials are very important to demonstrate the effectiveness toward pathogens and pests in different environments and growing contexts, and a flexibility could be required in suggested dosages to avoid that applying basic substances at suggested rated can lead to a lack of or poor effectiveness and then the disaffection of users toward these innovative compounds, and this is in contrast with the requirements of finding solutions alternatives to the application of synthetic pesticides keeping the standard quality and quantity of the production, which is one of the drivers of the Farm-to=Fork Strategy of European Green Deal. Moreover, the diluent allowed for basic substance, up to now concretely restricted to water, may be another substance. In this case, vinegar has just been authorized for chitosan. Finally, increasing the demand from growers and competition among companies can lead to the reduction of costs of the treatments that, nowadays, are often higher than standard treatments.

4. Materials and Methods

4.1. Collection of Data

A systematic literature search from 2009 to 2021 was performed using the database of Scopus with the keywords 'basic substance' and 'basic substances'. In the EU, several retailers request an amount of pesticide residue on fruit and vegetables below the legal limit (MRL), and data on some protocols were collected through companies and plant doctors.

4.2. Legislation

Basic substance criteria are defined by article 23 of Regulation (EC) No. 1107/2009, cited in introduction. By way of derogation from Article 4 of this regulation, a basic substance is approved when all relevant evaluations conducted in accordance with other Community legislation, governing other uses of this substance, showing that it has neither an immediate or delayed harmful effect on human or animal health nor any unacceptable influence on the environment. Active substances that could be defined as 'foodstuff' are intrinsically considered as basic substances, following Article 2 of Regulation (EC) No. 178/2002. Basic substances shall be approved in accordance with paragraphs 2–6 of regulation (EC) No. 1107/2009 and by way of derogation from Article 5, the approval shall be for an unlimited period. By way of derogation from Article 7 of Regulation (EC) No. 1107/2009, an application for approval of a basic substance can be made by a Member State or any interested party. At the end of the evaluation process, basic substances shall be listed separately in the Regulation referred to in Article 13(4). The Commission may review the approval of an active substance at any time. It may take into account the request of a Member State to review the approval. Article 28 of Regulation (EC) No. 1107/2009 set the absence of marketing authorizations and usages allowance for basic substances. However, no formal authorization is required as long as the product contains exclusively basic substances (see corresponding Review Report) [49,50].

4.3. Approval Process

The approval process of a basic substance starts with a request for approval (Figure 3). The applicant estimates if the substance concerned fulfil all criteria of basic substances category and then complete the BSAT, in English, to obtain a Basic Substance Application. Several guidance documents, such as the official SANCO guide or the teaching guide from the ITAB, have been published to help applicants to build basic substance application correctly [50]. For the transmission of the basic substance application, once completed, the file should be sent to the DGSanté, representing the European Commission (EC). The Basic Substance Application can firstly be sent to national competent authorities for a preassessment and possibly a support. For example, in France, the Basic Substance Application can be sent to the Ministry of Agriculture (DGAI in France), who can ask for the National Authority' opinion and then transfer the file to the EC. Upon receipt of the Basic Substance Application, EC implements the approval procedure detailed in Article 23 of Regulation (EC) No. 1107/2009. Admissibility may be pronounced at any time, directly or after questions from DGSanté. It constitutes the real start of the application (black line in Figure 3). The first stage is based on the Basic Substance Application evaluation by Member States and EFSA as scientific assistance leading to a request for corrections and questions. The request is sent to the applicant, and his answers shall be sent back within one month to the EFSA. For decision and approval, at the end of the basic substance application evaluation, EFSA will deliver its opinion, append a comment, and send the basic substance application to the DG Health within 3 months for the final vote of Member States in the PAFF committee (Figure 3). Approval, if accorded, is effective at the date of the publication of an implementing Regulation modifying Regulation (EU) No. 540/2011 [8].

Basic Substance pathway




Figure 3. Approval process and timeline of a Basic Substance Application (BSA).

The period of examination of the basic substance application is established in paragraph 1 of article 37 of Regulation (EC) No. 1107/2009. It is said: ‘The Member State examining the application shall decide within 12 months of receiving it whether the requirements for authorization are met. Where the Member State needs additional information, it shall set a period for the applicant to supply it. In that case, the 12-month period shall be extended by the additional period granted by the Member State. That additional period shall

be a maximum of 6 months and shall cease at the moment when the additional information is received by the Member State. Where at the end of that period the applicant has not submitted the missing elements, the Member State shall inform the applicant that the application is inadmissible.' [10]. The maximum delay is therefore set at 18 months. However, although clearly defined, these steps are not so straightforward in many cases [51].

4.4. Extension of Uses Process

The request for an extension is somehow similar, except the need of support from corresponding agricultural sectors at the deposit step. Some extensions were voted after submission, some others were granted with admissibility and voted rapidly after; some later were following the full approval pathway, including admissibility, evaluation, outcome, full vote at PAFF Committee (appearance in Part A (lecture, discussion), C (proposal) and B (effective vote)). This latter process sometimes takes the same amount of time compared to a new approval, which is considered very excessive by the applicants, having an approved substance at the beginning of their request and only asking for one line sometimes in the Good Agricultural Practices (GAP) table.

4.5. Regulation Analysis

The EU Pesticides Database [52] was used to detect basic substances and their status (approved, nonapproved, pending, and modifications of Review Reports). Corresponding linked Implementing Regulations [20] attached to each active substance were found using the same method and cross-verified with Implementing Regulation (EU) 540/2011. The EU law database for Eur-Lex was also used to track each Implementing Regulation publication. Furthermore, EFSA documents were also compiled to extract decisions supportive analyses.

5. Conclusions

Searching for alternative products for crop protection is an important strategy for promoting more sustainable food systems. The use of basic substances is in line with the restriction on the application of chemical PPPs and the principles of the European Green Deal and SDGs, mostly renewables and with no MRL. There is relatively poor information about the effectiveness of basic substances as compared to synthetic pesticides and biological PPPs. A higher testing and validation of the use of basic substances as a phytosanitary measure can lead to further reduction of application of synthetic pesticides. In addition, searching for the most effective dosage of the basic substance is critical and an important question for phytosanitary consultants (the plant doctors that are opinion leaders in application of innovations in pest management), growers, stakeholder, and companies to avoid that their application at the recommended dose can lead to a lack of or poor effectiveness of these substances. For this reason, a flexibility might be required in the suggested dosage of basic substances approved to ensure good maintenance of the quality and quantity of production, which is one of the keys of the Farm to Fork Strategy of the European Green Deal. Moreover, a defined timeline for approval is basilar to have the chance to increase the number of basic substances available for growers, the scientific community, and the whole agricultural sector, with final benefits for the consumers.

6. Patents

All Implementing Regulations may be considered as patents but with free exploitation, since no Marketing Authorizations are needed for basic substances.

Supplementary Materials: The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/molecules27113484/s1>, Figure S1: Time needed for Basic Substance Application admissibility evaluation over time (bars) and tendency line (dotted line); Table S1: Total time of basic substance application process within admissibility to Implementing Regulation publication in months.

Author Contributions: Conceptualization, G.R. and P.A.M.; methodology, Y.O. and M.M.; resources, G.R., Y.O., Y.D., M.M. and P.A.M.; writing—original draft preparation, G.R., Y.O., Y.D., M.M. and P.A.M.; writing—review and editing, G.R., Y.D., M.M. and P.A.M.; supervision, G.R. and P.A.M.; and funding acquisition, G.R. and P.A.M. All authors have read and agreed to the published version of the manuscript.

Funding: For G.R. and M.M., this work was conducted within the framework of the PSR ZeroSprechi, PSR Vitinnova, PSR CleanSeed, and of PRIMA StopMedWaste projects, which are funded by PRIMA, a program supported by the European Union. For Authors Y.O., Y.D., and P.A.M., the French Ministry of Agriculture (CASDAR ‘4P’, ‘Contrat de branche Carie’, ‘Sweet’, and ‘HE’; Ecophyto ‘Usage’, ‘Biocontrol’ ‘INADOM’, and ‘PARMA’), French Ministry of Ecology (Project ‘PNPP’ CT0007807, ‘SubDOMEx’, and ‘Jussie’). All authors worked within the “Euphresco BasicS (Objective 2020-C-353) project. Thanks are expressed to Antonello Lepore, Gianni Ceredi and other technicians for providing data about pesticide residues requested by retailers.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Data is contained within the article or supplementary material.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. United Nation. Growing at a Slower Pace, World Population Is Expected to Reach 9.7 billion in 2050 and Could Peak at Nearly 11 billion around 2100. 2019. Available online: <https://www.un.org/development/desa/en/news/population/world-population-prospects-2019.html> (accessed on 10 December 2021).
2. FAO. 2021. Available online: <https://www.fao.org/news/story/en/item/1402920/icode/> (accessed on 1 December 2021).
3. Savary, S.; Willocquet, L.; Pethybridge, S.J.; Esker, P.; McRoberts, N.; Nelson, A. The global burden of pathogens and pests on major food crops. *Nat. Ecol. Evol.* **2019**, *3*, 430–439. [[CrossRef](#)] [[PubMed](#)]
4. Dietz-Pfeilstetter, A.; Mendelsohn, M.; Gathmann, A.; Klinkenbuß, D. Considerations and regulatory approaches in the USA and in the EU for dsRNA-based externally applied pesticides for plant protection. *Front. Plant Sci.* **2021**, *12*, 682387. [[CrossRef](#)] [[PubMed](#)]
5. Romanazzi, G.; Smilanick, J.L.; Feliziani, E.; Droby, S. Integrated management of postharvest gray mold on fruit crops. *Postharvest Biol. Technol.* **2016**, *113*, 69–76. [[CrossRef](#)]
6. Chèze, B.; David, M.; Martinet, V. Understanding farmers’ reluctance to reduce pesticide use: A choice experiment. *Ecol. Econ.* **2020**, *167*, 106349. [[CrossRef](#)]
7. Robin, D.; Marchand, P.A. Evolution of Directive (EC) No 128/2009 of the European parliament and of the council establishing a framework for Community action to achieve the sustainable use of pesticides. *JRS* **2019**, *7*, 1–7. [[CrossRef](#)]
8. EU. Commission Implementing Regulation No 540/2011 of 25 May 2011 implementing Regulation (EC) No 1107/2009 of the European parliament and of the council as regards the list of approved active substances. *OJ* **2011**, *L153*, 1–186.
9. Matyjaszczyk, E. Problems of implementing compulsory integrated pest management. *Pest Manag. Sci.* **2019**, *75*, 2063–2067. [[CrossRef](#)]
10. Robin, D.; Marchand, P.A. Biocontrol active substances: Evolution since the entry in vigour of Reg. 1107/2009. *Pest Manag. Sci.* **2019**, *75*, 950–959. [[CrossRef](#)]
11. EC. Commission Regulation No 1107/2009 of the European parliament and of the council of 21 October 2009 concerning the placing of plant protection products on the market and repealing council directives 79/117/EEC and 91/414/EEC. *OJ* **2009**, *L309*, 1–50.
12. Kowalska, J.; Roszkowski, S.; Krzysińska, J. Substancje podstawowe—Efektywne uzupełnienie metod ochrony upraw. Basic substances—An effective supplement to crop protection methods. *Prog. Plant Prot.* **2021**, *61*, 139–146. [[CrossRef](#)]
13. Marchand, P.A. Basic and low-risk substances under European Union pesticide regulations: A new choice for biorational portfolios of small and medium-sized enterprises. *J. Plant Prot. Res.* **2017**, *57*, 433–440. [[CrossRef](#)]
14. Charon, M.; Robin, D.; Marchand, P.A. The importance of substances without maximum residue limit (MRL) in integrated pest management (IPM). *Biotechnol. Agron. Soc. Environ.* **2019**, *23*, 22–29. [[CrossRef](#)]
15. EC. Regulation No 178/2002 of the European parliament and of the council of 28 January 2002 laying down the general principles and requirements of food law, establishing the European Food Safety Authority and laying down procedures in matters of food safety. *OJ* **2002**, *L31*, 1–24.
16. Euphresco. BasicS Project. 2021. Available online: <https://www.researchgate.net/project/EUPHRESCO-Basic-substances-as-an-environmentally-friendly-alternative-to-synthetic-pesticides-for-plant-protection-BasicS> (accessed on 1 December 2021).

17. Marchand, P.A.; Davillerd, Y.; Riccioni, L.; Sanzani, S.M.; Horn, N.; Matyjaszczyk, E.; Golding, J.; Roberto, S.R.; Mattiuz, B.-H.; Xu, D.; et al. BasicS, an euphresco international network on renewable natural substances for durable crop protection products. *Chron. Bioresour. Manag.* **2021**, *5*, 77–80.
18. Romanazzi, G.; Feliziani, E.; Sivakumar, D. Chitosan, a biopolymer with triple action on postharvest decay of fruit and vegetables: Eliciting, antimicrobial and film-forming properties. *Front. Microbiol.* **2018**, *9*, 2745. [CrossRef]
19. European Commission. *Working Document on the Procedure for Application of Basic Substances to be Approved in Compliance with Article 23 of Regulation (EC) No 1107/2009*; SANCO/10363/2012 rev. 10; European Commission: Brussels, Belgium, 25 January 2021.
20. EU. EU Pesticides Database (v.2.2) Search Active Substances, Safeners and Synergists (europa.eu). 2021. Available online: <https://ec.europa.eu/food/plant/pesticides/eu-pesticides-database/active-substances/?event=search.as> (accessed on 23 May 2022).
21. Matyjaszczyk, E. Plant protection means used in organic farming throughout the European Union. *Pest Manag. Sci.* **2018**, *74*, 505–510. [CrossRef]
22. Marchand, P.A. Basic substances under EU pesticide regulation: An opportunity for organic production? *Org. Farming* **2017**, *3*, 16–19. [CrossRef]
23. EU. Commission Implementing Regulation (EU) amending and correcting Implementing Regulation (EU) 2021/1165 authorising certain products and substances for use in organic production and establishing their lists. *Official J. Eur. Union* **2022**, in press.
24. Marchand, P.A.; Isambert, C.A.; Jonis, M.; Parveaud, C.-E.; Chevolon, M.; Gomez, C.; Lambion, J.; Ondet, S.J.; Aveline, N.; Molot, B.; et al. Evaluation des caractéristiques et de l'intérêt agronomique de préparations simples de plantes, pour des productions fruitières, légumières et viticoles économes en intrants. *Innov. Agron.* **2014**, *34*, 83–96.
25. CASDAR 4P, Protéger les Plantes Par les Plantes. 2014. Available online: <https://ecophytopic.fr/recherche-innovation/protoger/projet-4p> (accessed on 23 May 2022).
26. COPPEREPLACE. 2015. Available online: <https://coppereplace.com/fr/project-coppereplace/> (accessed on 23 May 2022).
27. Robin, N.; Bruyere, J. Traitements de semences: Contrôler la carie. In *Actes Journée Technique*; ITAB: Paris, France, 2012; pp. 31–40. Available online: <http://www.itab.asso.fr/downloads/actes%20suite/carie-actes2012.pdf> (accessed on 23 May 2022).
28. Arnault, I.; Bardin, M.; Ondet, S.; Furet, A.; Chovelon, M.; Kasprick, A.-C.; Marchand, P.; Clerc, H.; Davy, M.; Roy, G.; et al. Utilisation de micro-doses de sucres en protection des cultures. *Innov. Agron.* **2015**, *46*, 1–10.
29. Arnault, I.; Aveline, N.; Bardin, M.; Brisset, M.N.; Carriere, J.; Chovelon, M.; Delanoue, G.; Furet, A.; Frérot, B.; Lambion, J.; et al. Optimisation des stratégies de biocontrôle par la stimulation de l'immunité des plantes avec des applications d'infra-doses de sucres simples. *Innov. Agron.* **2021**, *82*, 411–423. [CrossRef]
30. Vidal, R.; Muchembled, J.; Deweer, C.; Tournant, L.; Corroyer, N.; Flammier, S. Évaluation de l'intérêt de l'utilisation d'huiles essentielles dans des stratégies de protection des cultures. *Innov. Agron.* **2018**, *63*, 1–20.
31. Compo Expert. Report 2015. In *LA PUGERE, 2011 Pear Psylla [Psylla piri], The TALC Efficiency Evaluation in a Preventive Control Strategy of the Pear Psylla Year*; Station d'experimentation La Pugere: Mallemort, France, 2015; pp. 1–68.
32. CA (Chambre d'Agriculture de l'Aude). *Réduction des coûts en viticulture. Produits Alternatifs: Lactosérum [Reducing Costs in Viticulture: Alternative Crop Protection: Lactoserum]*; Technical Report; Chambre d'Agriculture de l'Aude (CA11): Carcas-sonne, France, 2011; 123p.
33. Duriez, J.M. *Le phosphate di-ammonium, un attractant de la mouche de l'olive* Journées Techniques Intrants; ITAB: Paris, France, 2016. Available online: http://www.itab.asso.fr/downloads/jt-intrants-2016/10_duriez-afidol-pda.pdf (accessed on 23 May 2022).
34. Montag, J.; Schreiber, L.; Schönherr, J. The post-infection activity of hydrated lime against conidia of *Venturia inaequalis*. In Proceedings of the Ecofruit-12th International Conference on Cultivation Technique and Phytopathological Problems in Organic Fruit-Growing, Weinsberg, Germany, 31 January–2 February 2006.
35. Romanazzi, G.; Piancatelli, S.; D'Ignazi, G.; Moumni, M. Innovative approaches to grapevine downy mildew management on large and commercial scale. In Proceedings of the 9th International Workshop on Grapevine Downy and Powdery Mildew, Cremona, Italy, 20–22 July 2022.
36. Romanazzi, G.; Moumni, M. New challenges in preventing and managing fresh fruit loss and waste. In Proceedings of the VI International Symposium on Postharvest Pathology, Limassol, Cyprus, 29 May–2 June 2022.
37. D'Ortenzio, A.L.; Fava, G.; Mazzoni, S.; Acciarri, P.; Baronciani, L.; Ceredi, G.; Romanazzi, G. Postharvest application of natural compounds and biocontrol agents to manage brown rot of stone fruit. In Proceedings of the VI International Symposium on Postharvest Pathology, Limassol, Cyprus, 29 May–2 June 2022.
38. Piancatelli, S.; Moumni, M.; Binni, T.; Giardini, D.; Profili, R.; Napoleoni, D.; Morbidelli, M.; Fabbri, G.; Piersanti, G.; Nardi, S.; et al. Impiego di sostanze di origine naturale e a basso impatto ambientale nella protezione del cavolo cappuccio da seme. *Giornate Fitopatol.* **2022**; in press.
39. Park, D.W.; Yang, Y.S.; Lee, Y.U.; Han, S.J.; Kim, H.J.; Kim, S.H.; Kim, J.P.; Cho, S.J.; Lee, D.; Song, N.; et al. Pesticide residues and risk assessment from monitoring programs in the largest production area of leafy vegetables in South Korea: A 15-year study. *Foods* **2021**, *10*, 425. [CrossRef]
40. El-Nahhal, I.; El-Nahhal, Y. Pesticide residues in drinking water, their potential risk to human health and removal options. *J. Environ. Manag.* **2021**, *299*, 113611. [CrossRef]
41. Hejazi, M.; Grant, J.H.; Peterson, E. Trade impact of maximum residue limits in fresh fruits and vegetables. *Food Policy* **2022**, *106*, 102203. [CrossRef]

42. Romanazzi, G.; Feliziani, E.; Baños, S.B.; Sivakumar, D. Shelf life extension of fresh fruit and vegetables by chitosan treatment. *Crit. Rev. Food Sci. Nutr.* **2017**, *57*, 579–601. [[CrossRef](#)]
43. Romanazzi, G.; Sanzani, S.M.; Bi, Y.; Tian, S.; Gutierrez-Martinez, P.; Alkan, N. Induced resistance to control postharvest decay of fruit and vegetables. *Postharvest Biol. Technol.* **2016**, *122*, 82–94. [[CrossRef](#)]
44. Robin, D.; Marchand, P.A. Expansion of the low-risk substances in the framework of the European Pesticide Regulation (EC) No. 1107/2009. *Eur. J. Risk Regul.* **2022**, 1–20. [[CrossRef](#)]
45. Marchand, P.A. Basic substances: An opportunity for approval of low-concern substances under EU pesticide regulation. *Pest Manag. Sci.* **2015**, *71*, 1197–1200. [[CrossRef](#)]
46. Marchand, P.A. Basic substances under EC 1107/2009 phytochemical regulation: Experience with non-biocide and food products as biorationals. *J. Plant Prot. Res.* **2016**, *56*, 312–318. [[CrossRef](#)]
47. Lykogianni, M.; Bempelou, E.; Karamaouna, F.; Aliferis, K.A. Do pesticides promote or hinder sustainability in agriculture? The challenge of sustainable use of pesticides in modern agriculture. *Sci. Total Environ.* **2021**, *795*, 148625. [[CrossRef](#)] [[PubMed](#)]
48. Romanazzi, G.; Mancini, V.; Foglia, R.; Marcolini, D.; Kavari, M.; Piancatelli, S. Use of chitosan and other natural compounds alone or in different strategies with copper hydroxide for control of grapevine downy mildew. *Plant Dis.* **2021**, *105*, 3261–3268. [[CrossRef](#)] [[PubMed](#)]
49. EU. *Working Document on the Procedure for Application of Basic Substances to be Approved in Compliance with Article 23 of Regulation (EC) No. 1107/2009 SANCO/10363/2012*; (rev. 9 of 21 March 2014); EU: Brussels, Belgium, 2012.
50. EU. Commission Implementing Regulation (EU) 2022/456 of 21 March 2022 approving the basic substance chitosan in accordance with Regulation (EC) No 1107/2009 of the European Parliament and of the Council concerning the placing of plant protection products on the market, and amending the Annex to Implementing Regulation (EU) No 540/2011. *OJ* **2022**, *L93*, 138–141.
51. Marchand, P.A.; Carrière, J. *Guide Pédagogique: Constituer un dossier d'approbation de substance naturelle au règlement (CE) n° 1107/2009*; ITAB: Paris, France, 2012. [[CrossRef](#)]
52. Vekemans, M.C.; Marchand, P.A. The fate of the Biocontrol agents under the European phytopharmaceutical regulation: A hindering for approval botanicals as new active substances? *Environ. Sci. Pollut. Res.* **2020**, *27*, 39879–39887. [[CrossRef](#)]