

POSTER PRESENTATION

Open Access

Mutations in the dynein assembly factor PF22 (DNAAF3) cause primary ciliary dyskinesia with absent dynein arms

M Schmidts^{1*}, J Freshour², NT Loges³, A Dritsoula¹, D Antony¹, RA Hirst⁴, C O'Callaghan⁴, H Blau⁵, H Olbrich³, T Yagi⁶, H Mussaffi⁵, EMK Chung⁷, H Omran³, DR Mitchell², H Mitchison¹

From First International Cilia in Development and Disease Scientific Conference (2012) London, UK. 16-18 May 2012

The genetic disorder primary ciliary dyskinesia (PCD) arises from dysmotility of cilia in the respiratory tract, brain ventricles, oviduct and the embryonic node. Patients have chronic obstructive pulmonary disease, reduced fertility and situs abnormalities. PCD is genetically heterogeneous with 12 genes causing ~40% of all cases, two encoding proteins (KTU, LRRC50) involved in cytosolic axonemal dynein co-assembly. We have identified mutations in the *C19ORF51* gene located within a previously mapped PCD locus. *C19ORF51* encodes a protein orthologous to PF22, a Chlamydomonas protein involved in the cytoplasmic assembly of outer dynein arms preceding their import into the axoneme. Chlamydomonas pf22 cells display a disturbance in their cytoplasm of dynein heavy chain stability and the co-assembly of heavy with intermediate chains, both essential for dynein arm assembly. PF22 appears to act downstream of KTU and LRRC50 in the dynein preassembly pathway. *PF22* knockdown in zebrafish causes a loss of dynein arms, cilia dysmotility, and a typical ciliopathy phenotype with axis curvature, pronephric cysts, hydrocephalus and situs inversus. We propose the existence of a conserved multi-step pathway for formation of assembly-competent dynein complexes, and that *PF22* (now renamed *DNAAF3*, “dynein axonemal assembly factor 3”) mutations cause PCD with situs inversus due to deficient cytoplasmic dynein assembly which in turn leads to missing outer arms in the axoneme.

Author details

¹Molecular Medicine Unit, UCL Institute of Child Health, UK.
²Cell & Developmental Biology, SUNY Upstate Medical University, USA.

* Correspondence: m.schmidts@ich.ucl.ac.uk

¹Molecular Medicine Unit, UCL Institute of Child Health, UK
Full list of author information is available at the end of the article

³Department of Pediatrics and Adolescent Medicine, University Hospital Freiburg and University Hospital Muenster, Germany. ⁴Department of Infection Immunity and Inflammation, University of Leicester, UK. ⁵Pulmonary Unit, Schneider Children's Medical Center of Israel, Israel. ⁶Department of Cell Biology and Anatomy, Graduate School of Medicine, University of Tokyo, Japan. ⁷General and Adolescent Paediatric Unit, UCL Institute of Child Health, UK.

Published: 16 November 2012

doi:10.1186/2046-2530-1-S1-P101

Cite this article as: Schmidts *et al.*: Mutations in the dynein assembly factor PF22 (DNAAF3) cause primary ciliary dyskinesia with absent dynein arms. *Cilia* 2012 **1**(Suppl 1):P101.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

