

Ethyl 7-amino-1-cyclopropyl-6-fluoro-8-methoxy-4-oxo-1,4-dihydroquinoline-3-carboxylate monohydrate

Jia Pan,^a Li Yang,^a Zhi-Hua Mao^b and Ling-Ling Weng^{a*}

^aDepartment of Medicinal Chemistry, West China School of Pharmacy, Sichuan University, Chengdu 610041, People's Republic of China, and ^bThe Center for Testing and Analysis, Sichuan University, Chengdu 610064, People's Republic of China

Correspondence e-mail: lingling.weng@yahoo.com.cn


Received 17 January 2008; accepted 23 January 2008

Key indicators: single-crystal X-ray study; $T = 291\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.044; wR factor = 0.126; data-to-parameter ratio = 12.8.

In the title compound, $\text{C}_{16}\text{H}_{17}\text{FN}_2\text{O}_4\cdot\text{H}_2\text{O}$, the dihedral angle between the heterocyclic ring and the benzene ring is $5.77(9)^\circ$, that between the heterocycle and the ethoxy-carbonyl plane is $15.5(1)^\circ$, and that between the heterocyclic ring and the cyclopropane ring is $67.75(13)^\circ$. In the crystal structure, molecules are linked into a ribbon-like structure along the c axis by $\text{N}-\text{H}\cdots\text{O}$ and $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds.

Related literature

For general background, see: Fujita & Chiba (1998).



Experimental

Crystal data

$\text{C}_{16}\text{H}_{17}\text{FN}_2\text{O}_4\cdot\text{H}_2\text{O}$
 $M_r = 338.33$

Monoclinic, $P2_1/n$
 $a = 10.096(4)\text{ \AA}$

$b = 14.699(5)\text{ \AA}$
 $c = 11.028(6)\text{ \AA}$
 $\beta = 94.26(4)^\circ$
 $V = 1632.0(12)\text{ \AA}^3$
 $Z = 4$

Mo $K\alpha$ radiation
 $\mu = 0.11\text{ mm}^{-1}$
 $T = 291(2)\text{ K}$
 $0.45 \times 0.42 \times 0.39\text{ mm}$

Data collection

Enraf-Nonius CAD-4
diffractometer
Absorption correction: none
3157 measured reflections
3009 independent reflections

1741 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.007$
3 standard reflections
every 300 reflections
intensity decay: 0.8%

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.043$
 $wR(F^2) = 0.125$
 $S = 1.04$
3009 reflections
235 parameters

H atoms treated by a mixture of
independent and constrained
refinement
 $\Delta\rho_{\text{max}} = 0.20\text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.19\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N1—H1N1 \cdots O1W ⁱ	0.91 (3)	2.15 (3)	2.930 (3)	143 (2)
N1—H2N1 \cdots O4 ⁱⁱ	0.85 (3)	2.33 (3)	3.061 (3)	144 (2)
O1W—H1W \cdots O2 ⁱⁱⁱ	0.84 (3)	2.13 (3)	2.916 (3)	155 (3)
O1W—H2W \cdots O2	0.91 (3)	1.96 (3)	2.864 (3)	171 (3)

Symmetry codes: (i) $-x + 1, -y + 1, -z + 2$; (ii) $x, y, z + 1$; (iii) $-x + 1, -y + 1, -z + 1$.

Data collection: *DIFRAC* (Gabe & White, 1993); cell refinement: *DIFRAC*; data reduction: *NRCVAX* (Gabe *et al.*, 1989); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *SHELXL97*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CI2556).

References

- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Fujita, M. & Chiba, K. (1998). *Chem. Pharm. Bull.* **46**, 631–638.
- Gabe, E. J., Le Page, Y., Charland, J.-P., Lee, F. L. & White, P. S. (1989). *J. Appl. Cryst.* **22**, 384–387.
- Gabe, E. J. & White, P. S. (1993). *DIFRAC*. American Crystallographic Association Pittsburgh Meeting Abstract PA 104.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.

supplementary materials

Acta Cryst. (2008). E64, o527 [doi:10.1107/S1600536808002547]

Ethyl 7-amino-1-cyclopropyl-6-fluoro-8-methoxy-4-oxo-1,4-dihydroquinoline-3-carboxylate monohydrate

J. Pan, L. Yang, Z.-H. Mao and L.-L. Weng

Comment

Quinolone antibacterials were found several decades ago, and some excellent antibacterials have been developed and used widely now (Fujita & Chiba, 1998). An interest in search of more potent antibacterial agents led us to design and synthesis a new type of quinoline derivatives. The title compound is one of the key intermediates and we report here its crystal structure.

The pyridinone ring is planar to within ± 0.057 (2) Å (Fig. 1). The dihedral angle between the pyridine and benzene rings is 5.77 (9) $^\circ$, and that between the pyridine and carboxylate plane is 15.5 (1) $^\circ$. In the crystal structure, the molecules are linked into a ribbon like structure along the *c* axis (Fig. 2) by N—H \cdots O and O—H \cdots O hydrogen bonds (Table 1).

Experimental

Ethyl 7-azido-1-cyclopropyl-6-fluoro-8-methoxyl-4-oxo-1,4-dihydroquinoline-3- carboxylate (2 g, 5.8 mmol), 5% Pd/C (0.4 g) were suspended in methanol (20 ml) and the mixture was hydrogenated at 303 K for 6 h. The reaction mixture was then filtered and concentrated under vacuum. The residue obtained was purified by silica gel chromatography. Single crystals suitable for X-ray analysis were obtained by slow evaporation of a acetyl acetate-chloroform (1.2:1 *v/v*) solution at room temperature.

Refinement

The water H atoms were located in a difference Fourier map and refined isotropically. The remaining H atoms were placed in the calculated positions [C—H = 0.93 (aromatic) and 0.96 Å (methyl)] and refined in the riding-model approximation, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{aromatic-C})$ and $1.5U_{\text{eq}}(\text{methyl-C})$.

Figures




Fig. 1. The molecular structure of the title compound, showing the atomic numbering scheme. Displacement ellipsoids drawn at the 30% probability level. The dashed line indicates a hydrogen bond.



Fig. 2. Crystal packing of the title compound, viewed down the *a* axis. Hydrogen bonds are shown as dashed lines.

supplementary materials

Ethyl 7-amino-1-cyclopropyl-6-fluoro-8-methoxy-4-oxo-1,4-dihydroquinoline- 3-carboxylate monohydrate

Crystal data

C ₁₆ H ₁₇ FN ₂ O ₄ ·H ₂ O	$F_{000} = 712$
$M_r = 338.33$	$D_x = 1.377 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation
Hall symbol: -P 2yn	$\lambda = 0.71073 \text{ \AA}$
$a = 10.096 (4) \text{ \AA}$	Cell parameters from 24 reflections
$b = 14.699 (5) \text{ \AA}$	$\theta = 4.5\text{--}7.4^\circ$
$c = 11.028 (6) \text{ \AA}$	$\mu = 0.11 \text{ mm}^{-1}$
$\beta = 94.26 (4)^\circ$	$T = 291 (2) \text{ K}$
$V = 1632.0 (12) \text{ \AA}^3$	Block, yellow
$Z = 4$	$0.45 \times 0.42 \times 0.39 \text{ mm}$

Data collection

Enraf-Nonius CAD-4 diffractometer	$R_{\text{int}} = 0.007$
Radiation source: fine-focus sealed tube	$\theta_{\text{max}} = 25.4^\circ$
Monochromator: graphite	$\theta_{\text{min}} = 2.3^\circ$
$T = 291(2) \text{ K}$	$h = -12 \rightarrow 12$
$\omega/2\theta$ scans	$k = 0 \rightarrow 17$
Absorption correction: none	$l = -4 \rightarrow 13$
3157 measured reflections	3 standard reflections
3009 independent reflections	every 300 reflections
1741 reflections with $I > 2\sigma(I)$	intensity decay: 0.8%

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: mixed
$R[F^2 > 2\sigma(F^2)] = 0.043$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.125$	$w = 1/[\sigma^2(F_o^2) + (0.0627P)^2 + 0.1314P]$
$S = 1.04$	where $P = (F_o^2 + 2F_c^2)/3$
3009 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
235 parameters	$\Delta\rho_{\text{max}} = 0.20 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\text{min}} = -0.19 \text{ e \AA}^{-3}$
	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\text{sigma}(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
F1	0.51891 (13)	0.59935 (11)	1.01305 (12)	0.0667 (4)
O1	0.07128 (14)	0.68291 (10)	1.03507 (12)	0.0451 (4)
O2	0.33328 (14)	0.56826 (12)	0.58213 (13)	0.0519 (4)
O3	-0.04405 (16)	0.62649 (11)	0.41364 (13)	0.0552 (5)
O4	0.16963 (18)	0.62837 (18)	0.37733 (15)	0.1007 (9)
N1	0.3174 (3)	0.65375 (16)	1.14712 (18)	0.0514 (5)
H1N1	0.385 (3)	0.6210 (17)	1.184 (2)	0.068 (9)*
H2N1	0.248 (3)	0.6504 (18)	1.186 (2)	0.071 (9)*
N2	0.01966 (16)	0.64366 (12)	0.77689 (13)	0.0374 (4)
C1	0.3794 (2)	0.59542 (15)	0.83243 (19)	0.0436 (5)
H1	0.4507	0.5774	0.7894	0.052*
C2	0.3970 (2)	0.61005 (15)	0.95442 (19)	0.0433 (5)
C3	0.2938 (2)	0.63708 (14)	1.02489 (17)	0.0397 (5)
C4	0.1693 (2)	0.65127 (13)	0.96495 (17)	0.0352 (5)
C5	0.1457 (2)	0.63418 (13)	0.83915 (17)	0.0333 (5)
C6	0.2533 (2)	0.60753 (13)	0.77155 (17)	0.0358 (5)
C7	0.0710 (3)	0.78037 (18)	1.0445 (3)	0.0786 (9)
H7A	0.1564	0.8008	1.0778	0.118*
H7B	0.0041	0.7990	1.0968	0.118*
H7C	0.0523	0.8064	0.9653	0.118*
C8	0.2389 (2)	0.59719 (14)	0.63888 (17)	0.0378 (5)
C9	0.1107 (2)	0.62227 (14)	0.58234 (17)	0.0401 (5)
C10	0.0092 (2)	0.64129 (14)	0.65419 (17)	0.0397 (5)
H10	-0.0738	0.6536	0.6152	0.048*
C11	-0.1031 (2)	0.65083 (17)	0.83913 (18)	0.0449 (6)
H11	-0.1233	0.7109	0.8716	0.054*
C12	-0.2192 (2)	0.5947 (2)	0.7935 (2)	0.0723 (9)
H12A	-0.2086	0.5554	0.7242	0.087*
H12B	-0.3070	0.6208	0.7971	0.087*
C13	-0.1436 (2)	0.5712 (2)	0.9108 (2)	0.0627 (7)
H13A	-0.1857	0.5830	0.9854	0.075*
H13B	-0.0873	0.5177	0.9126	0.075*
C14	0.0853 (2)	0.62535 (17)	0.44863 (19)	0.0508 (6)

supplementary materials

C15	-0.0802 (3)	0.6314 (2)	0.28436 (19)	0.0693 (8)
H15A	-0.0528	0.5763	0.2449	0.083*
H15B	-0.0360	0.6826	0.2493	0.083*
C16	-0.2257 (2)	0.64228 (17)	0.2653 (2)	0.0589 (7)
H16A	-0.2687	0.5904	0.2979	0.088*
H16B	-0.2506	0.6471	0.1798	0.088*
H16C	-0.2523	0.6964	0.3058	0.088*
O1W	0.5509 (2)	0.44950 (16)	0.65115 (16)	0.0715 (6)
H1W	0.606 (3)	0.446 (2)	0.598 (3)	0.091 (11)*
H2W	0.486 (3)	0.488 (2)	0.621 (3)	0.097 (11)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
F1	0.0396 (8)	0.1032 (12)	0.0559 (8)	0.0081 (7)	-0.0058 (6)	-0.0147 (8)
O1	0.0457 (9)	0.0567 (10)	0.0346 (7)	0.0092 (8)	0.0143 (6)	-0.0021 (7)
O2	0.0422 (9)	0.0766 (11)	0.0388 (8)	0.0138 (8)	0.0163 (7)	-0.0026 (7)
O3	0.0469 (10)	0.0891 (13)	0.0303 (8)	0.0030 (9)	0.0083 (7)	0.0013 (7)
O4	0.0523 (12)	0.215 (3)	0.0373 (10)	0.0254 (13)	0.0193 (9)	0.0150 (12)
N1	0.0482 (13)	0.0683 (15)	0.0378 (11)	0.0021 (11)	0.0036 (10)	-0.0036 (10)
N2	0.0332 (10)	0.0504 (11)	0.0298 (9)	0.0072 (8)	0.0109 (7)	0.0037 (7)
C1	0.0340 (12)	0.0527 (14)	0.0451 (13)	0.0016 (10)	0.0109 (10)	-0.0051 (10)
C2	0.0337 (12)	0.0528 (14)	0.0428 (12)	0.0019 (10)	-0.0008 (10)	-0.0034 (10)
C3	0.0449 (13)	0.0402 (12)	0.0344 (11)	-0.0043 (10)	0.0063 (9)	0.0002 (9)
C4	0.0364 (12)	0.0368 (11)	0.0340 (10)	0.0013 (9)	0.0121 (9)	0.0001 (9)
C5	0.0331 (11)	0.0336 (11)	0.0341 (10)	0.0011 (9)	0.0075 (8)	0.0031 (8)
C6	0.0337 (12)	0.0373 (12)	0.0373 (11)	-0.0026 (9)	0.0085 (9)	-0.0001 (8)
C7	0.097 (2)	0.0622 (18)	0.0805 (19)	0.0211 (17)	0.0311 (17)	-0.0148 (15)
C8	0.0384 (12)	0.0414 (12)	0.0355 (11)	-0.0007 (10)	0.0145 (9)	-0.0008 (9)
C9	0.0378 (12)	0.0518 (14)	0.0315 (11)	0.0038 (10)	0.0093 (9)	0.0018 (9)
C10	0.0350 (12)	0.0519 (14)	0.0327 (10)	0.0079 (10)	0.0069 (9)	0.0078 (9)
C11	0.0320 (12)	0.0659 (15)	0.0385 (11)	0.0082 (11)	0.0129 (9)	0.0041 (10)
C12	0.0403 (15)	0.130 (3)	0.0480 (14)	-0.0086 (15)	0.0124 (12)	0.0057 (15)
C13	0.0492 (15)	0.094 (2)	0.0469 (13)	-0.0127 (14)	0.0168 (11)	0.0119 (13)
C14	0.0444 (14)	0.0763 (17)	0.0331 (11)	0.0094 (12)	0.0123 (10)	0.0023 (11)
C15	0.0627 (18)	0.114 (2)	0.0311 (12)	0.0122 (16)	0.0048 (12)	-0.0030 (13)
C16	0.0602 (17)	0.0649 (17)	0.0508 (14)	0.0107 (13)	-0.0012 (12)	-0.0041 (12)
O1W	0.0600 (13)	0.1157 (18)	0.0396 (9)	0.0301 (12)	0.0077 (9)	-0.0035 (10)

Geometric parameters (\AA , $^\circ$)

F1—C2	1.356 (2)	C7—H7B	0.96
O1—C4	1.381 (2)	C7—H7C	0.96
O1—C7	1.436 (3)	C8—C9	1.442 (3)
O2—C8	1.252 (2)	C9—C10	1.371 (3)
O3—C14	1.334 (3)	C9—C14	1.479 (3)
O3—C15	1.447 (3)	C10—H10	0.93
O4—C14	1.202 (3)	C11—C13	1.486 (3)
N1—C3	1.373 (3)	C11—C12	1.490 (3)

N1—H1N1	0.91 (3)	C11—H11	0.98
N1—H2N1	0.85 (3)	C12—C13	1.492 (4)
N2—C10	1.350 (2)	C12—H12A	0.97
N2—C5	1.408 (3)	C12—H12B	0.97
N2—C11	1.465 (3)	C13—H13A	0.97
C1—C2	1.361 (3)	C13—H13B	0.97
C1—C6	1.405 (3)	C15—C16	1.477 (3)
C1—H1	0.9300	C15—H15A	0.97
C2—C3	1.403 (3)	C15—H15B	0.97
C3—C4	1.391 (3)	C16—H16A	0.96
C4—C5	1.413 (3)	C16—H16B	0.96
C5—C6	1.418 (3)	C16—H16C	0.96
C6—C8	1.467 (3)	O1W—H1W	0.84 (3)
C7—H7A	0.96	O1W—H2W	0.91 (3)
C4—O1—C7	112.45 (17)	N2—C10—C9	125.34 (19)
C14—O3—C15	117.11 (18)	N2—C10—H10	117.3
C3—N1—H1N1	114.2 (16)	C9—C10—H10	117.3
C3—N1—H2N1	113.3 (18)	N2—C11—C13	117.9 (2)
H1N1—N1—H2N1	112 (2)	N2—C11—C12	118.3 (2)
C10—N2—C5	119.16 (17)	C13—C11—C12	60.20 (17)
C10—N2—C11	117.75 (17)	N2—C11—H11	116.3
C5—N2—C11	123.00 (15)	C13—C11—H11	116.3
C2—C1—C6	119.99 (19)	C12—C11—H11	116.3
C2—C1—H1	120.0	C11—C12—C13	59.79 (16)
C6—C1—H1	120.0	C11—C12—H12A	117.8
F1—C2—C1	120.05 (19)	C13—C12—H12A	117.8
F1—C2—C3	117.01 (18)	C11—C12—H12B	117.8
C1—C2—C3	122.9 (2)	C13—C12—H12B	117.8
N1—C3—C4	121.6 (2)	H12A—C12—H12B	114.9
N1—C3—C2	120.8 (2)	C11—C13—C12	60.01 (16)
C4—C3—C2	117.47 (18)	C11—C13—H13A	117.8
O1—C4—C3	116.36 (17)	C12—C13—H13A	117.8
O1—C4—C5	122.29 (18)	C11—C13—H13B	117.8
C3—C4—C5	121.36 (18)	C12—C13—H13B	117.8
N2—C5—C4	122.81 (18)	H13A—C13—H13B	114.9
N2—C5—C6	118.12 (17)	O4—C14—O3	122.4 (2)
C4—C5—C6	119.06 (18)	O4—C14—C9	125.0 (2)
C1—C6—C5	119.07 (18)	O3—C14—C9	112.52 (18)
C1—C6—C8	118.85 (18)	O3—C15—C16	108.7 (2)
C5—C6—C8	121.99 (18)	O3—C15—H15A	109.9
O1—C7—H7A	109.5	C16—C15—H15A	109.9
O1—C7—H7B	109.5	O3—C15—H15B	109.9
H7A—C7—H7B	109.5	C16—C15—H15B	109.9
O1—C7—H7C	109.5	H15A—C15—H15B	108.3
H7A—C7—H7C	109.5	C15—C16—H16A	109.5
H7B—C7—H7C	109.5	C15—C16—H16B	109.5
O2—C8—C9	124.20 (18)	H16A—C16—H16B	109.5
O2—C8—C6	120.70 (19)	C15—C16—H16C	109.5
C9—C8—C6	115.09 (17)	H16A—C16—H16C	109.5

supplementary materials


C10—C9—C8	119.22 (18)	H16B—C16—H16C	109.5
C10—C9—C14	119.2 (2)	H1W—O1W—H2W	107 (3)
C8—C9—C14	121.54 (18)		

Hydrogen-bond geometry (\AA , $^\circ$)

$D\text{—H}\cdots A$	$D\text{—H}$	$\text{H}\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
N1—H1N1…O1W ⁱ	0.91 (3)	2.15 (3)	2.930 (3)	143 (2)
N1—H2N1…O4 ⁱⁱ	0.85 (3)	2.33 (3)	3.061 (3)	144 (2)
O1W—H1W…O2 ⁱⁱⁱ	0.84 (3)	2.13 (3)	2.916 (3)	155 (3)
O1W—H2W…O2	0.91 (3)	1.96 (3)	2.864 (3)	171 (3)

Symmetry codes: (i) $-x+1, -y+1, -z+2$; (ii) $x, y, z+1$; (iii) $-x+1, -y+1, -z+1$.

Fig. 1



supplementary materials

Fig. 2

