



■ Original Article

# The Effect of Access to Primary Care Physicians on Avoidable Hospitalizations: A Time Series Study in Rural Areas of Tehran Province, Iran

Arash Rashidian<sup>1</sup>, Sedigheh Salavati<sup>2,\*</sup>, Hanan Hajimahmoodi<sup>3</sup>

<sup>1</sup>Department of Health Management and Economics, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

<sup>2</sup>Department of Public Health, School of Nursery, Maragheh University of Medical Sciences, Maragheh, Iran

<sup>3</sup>Department of Family Physician Program, Iran Health Insurance Organization, Tehran, Iran

**Background:** Avoidable hospitalizations (AHs) are defined as hospitalizations that could have been prevented through timely and effective services. AHs are, therefore, an indicator used to evaluate the access and effectiveness of primary health care services.

**Methods:** A retrospective time-series study spanning 8 years (2006–2013) was conducted to determine the relationship between AHs and gender, age, and access to primary health care physicians in rural areas in Tehran province, the capital of Iran. The total number of avoidable hospitalizations was 22,570; logistic regression was estimated for each year separately.

**Results:** Total hospitalizations and AHs increased during the study period, especially during the first 3 years of the study. AHs, as a percentage of total hospitalizations, did not change significantly throughout the study years. This value was 22.3% during the first year of study and varied between 17% and 19.6% from 2007 to 2013. No statistically significant relationship was seen between AH occurrence and access to a physician during the study years.

**Conclusion:** Increasing access to primary health care physicians cannot necessarily result in decreased AHs. Considering the factors influencing AHs while designing and implementing the family physicians program is important to achieve the expected results regarding the effectiveness of primary health care services.

**Keywords:** Primary Health Care; Physicians; Avoidable Hospitalization; Rural Health; Health Insurance

Received: March 26, 2019, Revised: July 8, 2019, Accepted: August 30, 2019

\*Corresponding Author: Sedigheh Salavati <https://orcid.org/0000-0001-9206-1553>

Tel: +98-4137256901, Fax: +98-4137256901, E-mail: [ssalavati@mrqums.ac.ir](mailto:ssalavati@mrqums.ac.ir)

## INTRODUCTION

Avoidable hospitalizations are those hospitalizations that could have been prevented through timely and effective primary health care services.<sup>1)</sup> Adequate accessibility and quality of primary health care services reportedly result in decreased avoidable hospitalizations.<sup>2-5)</sup> Hospitalizations for chronic conditions such as diabetes, asthma, and hypertension, and acute diseases, such as pneumonia and complicated appendicitis, are considered as avoidable hospitalizations.<sup>1,6)</sup> Other factors, such as demographic characteristics, socioeconomic status, health status, and biology, can also affect hospitalizations.<sup>3,5,7,8)</sup> For instance, differences between men and women in biology, health status, and healthcare-seeking behaviors can affect hospitalizations.<sup>9-11)</sup> Age is a frequently reported factor significantly related to hospitalization, and its influence should be considered alongside gender.<sup>5,12)</sup> Simultaneously, exploring the relationship between the population's demographic characteristics, access to primary health care services, and avoidable hospitalizations can provide policymakers with a comprehensive picture of the effectiveness of policy interventions related to primary health care and evidence regarding the performance of first-level healthcare services.

In the Islamic Republic of Iran, public, private, quasi-public, and charity sectors provide healthcare services. Over 93% of second and third level healthcare services are provided by clinics, centers, and hospitals affiliated to the Iran Ministry of Health and Medical Education (MOHME), with other centers affiliated to the Social Security Organization and the private sector.<sup>13)</sup>

Primary health care networks provide the population with access to public primary health care services, free of charge.<sup>14)</sup> Actions towards universal health coverage, first introduced in 2005, resulted in the implementation of the rural family physician program in rural areas and cities with a population of less than 20,000. This reform led to primary care services—provided by family physicians—becoming available for rural residents.<sup>15)</sup> Some physicians and midwives were added to the primary health care network over three years as part of the program that was intended to improve access to a more comprehensive, preventive, and outpatient care. Family physicians manage patients referred to them by the “behvarz,” a community health worker at first-line of service that provides and offers primary health care services, including vaccinations, prenatal and postnatal care, growth monitoring, family planning, management of common infectious and chronic conditions, and environmental health services.

Access to primary health care physicians is unequal across rural areas and depends on an area's geographic distance from the main village. The main villages with rural health care centers have a resident primary health care physician. However, in smaller or farther villages, typically those with health houses instead of healthcare centers, resident physicians are lacking, and primary health care physicians provide services once or twice a month, with an accompanying health team. Residents of satellite villages are forced to travel to villages with health houses to use their primary health care services. Remote villag-

es have the lowest access to primary health care physicians, and can only use the services of physicians who travel to these villages once a month. Therefore, avoidable hospitalizations in the rural areas of Iran's health system can be an acceptable indicator of access to primary health care physicians and the effectiveness of their services.

The effect of access to family physicians on total and avoidable hospitalizations in Iran has been explored in two previous studies,<sup>16,17)</sup> but there have been no studies on the differences in the impact of the family physician reform on avoidable hospitalizations in the first year of the program compared to the following years. This study aimed to simultaneously explore the relationship between access to primary health care physicians, demographic characteristics, and avoidable hospitalizations among the rural population in different years following the implementation of the family physician program.

## METHODS

A retrospective time-series study was used to determine the relationship between avoidable hospitalizations and age, gender, and access to a primary health care physician. Data on rural resident's hospitalizations during the 8 years following the family physician program implementation (2006–2013) were gathered in the rural areas of Tehran province, the capital of Iran. The list of diagnostic codes related to avoidable hospitalizations had not been previously determined in any study in Iran; therefore, the literature was reviewed to determine the avoidable hospitalization codes.<sup>5,18)</sup> Experts were consulted regarding the correctness of these codes and ways to avoid admissions falling in the category of avoidable hospitalizations. These diagnostic codes were then used as markers of avoidable hospitalizations in other studies in Iran.<sup>16,17)</sup> The avoidable hospitalizations' diagnostic codes used in this study are presented in Appendix 1 and are considered as determinants of the effectiveness and accessibility of primary health care services.

### 1. Data Sources

Our data, including the diagnostic codes of avoidable hospitalizations based on the 10th revision of the International Statistical Classification of Diseases, gender, age, and address of the hospitalized population and their degree of access to a primary health care physician, were gathered using the various databases of Iran Health Insurance Organization. The individual's national number was the common factor that allowed us to link the data in these different databases. The Primary Health Care Networks Development Plan of MOHME was used to determine the level of physician access in each village. Villages were divided into four categories based on their level of family physician access. Villages with rural health care centers had the highest physician access, followed by villages with health houses and satellite villages. Remote villages had the lowest level of physician access.

### 2. Analysis

For a descriptive investigation of avoidable hospitalizations between

2006 and 2013, the percentage of avoidable hospitalizations (out of total hospitalizations) was calculated. This was calculated based on age groups, gender groups, and four different categories of access to a primary physician. Trend series were plotted using Microsoft Excel software version 2016 (Microsoft Corp., Redmond, WA, USA).

To explore the relationship between demographic characteristics, access to a primary health care physician, and avoidable hospitalization, binary logistic regression was estimated separately for each year. We included gender, age, and access to a physician as independent variables and hospitalization as the categorical dependent variable, in the model. The observed outcome for this dependent variable has two possible types (happen or not happens avoidable hospitalization). Some diagnostic tests were conducted after developing the model. The link test was used to ensure that there was no specification error in the estimated model. Additionally, Hosmer-Lemeshow test was used to test the model's goodness of fit. Collinearity among the various dependent variables was also explored. All analyses were conducted in STATA software ver. 13.0 (Stata Corp., College Station, TX, USA).

## RESULTS

Total and avoidable hospitalizations increased during the study period, especially during the first 3 years of study. However, avoidable hospitalizations, as a percentage of total hospitalizations, did not change significantly throughout the study years. This value was 22.3% during the first year of study and varied between 17% and 19.6% from 2007 to 2013 (Figure 1).

Results show that among the four categories of villages, remote villages had the highest variation in the percentage of avoidable hospitalizations during the study period. It is necessary to mention that the populations of these villages are considerably lower than in other rural areas; therefore, small changes in hospitalizations can have a large effect on the shape of the figure trend. Figure 2 presents the percentage of avoidable hospitalizations among total hospitalizations in different rural areas.

Comparing the percentage of avoidable hospitalizations between men and women showed that this value is higher among men. During the study years, this percentage remained relatively stable among men, while showing a decrease among women. This decrease began one year after the implementation of the family physician program.




Figure 1. Avoidable and unavoidable hospitalizations in rural Tehran (2006–2013).




Figure 2. Comparison of the avoidable hospitalization rate between different rural areas in Tehran province.

However, this percentage remained stable for 8 years after the reform implementation (Figure 3).

Figure 4 displays the comparison of the percentages of avoidable hospitalizations among different age groups. The highest percentage of avoidable hospitalizations is seen in the >70 age group, while it is lowest in people aged between 10 and 30 years.

Table 1 presents the logistic regression analysis for each year. In this model, the dependent variable is the occurrence of avoidable hospitalizations, and the independent variables are age, gender, and level of access to a primary health care physician. No statistically significant relationship was found between avoidable hospitalization occurrence and physician access during the study years. Throughout the study, the probability of avoidable hospitalizations in the >70 years age group was higher as compared to 10–50 years. In some time periods, the probability of avoidable hospitalizations among those aged >70 years was higher than the 1–10 years and 50–70 years age groups. Exploring the relationship between gender and avoidable hospitalizations showed that the probability of avoidable hospitalization among women was lower than men in the years 2009, 2010, and 2012 (Table 1).

### DISCUSSION

This study explored the relationship between access to a primary health care physician and avoidable hospitalization among the rural population during the 7 years following the implementation of the rural family physician program. No significant relationship was found during the study years. Simultaneously, the relationship between demographic characteristics and avoidable hospitalization was investigated, which showed that avoidable hospitalizations were significantly higher among >70 years age group than younger age groups throughout the study. Avoidable hospitalizations were also higher among men than women; however, this finding was only significant in three of the study years.

Another study conducted in Tehran province to explore the relation-



Figure 3. Comparison of the avoidable hospitalization rate between males and females in rural areas of Tehran province.

ship between family physician access and avoidable hospitalization following the rural family physician program showed that access to a physician is not related to avoidable hospitalizations.<sup>17</sup> In our study, this relationship was explored over several years, however, the results had not been segregated based on year in the previous study. Both studies yielded similar results.

Theories suggest that health systems that rely more on primary health care and primary care physicians result in better health outcomes for the population and also decrease the costs of unnecessary and avoidable hospitalizations.<sup>19</sup> However, based on the results of these two studies, it appears that the gate-keeping function of the primary health care physician aimed at preventing unnecessary services such as avoidable hospitalization has not been implemented accurately in the family physician program. Therefore, investigating the flaws in the implementation of this program that led to this effect may be worthwhile.

Additionally, our results differ from other studies that evaluated the effects of the rural family physician program on total hospitalizations across Iran and showed a significant increase in both short- and long-term hospitalizations.<sup>20</sup> This difference in results may be because the nationwide study did not separate the effect of the family physician program on avoidable and referral sensitive hospitalizations, or due to the previous study's quasi-experimental design.

Besides, our results are in contrast with other before-and-after studies conducted in one of the counties in Tehran. Study results at the county level revealed that increasing access to primary care physicians resulted in a decrease in avoidable hospitalizations.<sup>16</sup> Therefore, a difference exists in the relationship between access to a primary care physician and avoidable hospitalizations at the province and county




Figure 4. Comparison of the avoidable hospitalization rate between different age groups in rural areas of Tehran province.

**Table 1.** Odds ratio and related confidence intervals of avoidable hospitalization based on age, gender, and access to a primary health care physician

Variable	Year									
	2006	2007	2008	2009	2010	2011	2012	2013		
<b>Gender</b>										
Female	1.34 (0.95–1.89)	0.88 (0.68–1.14)	0.89 (0.72–1.1)	0.75* (0.62–0.9)	0.77* (0.63–0.94)	1.05 (0.87–1.26)	0.82* (0.66–0.99)	0.83 (0.68–0.01)		
Male	-	-	-	-	-	-	-	-		
<b>Age (y)</b>										
1–10	0.56* (0.32–0.99)	0.66* (0.43–0.99)	0.7* (0.5–0.96)	0.08 (0.58–1.2)	0.94 (0.68–1.3)	0.96 (0.71–1.31)	0.84 (0.61–1.16)	0.56* (0.4–0.78)		
10–20	0.12* (0.05–0.33)	0.09* (0.04–0.21)	0.09* (0.05–0.19)	0.12* (0.06–0.22)	0.15* (0.09–0.25)	0.18* (0.1–0.33)	0.25* (0.14–0.44)	0.13* (0.07–0.23)		
20–30	0.17* (0.08–0.34)	0.11* (0.07–0.2)	0.09* (0.06–0.16)	0.17* (0.11–0.26)	0.15* (0.1–0.22)	0.18* (0.12–0.28)	0.21* (0.14–0.32)	0.09* (0.05–0.15)		
30–40	0.19* (0.1–0.38)	0.1* (0.05–0.19)	0.16* (0.1–0.26)	0.32* (0.22–0.47)	0.23* (0.15–0.34)	0.23* (0.16–0.34)	0.22* (0.14–0.33)	0.19* (0.12–0.29)		
40–50	0.44* (0.24–0.08)	0.49* (0.32–0.75)	0.43* (0.3–0.62)	0.51* (0.36–0.72)	0.49* (0.34–0.7)	0.55* (0.4–0.75)	0.54* (0.39–0.74)	0.39* (0.28–0.54)		
50–60	1.16 (0.67–2)	0.62* (0.42–0.93)	0.8 (0.57–1.13)	0.92 (0.66–1.28)	0.78 (0.55–1.2)	0.82 (0.61–1.1)	0.93 (0.68–1.27)	0.54* (0.4–0.72)		
60–70	0.95 (0.53–1.74)	0.62* (0.4–0.97)	0.63* (0.42–0.96)	1.05 (0.73–1.5)	1.05 (0.72–1.54)	0.1 (0.72–1.38)	0.91 (0.65–1.28)	0.57* (0.4–0.81)		
>70	-	-	-	-	-	-	-	-		
<b>Access</b>										
Remote villages	1.39 (0.7–2.78)	0.52 (0.25–1.06)	1.41 (0.9–2.24)	1.46 (0.94–2.27)	1.18 (0.79–1.77)	1.85 (1.22–2.82)	0.6 (0.34–1.06)	0.55 (0.3–1.02)		
Satellite villages	1.07 (0.58–1.95)	1.12 (0.73–1.71)	1.03 (0.71–1.49)	1.38 (0.1–1.9)	0.89 (0.62–1.26)	1.1 (0.72–1.43)	1.12 (0.78–1.61)	0.92 (0.61–1.39)		
Villages with health houses	1.17 (0.79–1.74)	0.85 (0.64–1.14)	1.26 (0.1–1.6)	1.12 (0.9–1.39)	0.92 (0.75–1.14)	1.1 (0.9–1.36)	0.95 (0.77–1.16)	0.9 (0.72–1.13)		
Villages with health care centers	-	-	-	-	-	-	-	-		
<b>Diagnostic tests statistics</b>										
Likelihood-ratio chi-square (11)	85.7	168.04	24.23	221.73	225.58	197.37	170.16	198.82		
P-value	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001		
Hosmer-Lemeshow chi-square (8)	6.73	11.63	8.62	7.12	4.85	8.98	8.6	4.04		
P-value >chi-square	0.56	0.16	0.37	0.52	0.77	0.34	0.37	0.85		

Values are presented as odds ratio (95% confidence interval), unless otherwise stated.  
\*P<0.05.

level. This may be due to the differences in the demographic, social, and economic characteristics of the rural population in the studied county in comparison with the whole of Tehran province. Therefore, it is better to consider other factors that may influence this issue.

Studies exploring the relationship between avoidable hospitalizations and access to a primary care physician in other countries show different results. In Ukraine, pediatric hospitalization rates for some preventable conditions were positively related to the availability of a primary care physician.<sup>21)</sup> In Canada, after adjusting for age, sex, and comorbidity, an increased rate of emergency department use was associated with the lack of a primary physician among people aged 65 years and older.<sup>22)</sup> In the United States, physician supply was not associated with avoidable hospitalizations in rural areas.<sup>23)</sup> In another study conducted in the United States, physician supply and distribution had little effect on avoidable hospitalization rates.<sup>24)</sup> Based on these studies, the effect of access to a primary health care physician on avoidable hospitalizations appears to be different in various health care systems with different financial systems, different payment methods to providers, and different populations under study.

The Health Information System in Iran does not hold data on the presence of simultaneous diseases, patient's disease history, their income and job situation, or their economic classification. Considering these variables alongside demographic variables can present a more comprehensive picture in determining the relationship between access to a physician and hospitalizations. However, it can be said that rural areas with lower access to physicians have a lower economical and welfare status in this study, although this cannot be claimed confidently.

In our study, we did not consider the structural and organizational aspects of the primary health care physician program through which the physicians provide services when exploring the relationship between physician access and hospitalization. However, according to the results of other studies, these characteristics may affect avoidable hospitalizations. For example, in Sweden, general practitioner visit rates were negatively associated with hospitalization rates, considering the country's socioeconomic status health care structure.<sup>25)</sup> In Taiwan, which has a system with higher access to health care, continuity of care resulted in decreased avoidable hospitalizations.<sup>26)</sup> In Canada, high continuity of care was associated with reduced odds of ambulatory care-sensitive hospitalizations controlling for demographic and self-reported, health-related measures.<sup>27)</sup> Based on a systematic review, a powerful primary health care system with access to primary care physicians and long associations between the physician and patient reduce the probability of avoidable hospitalization.<sup>3)</sup>

Therefore, health policymakers need to consider the structural and organizational aspects (such as disease management guideline development, continuity of care by physicians, types of service providers, etc.) when developing similar programs. Examples include the urban family physician program, which is the current agenda in Iran. This may help health policymakers to decrease costs and hospitalizations.

In our study, we found that elderly people (>70 years) were more

likely to undergo avoidable hospitalizations. Our results were consistent with other studies. In the United States, avoidable hospitalizations in people younger than 65 years were significantly lower than older age groups.<sup>28)</sup> In Italy, avoidable hospitalizations were higher among the elderly and increased with lower levels of access to primary care physicians.<sup>29)</sup> Higher avoidable hospitalizations among the elderly may be reasonable, given the higher rates of chronic conditions among this age group. Therefore, access to a primary care physician can reduce the need for hospitalization among older adults. Health policymakers should, therefore, pay special attention to provide the elderly with primary health care services to prevent future costs related to avoidable hospitalizations. This issue is of special importance in rural areas, where access to higher levels of health care is more complex.

In our study, a statistically significant difference was found in hospitalization among the sexes, with avoidable hospitalizations being higher in men compared to women throughout the study. A previous study in Tehran assessing the relationship between hospitalization, demographic characteristics, and access to physicians reported higher avoidable hospitalizations among men. However, these relationships were not simultaneously explored based on a time-series.<sup>17)</sup> In Norway, elderly women were hospitalized less frequently than men, while elderly women in Finland were more likely to use all health care services than men, except for hospital care services.<sup>30)</sup> In both Spain and Italy, avoidable hospitalization rates among men were reportedly higher than women.<sup>1,11)</sup>

Previous studies generally report higher use of health care services among females during their reproductive years. This is later reversed, with elderly men using health care services more than women. These results could be explained by differences in health status, care-seeking behavior, and care provisions among men and women. Their working time as farmers could be another cause of higher avoidable hospitalizations among rural men in comparison with women in this study. Time limitations imposed on males by their jobs in farming implies that they have less time to access services provided by primary health care physicians.

Our study provides valuable evidence that can be used when developing similar reforms in Iran and other developing countries. Implementation of the urban family physician program is currently on the agenda, and our results suggest that policymakers should pay greater attention to the urban population's demographic characteristics and their possible effects on the patients' use of physician and hospital services.

In conclusion, data analysis over several years following the primary health care reforms revealed that increasing access to primary health care physicians cannot, by itself, result in decreased avoidable hospitalizations. Therefore, policymakers need to consider the different factors and aspects of structure and organization when developing and implementing new interventions in primary health care to promote the performance of this level of health service provision.

## CONFLICT OF INTEREST

No potential conflict of interest relevant to this article was reported.

## ACKNOWLEDGMENTS

This research was supported by Tehran University of Medical Sciences and provided with a health services grant (no., 30317). Also we thank the Iranian Health Insurance Organization for their support in providing us with the study data.

## ORCID

Arash Rashidian: <https://orcid.org/0000-0002-4005-5183>

Sedigheh Salavati: <https://orcid.org/0000-0001-9206-1553>

Hanan Hajimahmoodi: <https://orcid.org/0000-0002-9520-4692>

## REFERENCES

- Rosano A, Loha CA, Falvo R, van der Zee J, Ricciardi W, Guasticchi G, et al. The relationship between avoidable hospitalization and accessibility to primary care: a systematic review. *Eur J Public Health* 2013;23:356-60.
- Health Services Delivery Programme, Division of Health Systems and Public Health. Assessing health services delivery performance with hospitalizations for ambulatory care sensitive conditions. Copenhagen: World Health Organization Regional Office for Europe; 2016.
- Van Loenen T, van den Berg MJ, Westert GP, Faber MJ. Organizational aspects of primary care related to avoidable hospitalization: a systematic review. *Fam Pract* 2014;31:502-16.
- Gao J, Moran E, Li YF, Almenoff PL. Predicting potentially avoidable hospitalizations. *Med Care* 2014;52:164-71.
- Muenchberger H, Kendall E. Determinants of avoidable hospitalization in chronic disease: development of a predictor matrix. Meadowbrook: Griffith University; 2008.
- Purdy S, Griffin T, Salisbury C, Sharp D. Ambulatory care sensitive conditions: terminology and disease coding need to be more specific to aid policy makers and clinicians. *Public Health* 2009;123:169-73.
- Ansari Z. The concept and usefulness of ambulatory care sensitive conditions as indicators of quality and access to primary health care. *Aust J Prim Health* 2007;13:91-110.
- Caminal J, Starfield B, Sanchez E, Casanova C, Morales M. The role of primary care in preventing ambulatory care sensitive conditions. *Eur J Public Health* 2004;14:246-51.
- Mustard CA, Kaufert P, Kozyrskyj A, Mayer T. Sex differences in the use of health care services. *N Engl J Med* 1998;338:1678-83.
- Macintyre S, Hunt K, Sweeting H. Gender differences in health: are things really as simple as they seem? *Soc Sci Med* 1996;42:617-24.
- Redondo-Sendino A, Guallar-Castillon P, Banegas JR, Rodriguez-Artalejo F. Gender differences in the utilization of health-care services among the older adult population of Spain. *BMC Public Health* 2006;6:155.
- Guo L, MacDowell M, Levin L, Hornung RW, Linn S. How are age and payors related to avoidable hospitalization conditions? *Manag Care Q* 2001;9:33-42.
- Rashidian A, Moradi G, Takian A, Sakha MA, Salavati S, Faraji O, et al. Effects of the Health Transformation Plan on caesarean section rate in the Islamic Republic of Iran: an interrupted time series. *East Mediterr Health J* 2019;25:254-61.
- Shadpour K. Primary health care networks in the Islamic Republic of Iran. *East Mediterr Health J* 2000;6:822-5.
- Takian A, Doshmangir L, Rashidian A. Implementing family physician programme in rural Iran: exploring the role of an existing primary health care network. *Fam Pract* 2013;30:551-9.
- Salavati S, Rashidian A, Emamgholipour S, Varahrami V. The impact of rural health insurance and the family physician program on hospitalizations, a beforeafter study at the county level conducted in Tehran province, Iran. *Med J Islam Repub Iran* 2018;32:32.
- Salavati S, Rashidian A, Hajimahmoodi H, Emamgholipour S, Varahrami V, Moez EK. The impact of the rural family physician and health insurance reforms in Iran on total and avoidable hospitalizations. *Prim Health Care Res Dev* 2018.
- Freund T, Campbell SM, Geissler S, Kunz CU, Mahler C, Peters-Klimm F, et al. Strategies for reducing potentially avoidable hospitalizations for ambulatory care-sensitive conditions. *Ann Fam Med* 2013;11:363-70.
- Atun R. What are the advantages and disadvantages of restructuring a health care system to be more focused on primary care services. Copenhagen: World Health Organization Regional Office for Europe; 2004.
- Rashidian A, Salavati S, Hajimahmoodi H, Kheirandish M. Does rural health system reform aimed at improving access to primary health care affect hospitalization rates?: an interrupted time series analysis of national policy reforms in Iran. *J Health Serv Res Policy* 2019;24:73-80.
- Oleksiuk O. Avoidable hospitalization of children, morbidity and health care supply. *Hygeia Public Health* 2013;48:234-7.
- Ionescu-Itu R, McCusker J, Ciampi A, Vadeboncoeur AM, Roberge D, Larouche D, et al. Continuity of primary care and emergency department utilization among elderly people. *CMAJ* 2007;177:1362-8.
- Laditka JN, Laditka SB, Probst JC. More may be better: evidence of a negative relationship between physician supply and hospitalization for ambulatory care sensitive conditions. *Health Serv Res* 2005;40:1148-66.
- Krakauer H, Jacoby I, Millman M, Lukomnik JE. Physician impact on hospital admission and on mortality rates in the Medicare population. *Health Serv Res* 1996;31:191-211.
- Lindstrom K, Engstrom S, Bengtsson C, Borgquist L. Determinants of hospitalisation rates: does primary health care play a role? *Scand J Prim Health Care* 2003;21:15-20.
- Cheng SH, Chen CC, Hou YF. A longitudinal examination of continuity of care and avoidable hospitalization: evidence from a universal coverage health care system. *Arch Intern Med* 2010;170:1671-7.
- Menec VH, Sirski M, Attawar D, Katz A. Does continuity of care with a family physician reduce hospitalizations among older adults? *J Health Serv Res Policy* 2006;11:196-201.
- Culler SD, Parchman ML, Przybylski M. Factors related to potentially preventable hospitalizations among the elderly. *Med Care* 1998;36:804-17.
- Rizza P, Bianco A, Pavia M, Angelillo IF. Preventable hospitalization

- and access to primary health care in an area of Southern Italy. *BMC Health Serv Res* 2007;7:134.
30. Suominen-Taipale AL, Martelin T, Koskinen S, Holmen J, Johnsen R. Gender differences in health care use among the elderly population in areas of Norway and Finland: a cross-sectional analysis based on the HUNT study and the FINRISK Senior Survey. *BMC Health Serv Res* 2006;6:110.



**Appendix 1.** List of avoidable hospitalization diagnostic codes which were used in the study (based on the 10th revision of the International Statistical Classification of Diseases)

Diseases classification	Avoidable diagnostic codes
Angina	I20, I25
Asthma	J45, J46
Cellulitis	L03, L04, L88
Congestive heart failure	I50, J81
Convulsions and epilepsy	G40, G41, R56, O15
Chronic obstructive pulmonary disease	J20, J41, J42, J43, J47, J12, J15, J16, J21, J22, J44
Dehydration and gastroenteritis	E86, A09, A04
Ear, nose and throat infections	H66, H67, J02, J03, J06,
Gangrene	R02
Hypertension	I10, I11
Influenza and pneumonia	J10, J11, J13, J14, J18
Nutritional deficiency	E40, E41, E42, E43
Other vaccine-preventable diseases	B26, A35, A36, A37, A80, B05, B06
Pelvic inflammatory disease	N70, N73, N74
Pyelonephritis	N10, N11, N12
Alcohol-related diseases	F10
Dyspepsia and other stomach function disorders	K30
Gastro esophageal reflux disease	K21
Migraine/acute headache	G43