

# Analytical determination, antioxidant and anti-inflammatory activities of Bhamrung-Lohit a traditional Thai medicine

Chitralada Panchakul<sup>1</sup>, Pakakrong Thongdeeying<sup>2,3</sup>, Arunporn Itharat<sup>2,3,\*</sup>, Weerachai Pipatrattanaseree<sup>4</sup>, Chonthicha Kongkwamcharoen<sup>5</sup>, and Neal M. Davies<sup>6</sup>

<sup>1</sup>Faculty of Medicine, Thammasat University (Rangsit Campus), Pathumthani, 12120, Thailand.

<sup>2</sup>Department of Applied Thai Traditional Medicine, Faculty of Medicine, Thammasat University (Rangsit Campus), Pathumthani, 12120 Thailand.

<sup>3</sup>Center of Excellence in Applied Thai Traditional Medicine Research, Thammasat University (Rangsit Campus), Pathumthani, 12120, Thailand.

<sup>4</sup>Regional Medical Science Center 12 Songkhla, Department of Medical Sciences, Ministry of Public Health, Songkhla, 90100, Thailand.

<sup>5</sup>Graduate School of Applied Thai Traditional Medicine, Faculty of Medicine, Thammasat University, Pathum Thani 12120, Thailand.

<sup>6</sup>Faculty of Pharmacy and Pharmaceutical Sciences, Katz Centre for Pharmacy and Health Research, University of Alberta, Edmonton, Alberta T6G2E1, Canada.

---

## Abstract

**Background and purpose:** Bhamrung-Lohit (BRL) remedy is a traditional Thai medicine (TTM). There are few reports of biological activity, the activity of its constituent plants, or quantitative analytical methods for the content of phytochemicals. In this study, we investigated antioxidant, anti-inflammatory activity, and total phenolic and flavonoid content and validated a new analytical method for BRL.

**Experimental approach:** Antioxidant activity was evaluated by a 2,2-diphenyl-1-picrylhydrazyl (DPPH) and 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid (ABTS) radical scavenging. The cellular antioxidant activity was evaluated by inhibition of the superoxide anion ( $O_2^{\bullet-}$ ) production from HL-60 cells and anti-inflammatory activity by inhibition of nitric oxide production in RAW264.7 cells. The total phenolic and flavonoid contents were analyzed using the Folin-Ciocalteu method and an aluminum chloride colorimetric assay, respectively. Validated analytical procedures were conducted according to International Conference on Harmonization (ICH) guidelines.

**Findings/Results:** An ethanolic extract of BRL exerted potent DPPH radical scavenging activity and moderate antioxidant and anti-inflammatory activity. *Caesalpinia sappan* exerted the greatest effect and the highest content of total phenolics and flavonoids. The HPLC method validated parameters that complied with ICH requirements. Each peak showed selectivity with a baseline resolution of 2.0 and precision was less than 2.0% CV. The linearity of all compounds was  $> 0.999$  and the recovery % was within 98.0%-102.0%. The validated results demonstrated specificity/selectivity, linearity, precision, and accuracy with appropriate LOD and LOQ.

**Conclusion and implication:** BRL remedy, a TTM demonstrated antioxidant and anti-inflammatory properties. This study is the first report on the biological activity and the validation of an HPLC method for BRL remedy.

**Keywords:** Anti-inflammation; Antioxidant; Bhamrung-Lohit; HPLC; Method validation.

---

## INTRODUCTION

Oxidative stress occurs in disease and pathophysiology when redox reactions in the body are imbalanced. Reactive oxygen species (ROS) are also released in the body and can act

as signaling molecules. Overproduction of ROS and defects in the homeostasis of antioxidants can cause an imbalance in the redox system causing oxidative stress in cells and tissues.

---

\*Corresponding author: A. Itharat

Tel: +66-29269749, Fax: +66-29269705

Email: iarunporn@yahoo.com

Red blood cells (RBCs) are integral to the cardiovascular system and help maintain oxidative balance in the blood (1). Hemoglobin with the RBCs initiates oxidative stress by its auto-oxidation which produces the superoxide radical leading to RBC damage (2). Inflammation is an important aspect of oxidative stress (3). Nitric oxide (NO) is involved in redox reactions and is pro-inflammatory in the body, when in excess NO reacts with the superoxide anion ( $O_2^{\bullet-}$ ) producing a peroxynitrite radical ( $ONOO^{\bullet}$ ) which reacts with enzymes, DNA, and cell membranes and causes irreversible damage leading to tissue damage and cell death (4).

Various cellular antioxidants in the cytosol including superoxide dismutase, and glutathione peroxidase reduce the overproduction of ROS and oxidative balance is maintained (5).

The functional use of antioxidant supplements is to provide beneficial effects in the amelioration of oxidative stress (6). These antioxidants can be medicinally synthesized or naturally occurring xenobiotics. All antioxidants can have potential side effects including allergic reactions, and gastrointestinal tract irregularities (7-10). The use of plants and traditional herbal remedies may provide an alternative source of new phytochemicals that impart antioxidant properties.

**Table 1.** The ethnobotanical data of Bhamrung-Lohit remedy and its plant components.

Plant species	Family	Plant part	Voucher specimen number	Proportion (%w/w)	Taste	Traditional use
<i>Alyxia reinwardtii</i>	Apocynaceae	Peel	SKP 013 01 18 01	1.41	Sweet	Tonic, anti-dizziness (49)
<i>Amomum verum</i>	Zingiberaceae	Fruit	SKP 206 01 20 01	1.41	Spicy	Antiemetic, carminative (50)
<i>Anaxagorea luzonensis</i>	Annonaceae	Heartwood	SKP 011 01 12 01	2.82	Spicy and astringent	Blood tonic, tonic, paregoric (51)
<i>Anethum graveolens</i>	Apiaceae	Dry fruit	SKP 199 01 07 01	1.41	Bitter and spicy	Tonic, carminative (50,52)
<i>Angelica dahurica</i>	Apiaceae	Root	SKP 199 01 04 01	1.41	Spicy and bitter	Tonic, antipyretic, paregoric, relieve asthma (49)
<i>Angelica sinensis</i>	Apiaceae	Root	SKP 199 01 19 01	1.41	Sweet and spicy	Antipyretic, gynecological (49,53,54)
<i>Aquilaria crassna</i>	Thymelaeaceae	Heartwood	SKP 193 01 03 01	1.41	Spicy and bitter	Blood tonic, tonic, cardi tonic (49)
<i>Arcangelisia flava</i>	Menispermaceae	Stem	SKP 114 11 06 01	2.82	Spicy and astringent	Carminative, anti-diarrhea, blood tonic (55)
<i>Artemisia annua</i>	Compositae	Aerial parts	SKP 051 01 01 01	1.41	Bitter	Antipyretic, relieve cough (56)
<i>Atractylodes lancea</i>	Compositae	Rhizome	SKP 051 01 12 01	1.41	Sweet and bitter	Tonic, oral prophylaxis, antipyretic (50)
<i>Avicennia marina</i>	Acanthaceae	Heartwood	SKP 213 01 13 01	1.41	Salty and bitter	Carminative, gynecological (57)
<i>Bixa orellana</i>	Bixaceae	Flower	SKP 026 02 15 01	14.08	Sweet and mellow	Blood tonic, Treatment of anemia (55)
<i>Caesalpinia sappan</i>	Leguminosae	Heartwood	SKP 072 03 19 01	14.08	Bitter and astringent	Blood tonic, treatment of menstruation pain (50)(56)
<i>Cinnamomum verum</i>	Lauraceae	Peel	SKP 096 03 03 01	1.41	Spicy and sweet	Carminative, relieve dyspepsia (58)
<i>Cuminum cyminum</i>	Apiaceae	Dry fruit	SKP 199 03 03 01	1.41	Spicy	Carminative (50)

<i>Dactylopius coccus</i>	Dactylopiidae	Lac	SKP 215 12 12 01	5.63	Astringent	Blood tonic (50,59)
<i>Dracaena cochinchinensis</i>	Asparagaceae	Heartwood	SKP 065 04 12 01	1.41	Bitter	Cardiotonic, anti-inflammatory (50)
<i>Foeniculum vulgare</i>	Apiaceae	Dry fruit	SKP 199 06 22 01	1.41	Sweet and spicy	Tonic, carminative (50)
<i>Lepidium sativum</i>	Brassicaceae	Dry fruit	SKP 057 12 19 01	1.41	Spicy and bitter	Gastrointestinal agent (60)
<i>Ligusticum sinense</i>	Apiaceae	Rhizome	SKP 199 12 19 01	1.41	Bitter	Carminative, blood tonic, gynecology (50)
<i>Mammea siamensis</i>	Guttiferae	Flower	SKP 083 13 19 01	2.82	Bitter	Cardiotonic, tonic (49,50)
<i>Mesua ferrea</i>	Calophyllaceae	Flower	SKP 083 13 06 01	2.82	Bitter	Blood tonic, cardiotonic, antipyretic (49,50)
<i>Mimusops elengi</i>	Sapotaceae	Flower	SKP 171 13 05 01	2.82	Sweet and astringent	Cardiotonic, antipyretic (50,61)
<i>Myristica fragrans</i>	Myristicaceae	Aril	SKP 121 13 06 01	1.41	Spicy	Blood tonic, tonic, carminative (50)
<i>Myristica fragrans</i>	Myristicaceae	Seed	SKP 121 13 06 01	1.41	Bitter and astringent	Tonic, carminative, antiemetic (50)
<i>Nelumbo nucifera</i>	Nelumbonaceae	Pollen	SKP 125 14 14 01	2.82	Astringent	Cardiotonic, tonic (50)
<i>Nigella sativa</i>	Ranunculaceae	Dry fruit	SKP 160 14 19 01	1.41	Spicy and bitter	Antiemetic, carminative (50)
<i>Piper retrofractum</i>	Piperaceae	Flower	SKP 146 16 18 01	2.82	Spicy and bitter	Promote blood circulation, antiemetic (50)
<i>Piper ribesoides</i>	Piperaceae	Vine	SKP 146 16 14 01	2.82	Spicy	Carminative (50)
<i>Piper sarmentosum</i>	Piperaceae	Root	SKP 146 16 19 01	2.82	Spicy	Carminative (50)
<i>Plumbago indica</i>	Plumbaginaceae	Root	SKP 148 16 09 01	2.82	Spicy	Promote blood circulation, gynecology, carminative (50)
<i>Senna garrettiana</i>	Leguminosae	Heartwood	SKP 072 19 07 01	1.41	Bitter	Carminative, gynecology (55)
<i>Syzygium aromaticum</i>	Myrtaceae	Flower	SKP 123 19 01 01	1.41	Spicy	Paregoric of tooth and pyorrhea, carminative, promote blood circulation (50)
<i>Terminalia bellirica</i>	Combretaceae	Fruit	SKP 049 20 02 01	1.41	Sour and astringent	Laxative, carminative, relieve cough (62)
<i>Terminalia chebula</i>	Combretaceae	Fruit	SKP 045 20 03 01	1.41	Astringent	Laxative, anti-diarrhea, relieve cough (55)
<i>Terminalia citrina</i>	Combretaceae	Fruit	SKP 049 20 03 01	1.41	Bitter and astringent	Laxative, antipyretic (63)
<i>Urceola rosea</i>	Apocynaceae	Stem	SKP 013 05 18 01	2.82	Astringent	Carminative (64)
<i>Zingiber officinale</i>	Zingiberaceae	Rhizome	SKP 206 26 15 01	2.82	Spicy	Carminative, antiemetic (56)

In traditional Thai medicine (TTM), the herbal remedy “Bhamrung-Lohit” (BRL) is listed in the Thailand National List of Essential Medicine (NLEM) since 2013 for nourishing blood and treating extensive menstrual blood loss in women (the Thai word “Bhamrung” means nourishment and “Lohit” means blood). BRL remedy consists of 38 plant ingredients which are combined in different proportions according to Table 1. *Caesalpinia sappan* and *Bixa orellana* are the two major plant ingredients in the remedy (11).

This study aimed to investigate the biological activity related to antioxidant properties of BRL remedy and its plant components as follows: chemical-based antioxidant activities by DPPH and ABTS•+ scavenging activity, cellular-based antioxidant activity through inhibition of O<sub>2</sub>•- from the promyelocyte cell, HL-60 cell assay and anti-inflammatory activity by inhibition of NO production from the macrophage RAW264.7 cells. A quantitative high-performance liquid chromatography (HPLC) method for determination of the major analytes brazilin, ellagic acid, eugenol, piperine, and myristicin in ethanolic extract of BRL remedy was developed and validated. This study is the first report of antioxidant activity, anti-inflammatory activity, and quantitative HPLC methodology. The study provides helpful information for further study of BRL remedy and the validated method can be used as a quality control standard for further development of BRL remedy extract.

## MATERIALS AND METHODS

### *Plant materials and preparation of BRL extract*

All plant materials were purchased from Charernsuka Osot Pharmacy (Nakornpathom, Thailand). The plant specimens were compared to the reference plant standards deposited at the Herbarium of Southern Center of Thai Medicinal Plants at the Faculty of Pharmaceutical Science, Prince of Songkhla University, Songkhla Province, Thailand. The voucher IDs are shown in Table 1. Plant materials were washed, dried, and cut into small pieces, weighed according to proportions and

mixed homogeneously. Each plant ingredient and the BRL remedy were macerated with 95% ethanol for three days. The extract was filtered and evaporated at 45 °C under a vacuum. The residue of BRL was then re-extracted an additional two times. All three extracts were combined and dried in a vacuum oven at 45 °C.

### *Chemicals and reagents*

Brazilin, ellagic acid, eugenol, myristicin, piperine, gallic acid, and propyl gallate (purity > 98%) were purchased from Sigma-Aldrich (MO, USA). Folin-Ciocalteu's phenol reagent, 2,2-diphenyl-1-picrylhydrazyl (DPPH), and butylated hydroxytoluene (BHT) were purchased from Fluka (Munich, Germany). Dimethyl sulfoxide (DMSO), acetonitrile, and phosphoric acid were purchased from RCI Lab Scan (Bangkok, Thailand). Purified water was prepared by Milli Q<sup>®</sup> system from Millipore (Bedford, MA, USA).

Murine macrophage leukemia cell line (RAW 264.7: ATCC<sup>®</sup> TIB-71<sup>™</sup>) and Human promyelocytic leukemia cells (HL-60) were purchased from American Type Culture Collection (VA, USA). Roswell Park Memorial Institute medium 1640 (RPMI-1640), fetal bovine serum (FBS), penicillin-streptomycin, phosphate buffer saline (PBS), and trypsin-EDTA were purchased from Biochrom (MA, Germany). Lipopolysaccharide (LPS), 3-(4,5-dimethyl-2-thiazolyl)-2,5 diphenyl-2H-tetrazolium bromide (MTT) and nitroblue tetrazolium (NBT) were purchased from Sigma-Aldrich. (MO, USA).

### *DPPH radical scavenging activity*

The antioxidant activity assessed by DPPH assay was conducted according to the modified method of Yamasaki *et al.* (12). Briefly, the stock solution of the extract was dissolved in absolute ethanol and also diluted to obtain various sample solutions at different concentrations. The sample solution (100 µL) was added with an equal volume of  $6 \times 10^{-5}$  M DPPH (in absolute ethanol) and kept in darkness at room temperature for 30 min. The absorbance of the reaction was measured at 520 nm. BHT was used as a positive control. The scavenging activity of the sample is the ability to reduce the color intensity of DPPH.

Inhibition (%) was calculated using the following equation:

$$\text{Inhibition (\%)} = \frac{\text{Abs. control} - \text{Abs. sample}}{\text{Abs. control}} \times 100 \quad (1)$$

#### **ABTS●+ radical scavenging activity**

ABTS●+ radical scavenging assay was conducted according to the method of Re *et al.* (13). Briefly, sample solutions of each extract were prepared at various concentrations in absolute ethanol. ABTS●+ radical solution was prepared by dissolving potassium persulfate with distilled water to produce the radical solution at a concentration of 2.45 mM. The sample solution (20 µL) was mixed with ABTS●+ solution (180 µL) and incubated at room temperature for 6 min. The absorbance of the solution reaction was measured by a microplate reader at a wavelength of 734 nm. The percentage of ABTS●+ scavenging activity was calculated. BHT was used as a positive control. The calculation of percent scavenging activity was calculated using the equation (1).

#### **Cellular antioxidant by NBT dye reduction assay**

The cellular antioxidant of ethanolic extract of BRL remedy was evaluated by inhibition of O<sub>2</sub>●- in differentiated HL-60 cells according to the method of Surarit *et al.* (14). Briefly, human promyelocytic leukemia cells (HL-60) were cultured in RPMI 1640 supplemented with 10% heated FBS, 50 IU/mL penicillin and 50 µg/mL streptomycin. HL-60 cells were seeded at 10,000 cells and incubated to obtain morphologically and functionally mature granulocytes by culturing in DMSO 1.3% in RPMI for 7 days at 37 °C, 5% CO<sub>2</sub> and 95% humidity. The stock solution of the extract was prepared in DMSO. The extract was diluted in a medium to produce working solutions at various concentrations. The concentration of DMSO in the treated cell was not more than 0.2% v/v. Propyl gallate was used as a positive control.

The differentiated HL-60 was dissolved in Hanks' balanced salt solution in a 50 mL centrifuge tube to produce a concentration of  $1 \times 10^6$  cells. Various concentrations of the extract (500 µL) were added into each tube and

incubated in an incubator at 37 °C with 5% CO<sub>2</sub> and 95% humidity for 15 min. After that, phorbol 12-myristate 13-acetate solution was added (final concentration of 250 ng/mL) and an aliquot of 250 µL of NBT solution (1.25 mg/mL) was added. Then, the mixture was incubated for 60 min and then, 2 mL of 1 N HCl was added. After mixing and centrifugation, the supernatant was removed and the produced NBT formazan was dissolved in 300 µL DMSO and the solution was transferred into 96 well plates. The plate was measured at 572 nm using a microplate reader. The inhibition of each concentration of the extract against superoxide formation measured by NBT reduction was calculated using equation (1).

#### **Anti-inflammatory activity by inhibition of nitric oxide production from RAW 264.7 cells and cytotoxicity by MTT assay**

Anti-inflammatory activity by inhibition of nitric oxide production in RAW264.7 cells was determined according to the method of Tewtrakul and co-workers with slight modifications (15). Briefly, RAW264.7 cells were cultured in DMEM medium with 10% FBS, penicillin (100 units/mL) and streptomycin (100 µg/mL) and incubated at 37 °C under 5% CO<sub>2</sub>. The cells were seeded into a 96-well plate and incubated for 24 h. The cells were treated with LPS at a final concentration of 5 ng/mL. A sample solution was added and incubated for 24 h. The supernatant was transferred to a new 96-well plate and Griess's reagent was added. The plate optical density (OD) was measured with a microplate reader at the wavelength of 570 nm. The viability of treated cells was determined by MTT solution. The OD was measured at 570 nm. The % inhibition and % toxicity were calculated.

#### **Determination of total phenolic content**

The total phenolic content of the extracts was determined by the modified Folin-Ciocalteu method (16). A 20 µL aliquot of the extracts was mixed with 100 µL of Folin-Ciocalteu's reagent and 80 µL of sodium carbonate solution. The plate was mixed and allowed to stand at room temperature to develop color for 30 min. The absorbance of the

reaction was measured at 765 nm. Total phenolic content was calculated from a calibration curve of gallic acid standard solutions (ranging from 2.5 to 100 µg/mL) and expressed as mg gallic acid equivalents (GAE)/g of dried plant.

#### **Determination of total flavonoid content**

Total flavonoid content was measured by the aluminum chloride colorimetric assay (17). Briefly, an aliquot (1 mL) of the extracts or standard solution of quercetin (20, 40, 60, 80, 100, 120, and 240 mg/L) was added to a volumetric flask. To the flask were added 75 µL of 5% NaNO<sub>2</sub> and 150 µL of 10% AlCl<sub>3</sub>. After 5 min, 500 µL of 1 M NaOH was added and the volume was adjusted by H<sub>2</sub>O. The solution was mixed well and the absorbance was measured against the prepared reagent blank at 510 nm. The concentration of total flavonoid content in the test samples was calculated from the calibration plot and expressed as mg quercetin equivalent/g of dried plant.

#### **HPLC analysis**

The analytical method for determination of anti-inflammatory markers in BRL ethanolic extract was carried out by using an HPLC system (Shimadzu, Japan) consisting of a solvent degasser (G1322A), an autosampler (G1329A), a quaternary solvent pump (G1311A), a photodiode array detector (G1315D), and a column oven (G1316A). The chromatographic data were processed by the Chemstation® software revision B.04.01 SP1.

#### **Chromatographic conditions**

All markers were separated along a C18 HPLC column (5 µm, 4.6 mm × 250 mm;

Zorbax®, Agilent, USA) protected by a C18 guard cartridge (5 µm, 4.6 mm × 3 mm; Zorbax®, Agilent, USA). The mobile phase consisted of acetonitrile (A) and 0.1% v/v phosphoric acid in water (B). The gradient elution of the mobile phase was programmed and presented in Table 2.

The flow rate was set at 1.0 mL/min. The operating temperature was maintained at room temperature. Ten µL of sample and standard solutions were injected into the chromatographic system and the anti-inflammatory markers were detected at a wavelength of 210 nm (brazilin, eugenol and myristicin), 254 nm (ellagic acid), and 320 nm (piperine).

#### **Preparation of standard and sample solution**

Each stock standard solution of brazilin, ellagic acid, eugenol, myristicin, and piperine was prepared at a concentration of 1.0 mg/mL in methanol. The working standard mixtures were prepared in methanol according to the concentration from the linear range of each standard. For sample solutions, the ethanolic of BRL extract was dissolved in methanol to produce a sample solution at the concentration of 10 mg/mL.

#### **Validation of the HPLC method**

The developed HPLC method was validated as per International Conference on Harmonization (ICH 2005) guidelines (18). The validation parameters included selectivity, linearity, accuracy, precision, the limit of detection (LOD), and the limit of quantitation (LOQ). All validation parameters passed the specification limits.

**Table 2.** Mobile phase gradient elution program for developed analytical assay. The mobile phase consisted of acetonitrile (A) and 0.1% v/v phosphoric acid in water (B).

Time (min)	Mobile phase composition (%)
0-5	5 A: 95 B
5-45	5 A: 95 B – 50 A: 50 B
45-60	50 A: 50 B – 95 A: 5 B
60-65	95 A: 5 B – 100 A: 0 B
65-70	5 A: 95 B

**Statistical analysis**


The results of the anti-inflammatory activities were reported as mean ± standard error of the mean (SEM) from three replicated experiments. IC<sub>50</sub> values were calculated by using regression analysis using the GraphPad Prism software (CA, USA). Mean differences among groups were analyzed by Student’s t-test. Statistical analysis was conducted by using GraphPad Prism software (CA, USA). The results of the HPLC analysis were expressed as mean ± SD. The hierarchical cluster analysis (HCA) and heat map were analyzed using Heatmap Illustrator (HemI) version 1.0 software (19). Statistical

comparisons between groups will be calculated via independent sample t-test and one-way analysis of variance (ANOVA) followed by the LSD post hoc test. Differences were considered significant at *P* < 0.05.

**RESULTS**

**Antioxidant activity and Anti-inflammatory activity by inhibition of nitric oxide production**

The results of DPPH, ABTS•+ radical scavenging activities, the cellular antioxidant by NBT assay, and anti-inflammatory activity by inhibition of NO production are shown in Table 3 and Fig. 1.



**Fig. 1.** Heat map and hierarchical cluster analysis of plant and Bhamrung-Lohit remedy extracts. The color scale bar showed a range of IC<sub>50</sub>, the red color bar represents more potent activity (0-10 µg/mL) incrementing to a blue color bar which represents weak activity (90-100 µg/mL).

**Table 3.** Antioxidant and anti-inflammatory activities of ethanolic extract of Bhamrung-Lohit remedy and its plant ingredients. Data are expressed as mean  $\pm$  SEM; n = 3.

Plant genus and species	Antioxidant (EC <sub>50</sub> ; $\mu$ g/mL)			Inhibitory activity of nitric oxide (IC <sub>50</sub> ; $\mu$ g/mL)
	DPPH	ABTS	NBT	
<i>Artemisia annua</i>	72.64 $\pm$ 3.48	> 100	> 100	64.11 $\pm$ 5.97
<i>Aquilaria crassna</i>	84.85 $\pm$ 3.85	32.81 $\pm$ 3.73	> 100	2.33 $\pm$ 2.33
<i>Angelica dahurica</i>	71.06 $\pm$ 3.18	84.15 $\pm$ 6.28	> 100	36.47 $\pm$ 1.92
<i>Arcangelisia flava</i>	43.95 $\pm$ 0.58	28.93 $\pm$ 2.61	> 100	91.13 $\pm$ 5.50
<i>Anethum graveolens</i>	64.26 $\pm$ 1.73	> 100	> 100	95.27 $\pm$ 2.69
<i>Atractylodes lancea</i>	90.17 $\pm$ 0.64	> 100	> 100	15.66 $\pm$ 1.45
<i>Anaxagorea luzonensis</i>	5.23 $\pm$ 0.34	5.15 $\pm$ 0.66	> 100	11.26 $\pm$ 0.8
<i>Avicennia marina</i>	25.19 $\pm$ 0.87	30.37 $\pm$ 1.8	> 100	7.59 $\pm$ 2.12
<i>Alyxia reinwardtii</i>	> 100	58.85 $\pm$ 4.71	> 100	89.93 $\pm$ 3.29
<i>Angelica sinensis</i>	56.37 $\pm$ 1.44	75.12 $\pm$ 5.87	> 100	14.24 $\pm$ 1.41
<i>Artemisia verum</i>	> 100	> 100	> 100	64.75 $\pm$ 4.47
<i>Bixa orellana</i>	8.32 $\pm$ 0.48	18.36 $\pm$ 1.56	> 100	57.62 $\pm$ 2.8
<i>Cuminum cyminum</i>	48.5 $\pm$ 4.52	> 100	> 100	27.19 $\pm$ 1.08
<i>Caesalpinia sappan</i>	4.51 $\pm$ 0.14	4.98 $\pm$ 0.18	12.07 $\pm$ 0.44	7.57 $\pm$ 1.46
<i>Cinnamomum verum</i>	16.55 $\pm$ 2.22	55.85 $\pm$ 1.41	> 100	21.19 $\pm$ 6.58
<i>Dactylopius coccus</i>	> 100	> 100	> 100	64.99 $\pm$ 3.4
<i>Dracaena cochinchinensis</i>	17.36 $\pm$ 0.64	7.55 $\pm$ 0.45	> 100	42.21 $\pm$ 0.57
<i>Foeniculum vulgare</i>	64.08 $\pm$ 3.56	> 100	> 100	86.30 $\pm$ 3.40
<i>Lepidium sativum</i>	> 100	> 100	> 100	> 100
<i>Ligusticum sinense</i>	52.23 $\pm$ 2.53	63.18 $\pm$ 4.79	> 100	46.67 $\pm$ 3.29
<i>Mimusops elengi</i>	55.84 $\pm$ 1.01	75.6 $\pm$ 5.83	> 100	> 100
<i>Mesua ferrea</i>	31.88 $\pm$ 1.93	67.91 $\pm$ 1.93	> 100	27.46 $\pm$ 0.78
<i>Myristica fragrans (Arail)</i>	21.27 $\pm$ 0.42	18.4 $\pm$ 1.57	> 100	81.81 $\pm$ 1.98
<i>Myristica fragrans (Nutmeg)</i>	21.91 $\pm$ 0.78	14.98 $\pm$ 1.29	> 100	> 100
<i>Mammea siamensis</i>	13.22 $\pm$ 1.13	28 $\pm$ 2.93	> 100	64.49 $\pm$ 4.19
<i>Nelumbo nucifera</i>	> 100	> 100	> 100	> 100
<i>Nigella sativa</i>	> 100	> 100	> 100	67.17 $\pm$ 3.17
<i>Plumbago indica</i>	80.92 $\pm$ 5.99	> 100	> 100	22.58 $\pm$ 2.6
<i>Piper retrofractum</i>	> 100	> 100	> 100	21.40 $\pm$ 4.40
<i>Piper ribesiodes</i>	22.82 $\pm$ 1.9	17.61 $\pm$ 1.89	> 100	42.38 $\pm$ 5.59
<i>Piper sarmentosum</i>	95.66 $\pm$ 1.86	> 100	> 100	66.48 $\pm$ 3.95
<i>Syzygium aromaticum</i>	7.2 $\pm$ 0.15	2.94 $\pm$ 0.22	> 100	3.06 $\pm$ 0.67
<i>Senna garrettiana</i>	4.87 $\pm$ 0.33	4.46 $\pm$ 0.95	10.84 $\pm$ 1.14	98.64 $\pm$ 0.04
<i>Terminalia bellirica</i>	3.84 $\pm$ 0.35	5.94 $\pm$ 0.59	23.81 $\pm$ 0.56	68.83 $\pm$ 0.14
<i>Terminalia chebula</i>	5.37 $\pm$ 0.58	8.51 $\pm$ 0.69	78.16 $\pm$ 1.49	93.49 $\pm$ 2.24
<i>Terminalia citrina</i>	2.42 $\pm$ 0.27	3.93 $\pm$ 0.3	49.15 $\pm$ 2.18	83.90 $\pm$ 6.10
<i>Urceola rosea</i>	5.02 $\pm$ 0.16	7.64 $\pm$ 0.52	> 100	> 100
<i>Zingiber officinale</i>	11.83 $\pm$ 0.71	10.86 $\pm$ 1.43	> 100	9.29 $\pm$ 0.13
<b>Bhamrung-Lohit remedy</b>	6.01 $\pm$ 0.21	37.77 $\pm$ 2.48	84.94 $\pm$ 7.47	57.63 $\pm$ 4.01
<b>Butylated hydroxytoluene</b>	13.78 $\pm$ 0.23	N/A	N/A	N/A
<b>Trolox</b>	N/A	5.55 $\pm$ 0.70	N/A	N/A
<b>Propyl gallate</b>	N/A	N/A	19.52 $\pm$ 0.31	N/A
<b>Prednisolone</b>	N/A	N/A	N/A	0.44 $\pm$ 0.04

N/A, Not applicable.

The HCA (Fig. 1) clustered the plant constituents into two major groups, the low-activity group (group I) and the moderate to high-activity group (group II). Each sub-group in group II expressed its characteristic on the tested activities. BRL remedy showed potent activity on DPPH radical scavenging and exerted moderate activity on ABTS•+ scavenging activity, cellular antioxidant by inhibition of superoxide anion production (NBT assay), and inhibition of NO

production. For the plant ingredients of the BRL remedy, *C. sappan* exerted potent activity on every tested activity with EC<sub>50</sub> and IC<sub>50</sub> less than 20  $\mu$ g/mL. The sub-group containing *C. sappan*, *Senna garrettiana*, *Terminalia bellirica*, and *Tholymis citrina* performed inhibitory effects on all tested activities. The sub-group containing *Anaxagorea luzonensis*, *Syzygium aromaticum*, *Zingiber officinale*, and *Acaryochloris marina* showed potent activity on DPPH, ABTS•+ and



inhibition of NO production. With regard to antioxidant activity, *C. sappan* and *S. garrettiana* exerted potent activity more than the positive controls in all tested antioxidant activity methodology. *A. luzonensis*, *S. Aromaticum*, and *T. citrina* exerted potent activity greater than the positive controls on DPPH and ABTS•+ scavenging activity. For inhibitory activity on NO production, several extracts showed potent activity with IC<sub>50</sub> less than 20 µg/mL including *Aguilaria crassna*, *Atractylodes lancea*, *A. luzonensis*, *Acaryochloris marina*, *Angelica sinensis*, *C. sappan*, *S. Aromaticum*, and *Z. officinale*. The BRL remedy showed inhibitory activity with an IC<sub>50</sub> value of 57.63 ± 4.01 µg/mL.

All plant constituents exerted significantly less anti-inflammatory activity than the positive control (prednisolone).

#### **Total phenolic content and total flavonoid content**

As shown in Table 4, *C. sappan* showed the highest content of total phenolic followed by *S. garrettiana* and *T. citrina*, respectively. For total flavonoid content, *C. sappan* also showed the highest content followed by *S. garrettiana* and *A. luzonensis*. BRL remedy showed total phenolic content and total flavonoid content of 207.93 ± 7.40 mgGAE/g) and 457.63 ± 11.25 mg quercetin equivalent/g, respectively.

**Table 4.** Total flavonoid content of ethanolic extract of Bhamrung-Lohit remedy and its plant components. Data are expressed as mean ± SEM; n = 3.

Plant genus and species	Total phenolic content (mg GAE/g)	Total flavonoid content (mg QuE/g)
<i>Artemisia annua</i>	50.26 ± 1.41	89.97 ± 1.09
<i>Aquilaria crassna</i>	104.06 ± 5.39	148.25 ± 7.19
<i>Angelica dahurica</i>	68.88 ± 2.38	142.47 ± 3.28
<i>Arcangelisia flava</i>	103.30 ± 1.54	176.85 ± 2.04
<i>Anethum graveolens</i>	24.36 ± 2.30	106.85 ± 5.16
<i>Atractylodes lancea</i>	33.88 ± 1.77	238.72 ± 7.66
<i>Anaxagorea luzonensis</i>	679.24 ± 18.72	1,094.82 ± 54.38
<i>Avicennia marina</i>	97.22 ± 8.57	211.85 ± 0.78
<i>Alyxia reinwardtii</i>	51.67 ± 0.57	156.85 ± 4.54
<i>Angelica sinensis</i>	59.88 ± 2.29	129.66 ± 0.78
<i>Artemisia verum</i>	36.42 ± 3.00	143.57 ± 1.88
<i>Bixa orellana</i>	168.13 ± 5.50	236.85 ± 6.10
<i>Cuminum cyminum</i>	39.19 ± 2.74	192.00 ± 0.94
<i>Caesalpinia sappan</i>	759.88 ± 46.44	2,535.91 ± 1.72
<i>Cinnamomum verum</i>	62.32 ± 5.01	431.38 ± 4.38
<i>Dactylopius coccus</i>	7.61 ± 0.2	95.60 ± 4.54
<i>Dracaena cochinchinensis</i>	373.06 ± 18.86	327.47 ± 13.91
<i>Foeniculum vulgare</i>	25.35 ± 2.02	92.16 ± 2.97
<i>Lepidium sativum</i>	13.83 ± 1.08	104.35 ± 5.79
<i>Ligusticum sinense</i>	53.17 ± 1.47	157.32 ± 4.07
<i>Mimusops elengi</i>	24.65 ± 2.10	75.13 ± 2.82
<i>Mesua ferrea</i>	66.33 ± 5.66	177.79 ± 2.35
<i>Myristica fragrans (Arail)</i>	86.24 ± 3.50	336.85 ± 0.16
<i>Myristica fragrans (Nutmeg)</i>	101.12 ± 3.45	291.07 ± 7.82
<i>Mammea siamensis</i>	98.73 ± 10.81	310.28 ± 7.03
<i>Nelumbo nucifera</i>	25.64 ± 0.64	181.07 ± 3.44
<i>Nigella sativa</i>	9.50 ± 0.44	77.79 ± 1.10
<i>Plumbago indica</i>	11.99 ± 0.16	150.13 ± 5.32
<i>Piper retrofractum</i>	22.69 ± 2.56	271.85 ± 1.10
<i>Piper ribesoides</i>	151.44 ± 5.08	490.13 ± 12.50
<i>Piper sarmentosum</i>	36.65 ± 0.84	259.50 ± 6.56
<i>Syzygium aromaticum</i>	548.98 ± 14.39	846.06 ± 3.75
<i>Senna garrettiana</i>	754.85 ± 10.62	1,364.04 ± 30.16
<i>Terminalia bellirica</i>	419.14 ± 2.16	135.60 ± 4.85
<i>Terminalia chebula</i>	297.18 ± 9.36	98.41 ± 4.53
<i>Terminalia citrina</i>	694.81 ± 3.27	187.31 ± 8.75
<i>Urceola rosea</i>	488.06 ± 22.50	1,241.07 ± 19.68
<i>Zingiber officinale</i>	197.64 ± 4.90	780.75 ± 19.69
<b>Bhamrung-Lohit remedy</b>	<b>207.93 ± 7.40</b>	<b>457.63 ± 11.25</b>

GAE, Gallic acid equivalents; QuE, quercetin equivalent.

## Validation of the HPLC method

### Specificity/selectivity

A new analytical method for the simultaneous determination of brazilin, ellagic acid, eugenol, myristicin, and piperine in ethanol extract of BRL remedy was developed and validated. As shown in Fig. 2, the chromatogram of the sample was compared to the chromatogram of the standard solution. The retention time of brazilin, ellagic acid, eugenol, piperine, and myristicin in the sample solution were 19.17, 23.8, 42.1, 49.0, and 52.1 min, respectively. The UV spectrum of the analyzed peaks in the chromatogram of the sample solution corresponded to the respective standards in the chromatogram of the standard solution (Fig. 2). These results indicated the specificity and selectivity of this method.

### Linearity and ranges

The linearity of the developed method was determined by the coefficient of determination ( $R^2$ ) calculated from the constructed calibration curves of each marker. As shown in Table 5, the  $R^2$  values of brazilin, ellagic acid, eugenol, piperine, and myristicin were 0.9996, 0.9991, 0.9996, 0.9991, and 0.9999, respectively. The linear range of each marker varied relating to their contents in the BRL extract and their absorptivity at the detected wavelength.

### LODs and LOQs


LODs and LOQs of the markers were analyzed according to the visualization shown in Table 5. The LODs of brazilin, ellagic acid, eugenol, piperine, and myristicin were 0.1, 0.5, 0.5, 1.0, and 0.2  $\mu\text{g/mL}$ , respectively. The LOQs were 6.25, 1.25, 2.0, 5.0, and 1.0  $\mu\text{g/mL}$  respectively.




### Accuracy and precision





The accuracy of the method was presented as % recovery of each compound shown in Table 6. The % recovery of all standards was within 97.62% - 101.57%. The precision of the analytical method, both intra-run and inter-run, was performed as % CV which was less than 1.99.


### Content of the markers in ethanolic extract of BRL remedy

The ethanolic extract of the BRL remedy was analyzed for the content of brazilin, ellagic acid, eugenol, myristicin and piperine by using the developed and validated HPLC method. The results found the amount of brazilin, ellagic acid, eugenol, piperine and myristicin were  $120.23 \pm 4.90$ ,  $2.03 \pm 0.07$ ,  $12.27 \pm 0.19$ ,  $4.66 \pm 0.12$  and  $0.77 \pm 0.17$  mg/g of extract, respectively.









**Fig. 2.** The HPLC chromatogram and UV spectrum of the mixed standard solution and BRL sample solution. (A, C, E) Chromatograms of standard solution detecting at the wavelength of 210, 254, and 320, respectively; (B, D, F) chromatograms of BRL sample solution detecting at the wavelength of 210, 254, and 320, respectively. Comparative UV-spectrum of peaks in standard and sample; (G) brazilin, (H) eugenol, (I) myristicin, (J) ellagic acid, and (K) piperine. BRL, Bhamrung-Lohit.

**Table 5.** Linearity, LODs, and LOQs of brazilin, ellagic acid, eugenol, piperine and myristicin in Bhamrung-Lohit extract.

Compounds	Validation parameters			
	Range (µg/mL)	Linearity	LOD (µg/mL)	LOQ (µg/mL)
<b>Brazilin</b>	500-1800	$Y = 11.621X - 988.790; r^2 = 0.9996$	0.1	0.325
<b>Ellagic</b>	5-40	$Y = 158.990X - 9.785; r^2 = 0.9991$	0.5	1.25
<b>Eugenol</b>	25-270	$Y = 59.468X - 571.650; r^2 = 0.9996$	0.5	2.5
<b>Piperine</b>	15-60	$Y = 47.834X - 6.381; r^2 = 0.9991$	0.2	1.0
<b>Myristicin</b>	1-25	$Y = 191.950X - 78.762; r^2 = 0.9999$	0.2	1.0

LOD, Limit of detection; LOQ, limit of quantification.

**Table 6.** Accuracy of the analytical method for determination of brazilin, ellagic acid, eugenol, piperine, and myristicin in Bhamrung-Lohit extract

Compounds	Spiked concentration (µg/mL)	Intra-run (n = 3)			Inter-run (n = 9)		
		Concentration found (µg/mL; mean ± SD)	Recovery (%)	CV (%)	Concentration found (µg/mL; mean ± SD)	Recovery (%)	CV (%)
<b>Brazilin</b>	600	606.23 ± 11.85	100.37 - 101.04	0.96	589.49 ± 10.39	99.75 - 100.49	1.74
	900	903.88 ± 10.73		0.69	904.39 ± 7.31		0.81
	1100	1104.17 ± 4.75		0.31	1101.66 ± 18.51		1.68
<b>Ellagic acid</b>	10	9.76 ± 0.09		0.60	9.84 ± 0.17		1.70
	20	20.31 ± 0.38	97.62 - 101.57	0.68	20.01 ± 0.40	98.37 - 100.21	1.99
	25	24.61 ± 0.31		1.11	24.96 ± 0.36		1.43
<b>Eugenol</b>	50	50.23 ± 0.29		0.75	50.54 ± 0.66		1.31
	150	150.31 ± 1.02	99.89 - 100.47	1.26	149.92 ± 1.19	99.19 - 100.09	0.80
	200	199.79 ± 3.83		1.35	198.37 ± 2.83		1.43
<b>Piperine</b>	20	20.19 ± 0.16		0.34	20.00 ± 0.37		1.83
	30	30.25 ± 0.37	100.51 - 100.97	0.60	29.87 ± 0.40	99.38 - 99.98	1.34
	35	35.18 ± 0.62		0.25	34.78 ± 0.69		1.98
<b>Myristicin</b>	5	4.91 ± 0.05		0.66	4.97 ± 0.07		1.49
	15	15.09 ± 0.29	98.35 - 101.37	0.74	14.88 ± 0.25	99.20 - 99.86	1.66
	20	20.27 ± 0.31		1.12	19.97 ± 0.35		1.73

## DISCUSSION

Oxidative stress plays a significant role as a risk factor for diseases including cancer and cardiovascular disease (6). In the cardiovascular system, hemoglobin within red blood cells plays a key role in the initiating of oxidative stress by its auto-oxidation which produces superoxide radicals including  $O_2^{\bullet-}$  and  $H_2O_2$  (2). In inflammatory processes, the upregulation and stimulation of NO which reacts with superoxide radicals leads to damage from the ONOO $\bullet$  radical (2). Therefore, in this study, evaluating the effects of BRL remedy and its components on both antioxidant and anti-inflammatory activity were assessed.

The ethanolic extract of BRL exerted significant antioxidant activity as measured with the DPPH scavenging activity and also showed significant activity on ABTS $\bullet+$  radical scavenging, as well as a cellular antioxidant activity through inhibition of  $O_2^{\bullet-}$  and concurrent anti-inflammatory activity by inhibition of NO production. This study is the first report of BRL extract demonstrating both *in vitro* antioxidant and anti-inflammatory activity.

Concerning the plant components of BRL, *C. sappan* exerted potent activity. *S. garrettiana* also exerted potent activity in all tested antioxidant systems. *T. bellirica* and *T. citrina* exerted potent activity on DPPH and ABTS $\bullet+$  radical scavenging but exerted less pronounced effects on cellular antioxidant and inhibitory activity on NO production. For the sub-group containing *A. luzonensis*, *S. aromatica*, *Z. Offinale*, and *A. marina*. All four plants were inactive in their inhibitory activity on  $O_2^{\bullet-}$  production determined by the NBT assay. *A. luzonensis*, *S. aromatica*, and *Z. officinale* exerted potent DPPH and ABTS $\bullet+$  radical scavenging and anti-inflammatory activity while *A. Marina* exerted potent anti-inflammatory activity and demonstrated correspondingly less activity on DPPH and ABTS $\bullet+$  radical scavenging. Eight plant constituents in particular demonstrated noteworthy bioactivity findings. Firstly, *C. sappan* and its major antioxidant compound

(brazilin) have been previously reported to possess potent antioxidant and anti-inflammatory activity similar to our findings (20-23). For *S. garrettiana*, this study is the first report of DPPH and ABTS●+ radical scavenging, inhibitory activity on O<sub>2</sub>●- production in HL-60 cells and inhibitory activity on NO production in RAW264.7 cells. The major compound in *S. garrettiana* is trans-piceatannol, a stilbene, which has been reported to have antioxidant activity as it increased cellular antioxidant defense (24). For *T. bellirica* and *T. citrina*, Hazra and co-workers reported that *T. bellirica* exerted potent DPPH radical scavenging similar to our study. Moreover, potent activity on the scavenging of O<sub>2</sub>●-, NO, and ONOO radicals have been reported (25). *T. bellirica* possesses potent anti-inflammatory activity through the inhibition of NO, cyclooxygenase-2, tumor necrosis factor alpha and interleukin-6 (26,27). The leaves of *T. citrina* exerted weak DPPH scavenging activity differing from our study in which the fruit was utilized (28). The bioactive constituents in *T. bellirica* and *T. citrina* are tannin, glucoside, pyranosides, triterpenoids, and acidic compounds (29). Ellagic acid in *Terminalia* spp. exerted anti-inflammatory effects by inhibition of NO production in RAW 264.7 (30). For *A. luzonensis*, few studies have been reported on activity related to antioxidation and anti-inflammatory effects and its phytochemicals. Chaiyasut and co-workers reported potent DPPH scavenging activity and high content of total phenolic compounds which is similar to our study (31). Gonda and co-workers reported the xanthone derivatives are constituents in *A. luzonensis* (32). *S. aromatica* and *Z. officinale* are common plants well known for their activity. *S. aromaticum* (clove) and its major constituent, eugenol, exerted potent antioxidant and anti-inflammatory activity (33-36). Perez-Roses and co-workers reported that eugenol exerted antioxidant activity by scavenging DPPH, NO, and H<sub>2</sub>O<sub>2</sub> (37). Yeh and co-workers reported that eugenol inhibited the expression of the inducible nitric oxide synthase from the LPS-induced macrophages. It also reduced the tumor necrosis factor alpha and IL-1β as well as the nuclear factor kappa B, extracellular signal-

regulated kinase 1/2, and p38 mitogen-activated protein kinases signaling pathways (38). For *Z. officinale*, several studies reported its antioxidant and anti-inflammatory activity both *in vitro* and *in vivo* (39-41). Kim and co-workers reported that 6-gingerol, an active compound in *Z. officinale*, exerted potent activity both *in vivo* and *in vitro*, and exerted strong anti-inflammatory activity (42). For *A. marina*, several studies reported its potential antioxidant activity both *in vitro* (43) and *in vivo* (44). Lincy and co-workers reported that *A. marina* has a high content of total phenolics and flavonoids imparting antioxidant properties (43). For the phytochemicals, *A. marina* contains various constituents, including naphthalene derivatives, flavones, iridoid glucosides, phenylpropanoid glycosides, diterpenoid glucosides, flavonoids, terpenoids and steroids (45). Avicequinone C has been isolated from the methanolic extract of *A. Marina* (46).

With regard to the analyte markers analyzed by HPLC, brazilin is the major compound in *C. Sappan* (22), ellagic acid from *Terminalia* spp. (30), eugenol from *S. aromatica* (36), piperine from *Piper* spp (47). and myristicin from *M. Fragrans* (48). The present study aimed at developing and validating an HPLC method for the determination of the main major constituents of BRL extract. The method can be used as a quality control method for further development of BRL extract into health products. There are no previous analytical methods reported for quality control of the BRL remedy. The current study is the first study presenting validation results of an HPLC method for the determination of the major marker constituent brazilin, ellagic acid, eugenol, piperine, and myristicin in BRL remedy.

An HPLC method for simultaneous quantitative analysis of brazilin, ellagic acid, eugenol, piperine, and myristicin in the ethanolic extract of BRL remedy was developed and validated. In the present study, 0.1% v/v phosphoric acid was used as a component in the mobile phase to suppress the ionization of phenolic compounds, such as brazilin and ellagic acid, causing a reduction in peak tailing. All marker analytes were

separated with the C18 column with a resolution value of more than 2.0. The linear range of the calibration curve of each marker varied depending on its content in BRL extract and absorptivity. Each marker was detected at its appropriated maximum detection wavelength as follows: 210 nm for brazilin, eugenol, and myristicin, 254 nm for ellagic acid, and 320 nm for piperine. The developed HPLC method resulted in excellent specificity for the determination of the markers in BRL ethanol extract. The calibration curves showed excellent linearity with  $R^2 > 0.999$  at the analyzed concentration ranges. The LODs for all markers ranged between 0.1 – 1.0  $\mu\text{g/mL}$  and the LOQs ranged between 1.0 – 6.25  $\mu\text{g/mL}$ . The accuracy of each marker ranged between 97.62% – 101.57%. The intra-run and inter-run precisions presented as %CV were less than 1.99%. The method validation demonstrated excellent results meeting ICH criteria for quality control of the BRL extract.

## CONCLUSION

BRL remedy, a TTM remedy is being used in patients, particularly in Thailand and neighboring nations. In this study, an ethanol extract of BRL remedy exerted antioxidant and anti-inflammatory activity as measured *in vitro* by DPPH and ABTS $\bullet$ + radical scavenging activity, inhibitory activity on  $\text{O}_2\bullet$ - production and inhibitory activity on NO production. This study is the first report of a BRL extract on its anti-inflammatory activity. For quality control of the ethanolic extract of BRL, a developed HPLC method for simultaneous quantification of the major markers was validated. The validated results showed good selectivity, linearity, precision, and accuracy with appropriate LOD and LOQ. This study is the first report of validation of an HPLC method of BRL for use in quality control of the BRL extract.

## Acknowledgments

The authors would like to thank the Thailand Science Research and Innovation Fundamental Fund, Center of Excellence in Applied Thai Traditional Medicine Research (CEATMR) and Bualuang ASEAN Chair Professorship,

Faculty of Medicine, Thammasat University, for financial support and the use of laboratory facilities.

## Conflict of interest statement

The authors declared no conflicts of interest in this study.

## Authors' contribution

A. Itharat conceived and supervised the project; C. Panchakul, W. Pipatrattanaseree, and P. Thongdeeying carried out the experiments and wrote the manuscript with support from A. Itharat; C. Kongkwamcharoen revised the manuscript; N.M. Davies, the Bualuang ASEAN Chair Professor, revised, reviewed, and analyzed the data and provided scientific insights and revisions to the manuscript. The final version of the manuscript was approved by all authors.

## REFERENCES

1. Gwozdziński K, Pieniazek A, Gwozdziński L. Reactive oxygen species and their involvement in red blood cell damage in chronic kidney disease. *Oxid Med Cell Longev*. 2021;2021:6639199. DOI: 10.1155/2021/6639199.
2. Motterlini R, Foresti R, Vandegriff K, Intaglietta M, Winslow R.M. Oxidative-stress response in vascular endothelial cells exposed to acellular hemoglobin solutions. *Am J Physiol* 1995;269(2 Pt 2):H648-55. DOI: 10.1152/ajpheart.1995.269.2.H648.
3. Zuo L, Prather E, Stetskiy M, Garrison DE, Meade JR, Peace TI, Zhou T. Inflammaging and Oxidative Stress in Human Diseases: From Molecular Mechanisms to Novel Treatments. *Int J Mol Sci*. 2019;20(18):4472. DOI: 10.3390/ijms20184472.
4. Wang C, Schuller Levis GB, Lee EB, Levis WR, Lee DW, Kim BS, Park SY, Park E. Platycodin D and D3 isolated from the root of *Platycodon grandiflorum* modulate the production of nitric oxide and secretion of TNF-[alpha] in activated RAW 264.7 cells *International Immunopharmacology* 2004;4(8): 1039-1049. DOI: 10.1016/j.intimp.2004.04.005.
5. Kurutas EB. The importance of antioxidants which play the role in cellular response against oxidative nitrosative stress: current state. *Nutr J*. 2016;5(1):71. DOI: 10.1186/s12937-016-0186-5.
6. Bjelakovic G, Nikolova D, Gluud LL, Simonetti RG, Gluud C. Antioxidant supplements for prevention of mortality in healthy participants and patients with various diseases. The Cochrane database of systematic reviews, 2012;2012(3), Cochrane Database Syst Rev 2008;(2):CD007176. DOI: 10.1002/14651858.CD007176.



7. Jeong SH, Kim BY, Kang HG, Ku HO, Cho JH. Effects of butylated hydroxyanisole on the development and functions of reproductive system in rats. *Toxicology* 2005;208(1):49-62. DOI: 10.1016/j.tox.2004.11.014.
8. Engin AB, Bukan N, Kurukahvecioglu O, Memis L, Engin A. Effect of butylated hydroxytoluene (E321) pretreatment versus l-arginine on liver injury after sub-lethal dose of endotoxin administration. *Environ Toxicol Pharmacol* 2011;32(3):457-464. DOI: 10.1016/j.etap.2011.08.014. Epub 2011 Sep 10.
9. Botterweck A, Verhagen H, Goldbohm R, Kleinjans J, Brandt PV. Intake of butylated hydroxyanisole and butylated hydroxytoluene and stomach cancer risk: results from analyses in the Netherlands Cohort Study. *Food Chem Toxicol.* 2000;38(7):599-605. DOI: 10.1016/s0278-6915(00)00042-9
10. Randhawa S, Bahna SL. Hypersensitivity reactions to food additives. *Curr Opin Allergy Clin Immunol.* 2009;9(3):278-283. DOI: 10.1097/ACI.0b013e32832b2632.
11. National Drug System Development Committee, "National list of essential medicines: herbal medicine list," National Drug System Development Committee, Thailand; 2012: 56-57. Available from: kpo.moph.go.th/webkpo/tool/Thaimed2555.pdf.
12. Yamasaki K, Hashimoto A, Kokusunya Y, Miyamoto T, Sato T. Electrochemical method for estimating the antioxidative effects of methanol extracts of crude drugs. *Chem Pharm Bull (Tokyo).* 1994;42(8):1663-1665. DOI: 10.1248/cpb.42.1663.
13. Re R, Pellegrini N, Proteggente A, Pannala A, Yang M, Rice-Evans C. Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radic Biol Med.* 1999;26(9-10):1231-1237. DOI: 10.1016/s0891-5849(98)00315-3.
14. Surarit W, Jansom C, Lerdvuthisophon N, Kongkham S, Hansakul P. Evaluation of antioxidant activities and phenolic subtype contents of ethanolic bran extracts of Thai pigmented rice varieties through chemical and cellular assays. *Int J Food Sci Tech.* 2015;50(4):990-998. DOI: 10.1111/IJFS.12703.
15. Makchuchit S, Rattarom R, Itharat A. The anti-allergic and anti-inflammatory effects of Benjakul extract (a Thai traditional medicine), its constituent plants and its some pure constituents using *in vitro* experiments. *Biomed Pharmacother.* 2017;89:1018-1026. DOI: 10.1016/j.biopha.2017.02.066.
16. Folin O, Ciocalteu V. On tyrosine and tryptophan determination in proteins. *J Biol Chem.* 1927;73(2):627-650. DOI: 10.1016/S0021-9258(18)84277-6.
17. Zhishen J, Mengcheng T, Jianming W. The determination of flavonoid contents in mulberry and their scavenging effects on superoxide radicals. *Food Chem.* 1999;64(4):555-559. DOI: 10.1016/S0308-8146(98)00102-2.
18. ICH Harmonised Tripartite Guideline, "Validation of analytical procedures: text and methodology Q2(R1)," Proceedings of the International Conference on Harmonisation of Technical Requirements for Registration of Pharmaceuticals for Human, ICH Harmonised Tripartite Guideline, Geneva, Switzerland; 2005. pp:6-13. Available from: [https://database.ich.org/sites/default/files/Q2\\_R1\\_Guideline.pdf](https://database.ich.org/sites/default/files/Q2_R1_Guideline.pdf).
19. Deng W, Wang Y, Liu Z, Cheng H, Xue Y. HemI: a toolkit for illustrating heatmaps. *PLoS One.* 2014;9(11):111988,1-5. DOI: 10.1371/journal.pone.0111988.
20. Batubara I, Mitsunaga T, Ohashi H. Screening antiacne potency of Indonesian medicinal plants: antibacterial, lipase inhibition, and antioxidant activities. *J Wood Sci.* 2009;55:230-235. DOI: 10.1007/s10086-008-1021-1.
21. Sasaki Y, Hosokawa T, Nagai M, Nagumo S. *In vitro* study for inhibition of NO production about constituents of Sappan lignum. *Bio Pharm Bull.* 2007;30(1):193-196. DOI: 10.1248/BPB.30.193.
22. Hu CM, Liu YH, Cheah KP, Li JS, Kei Lam CS, Yu WY, *et al.* Heme oxygenase-1 mediates the inhibitory actions of brazilin in RAW264.7 macrophages stimulated with lipopolysaccharide.85, 2009. *J Ethnopharmacol.* 2009;121(1):79-85. DOI: 10.1016/j.jep.2008.09.030.
23. Wu SQ, Otero M, Unger FM, Mary B, Goldring MB, Phrutivorapongkul A, *et al.* Anti-inflammatory activity of an ethanolic *Caesalpinia sappan* extract in human chondrocytes and macrophages. *J Ethnopharmacol.* 2011;138(2):364-372. DOI: 10.1016/j.jep.2011.09.011
24. da Costa FP, Puty B, Nogueira LS, Mitre GP, Santos S, Teixeira B, *et al.* Piceatannol increases antioxidant defense and reduces cell death in human periodontal ligament fibroblast under oxidative stress. *Antioxidants (Basel).* 2019;9(1):16,1-14. DOI: 10.3390/antiox9010016.
25. Hazra B, Sarkar R, Biswas S, Mandal N. Comparative study of the antioxidant and reactive oxygen species scavenging properties in the extracts of the fruits of *Terminalia chebula*, *Terminalia bellerica* and *Emblca officinalis*. *BMC Complement Altern Med.* 2010;10:20,1-15. DOI: 10.1186/1472-6882-10-20.
26. Jayesh K, Karishma R, Vysakh A, Gopika P, Latha MS. *Terminalia bellirica* (Gaertn.) Roxb fruit exerts anti-inflammatory effect via regulating arachidonic acid pathway and pro-inflammatory cytokines in lipopolysaccharide-induced RAW 264.7 macrophages. *Inflammopharmacology.* 2020;28:265-274. DOI: 10.1007/s10787-018-0513-x.
27. Nampoothiri SV, Prathapan A, Cherian OL, Raghu KG, Venugopalan VV, Sundaresan A. *In vitro* antioxidant and inhibitory potential of *Terminalia bellerica* and *Emblca officinalis* fruits against LDL oxidation and key enzymes linked to type 2 diabetes. *Food Chem Toxicol.* 2011;49(1):125-131. DOI: 10.1016/j.fct.2010.10.006. Epub 2010 Oct 14.
28. Rajkumari J, Dyavaiah M, Sudharshan SJ, Busi S. Evaluation of *in vivo* antioxidant potential of

- Syzygium jambos* (L.) Alston and *Terminalia citrina* Roxb. towards oxidative stress response in *Saccharomyces cerevisiae*. J Food Sci Technol. 2018;55(11):4432-4439.  
DOI: 10.1007/s13197-018-3355-z.
29. Fahmy NM, Al-Sayed E, Singab AN. Genus *Terminalia*: a phytochemical and biological review. Med Aromat Plants. 2015;4:1000218,1-21.  
DOI: 10.4172/2167-0412.1000218.
  30. Tanaka M, Kishimoto Y, Sasaki M, Sato A, Kamiya T, Kondo K, et al. *Terminalia bellirica* (Gaertn.) Roxb. extract and gallic acid attenuate LPS-induced inflammation and oxidative stress via MAPK/NF- $\kappa$ B and Akt/AMPK/Nrf2 pathways. Oxid Med Cell Longev. 2018;2018:9364364,1-15.  
DOI: 10.1155/2018/9364364.
  31. Chaiyasut C, Kesika P, Chaiyasut K, Sittiyuno P, Peerajan S, Sivamaruth BS. Total phenolic content and free radical scavenging activity of representative medicinal plants of Thailand. Asian J Pharm Clin Res. 2017;10(11):137-141.  
DOI: 10.22159/ajpcr.2017.v10i11.20741.
  32. Gonda R, Takeda T, Akiyama T. Studies on the constituents of *Anaxagorea luzonensis* A. GRAY. Chem Pharm Bull (Tokyo). 2000;48(8):1219-1222.  
DOI: 10.1248/cpb.48.1219.
  33. Gülçin I, Şat IG, Beydemir S, Elmastaş M, Küfrevioğlu OI. Comparison of antioxidant activity of clove (*Eugenia caryophyllata* Thunb) buds and lavender (*Lavandula stoechas* L.). Food Chem. 2004;87(3):393-400.  
DOI: 10.1016/j.foodchem.2003.12.008.
  34. Bamdad F, Kadivar M, Karamat J. Evaluation of phenolic content and antioxidant activity of Iranian caraway in comparison with clove and BHT using model systems and vegetable oil. Int J Food Sci Technol. 2006;40:20-27.  
DOI: 10.1111/j.1365-2621.2006.01238.x.
  35. Rodrigues TG, Fernandes A Jr, Sousa JP, Bastos JK, Sforcin JM. *In vitro* and *in vivo* effects of clove on pro-inflammatory cytokines production by macrophages. Nat Prod Res. 2009;23(4):319-326.  
DOI: 10.1080/14786410802242679.
  36. Haro-González JN, Castillo-Herrera GA, Martínez-Velázquez M, Espinosa-Andrews H. Clove essential oil (*Syzygium aromaticum* L. Myrtaceae): extraction, chemical composition, food applications, and essential bioactivity for human health. Molecules. 2021;26(21):6387.  
DOI: 10.3390/molecules26216387..
  37. Pérez-Rosés R, Risco E, Vila R, Peñalver P, Cañigueral S. Biological and nonbiological antioxidant activity of some essential oils. J Agric Food Chem. 2016;64(23):4716-4724.  
DOI: 10.1021/acs.jafc.6b00986.
  38. Yeh JL, Hsu JH, Hong YS, Wu JR, Liang JC, Wu BN, et al. Eugenol and glyceryl-isoegenol suppress LPS-induced iNOS expression by down-regulating NF-kappaB AND AP-1 through inhibition of MAPKS and AKT/IkappaBalpha signaling pathways in macrophages. Int J Immunopathol Pharmacol. 2011;24(2):345-356.  
DOI: 10.1177/039463201102400208.
  39. Mashhadi NS, Ghiasvand R, Askari G, Hariri M, Darvishi L, Mofid MR. Anti-oxidative and anti-inflammatory effects of ginger in health and physical activity: review of current evidence. Int J Prev Med. 2013;4(1):S36-S42.  
PMID: 23717767.
  40. Jagetia GC, Baliga MS, Venkatesh P, Ulloor JN. Influence of ginger rhizome (*Zingiber officinale* Rosc) on survival, glutathione and lipid peroxidation in mice after whole-body exposure to gamma radiation. Radiat Res 2003;160(5):584-592.  
DOI: 10.1667/rr3057.
  41. Haksar A, Sharma A, Chawla R, Kumar R, Arora R, Singh S, et al. *Zingiber officinale* exhibits behavioral radioprotection against radiation-induced CTA in a gender-specific manner. Pharmacol Biochem Behav. 2006;84(2):179-188.  
DOI: 10.1016/j.pbb.2006.04.008.
  42. Kim JK, Kim Y, Na KM, Surh YJ, Kim TY. [6]-Gingerol prevents UVB-induced ROS production and COX-2 expression *in vitro* and *in vivo*. Free Radic Res. 2007;41(5):603-614.  
DOI: 10.1080/10715760701209896.
  43. Lincy MP, Paulpriya K, Mohan VR. *In vitro* antioxidant activity of *Avicennia marina* (Forssk) Vierh pneumatophore (Avicenniaceae) Sci Res Rep. 2013;3(2):106-114.
  44. Okla MK, Alamri SA, Alatar AA, Hegazy AK, Al-Ghamdi AA, Ajarem JS, et al. Antioxidant, hypoglycemic, and neurobehavioral effects of a leaf extract of *Avicennia marina* on autoimmune diabetic mice. Evid Based Complement Alternat Med. 2019;2019:1263260,1-8.  
DOI: 10.1155/2019/1263260.
  45. Zhu F, Chen X, Yuan Y, Huang M, Sun H, Xiang W. The chemical investigations of the mangrove plant *Avicennia marina* and its endophytes. Open Nat Prod J. 2009;2:24-32.  
DOI: 10.2174/1874848100902010024].
  46. Jain R, Monthakantirat O, Tengamnuay P, De-Eknamkul W. Avicenninone C isolated from *Avicennia marina* exhibits 5 $\alpha$ -reductase-type 1 inhibitory activity using an androgenic alopecia relevant cell-based assay system. Molecules. 2014;19(5):6809-6821.  
DOI: 10.3390/molecules19056809.
  47. Meghwal M, Goswami TK. *Piper nigrum* and piperine: an update. Phytother Res. 2013;27(8):1121-1130.  
DOI: 10.1002/ptr.4972.
  48. Lee JY, Park W. Anti-inflammatory effect of myristicin on RAW 264.7 macrophages stimulated with polyinosinic-polycytidylic acid. Molecules. 2011;16(8):7132-7142.  
DOI: 10.3390/molecules16087132.
  49. Chayan P, Wichien J. Handbook of Thai traditional pharmacy. Vol 5. Amarin Publishing House: Bangkok, Thailand; 2004. P. 234-239
  50. Chayan P, Manamas C, Wichien J. Description of the Narai's drug. Amarin Publishing House: Bangkok Thailand; 1999. P. 150-165

51. Tep-Areenan P, Sawasdee P. The vasorelaxant effects of *Anaxagorea luzonensis* A. grey in the rat aorta. Int J Pharmacol. 2011;7(1):119-124. DOI: 10.3923/IJP.2011.119.124.
52. Karnick CR. Pharmacopeial standards of herbal plants. Vol 1. Sri Satguru Publications; 1994. P.38
53. Soonthorncharoennon N, Ruangwiset N. Quality of Thai pharmaceuticals from research to sustainable development. Vol. 1. Bangkok: Concept Medicus; 2008. P. 92.
54. World Health Organization. WHO monographs on selected medicinal plants. Vol 2. World Health Organization; 2002. P. 15-23.55. Department of Pharmacy and Botany Faculty of Pharmacy Mahidol University Encyclopedia of Herbs. Vol 1. Herbs Suan Sirirukhachat; 2000. P. 48-60.
55. Department of Pharmacy and Botany Faculty of Pharmacy Mahidol University Encyclopedia of Herbs. Herbs Suan Sirirukhachat. Vol 1; 2000.
56. China Pharmacopoeia Commission. Pharmacopoeia of the People's Republic of China. Vol 1. People Republic of China; 2005. P. 18.
57. Thaikasetsart (Internet), Samae Talay. 2020, Available from: www.thaikasetsart.com.
58. Soonthorncharoennon N, Ruangwiset N. Quality of Thai pharmaceuticals from research to sustainable development. Vol. 1. Bangkok: Concept Medicus; 2008. P. 371.
59. Thaicrudedrug [Internet]. Khrang; [cited 2022 Aug 4]. Available from: <http://www.thaicrudedrug.com/main.php?action=viewpage&pid=198>.
60. Soonthorncharoennon N, Ruangwiset N. Quality of Thai pharmaceuticals from research to sustainable development. Vol. 1. Bangkok: Concept Medicus; 2008. P. 230.
61. Aromdee C, Vorarat S, Benjamapriyagoon S. Physicochemical properties of Pikul Thai J Pharm Sci; 2005.
62. Soonthorncharoennon N, Ruangwiset N. Quality of Thai pharmaceuticals from research to sustainable development. Vol. 1. Bangkok: Concept Medicus; 2008. P. 323.
63. Thaicrudedrug (Internet). Samodeengu. 2022. Available from: <http://www.thaicrudedrug.com/main.php?action=viewpage&pid=131>
64. Nguyen Ngoc H, Nghiem DT, Pham TLG, Stuppner H, Ganzera M. Phytochemical and analytical characterization of constituents in *Urceola rosea* (Hook. & Arn.) D.J. Middleton leaves. J Pharm Biomed Anal. 2018;149:66-69. DOI: 10.1016/j.jpba.2017.10.031.