

Steamed Ginger Extract Exerts Anti-inflammatory Effects in *Helicobacter pylori*-infected Gastric Epithelial Cells through Inhibition of NF- κ B

Moon-Young Song¹, Da-Young Lee¹, Sang-Yong Park², Seul-A Seo², Jeong-Seung Hwang², Soo-Hyeon Heo², Eun-Hee Kim¹

¹Institute of Pharmaceutical Sciences, College of Pharmacy, CHA University, Seongnam, ²SD Biotechnologies Co., Ltd., Seoul, Korea

J Cancer Prev 2021;26(4):289-297
<https://doi.org/10.15430/JCP.2021.26.4.289>

The authors wish to notify readers of the following change: In page 290 - sentence in the "Materials and Methods":

<Before correction>

MATERIALS AND METHODS

Preparation of steamed ginger extract (GGE03)

GGE03 was prepared as previously published [5]. Briefly, ginger was washed three times with distilled water, dried at 50°C for 30 hours, and then steamed at: 2 to 2.5 kgf/cm², 97°C, for 2 hours.

<After correction>

MATERIALS AND METHODS

Preparation of steamed ginger extract (GGE03)

GGE03 was prepared as previously published [5]. Briefly, ginger was washed three times with distilled water, steamed (2 to 2.5 kgf/cm², 97°C) for 2 hours, and then dried at 50°C for 30 hours.

We apologize for any inconvenience our mistake may have caused.

Correspondence to Eun-Hee Kim, E-mail: ehkim@cha.ac.kr, <https://orcid.org/0000-0002-8523-0440>



This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Copyright © 2022 Korean Society of Cancer Prevention