

RESEARCH ARTICLE

Wolbachia effects on Rift Valley fever virus infection in *Culex tarsalis* mosquitoes

Brittany L. Dodson¹ , Elizabeth S. Andrews² ^{¶a}, Michael J. Turell² ^{¶b}, Jason L. Rasgon^{1,3,4*}


1 Department of Entomology, Pennsylvania State University, University Park, PA, United States of America, **2** Virology Division, United States Army Medical Research Institute of Infectious Diseases, Fort Detrick, MD, United States of America, **3** Center for Infectious Disease Dynamics, Pennsylvania State University, University Park, PA, United States of America, **4** Huck Institutes of the Life Sciences, Pennsylvania State University, University Park PA, United States of America

 These authors contributed equally to this work.

^{¶a} Current address: Vector-Borne Disease Section, Division of Communicable Diseases, Center for Infectious Diseases, California Department of Public Health, Sacramento, CA, United States of America

^{¶b} Current address: VectorID LLC, Frederick, MD, United States of America

* jlr54@psu.edu



 OPEN ACCESS

Citation: Dodson BL, Andrews ES, Turell MJ, Rasgon JL (2017) *Wolbachia* effects on Rift Valley fever virus infection in *Culex tarsalis* mosquitoes. PLoS Negl Trop Dis 11(10): e0006050. <https://doi.org/10.1371/journal.pntd.0006050>

Editor: Rhoel Ramos Dinglasan, University of Florida, UNITED STATES

Received: May 19, 2017

Accepted: October 18, 2017

Published: October 30, 2017

Copyright: This is an open access article, free of all copyright, and may be freely reproduced, distributed, transmitted, modified, built upon, or otherwise used by anyone for any lawful purpose. The work is made available under the [Creative Commons CC0](https://creativecommons.org/licenses/by/4.0/) public domain dedication.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: This study was funded by NIH grants R01AI116636 and R21AI128918 to JLR. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

Abstract

Innovative tools are needed to alleviate the burden of mosquito-borne diseases, and strategies that target the pathogen are being considered. A possible tactic is the use of *Wolbachia*, a maternally inherited, endosymbiotic bacterium that can (but does not always) suppress diverse pathogens when introduced to naive mosquito species. We investigated effects of somatic *Wolbachia* (strain *wAlbB*) infection on Rift Valley fever virus (RVFV) in *Culex tarsalis* mosquitoes. When compared to *Wolbachia*-uninfected mosquitoes, there was no significant effect of *Wolbachia* infection on RVFV infection, dissemination, or transmission frequencies, nor on viral body or saliva titers. Within *Wolbachia*-infected mosquitoes, there was a modest negative correlation between RVFV body titers and *Wolbachia* density, suggesting that *Wolbachia* may slightly suppress RVFV in a density-dependent manner in this mosquito species. These results are contrary to previous work in the same mosquito species, showing *Wolbachia*-induced enhancement of West Nile virus infection rates. Taken together, these results highlight the importance of exploring the breadth of pathogen modulations induced by *Wolbachia*.

Author summary

An integrated vector management program utilizes several practices, including pesticide application and source reduction, to reduce mosquito populations. However, mosquitoes are developing resistance to some of these methods and new control approaches are needed. A novel technique involves the bacterium *Wolbachia* that lives naturally in many insects. *Wolbachia* can be transferred to uninfected mosquitoes and can block pathogen transmission to humans, although in some circumstances pathogen enhancement has been observed. Additionally, *Wolbachia* is maternally inherited, allowing it to spread

quickly through uninfected field populations of mosquitoes. We studied the impacts of *Wolbachia* on Rift Valley fever virus (RVFV) in the naturally uninfected mosquito, *Culex tarsalis*. *Wolbachia* had no effect on the frequencies at which *Culex tarsalis* became infected with or transmitted RVFV. However, when we analyzed the relationship between *Wolbachia* densities and RVFV titers, we determined that high densities of *Wolbachia* were associated with no virus infection or low levels of virus, suggesting that *Wolbachia* might suppress RVFV at high densities. These results contrast with our previous study that showed *Wolbachia* enhances West Nile virus infection in *Culex tarsalis*. Together, these studies highlight the importance of studying *Wolbachia* effects on a variety of pathogens so that control methods that use *Wolbachia* are not impeded by unintended or off-target effects.

Introduction

Globally, mosquito-borne diseases are a major health burden. To decrease mosquito populations, control programs often use integrated vector management practices including adulticide and larvicide application, source reduction, and biological control [1]. However, these mosquito control methods are losing efficacy due to increasing insecticide resistance and changes in mosquito behavior [2–4]. With these concerns, novel and sustainable control methods are under investigation, including strategies that target the pathogen [5,6]. *Wolbachia* is a maternally-inherited endosymbiotic bacterium that infects a large number of insects and other invertebrates [7]. Infection by *Wolbachia* can cause broad effects on host physiology. For example, natural *Wolbachia* infections in fruit flies protect against pathogen-induced mortality [8,9]. When experimentally transferred to uninfected mosquitoes, *Wolbachia* can suppress infection or transmission of viruses, *Plasmodium* parasites, and filarial nematodes [10–13]. *Wolbachia* also manipulates host reproduction in ways that allow it to spread through and persist in insect populations [14].

Investigations using *Wolbachia*-infected mosquitoes as a control method for dengue virus are underway [15], and field trials in Australia have indicated that *Wolbachia* can spread to near-fixation in naturally uninfected populations of *Aedes aegypti* mosquitoes [16,17]. These *Wolbachia*-infected *Ae. aegypti* populations can persist years after release, and mosquitoes retain the dengue virus-blocking phenotype [18]. Similar field experiments are being conducted in several other countries, but not all have reported successful replacement of the uninfected population with *Wolbachia*-infected mosquitoes [19].

The effects of *Wolbachia*-induced pathogen interference may differ depending on mosquito species, *Wolbachia* strain, pathogen type, and environment conditions [20–22]. For example, in *Anopheles gambiae*, transient somatic infection of the *Wolbachia* strain *wAlbB* inhibits *Plasmodium falciparum* but enhances *Plasmodium berghei* parasites [22,23]. Enhancement phenotypes have been observed in *Anopheles*, *Culex*, and *Aedes* mosquitoes, and across several malaria species and virus families [20,22,24–27]. Thus, it is important to examine the range of *Wolbachia*-induced phenotypes so that efficacy of disease control efforts using *Wolbachia*-induced pathogen interference are not impeded.

Previous work has demonstrated that somatic *Wolbachia* (strain *wAlbB*) infections in *Culex tarsalis* (Yolo strain) enhanced West Nile virus (WNV) infection rates [24]. To better understand the range of *Wolbachia*-induced phenotypes, we investigated the effects of *wAlbB* on Rift Valley fever virus (RVFV) infection in *Cx. tarsalis*. RVFV is a member of the genus *Phlebovirus* in the family *Bunyaviridae* and predominately affects domestic ruminants, causing severe

economic losses in the livestock industry and human morbidity in Africa and the Middle East [28–30]. Additionally, models and laboratory studies have suggested the United States may have environmental conditions and mosquito vectors that would permit RVFV introduction and invasion [31–34]. *Cx. tarsalis* are abundant in the western U.S. and are highly competent laboratory vectors for RVFV [33–35]. We assessed the ability of *Wolbachia* to affect RVFV infection, dissemination, and transmission within *Cx. tarsalis* at two time points and evaluated relationships between viral titer and *Wolbachia* density in mosquitoes.

Materials and methods

Ethics statement

Mosquitoes were maintained on commercially obtained anonymous human blood using a membrane feeder (Biological Specialty Corporation, Colmar, PA). RVFV experiments were performed under biosafety-level 3 (BSL-3) and arthropod-containment level 3 (ACL3) conditions.

Research at the U.S. Army Medical Research Institute of Infectious Diseases (USAMRIID) was conducted under an Institutional Animal Care and Use Committee (IACUC) approved protocol in compliance with the Animal Welfare Act, PHS Policy, and other federal statutes and regulations relating to animals and experiments involving animals. The facility where this research was conducted is accredited by the Association for Assessment and Accreditation of Laboratory Animal Care, International and adheres to the principles stated in the *Guide for the Care and Use of Laboratory Animals*, National Research Council, 2011. The USAMRIID IACUC specifically approved this study.

Mosquitoes and *Wolbachia*

The *Culex tarsalis* colony used for all experiments was derived from field mosquitoes collected in Yolo County, CA in 2009. Mosquitoes were reared and maintained at 27°C ± 1°C, 12:12 hr light:dark diurnal cycle at 80% relative humidity in 30×30×30 cm cages. The *wAlbB* *Wolbachia* strain was purified from *An. gambiae* Sua5B cells according to published protocols [36]. *Wolbachia* viability and density was assessed using the LIVE/DEAD BacLight Bacterial Viability Kit (Invitrogen, Carlsbad, CA) and a hemocytometer. The RVFV vector competence experiment was replicated three times with different hamsters, and *wAlbB* concentrations for those replicates as follows: replicate one, 2.5×10^9 bacteria/ml; replicate two, 2.5×10^9 bacteria/ml; replicate three, 5.0×10^9 bacteria/ml.

Two- to 4-day-old adult female *Cx. tarsalis* were anesthetized with CO₂ and intrathoracically injected with approximately 0.1 μl of either suspended *wAlbB* or Schneider's insect media (Sigma Aldrich, Saint Louis, MO) as a control. Mosquitoes were provided with 10% sucrose *ad libitum* and maintained at 27°C in a growth chamber.

Vector competence for RVFV

RVFV strain ZH501 was isolated from the blood of a fatal human case in Egypt in 1977 [37]. Adult female Syrian hamsters were inoculated intraperitoneally with 0.2 ml of a suspension containing RVFV in diluent (10% heat-inactivated fetal bovine serum in Medium 199 with Earle's salts [Invitrogen], sodium bicarbonate, and antibiotics) containing approximately 10⁵ plaque-forming units (PFU) per ml of RVFV. Approximately 28–30 hr post-inoculation, infected hamsters were anesthetized with a suspension of ketamine, acepromazine, and xylazine. A single viremic hamster was placed across two 3.8-liter cardboard cages containing either *Wolbachia*-infected *Cx. tarsalis* or control-injected *Cx. tarsalis*, treatments to which the experimenter was blinded. Mosquitoes were allowed to feed for one hour. After this period,

hamsters were removed, a blood sample taken to determine viremia, and hamsters were euthanized.

After feeding, mosquitoes were anesthetized with CO₂ and examined for feeding status; partially or non-blood fed females were discarded. For all replicates, one blood fed mosquito from each treatment was sampled to test for input viral titers. Mosquitoes were sampled at 7 and 14 days post-blood feeding. On day 7, mosquitoes were anesthetized with CO₂ and had their legs removed; each set of legs was placed into one 2-ml microcentrifuge tube containing 1 ml of mosquito diluent (20% heat-inactivated fetal bovine serum [FBS] in Dulbecco's phosphate-buffered saline, 50 µg/ml penicillin streptomycin, and 2.5 µg/ml fungizone). Bodies were placed separately into 2-ml microcentrifuge tubes (Eppendorf, Hauppauge, NY) containing 1 ml of mosquito diluent. On day 14, bodies and legs were collected in the same manner as day 7, except that prior to placing bodies into microcentrifuge tubes, saliva was collected from mosquitoes by positioning the proboscis of each mosquito into a capillary tube containing approximately 10 µl of a 1:1 solution of 50% sucrose and FBS. After 30 minutes, the contents were expelled in individual microcentrifuge tubes containing 0.3 ml of mosquito diluent. A 5 mm stainless steel bead (Qiagen, Valencia, CA) was placed into all microcentrifuge tubes that contained mosquito bodies and legs, homogenized in a mixer mill (Retsch, Haan, Germany) for 30 seconds at 24 cycles per second, and centrifuged for 1 minute at 8000 rpm. All mosquito bodies, legs, and saliva were stored at -80°C until assayed.

Samples were tested for RVFV infectious particles by plaque assay on Vero cells according to previous published protocols [38]. Serial dilutions were prepared for all mosquito body, leg, and saliva samples. One hundred microliters of each dilution was inoculated onto Vero cell culture monolayers. Inoculated plates were incubated at 37°C for 1 hr and an agar overlay was added (1X EBME, 0.75% agarose, 7% FBS, 1% penicillin streptomycin, and 1% nystatin). Plates were incubated at 37°C for 3 days and then a second overlay (1X EBME, 0.75% agarose, and 4% neutral red) was added. Plaques were counted 24 hr after application of the second overlay and titers calculated.

Quantitative real-time PCR of *Wolbachia* density

To evaluate relationships between *Wolbachia* density and RVFV titer, we measured *wAlbB* levels in individual mosquitoes. DNA was extracted from 200 µl of mosquito body homogenate using the DNeasy blood and tissue kit (Qiagen) and used as template for qPCR on a Rotor-Gene Q (Qiagen) with the PerfeCta SYBR FastMix kit (Quanta Biosciences, Beverly, MA) or on ABI 7500 with Power SYBR green master mix (Applied Biosystems, Foster City, CA). The qPCR assays were performed in 10µl reactions and amplification was carried out using a standardized program at 95°C for 5 min, 40 cycles of 95°C for 10 sec, 60°C for 15 sec, and 72°C for 10 sec. *Wolbachia* DNA was amplified with primers Alb-GF and Alb-GR [39] and was normalized to the *Cx. tarsalis* actin gene by using qGene software [24,40]. qPCRs were performed in duplicate.

Statistical analyses

Infection, dissemination, and transmission rates were compared between *Wolbachia*-infected and control *Cx. tarsalis*, and between replicates with Fisher's exact tests. Due to violations of assumptions needed for parametric tests, Mann-Whitney U was used to compare the following data sets: RVFV body titers between *Wolbachia*-infected and control mosquitoes, RVFV body titers between RVFV-positive saliva and RVFV-negative saliva, RVFV body titers over time, and *Wolbachia* density over time. Unpaired t-tests were used to analyze data that passed normality tests, including the comparison of RVFV saliva titers between *Wolbachia*-infected and control mosquitoes. To determine relationships between *Wolbachia* density and RVFV body

titer, the Spearman rank correlation test was used, as assumptions for Pearson correlation were violated. All statistical analyses were performed in GraphPad Prism version 7 for Windows (GraphPad Software, San Diego, CA).

Results

Vector competence for RVFV

After all hamster feeds (i.e., replicates), one blood fed mosquito from each treatment was tested for input RVFV titers on the day of blood feeding. These day 0 RVFV titer results for *Wolbachia*-infected *Cx. tarsalis* were as follows: replicate 1, 2.50×10^2 PFU/ml; replicate 2, 7.00×10^6 PFU/ml; replicate 3, 1.00×10^2 PFU/ml. Day 0 results for control *Cx. tarsalis* were as follows: replicate 1, 5.00×10^2 PFU/ml; replicate 2, 1.05×10^7 PFU/ml; replicate 3, 1.00×10^2 PFU/ml. Viremias in the three hamsters were 10^4 , 10^9 , and 10^3 PFU/ml, respectively.

To determine RVFV vector competence of *Wolbachia*-infected and *Wolbachia*-uninfected *Cx. tarsalis*, we examined frequencies of RVFV-positive bodies (Fig 1A), legs (Fig 1B), and saliva (Fig 1C). Infection rate is the proportion of mosquito bodies that contained infectious RVFV. Dissemination and transmission rates are the proportion of infected mosquitoes with RVFV positive legs and saliva, respectively. Three replicate experiments were performed, and individual data from those experiments are available in S1 Table. Hamster viremia in replicate three was low and resulted in low mosquito infection rates. Replicate two infection frequencies were significantly higher than replicate one for both treatments and at both day 7 and day 14 ($P < 0.0001$) (S1 Table). However, Fig 1 and S1 Table show that across replicates and time points, *Wolbachia*-infected *Cx. tarsalis* infection, dissemination, and transmission rates did not differ significantly from *Wolbachia*-uninfected *Cx. tarsalis*. Thus, the data were pooled for further analysis.

RVFV body (Fig 2A) and saliva titers (Fig 2B) were determined for *Wolbachia*-infected and control *Cx. tarsalis*. There were no significant differences in RVFV body titer or saliva titer between *Wolbachia*-infected and control *Cx. tarsalis* at either day 7 (Fig 2A) or day 14 (Fig 2B). Additionally, when replicate data were pooled, both *Wolbachia*-infected and uninfected *Cx. tarsalis* that had higher RVFV body titers were more likely to have RVFV-positive saliva (S1 Fig).

Quantitative real-time PCR of *Wolbachia* (wAlbB) density

We performed qPCR from the DNA of each mosquito and determined *Wolbachia* density as an expression normalized to a reference gene, actin [40]. We used all samples that were positive for *Wolbachia* to analyze relationships between *Wolbachia* density and RVFV body titer; we combined data from all replicate experiments (Fig 3). Overall, there was a moderate, negative correlation between *Wolbachia* density and RVFV body titer at both day 7 and 14 (Fig 3A and 3B). Replicate two, the replicate with the highest infection rates, did not have significant correlations between *Wolbachia* density and RVFV body titer at either day 7 ($n = 14$, $r = 0.0516$, $p = 0.062$) or day 14 ($n = 27$, $r = 0.112$, $p = 0.577$) (raw data available S2 Table). *Wolbachia* density was also compared across time; *Wolbachia* concentration at day 14 was significantly higher than at day 7, consistent with *Wolbachia* replication in mosquitoes (S3 Fig).

Discussion

Wolbachia infection can have varied effects on viruses and parasites transmitted by mosquitoes. These effects can include moderate to complete pathogen inhibition, as well as pathogen




Fig 1. Effects of *Wolbachia* infection on RVFV vector competence frequencies in *Cx. tarsalis*. RVFV infection 7 and 14 days post-feeding (A), dissemination 7 and 14 days post-feeding (B), and transmission rates 14 days post-feeding (C) were compared between *Wolbachia*-infected and control *Cx. tarsalis*. Bars represent data pooled from three replicates. Error bars denote binomial confidence intervals. See S1 Table for replicate-specific analyses.

<https://doi.org/10.1371/journal.pntd.0006050.g001>

enhancement [17,22,24,41,42]. In a previous study, we found that *Wolbachia* strain *wAlbB* enhanced WNV infection frequency in *Cx. tarsalis* [24], although in that study, viral infection titers were not measured. To understand how widespread the *Wolbachia*-induced enhancement phenotype is in *Cx. tarsalis*, we studied *wAlbB* effects on RVFV, an important arthropod-borne virus with potential to invade the United States [43,44]. In contrast to our previous results, we found that *wAlbB* did not affect RVFV body or saliva titers, nor RVFV infection, dissemination, or transmission frequencies in *Cx. tarsalis*.

Wolbachia-mediated effects on pathogens may depend on *Wolbachia* density. Several studies have reported that high densities of *Wolbachia* are more likely than low densities to block viruses in *Drosophila* spp. and mosquitoes [45–48]. Similarly, we found a moderate, negative correlation between RVFV body titer and *Wolbachia* density. High *Wolbachia* levels were associated with RVFV negative mosquitoes or those with low RVFV body titers. The low numbers of mosquitoes at the high *Wolbachia* densities may explain why we did not see a *Wolbachia* effect on population level vector competence measures. However, our correlation data suggests that in this system, *Wolbachia* may suppress RVFV in a density-dependent manner.

In this *Cx. tarsalis*-*wAlbB* system, we have reported different effects of *Wolbachia* on vector competence for WNV and RVFV [24]. Other studies have found similar differences in *Wolbachia* phenotypes and suggested they may depend on various factors including environmental conditions, and pathogen type [20,49]. RVFV and WNV belong to different virus families and could interact with the mosquito host environment and *Wolbachia* in different ways. For example, a recent study suggested that the mosquito JAK/STAT pathway may not have the




Fig 2. Comparison of RVFV body and saliva titers between *Wolbachia*-infected and control *Cx. tarsalis*. At both 7 and 14 days post-blood meal, there were no significant differences in RVFV body titers (A) or saliva titers (B) of *Wolbachia*-infected *Cx. tarsalis* compared to control *Cx. tarsalis*. All replicates are combined in this figure; separate replicates are provided in supplementary materials (S2 Fig). Bars represent medians and bolded numbers above the data points denote sample sizes.

<https://doi.org/10.1371/journal.pntd.0006050.g002>




Fig 3. Correlation between RVFV body titer and *Wolbachia* levels in *Cx. tarsalis*. *Wolbachia* levels were normalized to the host gene actin. Normalized *Wolbachia* levels and RVFV body titer for each mosquito were plotted and analyzed with the Spearman rank correlation test to determine relationships. There was a moderate, negative correlation between RVFV body titer and *Wolbachia* levels at both day 7 ($n = 46$) (A) and day 14 ($n = 54$) (B) post-blood feeding (Fig 3). Data for all replicates were combined; see S2 Table for replicate-specific raw data.

<https://doi.org/10.1371/journal.pntd.0006050.g003>

same antiviral effects on closely related viruses [50]. Although the mosquitoes in our two studies have the same genetic background, they were reared in separate facilities and may have different microbiomes that may explain differences in vector competence [51].

While artificial feeding methods such as membrane feeders or pledgets allow one to easily standardize viral titers, previous studies on mosquito RVFV and other virus infections demonstrated significantly reduced mosquito infectivity using an artificial feeding system compared to a live infected animal [52, 53]. We therefore decided to use live hamsters infected with a highly virulent and epidemiologically relevant viral strain in our experiments. However, this more biologically relevant choice comes with a trade-off; it is impossible to standardize the viral titer in a live animal infection model. While all hamsters were infected with approximately the same amount of virus, they varied in their response to the infection, resulting in significant variation in final viral blood titers across experimental replicates that reduced statistical power.

Our study was performed with an adult microinjection model that generates mosquitoes transiently infected with *Wolbachia*. It remains to be seen whether this model reflects relationships between *Wolbachia* and viruses in *Cx. tarsalis* in a stable infection system. However, a recent study showed that both stable and transient *wAlbB* infections in *Ae. aegypti* produced similar results [45]. This suggests that our transient infection model may correlate with a stable infection in *Cx. tarsalis*.

Despite relatively modest measurable effects in these experiments, our results underscore the necessity of studying diverse *Wolbachia*-host-pathogen systems. It is becoming increasingly clear that one cannot extrapolate results of one system of interactions across all systems; every *Wolbachia*-host-pathogen system must be individually examined. Future studies should seek to understand the mechanisms underlying variation of *Wolbachia* protective effects across diverse mosquito species, viral/pathogen species, and *Wolbachia* strains.

Supporting information

S1 Table. Vector competence of *Cx. tarsalis* following a RVFV blood meal. RVFV infection, dissemination, and transmission frequencies were compared between *Wolbachia*-infected and

control mosquitoes. Replicates are displayed individually.
(XLSX)

S2 Table. Raw data for manuscript.
(XLSX)

S1 Fig. Comparison of RVFV body titers in *Cx. tarsalis* with virus present or absent in the saliva. RVFV body titers were compared between mosquitoes that tested positive or negative for RVFV in their saliva. For both *Wolbachia*-infected and control *Cx. tarsalis*, mosquitoes positive for RVFV in the saliva had significantly higher RVFV body titers compared to mosquitoes negative for virus in the saliva. There was no significant difference in RVFV body titer of transmitters between *Wolbachia*-infected and control mosquitoes ($p = 0.7692$). Data from three replicates were pooled and analyzed with Mann-Whitney U, and the bars represent medians.
(TIF)

S2 Fig. Comparison of RVFV body titers between treatments by replicate. RVFV body titers were compared between *Wolbachia*-infected and control mosquitoes for replicates 1 (A), 2 (B), and 3 (C). In all replicates, there were no significant differences in RVFV body titer between *Wolbachia*-infected and control mosquitoes. Data did not pass assumptions for normality and were analyzed with Mann-Whitney U, and sample sizes are denoted above data points.
(TIF)

S3 Fig. *Wolbachia* density over time. *Wolbachia* levels for each mosquito, determined by qPCR, were combined across all three replicates. *Wolbachia* levels were significantly higher at day 14 compared to day 7. Due to violations of normality, Mann-Whitney U was used for comparisons, bars are medians, and numbers above data points are sample sizes.
(TIF)

Acknowledgments

We thank the Cell Culture and Diagnostics Systems Divisions at the United States Army Medical Research Institute of Infectious Diseases (USAMRIID) for providing experimental materials and equipment. We also thank J. Hinson (USAMRIID) for providing technical assistance. The views of the authors ESA and MJT do not necessarily reflect the position of the Department of Defense or the Department of the Army.

Author Contributions

Conceptualization: Brittany L. Dodson, Michael J. Turell, Jason L. Rasgon.

Data curation: Jason L. Rasgon.

Formal analysis: Elizabeth S. Andrews, Michael J. Turell, Jason L. Rasgon.

Funding acquisition: Jason L. Rasgon.

Investigation: Brittany L. Dodson, Michael J. Turell.

Methodology: Brittany L. Dodson, Elizabeth S. Andrews, Jason L. Rasgon.

Project administration: Jason L. Rasgon.

Supervision: Michael J. Turell, Jason L. Rasgon.

Writing – original draft: Brittany L. Dodson, Elizabeth S. Andrews, Michael J. Turell, Jason L. Rasgon.

Writing – review & editing: Brittany L. Dodson, Michael J. Turell, Jason L. Rasgon.

References

1. Best Management Practices for Integrated Mosquito Management. In: American Mosquito Control Association [Internet]. 2009. Available: <http://www.mosquito.org/assets/Resources/PRTTools/Resources/bmpsformosquitomanagement.pdf>
2. Moiroux N, Gomez MB, Pennetier C, Elanga E, Djènontin A, Chandre F, et al. Changes in *Anopheles funestus* biting behavior following universal coverage of long-lasting insecticidal nets in Benin. *J Infect Dis*. 2012; 206: 1622–1629. <https://doi.org/10.1093/infdis/jis565> PMID: 22966127
3. Syed Z, Leal WS. Mosquitoes smell and avoid the insect repellent DEET. *Proc Natl Acad Sci*. 2008; 105: 13598–13603. <https://doi.org/10.1073/pnas.0805312105> PMID: 18711137
4. Ranson H, Lissenden N. Insecticide resistance in African *Anopheles* mosquitoes: A worsening situation that needs urgent action to maintain malaria control. *Trends Parasitol*. Elsevier Ltd; 2016; 32: 187–196. <https://doi.org/10.1016/j.pt.2015.11.010> PMID: 26826784
5. Read AF, Lynch PA, Thomas MB. How to make evolution-proof insecticides for malaria control. *PLoS Biol*. 2009; 7: e1000058. <https://doi.org/10.1371/journal.pbio.1000058> PMID: 19355786
6. Ricci I, Damiani C, Rossi P, Capone A, Scuppa P, Cappelli A, et al. Mosquito symbioses: from basic research to the paratransgenic control of mosquito-borne diseases. *J Appl Entomol*. 2011; 135: 487–493. <https://doi.org/10.1111/j.1439-0418.2011.01613.x>
7. Hilgenboecker K, Hammerstein P, Schlattmann P, Telschow A, Werren JH. How many species are infected with *Wolbachia*?—A statistical analysis of current data. *FEMS Microbiol Lett*. 2008; 281: 215–220. <https://doi.org/10.1111/j.1574-6968.2008.01110.x> PMID: 18312577
8. Teixeira L, Ferreira A, Ashburner M. The bacterial symbiont *Wolbachia* induces resistance to RNA viral infections in *Drosophila melanogaster*. *PLoS Biol*. 2008; 6: e1000002. <https://doi.org/10.1371/journal.pbio.1000002> PMID: 19222304
9. Hedges LM, Brownlie JC, O'Neill SL, Johnson KN. *Wolbachia* and virus protection in insects. *Science*. 2008; 322: 702. <https://doi.org/10.1126/science.1162418> PMID: 18974344
10. Kambris Z, Cook PE, Phuc HK, Sinkins SP. Immune activation by life-shortening *Wolbachia* and reduced filarial competence in mosquitoes. *Science*. 2009; 326: 134–136. <https://doi.org/10.1126/science.1177531> PMID: 19797660
11. Moreira LA, Iturbe-Ormaetxe I, Jeffery JA, Lu G, Pyke AT, Hedges LM, et al. A *Wolbachia* symbiont in *Aedes aegypti* limits infection with dengue, Chikungunya, and *Plasmodium*. *Cell*. 2009; 139: 1268–1278. <https://doi.org/10.1016/j.cell.2009.11.042> PMID: 20064373
12. Kambris Z, Blagborough AM, Pinto SB, Blagrove MSC, Godfray HCJ, Sinden RE, et al. *Wolbachia* stimulates immune gene expression and inhibits *Plasmodium* development in *Anopheles gambiae*. *PLoS Pathog*. 2010; 6: e1001143. <https://doi.org/10.1371/journal.ppat.1001143> PMID: 20949079
13. Andrews ES, Crain PR, Fu Y, Howe DK, Dobson SL. Reactive oxygen species production and *Brugia pahangi* survivorship in *Aedes polynesiensis* with artificial *Wolbachia* infection types. *PLoS Pathog*. 2012; 8. <https://doi.org/10.1371/journal.ppat.1003075> PMID: 23236284
14. Werren JH, Baldo L, Clark ME. *Wolbachia*: master manipulators of invertebrate biology. *Nat Rev Microbiol*. 2008; 6: 741–751. <https://doi.org/10.1038/nrmicro1969> PMID: 18794912
15. Bourtzis K, Dobson SL, Xi Z, Rasgon JL, Calvitti M, Moreira L a, et al. Harnessing mosquito-*Wolbachia* symbiosis for vector and disease control. *Acta Trop*. Elsevier B.V.; 2014; 132S: S150–S163. <https://doi.org/10.1016/j.actatropica.2013.11.004> PMID: 24252486
16. Hoffmann AA, Montgomery BL, Popovici J, Iturbe-Ormaetxe I, Johnson PH, Muzzi F, et al. Successful establishment of *Wolbachia* in *Aedes* populations to suppress dengue transmission. *Nature*. Nature Publishing Group; 2011; 476: 454–457. <https://doi.org/10.1038/nature10356> PMID: 21866160
17. Walker T, Johnson PH, Moreira LA, Iturbe-Ormaetxe I, Frentiu FD, McMeniman CJ, et al. The wMel *Wolbachia* strain blocks dengue and invades caged *Aedes aegypti* populations. *Nature*. Nature Publishing Group; 2011; 476: 450–453. <https://doi.org/10.1038/nature10355> PMID: 21866159
18. Frentiu FD, Zakir T, Walker T, Popovici J, Pyke AT, van den Hurk A, et al. Limited dengue virus replication in field-collected *Aedes aegypti* mosquitoes infected with *Wolbachia*. *PLoS Negl Trop Dis*. 2014; 8: e2688. <https://doi.org/10.1371/journal.pntd.0002688> PMID: 24587459
19. Nguyen TH, Nguyen H Le, Nguyen TY, Vu SN, Tran ND, Le TN, et al. Field evaluation of the establishment potential of wMelpop *Wolbachia* in Australia and Vietnam for dengue control. *Parasit Vectors*. 2015; 8: 563. <https://doi.org/10.1186/s13071-015-1174-x> PMID: 26510523

20. Murdock CC, Blanford S, Hughes GL, Rasgon JL, Thomas MB. Temperature alters *Plasmodium* blocking by *Wolbachia*. *Sci Rep*. 2014; 4: 3932. <https://doi.org/10.1038/srep03932> PMID: 24488176
21. van den Hurk AF, Hall-Mendelin S, Pyke AT, Frentiu FD, McElroy K, Day A, et al. Impact of *Wolbachia* on infection with chikungunya and yellow fever viruses in the mosquito vector *Aedes aegypti*. *PLoS Negl Trop Dis*. 2012; 6: e1892. <https://doi.org/10.1371/journal.pntd.0001892> PMID: 23133693
22. Hughes GL, Vega-Rodriguez J, Xue P, Rasgon JL. *Wolbachia* strain wAlbB enhances infection by the rodent malaria parasite *Plasmodium berghei* in *Anopheles gambiae* mosquitoes. *Appl Environ Microbiol*. 2012; 78: 1491–1495. <https://doi.org/10.1128/AEM.06751-11> PMID: 22210220
23. Hughes GL, Koga R, Xue P, Fukatsu T, Rasgon JL. *Wolbachia* infections are virulent and inhibit the human malaria parasite *Plasmodium falciparum* in *Anopheles gambiae*. *PLoS Pathog*. 2011; 7: e1002043. <https://doi.org/10.1371/journal.ppat.1002043> PMID: 21625582
24. Dodson BL, Hughes GL, Paul O, Matakachiero AC, Kramer LD, Rasgon JL. *Wolbachia* enhances West Nile virus (WNV) infection in the mosquito *Culex tarsalis*. Kittayapong P, editor. *PLoS Negl Trop Dis*. 2014; 8: e2965. <https://doi.org/10.1371/journal.pntd.0002965> PMID: 25010200
25. Baton LA, Pacidônio EC, Gonçalves DDS, Moreira LA. wFlu: Characterization and evaluation of a native *Wolbachia* from the mosquito *Aedes fluviatilis* as a potential vector control agent. *PLoS One*. 2013; 8: e59619. <https://doi.org/10.1371/journal.pone.0059619> PMID: 23555728
26. Zélé F, Nicot A, Berthomieu A, Weill M, Duron O, Rivero A, et al. *Wolbachia* increases susceptibility to *Plasmodium* infection in a natural system. *Proc R Soc*. 2014; 281. <https://doi.org/10.1098/rspb.2013.2837> PMID: 24500167
27. Graham RI, Grzywacz D, Mushobozi WL, Wilson K. *Wolbachia* in a major African crop pest increases susceptibility to viral disease rather than protects. *Ecol Lett*. 2012; 15: 993–1000. <https://doi.org/10.1111/j.1461-0248.2012.01820.x> PMID: 22731846
28. Bird BH, Ksiazek TG, Nichol ST, MacLachlan J. Zoonosis Update Rift Valley fever virus. *Vet Med Today*. 2009; 883–893.
29. Wilson M. Rift Valley fever virus ecology and the epidemiology of disease emergence. *Ann New York Acad Sci*. 1994; 15: 169–180.
30. Jupp PG, Kemp A, Grobbelaar A, Lema P, Burt FJ, Alahmed AM, et al. The 2000 epidemic of Rift Valley fever in Saudi Arabia: mosquito vector studies. *Med Vet Entomol*. 2002; 16: 245–252. PMID: 12243225
31. Konrad SK, Miller SN. A temperature-limited assessment of the risk of Rift Valley fever transmission and establishment in the continental United States of America. *Geospat Health*. 2012; 6: 161–170. <https://doi.org/10.4081/gh.2012.134> PMID: 22639118
32. Pfeffer M, Dobler G. Emergence of zoonotic arboviruses by animal trade and migration. *Parasit Vectors*. 2010; 3: 35. <https://doi.org/10.1186/1756-3305-3-35> PMID: 20377873
33. Turell MJ, Wilson WC, Bennett KE. Potential for North American mosquitoes (Diptera: Culicidae) to transmit Rift Valley fever virus. *J Med Entomol*. 2010; 47: 884–889. <https://doi.org/10.1603/ME10007> PMID: 20939385
34. Gargan TP, Clark GG, Dohm DJ, Turell MJ, Bailey CL. Vector potential of selected North American mosquito species for Rift Valley fever virus. *Am J Trop Med Hyg*. 1988; 38: 440–446. Available: <http://www.ncbi.nlm.nih.gov/pubmed/2895591> PMID: 2895591
35. Bohart RM and Washino R. Mosquitoes of California. 3rd ed. Berkeley: University of California Division of Agricultural Sciences; 1978.
36. Rasgon JL, Gamston CE, Ren X. Survival of *Wolbachia pipientis* in cell-free medium. *Appl Environ Microbiol*. 2006; 72: 6934–6937. <https://doi.org/10.1128/AEM.01673-06> PMID: 16950898
37. Meegan JM. The Rift Valley fever epizootic in Egypt 1977–78. *Trans R Soc Trop Med Hyg*. 1979; 73: 618–623.
38. Blow JA, Dohm DJ, Negley DL, Mores CN. Virus inactivation by nucleic acid extraction reagents. *J Virol Methods*. 2004; 119: 195–198. <https://doi.org/10.1016/j.jviromet.2004.03.015> PMID: 15158603
39. Ruang-areerate T, Kittayapong P. *Wolbachia* transinfection in *Aedes aegypti*: A potential gene driver of dengue vectors. *Proc Natl Acad Sci*. 2006; 103: 6–11.
40. Simon P. Q-Gene: processing quantitative real-time RT-PCR data. *Bioinformatics*. 2003; 19: 1439–1440. <https://doi.org/10.1093/bioinformatics/btg157> PMID: 12874059
41. Glaser RL, Meola MA. The native *Wolbachia* endosymbionts of *Drosophila melanogaster* and *Culex quinquefasciatus* increase host resistance to West Nile virus infection. *PLoS One*. 2010; 5: e11977. <https://doi.org/10.1371/journal.pone.0011977> PMID: 20700535
42. Aliota MT, Peinado SA, Velez ID, Osorio JE. The wMel strain of *Wolbachia* reduces transmission of Zika Virus by *Aedes aegypti*. *Sci Rep*. Nature Publishing Group; 2016; 6: 28792. <https://doi.org/10.1038/srep28792> PMID: 27364935

43. Barker CM, Niu T, Reisen WK, Hartley DM. Data-driven modeling to assess receptivity for Rift Valley fever virus. *PLoS Negl Trop Dis*. 2013; 7: e2515. <https://doi.org/10.1371/journal.pntd.0002515> PMID: 24244769
44. Konrad SK, Miller SN. Application of a degree-day model of West Nile virus transmission risk to the East Coast of the United States of America. *Geospat Health*. 2012; 7: 15–20. Available: <http://www.ncbi.nlm.nih.gov/pubmed/23242676> <https://doi.org/10.4081/gh.2012.100> PMID: 23242676
45. Joubert DA, O'Neill SL. Comparison of stable and transient *Wolbachia* infection models in *Aedes aegypti* to block dengue and West Nile viruses. *PLoS Negl Trop Dis*. 2017; 11: e0005275. <https://doi.org/10.1371/journal.pntd.0005275> PMID: 28052065
46. Joubert DA, Walker T, Carrington LB, De Bruyne JT, Kien DHT, Hoang NLT, et al. Establishment of a *Wolbachia* superinfection in *Aedes aegypti* mosquitoes as a potential approach for future resistance management. *PLoS Pathog*. 2016; 12: 1–19. <https://doi.org/10.1371/journal.ppat.1005434> PMID: 26891349
47. Osborne SE, Iturbe-Ormaetxe I, Brownlie JC, O'Neill SL, Johnson KN. Antiviral protection and the importance of *Wolbachia* density and tissue tropism in *Drosophila simulans*. *Appl Environ Microbiol*. 2012; 78: 6922–6929. <https://doi.org/10.1128/AEM.01727-12> PMID: 22843518
48. Martinez J, Longdon B, Bauer S, Chan Y-S, Miller WJ, Bourtzis K, et al. Symbionts commonly provide broad spectrum resistance to viruses in insects: A comparative analysis of *Wolbachia* strains. *PLoS Pathog*. 2014; 10: e1004369. <https://doi.org/10.1371/journal.ppat.1004369> PMID: 25233341
49. Hughes GL, Rivero A, Rasgon JL. *Wolbachia* can enhance *Plasmodium* infection in mosquitoes: Implications for malaria control? *PLoS Pathog*. 2014; 10: e1004182. <https://doi.org/10.1371/journal.ppat.1004182> PMID: 25187984
50. Jupatanakul N, Sim S, Angleró-Rodríguez YI, Souza-Neto J, Das S, Poti KE, et al. Engineered *Aedes aegypti* JAK/STAT pathway-mediated immunity to dengue virus. *PLoS Negl Trop Dis*. 2017; 11: e0005187. <https://doi.org/10.1371/journal.pntd.0005187> PMID: 28081143
51. Hegde S, Rasgon JL, Hughes GL. The microbiome modulates arbovirus transmission in mosquitoes. *Curr Opin Virol*. 2015; 15: 97–102. <https://doi.org/10.1016/j.coviro.2015.08.011> PMID: 26363996
52. Turell MJ. Reduced Rift Valley fever virus infection rates in mosquitoes associated with feedings. *Am J Trop Med Hyg*. 1988; 39: 597–602. PMID: 3207178
53. Weaver SC, Lorenz LH, Scott TW. Distribution of western equine encephalomyelitis virus in the alimentary tract of *Culex tarsalis* (Diptera: Culicidae) following natural and artificial blood meals. *J Med Entomol*. 1993; 30: 391–397. PMID: 8459416