

Association of retinopathy and intima media thickness of common carotid artery in type 2 diabetic patients

Ali Momeni, Mohamad Ali Dyani¹, Elnaz Ebrahimi², Morteza Sedehi³, Afsaneh Naderi⁴

Departments of Nephrology, ¹Radiology, ²Internal Medicine, ³Epidemiology and ⁴Ophthalmology, Shahrekord University of Medical Sciences, Shahrekord, Iran

Background: This study was carried out in order to evaluate the relationship between retinopathy and carotid intima-media thickness (CIMT). **Materials and Methods:** In a cross-sectional study, 154 diabetic patients who had a history of diabetic disease were evaluated in two equal groups of 77 patients with and without retinopathy, respectively. CIMT was evaluated in all of the patients. **Results:** Mean age of the patients was 59.65 ± 9.37 years. Mean CIMT of all patients was 0.84 ± 0.18 . CIMT of patients with retinopathy was significantly greater than patients without retinopathy ($P < 0.001$). CIMT also correlated with age, duration of diabetes, systolic blood pressure, blood urea nitrogen, and serum creatinine. **Conclusion:** CIMT may be used as a simple, available and noninvasive method for screening of macro and microvascular complication of diabetic patients.

Key words: Common carotid artery, diabetic retinopathy, intima-media thickness, sonography

How to cite this article: Momeni A, Dyani MA, Ebrahimi E, Sedehi M, Naderi A. Association of retinopathy and intima media thickness of common carotid artery in type 2 diabetic patients. *J Res Med Sci* 2015;20:393-6.

INTRODUCTION

Diabetes mellitus is the most common cause of end-stage renal disease, non-traumatic lower-limb amputation and blindness. Furthermore, diabetic retinopathy (DR) is the most important cause of visual loss worldwide. The prevalence of DR increases with duration of diabetes. Some other risk factors of DR development include poor glycemic control, type of diabetes, and the presence of associated disorders such as dyslipidemia, hypertension, smoking, pregnancy, and nephropathy.^[1,2] DR could present as non-proliferative retinopathy, proliferative retinopathy or macular edema, however, the most important cause of visual loss among these patients is macular edema.^[3,4] Retinopathy is essential for the diagnosis of diabetic nephropathy, but it presents in 90% and 60% of type 1 and 2 diabetic patients, respectively.^[5] Because the rate of progression of retinopathy may be rapid, and treatment can be beneficial for reduction of disease progression, it is important to screen diabetic patients regularly for the development of retinopathy, so in type 1 diabetes, after 5 years and in type 2 diabetes, at the time of diagnosis and then annually, retinopathy should be evaluated.^[6,7] One of the other complications of diabetes is generalized atherosclerosis which can be presented as ischemic heart disease, cerebrovascular

accident or peripheral vascular disease. Diabetic atherosclerosis can be detected by measurement of intima-media thickness (IMT) of common or internal carotid artery (CIMT). The CIMT was used to predict of cardiovascular outcomes in diabetic patients.^[8] The easy applicability and the noninvasive nature of B-mode ultrasonography make it suitable for using as a surrogate endpoint for measuring the atherosclerotic burden in people with cardiovascular risk factors.^[9] The mean CIMT values by the different incidences were reported as 1.26 ± 0.6 mm (transversal), 1.17 ± 0.54 mm (longitudinal anterolateral), and 1.18 ± 0.58 mm (longitudinal posterolateral).^[10] Increases in the thickness of the carotid IMT may be associated with an increased risk of myocardial infarction and stroke in old patients without a history of cardiovascular disease.^[11] Mean CIMT was reported a reliable marker of risk of ischemic stroke in type 2 diabetic patients and could be used as a simple noninvasive screening test for the assessment of atherosclerosis in these patients.^[12] Mean CIMT may be associated with age, systolic blood pressure (SBP), smoking, the ratio of low-density lipoprotein to high-density lipoprotein cholesterol, mean glycosylated hemoglobin value (HbA1c), and urinary albumin excretion rate.^[13] In type 2 diabetic patients, significant predictors of IMT progression included albuminuria, advanced

Address for correspondence: Dr. Mohamad Ali Dyani, Hajar Hospital, Shahrekord, Iran. E-mail: m.a.dyani@yahoo.com

Received: 08-06-2014; **Revised:** 21-01-2015; **Accepted:** 14-04-2015

age, male sex, smoking, and higher SBP.^[14] In diabetic patients without a history of clinical CVD, the presence of advanced stage of DR is associated with subclinical coronary artery disease.^[15,16] DR was shown as subclinical atherosclerosis marker, so retinopathy may warrant a more careful cardiovascular assessment even in the early stages of diabetes.^[17]

Intima-media thickness is a sensitive marker of early carotid atherosclerosis, so ultrasound CIMT measurement can be used to assess the cardiovascular risk and to determine indications for intensified diabetic treatment; on the other hand, DR is an early and reliable marker of microvascular disease, and probably diabetic nephropathy,^[18] so the aim of our study was evaluation of relationship between retinopathy and CIMT as two valuable noninvasive methods for early detection of micro and macrovascular complication of diabetes.

MATERIALS AND METHODS

In a cross-sectional study (2012), 154 diabetic patients who had a medical history and follow-up in Imam Ali Clinic of Shahrekord were enrolled in the study. They were divided into two equal groups of 77 patients that were case group (with retinopathy) and control group (without retinopathy). Diabetes is defined as fasting blood sugar equal or more than 126 mg/dL.^[19] DR is defined as a microvascular complication of diabetes that affect retinal arterioles and it has two forms as nonproliferative and proliferative retinopathy.^[20] Exclusion criteria were: Age lesser than 40, presence of cataract in the ophthalmologic exam that prevents retinopathy evaluation, noncooperative patients during the study. It has been explained to the participants that all information will be confidential and a written consent form were filled in by all patients. Common carotid intima-media thickness (CIMT) is defined as the largest distance

between the luminal intima interface and the medial adventitia interface that is located at 1 cm of the initiation of the common carotid artery^[21] that was measured by one sonographer in recombinant position and using Doppler sonography devices (Siemens, G50, Germany). Demographic criteria such as age, duration of diabetes, body mass index (BMI), SBP and diastolic blood pressure (DBP), and laboratory results including fasting blood sugar (FBS), Glycated hemoglobin (HbA1c), blood urea nitrogen (BUN), and creatinine (Cr) were checked. BMI was measured by formula (body weight [Kg]/Length [m²]) and laboratory tests were conducted by Biotechnica Instruments (BT 3000). Collected data were entered to SPSS (Statistical Package for the Social Sciences, version 19.0, SPSS Inc, Chicago, Ill, USA) and analysis was done by *t*-test, Chi-square test, and Pearson correlation. This study was the result of research project number of 986, which approved by the research committee of Shahrekord University of Medical Sciences. Data are presented as mean (standard deviation).

RESULTS

Mean age of the patients was 59.65 ± 9.37 years. Mean age of the patients with retinopathy (Group 1) was 62.5 ± 9.75 years and in the patients without retinopathy (Group 2) was 58 ± 10 years (*P* = 0.006). Mean CIMT of all patients was 0.84 ± 0.18 and in the female and male patients were 0.82 ± 0.16 mm and 0.88 ± 0.22 mm respectively (*P* = 0.03). In all of the patients, CIMT was associated with age, duration of diabetes (based on history), SBP, serum BUN, and Cr [Table 1]. The number of male and female in Group 1 was 34 (44.2%) and 43 (55.8%) and in Group 2 were 21 (27.3%) and 56 (72.7%), respectively. CIMT of the patients of Group 1 was significantly greater than Group 2 (*P* < 0.001). Mean HbA1c in Group 1 and 2 were 7.20 ± 1.21 and 7.04 ± 1.33, respectively (*P* = 0.39). Table 2 showed that in patients of Group 1, CIMT was correlated only with SBP (*P* = 0.01); however in the

Table 1: Comparison of different variables in two groups of the patients

Variables	Age (years)	BMI (kg/m ²)	SBP (mmHg)	DBP (mmHg)	BUN (mg/dL)	Cr (mg/dL)	HbA1c (%)	CIMT (mm)
Group 1	62.01±9.08	28.94±4.73	143.37±20.04	83.11±11.5	26.31±12.79	1.45±0.75	7.39±1.19	0.95±0.18
Group 2	57.29±9.84	29.63±4.07	134.94±19.03	84.35±15.6	21.13±8.79	1.1±0.36	7.01±1.29	0.73±0.11
<i>P</i>	0.002	0.33	0.008	0.57	0.004	<0.0001	0.39	0.0001

Group 1 = Diabetic patients with retinopathy; Group 2 = Diabetic patients without retinopathy. BMI = Body mass index; SBP = Systolic blood pressure; DBP = Diastolic blood pressure; BUN = Blood urea nitrogen; Cr = Creatinine; CIMT = Carotid intima-media thickness; HbA1c = Glycosylated hemoglobin

Table 2: Association of CIMT with different variables in the patients

Variable	Age	BMI	SBP	DBP	BUN	Cr	HbA1c	FBS
Patients with retinopathy								
Correlation coefficient	0.14	0.003	0.276	0.179	0.016	0.067	0.069	0.183
<i>P</i>	0.2	0.981	0.01	0.12	0.89	0.56	0.55	0.11
Patients without retinopathy								
Correlation coefficient	0.256	0.179	0.186	0.274	0.19	0.008	0.09	0.07
<i>P</i>	0.2	0.11	0.11	0.03	0.09	0.94	0.42	0.51

BMI = Body mass index; SBP = Systolic blood pressure; DBP = Diastolic blood pressure; BUN = Blood urea nitrogen; Cr = Creatinine; CIMT = Carotid intima-media thickness; HbA1c = Glycosylated hemoglobin; FBS = Fasting blood sugar

patients of Group 2, CIMT was correlated with DBP. To remove confounding bias effect, linear regression model was used and the correlation between DR and CIMT was illustrated.

DISCUSSION

We found a significant association between CIMT and retinopathy in type two diabetic patients. Prevalence of DR in type 2 diabetic patients was reported 34.6% in Yau *et al.* study with no difference in male and female,^[22] whereas Zhang *et al.* showed slightly more common in male patients.^[2] Besides, Harris *et al.* found a greater prevalence and severity of DR in non-hispanic and Mexican Americans with type 2 diabetes.^[23] There are a few studies about correlation of CIMT and retinopathy in diabetic patients, for example, Miyamoto in evaluation of 102 diabetic patients showed the significant correlation between retinopathy and common carotid artery thickness,^[24] also Torres *et al.* in a study on 173 patients with hypertension showed, significant and independent association of carotid intima-media thickness with arteriolar caliber of retina.^[25] In our study CIMT was associated with age, duration of diabetes, SBP, serum BUN, and Cr whereas Cardoso *et al.* found the correlation of CIMT with age, male sex, smoking status, and ambulatory blood pressure.^[26] Ogawa *et al.* in a study on 634 type 2 diabetic patients reported the positive correlation of CIMT and patients BMI, also he found correlation between maximum BMI and retinopathy, but we didn't find this association in the patients. The reason of this discrepancy may be due to different number of patients in two studies.^[27] Correlation of CIMT and HTN was reported in some studies,^[12,26] also in our study CIMT was correlated with SBP in patients with retinopathy and with DBP in patients without retinopathy, however, Alizadeh *et al.* in a study on 40 diabetic patients did not find these correlation.^[28] Similar to our results, in Ogawa *et al.* study also CIMT had not correlation with control of blood sugar.^[29] About association of HbA1c and CIMT we did not find a significant correlation. Furthermore similar result was found by Choi *et al.* in the study on 370 type 2 diabetic patients.^[30]

Small sample size is a limitation in our study, so we suggest more studies to be carried out using larger sample size. Therefore, the relationship between CIMT and other complications of diabetes such as microalbuminuria, macroalbuminuria or neuropathy need to be evaluated.

CONCLUSION

In diabetic patients, CIMT is a marker of atherosclerosis and macrovascular damage which has had a correlation with DR (as a potential reliable marker of microvascular damage), so we may use sonographic measurement of CIMT as a simple,

available, and noninvasive method for screening of macro and microvascular complications among diabetic patients.

ACKNOWLEDGMENTS

The research project number of the study was 986. We acknowledge all staff of Hajar Sonography Center and the patients for their cooperation in this study.

AUTHOR'S CONTRIBUTION

All authors have contributed in designing and conducting the study. All authors have assisted in preparation of the first draft of the manuscript or revising it critically for important intellectual content. All authors have read and approved the content of the manuscript and confirmed the accuracy or integrity of any part of the work.

REFERENCES

1. Davis MD, Fisher MR, Gangnon RE, Barton F, Aiello LM, Chew EY, *et al.* Risk factors for high-risk proliferative diabetic retinopathy and severe visual loss: Early Treatment Diabetic Retinopathy Study Report #18. *Invest Ophthalmol Vis Sci* 1998;39:233-52.
2. Zhang X, Saaddine JB, Chou CF, Cotch MF, Cheng YJ, Geiss LS, *et al.* Prevalence of diabetic retinopathy in the United States, 2005-2008. *JAMA* 2010;304:649-56.
3. McMeel JW, Trempe CL, Franks EB. Diabetic maculopathy. *Trans Sect Ophthalmol Am Acad Ophthalmol Otolaryngol* 1977;83:OP476-87.
4. Wilkinson CP, Ferris FL 3rd, Klein RE, Lee PP, Agardh CD, Davis M, *et al.* Proposed international clinical diabetic retinopathy and diabetic macular edema disease severity scales. *Ophthalmology* 2003;110:1677-82.
5. Fong DS, Aiello L, Gardner TW, King GL, Blankenship G, Cavallerano JD, *et al.* Retinopathy in diabetes. *Diabetes Care* 2004;27 Suppl 1:S84-7.
6. Frank RN. Diabetic retinopathy. *N Engl J Med* 2004;350:48-58.
7. Mizutani M, Kern TS, Lorenzi M. Accelerated death of retinal microvascular cells in human and experimental diabetic retinopathy. *J Clin Invest* 1996;97:2883-90.
8. Polak JF, Pencina MJ, Pencina KM, O'Donnell CJ, Wolf PA, D'Agostino RB Sr. Carotid-wall intima-media thickness and cardiovascular events. *N Engl J Med* 2011;365:213-21.
9. George JM, Bhat R, Pai KM, Arun S, Jeganathan J. The carotid intima media thickness: A predictor of the clinical coronary events. *J Clin Diagn Res* 2013;7:1082-5.
10. Casella IB, Presti C, Porta RM, Sabbag CR, Bosch MA, Yamazaki Y. A practical protocol to measure common carotid artery intima-media thickness. *Clinics (Sao Paulo)* 2008;63:515-20.
11. Nichols WW, Pepine CJ, O'Rourke MF. Carotid-artery intima and media thickness as a risk factor for myocardial infarction and stroke. *N Engl J Med* 1999;340:1762-3.
12. Kota SK, Mahapatra GB, Kota SK, Naveed S, Tripathy PR, Jammula S, *et al.* Carotid intima media thickness in type 2 diabetes mellitus with ischemic stroke. *Indian J Endocrinol Metab* 2013;17:716-22.
13. Nathan DM, Lachin J, Cleary P, Orchard T, Brillon DJ, Backlund JY, *et al.* Intensive diabetes therapy and carotid intima-media thickness in type 1 diabetes mellitus. *N Engl J Med* 2003;348:2294-303.
14. Polak JF, Backlund JY, Cleary PA, Harrington AP, O'Leary DH, Lachin JM, *et al.* Progression of carotid artery intima-media

- thickness during 12 years in the Diabetes Control and Complications Trial/Epidemiology of Diabetes Interventions and Complications (DCCT/EDIC) study. *Diabetes* 2011;60:607-13.
15. Kawasaki R, Cheung N, Islam FM, Klein R, Klein BE, Cotch MF, *et al.* Is diabetic retinopathy related to subclinical cardiovascular disease? *Ophthalmology* 2011;118:860-5.
 16. Marin-Sanabria EA, Kondoh T, Yamanaka A, Kohmura E. Ultrasonographic screening of carotid artery in patients with vascular retinopathies. *Kobe J Med Sci* 2005;51:7-16.
 17. Son JW, Jang EH, Kim MK, Kim IT, Roh YJ, Baek KH, *et al.* Diabetic retinopathy is associated with subclinical atherosclerosis in newly diagnosed type 2 diabetes mellitus. *Diabetes Res Clin Pract* 2011;91:253-9.
 18. Bartman W, Pierzchala K. Clinical determinants of carotid intima-media thickness in patients with diabetes mellitus type 2. *Neurol Neurochir Pol* 2012;46:519-28.
 19. Canadian Diabetes Association Clinical Practice Guidelines Expert Committee, Goldenberg R, Punthakee Z. Definition, classification and diagnosis of diabetes, prediabetes and metabolic syndrome. *Can J Diabetes* 2013;37 Suppl 1:S8-11.
 20. Frank RN. Diabetic retinopathy. *N Engl J Med* 2004; 350:48.
 21. Stein JH, Korcarz CE, Hurst RT, Lonn E, Kendall CB, Mohler ER, *et al.* Use of carotid ultrasound to identify subclinical vascular disease and evaluate cardiovascular disease risk: A consensus statement from the American Society of Echocardiography Carotid Intima-Media Thickness Task Force. Endorsed by the Society for Vascular Medicine. *J Am Soc Echocardiogr* 2008;21:93-111.
 22. Yau JW, Rogers SL, Kawasaki R, Lamoureux EL, Kowalski JW, Bek T, *et al.* Global prevalence and major risk factors of diabetic retinopathy. *Diabetes Care* 2012;35:556-64.
 23. Harris MI, Klein R, Cowie CC, Rowland M, Byrd-Holt DD. Is the risk of diabetic retinopathy greater in non-Hispanic blacks and Mexican Americans than in non-Hispanic whites with type 2 diabetes? A U.S. population study. *Diabetes Care* 1998;21:1230-5.
 24. Miyamoto M, Kotani K, Okada K, Fujii Y, Konno K, Ishibashi S, *et al.* The correlation of common carotid arterial diameter with atherosclerosis and diabetic retinopathy in patients with type 2 diabetes mellitus. *Acta Diabetol* 2012;49:63-8.
 25. Torres FS, Fuchs SC, Maestri MK, Fuchs FD, Oliveira MM, Moreira LB, *et al.* Association between carotid intima-media thickness and retinal arteriolar and venular diameter in patients with hypertension: A cross-sectional study. *Atherosclerosis* 2013;229:134-8.
 26. Cardoso CR, Marques CE, Leite NC, Salles GF. Factors associated with carotid intima-media thickness and carotid plaques in type 2 diabetic patients. *J Hypertens* 2012;30:940-7.
 27. Ogawa K, Ueda K, Sasaki H, Yamasaki H, Okamoto K, Wakasaki H, *et al.* History of obesity as a risk factor for both carotid atherosclerosis and microangiopathy. *Diabetes Res Clin Pract* 2004;66 Suppl 1:S165-8.
 28. Alizadeh A, Roudbari A, Heidarzadeh A, Babaei Jandaghi A, Bani Jamali M. Ultrasonic measurement of common carotid intima-media thickness in type 2 diabetic and non-diabetic patients. *Iran J Radiol* 2012;9:79-82.
 29. Ogawa Y, Uchigata Y, Iwamoto Y. Progression factors of carotid intima-media thickness and plaque in patients with long-term, early-onset type 1 diabetes mellitus in Japan: Simultaneous comparison with diabetic retinopathy. *J Atheroscler Thromb* 2009;16:821-8.
 30. Choi SW, Shin MH, Yun WJ, Kim HY, Lee YH, Kweon SS, *et al.* Association between hemoglobin A1c, carotid atherosclerosis, arterial stiffness, and peripheral arterial disease in Korean type 2 diabetic patients. *J Diabetes Complications* 2011;25:7-13.

Source of Support: Nil, **Conflict of Interest:** None declared.