

Contents lists available at [ScienceDirect](http://www.sciencedirect.com)

International Journal of Surgery Case Reports

journal homepage: www.casereports.com

Ambiguous genitalia–A social dilemma in Bangladesh: A case report

Mohiuddin Ahsanul Kabir Chowdhury^{a,*}, Rashidul Anwar^a, Arnab Saha^b^a International Centre for Diarrhoeal Diseases Research, 68, Shaheed Tajuddin Ahmed Sarani, Mohakhali, Dhaka, 1212, Bangladesh^b Interfaith Medical Center, 1545 Atlantic Ave, Brooklyn, NY 11213, United States

ARTICLE INFO

Article history:

Received 6 October 2017

Received in revised form

28 November 2017

Accepted 28 November 2017

Available online 8 December 2017

Keywords:

Disorder of sexual development (DSD)

Ambiguous genitalia

Gender assignment

Management

Bangladesh

ABSTRACT

INTRODUCTION: Ambiguous Genitalia is a form of birth defect in which sex of the newborn cannot be readily distinguishable because of atypical appearance of the external genitalia.

CASE REPORT–CLINICAL FINDINGS: The patient, an 8 months old baby, was identified as a female baby since birth; but, some senior members of their neighborhood raised confusion regarding sex differentiation of the child. On examination, there was enlarged clitoris or micropenis, bifid scrotum or labioscrotal fold, and separate presence of urethral and vaginal orifices. Right testis was found in the middle of right inguinal canal and left testis was found near the deep inguinal ring.

CASE REPORT–DIAGNOSIS, THERAPEUTIC INTERVENTION AND OUTCOME: The ultrasonogram showed a rudimentary uterus measuring 26.7×7.27 mm with no ovary and the right testis was found in the mid inguinal canal while the left testis was in the deep inguinal ring. Diagnostic endoscopic procedure showed normal appearing testes in deep inguinal ring, about 2 cm vaginal remnant with normal urethra and bladder. Hormonal study revealed low serum testosterone with normal DHT and high oestradiol while the karyotyping revealed 46XY. After 3 months of hormonal treatment, multiple surgical interventions were taken to correctly determine the sex of the child. Later on the patient was discharged with follow up advices.

CONCLUSION: 'Ambiguous genitalia' is considered as a taboo in Bangladesh; hence, the social awareness regarding this curable disorder is a timely need in Bangladesh.

© 2017 The Authors. Published by Elsevier Ltd on behalf of IJS Publishing Group Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Ambiguous genitalia may be a consequence of a disorder of sexual development resulting in a defect at birth where the external genitals do not have a typical resemblance to that of a boy or a girl [1]. The common causes of ambiguous genitalia in genetic females are congenital adrenal hyperplasia (CAH), maternal ingestion of substances with male hormone activity during pregnancy, male hormone producing tumour in female foetus or mother, whereas possible causes in genetic males are Leydig cell aplasia, 5-alpha-reductase deficiency, androgen insensitivity syndrome, and maternal ingestion of substances with female hormone activity during pregnancy [2,3]. Congenital Adrenal Hyperplasia seems to be more common in people of Hispanic, Slavic, Italian and European Jewish ancestry which is followed by mixed gonadal dysgenesis (MGD) [3]. The incidence of genital ambiguity is 1 per 4500, though some degree of male under-virilisation or female virilisation may be present in 2% of live births [4,5]. A study showed that the incidence is about 1 in 18,000 births in UK [3]. There have been insufficient data regarding incidences of ambiguous genitalia or other simi-

lar birth defects in the low and middle income countries (LMICs) including Bangladesh. Most cases of ambiguous genitalia can be diagnosed at birth by equivocal features of external genitalia [6]. The outcome of the treatment of ambiguous genitalia has been reported differently in different studies. A study showed that most of the participants were content with their body image. 90% of men and 83% of women had sexual experience with a partner. Both men and women were satisfied with their sexual function. [7]. For majority of cases of extreme genital ambiguity in 46,XY individuals, both male or female sex of rearing could lead to successful long-term outcome. [7]. However, another study showed that there has been unsuccessful sexual life of the treated patients [8]. A table of the literature reports on DSD with title, author and journal details is provided in Table 1 [7–18].

But, most of these reports are mainly from the western world. The literature review suggests that there is insufficient data regarding ambiguous genitalia from LMICs. This article portrays a report of a case of ambiguous genitalia which was managed at a tertiary level public health facility of Bangladesh. In this article, the case study has been reported according to the SCARE criteria [20].

2. Case report

In May 2014, an 8 months old baby hailing from a remote district of Bangladesh presented with the complaints of difficulty in identification of sex by the parents for the previous 02 months.

* Corresponding author at: Maternal and Child Health Division, icddr,b, 68, Shaheed Tajuddin Ahmed Sarani, Mohakhali, Dhaka-1212, Bangladesh.

E-mail addresses: mohiuddin.chowdhury@icddr.org, makchowdhury.icddr@gmail.com (M.A.K. Chowdhury).

Table 1
Literature reports on DSD and similar cases.

Authors	Title	Reference details
Georgette Beatriz De Paula et al.	408 Cases of Genital Ambiguity Followed by Single Multidisciplinary Team during 23 Years: Etiologic Diagnosis and Sex of Rearing	International Journal of Endocrinology [11]
V. Anantha Kumari, A. Vani	Ambiguous genitalia: a clinical and chromosomal study	International Journal of Research in Medical Sciences [14]
Carol AB Warren Thyen U. et al	Gender reassignment surgery in the 18th century: A case study Epidemiology and Initial Management of Ambiguous Genitalia at Birth in Germany	Sexualities [17] Hormone Research in Paediatrics [16]
Claude J. Migeon et al.	Ambiguous Genitalia With Perineo-scrotal Hypospadias in 46,XY Individuals: Long-Term Medical, Surgical, and Psychosexual Outcome	Paediatrics [7]
Hazel Glenn Beh, Milton Diamond	An Emerging Ethical and Medical Dilemma: Should Physicians Perform Sex Assignment Surgery on Infants with Ambiguous Genitalia?	Michigan Journal of Gender & Law [19]
Milton Diamond William George Reiner Alice Domurat Dreger	Paediatric management of ambiguous and traumatized genitalia Assignment of sex in neonates with ambiguous-genitalia “Ambiguous Sex”—or Ambivalent Medicine?: Ethical Issues in the Treatment of Intersexuality	The Journal of Urology [12] Current Opinion in Paediatrics [15] The Hastings Center Report [18]

According to the statement of the parents, the baby was delivered at home and they had identified their child as a female baby since birth. But, within the previous two months of reporting at this hospital, some senior members of their neighborhood raised confusion regarding sex differentiation of the child which ultimately drove them to the department of Paediatric Surgery of a tertiary medical college hospital for proper advice which is about 250 kilometers away from their residence. On examination, there was enlarged clitoris or micropenis, bifid scrotum or labioscrotal fold, and separate presence of urethral and vaginal orifices. Right testis was found in the middle of right inguinal canal and left testis was found near the deep inguinal ring. The ultrasonogram showed a rudimentary uterus measuring 26.7×7.27 mm with no ovary and the right testis was found in the mid inguinal canal and left testis was in the deep inguinal ring. Diagnostic endoscopic procedure showed normal appearing testes in deep inguinal ring, about 2 cm vaginal remnant with normal urethra and bladder. Hormonal study revealed low serum testosterone with normal DHT and high oestradiol while the karyotyping revealed 46XY.

According to the decision of medical board comprising experts from different fields Testosterone was given in injectable form for consecutive 3 months as a part of multiphasic intervention plan and parents were counselled and convinced to convert her into male sex. After the hormonal treatment first stage of surgical intervention was performed during which vaginal remnant was found to be capacious measuring more than 5 cm. Right testis was found to be softer in consistency and smaller in size. Biopsy was taken from the same gonad and orchiopexy was performed. In the second setting, separation and closure of vaginal orifice was performed after having the biopsy report which conferred that this testicular tissue have degenerative changes. During this phase chordee was corrected as well followed by placement of buccal mucosal strip for urethral plate. In the third and final setting of surgical intervention urethroplasty, left orchiopexy and scrotal reconstruction were performed. Later on the patient was discharged with follow up advices to contact endocrinologist.

3. Discussion

3.1. Standard treatment of DSD and its multidisciplinary treatment in the study settings

Infants born with the disease represent a neonatal medical emergency for physical, social and psychological reasons and demand a multidisciplinary team for the management [4,21,22]. The team must include aptly trained paediatric surgeons or urologist, paediatric endocrinologist or neonatologists with their support staff along with backing of geneticists, psychologists, ethi-

cists, biochemists, counsellors and gynaecologists. [21,23]. The tertiary level hospitals (commonly known as medical college hospitals) are therefore required to manage the cases of DSD. In Bangladesh, currently, there are 31 medical colleges which have such mix of health care providers, although the genre of geneticists is usually scarce in the country. Accurate diagnosis, gender assignment, medical and surgical treatment and sharing of medical information with parents and patients are four key issues for management of ambiguous genitalia [24]. The proper diagnostic procedures should exhibit external appearance, internal anatomy, genetic make-up, and hormonal profile [6]. History has been of paramount importance to establish the diagnosis and start the management [2,6,25]. In the case presented in this article, history was taken quite carefully. Some of the key issues during the child birth could not be explored because of the delayed presentation of the baby. There might have been recall bias as well regarding extracting history from the patient's parents. Initial investigations determine whether the child is an under-virilised male or a virilised female from which differential diagnoses are established and further investigations planned [4]. As the differential diagnosis and relative investigations will depend on the genetic sex, an urgent karyotype has to be carried out within earliest possible time [4]. A more conservative approach has been followed considering the economic condition of the patient although extensive investigations are advised to reach the definitive diagnosis by different scientific papers [2,22,24–26]. Minimum diagnostic tests with high sensitivity and specificity were performed in this case to attain the correct diagnosis. It is noteworthy to mention that the sample for karyotyping was sent to a laboratory situated in New Delhi, India since in Bangladesh, during that period, the service for karyotyping was not available. However, currently the investigation is performed in the only medical university of Bangladesh. The treatment constituted both the available ones i.e. hormonal therapy followed by surgical measures although the timing of the following steps such as gender assignment and surgery is still a topic of debate [21]. Gender assignment should be aimed at offering the best prospect for healthy puberty and sex life [1,6,22,24] and should be attempted after comprehensive clinical, genetic and biochemical investigation actively involving the parents with detailed discussion and explanation [6,16,22,25]. The consensus for the management of ambiguous genitalia suggests that if it is decided to assign male gender to the infants, 10–25 mg of testosterone enanthate or cypionate is given intramuscularly for once a month for consecutive 3 months to ensure that the penis responds to androgens or to improve the size of penis prior to surgery [27]. However, patients with 5-alpha reductase deficiency should receive dihydrotestosterone replacement. In case of unavailability, higher dose of testosterone can be given as a replacement to overcome the enzyme block [27].

The team followed the standardized protocol for the testosterone therapy in the current case. For patients to be raised as males, müllerian structures can be removed with the goal of preserving the vas deferens. Various types of hypospadias repair could also be done and gonads could be replaced with prostheses. If needed, pre-penile scrotum could be reconstructed. [25,27]. Through different stages of surgical interventions the multidisciplinary team followed the standard procedures according to specific indications as mentioned earlier. Although the patient was lost to follow up in the discussed case, there is a desperate need for prospective long term follow up of such patients in order to explore their sexuality, identity and psychosocial outcomes [28]. As suggested by different scientific works, the multidisciplinary team has been composed of experts from different specialties that included paediatric surgeon, paediatric urologist and endocrinologist [1,4,6,8,21,28]. It helped not only in the diagnosis of the underlying condition, but also in reconstruction of the external genitalia, timely gonadectomy, gonadal biopsies or orchiopexy. As a result of our findings, we would advocate diagnostic laparoscopy as the investigation of choice for the identification of Müllerian structures in children with complex DSD, as well as being a useful treatment modality. However, a larger prospective study is required to compare different modalities such as laparoscopy, MRI and ultrasonography for the evaluation of Müllerian structures in children with complex DSD [29].

3.2. DSD as a taboo in bangladeshi culture and its relationship with management process

In Bangladesh, ambiguous genitalia is an under discussed agendum for which researches have been conducted to a minimal extent. The reason behind the paucity of discussion is the conservative attitude of the native Bangladeshi people regarding sex and reproductive health rights [9]. Sex itself is a taboo in the country [30], let alone the disorders related to sex. Hence, DSD, as an important issue, has been neglected for years [10]. A most important element in the care of these patients is the psychological support with utmost sensitivity required by families at first hand and then the patient and thus, cultural hindrances regarding DSD also hamper the management process. The psychologists and counsellors can be the key role players in this context [16,26]. The proper care of these complex patients requires that the physician be a scientist as well as a clinician and a skilled technician [25].

4. Conclusion

It is crucial to have a decent knowledge about DSD to analyse the attitude of parents and to provide suitable counselling. Management of DSD needs consistent classification, adequate cytogenetic facilities and customized approach along with integrated team management. At present, the patients with ambiguous genitalia stand a far better chance of receiving a rapid diagnosis, appropriate replacement therapy, and functional surgical reconstruction than before. In-utero diagnoses and treatment at the pre-implantation stage may prove to be very important for the care of some of these patients, who may be potential candidates for gene replacement therapy. Extensive research and social awareness campaigns are also a timely need to document evidences and to free the society from the obscure darkness of the taboo of previously called 'intersex disorders'.

Conflicts of interest

There is no potential conflict of interest declared by the authors.

Funding

No source of funding.

Ethical approval

Since this was just was a case report of a rare disease, ethical approval was exempted by our institution.

Consent

The data collectors took consent from the patient's parents since the patient was not eligible to provide consent. An additional consent was taken to capture a photograph of the case.

The patient's parents gave consent for the publication of this case report.

Author contribution

MAKC contributed in all components of this study ranging from data collection to writing the manuscript. He coordinated the project. RA was actively engaged with the management process of the case and manuscript writing. AS was involved in active management of the case. All the authors reviewed the article.

Guarantor

Mohiuddin Ahsanul Kabir Chowdhury

References

- [1] P.A. Lee, C.P. Houk, S.F. Ahmed, et al., Consensus statement on management of intersex disorders, *Int. Consens. Paediatr.* 118 (2) (2006) e488–e500.
- [2] *Ambiguous Genitalia Symptoms & Causes, in Conditions + Treatments*, Boston Children's Hospital, Boston, MA, 2016.
- [3] J.M. Khalid, J.M. Oerton, C. Dezateux, et al., Incidence and clinical features of congenital adrenal hyperplasia, *Arch. Dis. Child.* 97 (2) (2012) 101–106.
- [4] A. Ogilvy-Stuart, C. Brain, Early assessment of ambiguous genitalia, *Arch. Dis. Child.* 89 (5) (2004) 401–407.
- [5] I.A. Hughes, C. Houk, S.F. Ahmed, et al., Consensus statement on management of intersex disorders, *Arch. Dis. Child.* (2006).
- [6] *Guidance on the Initial Evaluation of an Infant or an Adolescent with a Suspected Disorder of Sex Development*, Society for Endocrinology, 2011.
- [7] C.J. Migeon, A.B. Wisniewski, J.P. Gearhart, H.F. Meyer-Bahlburg, J.A. Rock, T.R. Brown, et al., Ambiguous genitalia with perineoscrotal hypospadias in 46, XY individuals: long-term medical, surgical, and psychosexual outcome, *Paediatrics* 110 (3) (2002).
- [8] H.G. Beh, M. Diamond, An emerging ethical and medical dilemma: should physicians perform sex assignment surgery on infants with ambiguous genitalia? *Michigan J. Gen. Law* 7 (1) (2001).
- [9] S. Akhter, *Knowledge Attitudes and Practices on Reproductive Health and Rights of Urban and Rural Women in Bangladesh*, 2007.
- [10] T.K. Chowdhury, M. Kabir, M.Z. Chowdhury, J.M. Hutson, T. Banu, The challenges in diagnosis and gender assignment in disorders of sex development presenting to a paediatric surgical unit in a developing country: the role of laparoscopy and simple tests for gender identity, *J. Paediatr. Urol.* 10 (6) (2014) 1255–1260.
- [11] G.B. De Paula, et al., <it>408 cases of genital ambiguity followed by single multidisciplinary team during 23 years: etiologic diagnosis and sex of rearing, *Int. J. Endocrinol.* (2016) 2016.
- [12] M. Diamond, Paediatric management of ambiguous and traumatized genitalia, *J. Urol.* 162 (3) (1999) 1021–1028.
- [13] N. Diamond-Smith, M. Sudhinaraset, D. Montagu, Clinical and perceived quality of care for maternal, neonatal and antenatal care in Kenya and Namibia: the service provision assessment, *Reprod. health* 13 (1) (2016) 92.
- [14] V.A. Kumari, A. Vani, Ambiguous genitalia: a clinical and chromosomal study, *Int. J. Res. Med. Sci.* 3 (12) (2017) 3743–3748.
- [15] W.G. Reiner, Assignment of sex in neonates with ambiguous-genitalia, *Curr. Opin. Paediatr.* 11 (4) (1999) 363–366.
- [16] U. Thyen, K. Lanz, P.M. Holterhus, O. Hiort, Epidemiology and initial management of ambiguous genitalia at birth in Germany, *Horm. Res.* 66 (4) (2006) 195–203.
- [17] C.A. Warren, Gender reassignment surgery in the 18th century: a case study, *Sexualities* 17 (7) (2014) 872–884.
- [18] A.D. Dreger, Ambiguous sex—or ambivalent medicine?: ethical issues in the treatment of intersexuality, *Hastings Cent. Rep.* 28 (3) (1998) 24–35.

- [19] H.G. Beh, M. Diamond, An Emerging Ethical and Medical Dilemma: Should Physicians Perform Sex Assignment Surgery on Infants with Ambiguous Genitalia? 2000.
- [20] R.A. Agha, et al., The SCARE statement: consensus-based surgical case report guidelines *Int. J. Surg.* 34 (2016) 180–186.
- [21] O. Ercan, S. Kutlug, O. Uysal, et al., Gender identity and gender role in DSD patients raised as females: a preliminary outcome study, *Front. Endocrinol. (Lausanne)* 4 (86) (2013).
- [22] S.F. Ahmed, M. Rodie, Investigation and initial management of ambiguous genitalia, *Best Practice Res. Clin. Endocrinol. Metab.* 24 (2) (2010) 197–218.
- [23] L. Rangescroft, Surgical management of ambiguous genitalia, *Arch. Dis. Child.* 88 (9) (2003) 799–801.
- [24] Guidelines on Paediatric Urology, European Association of Urology, 2013.
- [25] D.D. Federman, P.K. Donahoe, Ambiguous genitalia—etiology, diagnosis, and therapy, *Adv. Endocrinol. Metab.* 6 (1995) 91–116.
- [26] L.D. Mastrandrea, C.H. Albini, R.J. Wynn, et al., Disorders of sex development: management of gender assignment in a preterm infant with intrauterine growth restriction, *Case Rep. Med.* (2012).
- [27] K.W. Ashcraft, G.W. Holcomb, J.P. Murphy, D.J. Ostlie, *Ashcraft's Paediatric Surgery*, Saunders/Elsevier, 2010.
- [28] K.P. Kulkarni, I. Panigrahi, R. Das, S. Kaur, R.K. Marwaha, Paediatric disorders of sex development, *Indian J. Pediatr.* 76 (9) (2009) 956–958.
- [29] M. Steven, S. O'Toole, J.P.H. Lam, G.A. MacKinlay, S. Cascio, Laparoscopy versus ultrasonography for the evaluation of Mullerian structures in children with complex disorders of sex development, *Pediatr. Surg Int.* 28 (12) (2012) 1161–1164.
- [30] P. Nahar, M. Van Reeuwijk, R. Reis, Contextualising sexual harassment of adolescent girls in Bangladesh, *Reprod. Health Matters* 21 (41) (2013) 78–86.

Open Access

This article is published Open Access at [sciencedirect.com](https://www.sciencedirect.com). It is distributed under the [IJSCR Supplemental terms and conditions](#), which permits unrestricted non commercial use, distribution, and reproduction in any medium, provided the original authors and source are credited.