

Complete Genome Sequence of a Wild-Type Measles Virus Isolated during the Spring 2013 Epidemic in Germany

Konstantin M. J. Sparrer,^{a,b} Stefan Krebs,^{b,c} Gundula Jäger,^a Sabine Santibanez,^d Annette Mankertz,^d Helmut Blum,^{b,c} Karl-Klaus Conzelmann^{a,b}

Max von Pettenkofer-Institute,^a Gene Center,^b and Laboratory for Functional Genome Analysis,^c Ludwig-Maximilians-University, Munich, Germany; Robert-Koch Institute, National Reference Center Measles, Mumps, Rubella, Berlin, Germany^d

Measles virus induces an acute disease with rash and fever. Despite ongoing vaccination and elimination campaigns, the measles virus still sustains long-lasting transmission chains in Europe. Here we report the complete genome sequence of a wild-type measles virus isolated from a patient in Munich (MVi/Muenchen.DEU/19.13[D8]) during a German measles outbreak in 2013.

Received 10 February 2014 Accepted 3 April 2014 Published 17 April 2014

Citation Sparrer KMJ, Krebs S, Jäger G, Santibanez S, Mankertz A, Blum H, Conzelmann K-K. 2014. Complete genome sequence of a wild-type measles virus isolated during the spring 2013 epidemic in Germany. *Genome Announc.* 2(2):e00157-14. doi:10.1128/genomeA.00157-14.

Copyright © 2014 Sparrer et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/3.0/).

Address correspondence to Karl-Klaus Conzelmann, conzelma@lmb.uni-muenchen.de.

Infections caused by measles virus (MV), the prototype of the *Morbillivirus* genus of the *Paramyxoviridae* family, remain a serious threat to human health worldwide. In 2011, approximately 160,000 deaths were caused by measles, although MV elimination is sought globally (1–3). Infection with MV is typically accompanied by transient immune suppression abetting secondary infections. MV itself can cause encephalitis and rare severe sequelae, including measles inclusion-body encephalitis (MIBE) and subacute sclerosing panencephalitis (SSPE), resulting inevitably in the death of the patient (4).

MV wild-type (wt) infection chains have been recently observed all over Europe (5, 6). In 2013, 704 measles cases were reported in Upper Bavaria and 306 in Munich. MV was isolated from a throat swab from a patient with typical clinical symptoms using Vero-hSLAM cells (7). The virus was cultivated for 48 h until syncytium formation was visible. Whole-cell RNA was extracted from infected Vero-hSLAM cells by using the Qiagen RNeasy minikit (Quiagen), and 100 ng was used to prepare a strand-specific RNA-Seq library kit (NuGEN Encore complete RNA-Seq library system; NuGEN, Inc.) following the manufacturer's instruction. The resulting library was sequenced on an Illumina MiSeq instrument in paired-end mode with a read length of 250 nucleotides (nt).

Sequence reads were aligned at first to the sequence of the MV vaccine strain Schwarz (GenBank accession number AF266291.1) and for further refinement to that of a closely related MV isolate (MVi/Texas.USA/4.07, genotype D8; GenBank accession number JN635407.1). Only reads mapping to these MV strains were used for genome assembly using the *de novo* assembler Velvet (8). Coverage for an average nucleotide in the N gene was approximately 400 reads, and for the L gene, 50 reads. Nucleotides with a low coverage within the L gene and H-L gene border were sequenced using conventional reverse transcription and sequencing using specific primers (GATC, Constance, Germany). The nucleotide sequences of the 3' leader and 5' trailer regions were determined by 5' rapid amplification of cDNA ends (5'-RACE) using specific

primers binding in the leader and trailer sequences and terminal deoxynucleotidyl transferase (TdT) for addition of a poly(A) tail. PCR was carried out using a dT(18) primer with an EcoRI site, and the resulting DNA fragments were cloned into pCR3 vectors for sequencing of multiple clones. The presented full genome therefore represents an assembly of sequences obtained by the three techniques described above.

Genotyping according to the WHO protocol (9) using the 450 nucleotides of the 3'-end of the N gene classified the present wt strain as a D8 genotype, subtype Frankfurt-Main (MVs/Frankfurt Main.DEU/17.11-variant). The genome of MVi/Muenchen.DEU/19.13 follows the canonical 3'-N-P/V/C-M-F-H-L-5' gene arrangement and comprises 15,984 nucleotides and therefore obeys the rule of six (10, 11). It shares 99% identity with the complete genome of the MVi/Texas.USA/4.07 wt isolate and 96.55% identity with that of the MV strain Schwarz. As is typically seen for wt versus vaccine strain comparisons, the C protein sequence contains a fully conserved nuclear localization sequence (NLS) (12), while the H protein lacks the typical N481Y or S546G mutation required for CD46 binding (13, 14). MVi/Muenchen.DEU/19.13 therefore represents a typical wt MV circulating in Germany.

Nucleotide sequence accession number. The complete genome sequence of the MVi/Muenchen.DEU/19.13[D8] isolate is available at GenBank under the accession number [KJ410048](https://www.ncbi.nlm.nih.gov/nuccore/KJ410048).

ACKNOWLEDGMENTS

We acknowledge Nadin Zapf for technical assistance, Yusuke Yanagi for Vero-hSLAM cells, and Klaus Förstemann for providing primers and helpful suggestions.

This research was supported by DFG (GraKo 1202 and SFB 870).

REFERENCES

1. World Health Organization. 2013. Measles reports. World Health Organization, Geneva, Switzerland. http://www.who.int/mediacentre/news/notes/2013/measles_20130117/en/.
2. World Health Organization. 2013. Measles fact sheet no. 286. World

- Health Organization, Geneva, Switzerland. <http://www.who.int/mediacentre/factsheets/fs286/en/>.
3. World Health Organization. 2013. Measles surveillance data. World Health Organization, Geneva, Switzerland. http://www.who.int/immunization/monitoring_surveillance/burden/vpd/surveillance_type/active/measles_monthlydata/en/.
 4. Griffin DE. 2010. Measles virus-induced suppression of immune responses. *Immunol. Rev.* 236:176–189. <http://dx.doi.org/10.1111/j.1600-065X.2010.00925.x>.
 5. Mankertz A, Mihneva Z, Gold H, Baumgarte S, Baillot A, Helble R, Roggendorf H, Bosevska G, Nedeljkovic J, Makowka A, Hutse V, Holzmann H, Aberle SW, Cordey S, Necula G, Mentis A, Korukluoglu G, Carr M, Brown KE, Hubschen JM, Muller CP, Mulders MN, Santibanez S. 2011. Spread of measles virus D4-Hamburg, Europe, 2008–2011. *Emerg. Infect. Dis.* 17:1396–1401. <http://dx.doi.org/10.3201/eid1708.101994>.
 6. Mankertz A, Mulders MN, Shulga S, Kremer JR, Brown KE, Santibanez S, Muller CP, Tikhonova N, Lipskaya G, Jankovic D, Khetsuriani N, Martin R, Gavrillin E. 2011. Molecular genotyping and epidemiology of measles virus transmission in the World Health Organization European Region, 2007–2009. *J. Infect. Dis.* 204(Suppl 1):S335–S342. <http://dx.doi.org/10.1093/infdis/jir101>.
 7. Ono N, Tatsuo H, Hidaka Y, Aoki T, Minagawa H, Yanagi Y. 2001. Measles viruses on throat swabs from measles patients use signaling lymphocytic activation molecule (CDw150) but not CD46 as a cellular receptor. *J. Virol.* 75:4399–4401. <http://dx.doi.org/10.1128/JVI.75.9.4399-4401.2001>.
 8. Zerbino DR, Birney E. 2008. Velvet: algorithms for de novo short read assembly using de Bruijn graphs. *Genome Res.* 18:821–829. <http://dx.doi.org/10.1101/gr.074492.107>.
 9. World Health Organization. 2001. Nomenclature for describing the genetic characteristics of wild-type measles virus. *Wkly. Epidemiol. Rec.* 76:249–251.
 10. Kolakofsky D, Pelet T, Garcin D, Hausmann S, Curran J, Roux L. 1998. Paramyxovirus RNA synthesis and the requirement for hexamer genome length: the rule of six revisited. *J. Virol.* 72:891–899.
 11. Calain P, Roux L. 1993. The rule of six, a basic feature for efficient replication of Sendai virus defective interfering RNA. *J. Virol.* 67:4822–4830.
 12. Sparrer KM, Pfaller CK, Conzelmann KK. 2012. Measles virus C protein interferes with beta interferon transcription in the nucleus. *J. Virol.* 86:796–805. <http://dx.doi.org/10.1128/JVI.05899-11>.
 13. Lecouturier V, Fayolle J, Caballero M, Carabaña J, Celma ML, Fernandez-Muñoz R, Wild TF, Buckland R. 1996. Identification of two amino acids in the hemagglutinin glycoprotein of measles virus (MV) that govern hemadsorption, HeLa cell fusion, and CD46 downregulation: phenotypic markers that differentiate vaccine and wild-type MV strains. *J. Virol.* 70:4200–4204.
 14. Shibahara K, Hotta H, Katayama Y, Homma M. 1994. Increased binding activity of measles virus to monkey red blood cells after long-term passage in Vero cell cultures. *J. Gen. Virol.* 75(Pt 12):3511–3516. <http://dx.doi.org/10.1099/0022-1317-75-12-3511>.