

CASE REPORT

Open Access



Diagnostic and therapeutic challenges of treating opportunistic fungal cellulitis: a case series

Jed Paul¹, Mary M. Czech^{1,2*} , Ramya Balijepally¹ and Janice Wes Brown^{1,2,3}

Abstract

Background: Cellulitis is an infection most commonly caused by bacteria and successfully treated with antibiotics. However, certain patient populations, especially the immunocompromised, are at risk for fungal cellulitis, which can be misidentified as bacterial cellulitis and contribute to significant morbidity and mortality.

Case presentations: We describe three cases of opportunistic fungal cellulitis in immunosuppressed patients that were initially mistaken for bacterial infections refractory to antibiotic therapy. However, atypical features of cellulitis ultimately prompted further diagnostics to identify fungal cellulitis and allow initiation of appropriate antifungals. We discuss: (1) a 52-year-old male immunosuppressed hematopoietic cell transplant recipient with *Fusarium solani* cellulitis on his right lower extremity that was treated with amphotericin B and voriconazole with full resolution of the cellulitis; (2) a 70-year-old male lung transplant recipient with *Fusarium solani* cellulitis on his left lower extremity that ultimately progressed despite antifungals; and (3) a 68-year-old male with a history of kidney transplantation with suspected *Purpureocillium lilacinum* cellulitis on his left lower extremity ultimately treated with posaconazole with resolution of the skin lesions.

Conclusions: *Fusarium solani* and *Purpureocillium lilacinum* are important pathogens causing opportunistic fungal cellulitis. These cases remind providers to be vigilant for fungal cellulitis when skin and soft tissue infection does not adequately respond to antibiotics and atypical features of cellulitis are present.

Keywords: Fungal cellulitis, *Fusarium solani*, *Purpureocillium lilacinum*, Immunocompromised, Opportunistic infection

Background

Cellulitis is a common infection, with incidence estimated at around 200 cases per 100,000 person-years in the USA [1]. The vast majority of cases are caused by bacterial etiologies such as *Streptococcus* and *Staphylococcus*, and less commonly gram-negative bacteria, with over 95% of cases responding to antimicrobial monotherapy. However, in immunocompromised patients, and rarely in immunocompetent hosts, cellulitis may be due to fungi.

Because of the relative rarity of fungal skin and soft tissue infection, misidentification of these cases often leads to inappropriate treatment and significant morbidity or mortality [1, 2]. In this paper, we highlight three cases of fungal cellulitis that were initially mistaken for bacterial cellulitis in part due to modest response to antibacterial therapy, and discuss the importance of fungal consideration when atypical features of cellulitis are identified.

Case presentation

Case 1

A 52-year-old male with insulin-dependent type 2 diabetes (hemoglobin A1C 9.2%) and a history of B cell acute lymphoblastic leukemia, who received an allogeneic

*Correspondence: mmc16@stanford.edu

¹ Department of Medicine, Stanford University School of Medicine, 300 Pasteur Drive, Lane Building, L134 MC:5107, Stanford, CA 94305-5107, USA
Full list of author information is available at the end of the article



hematopoietic cell transplant (HCT) four years prior to presentation, presented with right lower extremity pain and erythema. HCT was complicated by chronic graft-versus-host disease of the skin and oral mucosa for which the patient was maintained on prednisone 10 mg daily and ruxolitinib 10 mg twice daily. Five days prior to presentation, he noticed a small, painful papule on the dorsal hallux of his right foot which became ulcerative with painful erythema that spread circumferentially over his calf. He was prescribed a course of clindamycin with transient improvement. On admission, in addition to the erythema and right hallux ulcer, he was noted to have an onychomycotic right first toe (Fig. 1). He was started on broad spectrum antimicrobial coverage with intravenous (IV) vancomycin 1.25 g twice daily, meropenem 1 g three times daily, and isavuconazonium 4 mg/kg daily with improvement in his dermatitis over the next several days. A wound culture from the ulcer grew *Fusarium solani*, with pathology showing dermal edema with interstitial neutrophilic and lymphohistiocytic infiltrates. Polymerase chain reaction testing and fungal culture of the onychomycotic toenail additionally showed *Fusarium solani*, suggesting that cellulitis resulted from extension of the infected toenail. For *Fusarium solani* cellulitis, isavuconazonium was discontinued and both liposomal amphotericin B 5 mg/kg daily and intravenous voriconazole 4 mg/kg twice daily after a 6 mg/kg loading dose were started on hospital day 8. He was discharged home on hospital day 16 with residual but improving erythema of the leg on an additional four weeks of liposomal

amphotericin B and voriconazole plus topical efinaconazole for his affected nail. He continued to improve over the ensuing months. Ultimately, he had only minimal scarring from his ulcer ten months after discharge alongside a persistently onychomycotic but improved toenail.

Case 2

A 70-year-old male with a history of scleroderma-induced interstitial pneumonia, who underwent bilateral lung transplantation, developed left lower extremity cellulitis two months following his transplant. His post-transplant course was complicated by prolonged hospitalization for chronic mechanical respiratory failure unrelated to allograft function and multiple cardiac arrests. He received induction immunosuppression with basiliximab, and at the time of cellulitis he was maintained on azathioprine 50 mg daily and prednisone 5 mg twice daily. He received 5 days of IV cefazolin with improvement, but the erythema returned after antibiotics were stopped. Four days later, he was started on IV vancomycin 750 mg daily and cefepime 1 g daily (renal dosing). However, his erythema persisted, and he developed a black, ulcerative lesion on the dorsum of his left foot, in the setting of multiple onychomycotic toenails (Fig. 2). After nine days on vancomycin and cefepime, skin biopsy cultures grew *Fusarium solani*. Corresponding pathology of the biopsy specimen showed deep dermal fungal elements. He was started on IV amphotericin B 5 mg/kg daily and IV voriconazole 4 mg/kg twice daily after two loading doses. Blood cultures, respiratory cultures, chest



Fig. 1 Patient 1's leg and foot on presentation (A, B), partial response after antibiotic therapy (C), and three months post-discharge after antifungal therapy (D)



Fig. 2 Patient 2's left foot when lesion was first noticed on post-transplant day 86 (A) and progression despite antibiotic and antifungal therapy on post-transplant day 103 (B) with spread to thigh (C)

computed tomography (CT) scan, and ophthalmologic exam were unrevealing for disseminated fungal infection. Although samples from the onychomycotic toenails were not evaluated, it is suspected that onychomycosis was due to *Fusarium*, and skin breakdown surrounding the nail allowed for fungal inoculation. Despite antifungal therapy, the patient developed progressive erythematous nodules extending proximally to his upper thigh, consistent with vascular spread of infection. He subsequently developed septic shock refractory to broad spectrum antibiotics and ongoing antifungals. He was transitioned to comfort care and expired on post-transplant day 113.

Case 3

A 68-year-old male with a history of Alport Syndrome, who received a living related donor kidney transplant 30 years prior to presentation, presented with a 9-day history of global left lower extremity swelling with distal erythema and pain. His immunosuppressive regimen was azathioprine 100 mg daily and tacrolimus 2 mg daily. He had a recent folliculitis in his left groin 3 weeks prior to presentation, treated with ten days of oral doxycycline and fourteen days of oral azithromycin. Two weeks following the folliculitis, he was admitted at an outside hospital with new fever, progressive left lower extremity erythema, swelling and pain consistent with cellulitis. Within 48 h of starting IV vancomycin and clindamycin for presumed Methicillin-resistant *Staphylococcus aureus* (MRSA) cellulitis, fevers resolved and erythema and swelling improved. However, after discharge, his cellulitis again worsened, and he developed exquisite leg

tenderness with the development of multiple 2–4 mm tender purplish-black papules, some with punctate central eschar. On presentation to our hospital, these skin changes were noted to occur in the setting of longstanding first toe onychomycosis (Fig. 3). He was started on IV cefepime 2 g twice daily (renal dosing) and daptomycin 800 mg daily with marked improvement in both his erythema and pain over the next few days. However, the violaceous papules showed no change, prompting a punch biopsy of one papule. Gram stain of the biopsy specimen was unrevealing for significant polymorphonuclear cells (PMNs) or organisms. Tissue culture was negative for both bacterial and fungal growth. However, given concern for biopsy sampling error, he was started on empiric oral itraconazole 200 mg twice daily for presumed fungal cellulitis. Subsequently, fungal polymerase chain reaction of the onychomycotic toenail identified *Purpureocillium lilacinum*, which was also believed to be the causative pathogen of the purplish-black cutaneous papules. There was no evidence of deep tissue infection on magnetic resonance imaging (MRI) of his leg, and the CT scan of his chest did not suggest evidence of disseminated fungal infection. His pain and swelling improved by hospital day four, and he was discharged home on hospital day seven with fourteen additional days of doxycycline 100 mg twice daily and levofloxacin 750 mg every other day (renal dosing), as well as a prolonged course of itraconazole 200 mg twice daily. On follow-up 15 days after discharge, his erythema, swelling, and pain had resolved. Following 12 weeks of itraconazole, the black papules had flattened and faded, at which time antifungal therapy



Fig. 3 Patient 3's leg on presentation (A), after partial response to IV antibiotics (B), and one month after discharge on systemic antifungal therapy (C)

was stopped. However, one month after discontinuation of itraconazole the papular lesions reappeared with new lesions in the same distribution of his left leg. He was retreated with posaconazole, and three months into therapy, the black papules resolved.

Discussion and conclusions

These cases demonstrate episodes of fungal cellulitis initially mistaken for pure bacterial cellulitis in immunosuppressed patients. In all cases, summarized in Table 1, patients were initially treated with antibacterial agents with at least partial response to therapy. However, when the episodes of cellulitis seemed refractory to appropriate antibiotics, and displayed atypical features of cellulitis, opportunistic infections were appropriately considered. Diagnosis ultimately required clinical suspicion with tissue biopsy, cultures, and pathology to diagnose fungal cellulitis. With an accurate diagnosis, patients were able to be started on targeted long term antifungal therapy.

Fungal cellulitis has been increasingly described in immunocompromised individuals [3–6]. Impaired functioning of cell-mediated immunity pathways, particularly those involving T cells, predispose immunocompromised patients to infection with opportunistic and environmental fungi [7, 8]. As described in these cases, *Fusarium solani* and *Paecilomyces lilacinus*, renamed to *Purpureocillium lilacinum*, are ubiquitous fungi found in soil and decaying vegetation [9, 10]. As with most fungi, infection commonly occurs via inhalation with subsequent dissemination to skin and other sites, or as direct inoculation,

often at a site of skin or mucosal barrier breakdown [5, 11]. Cutaneous infection with these fungi manifests in many forms, including painless papules, tender nodules, and necrotic ulcers, occasionally requiring surgical debridement in addition to systemic antifungal therapies [3, 5, 12, 13].

Diagnosis of fungal cellulitis poses a significant challenge to clinicians for multiple reasons. As seen in our cases, fungal infections may mimic or possibly co-occur with bacterial infections, and a partial response to antibiotic therapy can give a false impression of adequate antimicrobial coverage. This prolongs the treatment course and puts severely immunocompromised patients at risk for both more severe local disease as well as disseminated fungal infection. Furthermore, the broad spectrum of manifestations of skin and soft tissue infections, especially among the immunocompromised, leads to misidentification of fungal lesions. Finally, even if a biopsy is obtained, advanced testing modalities may be required, and results must be interpreted with caution and consideration for the clinical context. For example, patient 3's skin biopsy was largely unrevealing. However, the absence of PMNs in gram stain was highly suggestive of sampling error given clear evidence of inflammation on exam and the absence of neutropenia in the patient's peripheral blood [14]. Therefore, clinicians were prompted to evaluate the patient's onychomycotic toenail, which was positive for *P. lilacinum*. Though it is possible different organisms caused the skin and toenail infections, we suspect the scattered necrotic papules

Table 1 Summary of case reports

	Underlying condition	Immuno-suppression	Atypical features of cellulitis	Fungal Pathogen	Antifungal regimen	Outcome	Relevant citations for other related case reports
Patient 1	B cell ALL s/p allo-HSCT c/b CGVHD	Prednisone 10 mg daily, ruxolitinib 10 mg twice daily	Ulcerative lesion on dor- sum of foot unresponsive to antibiotics	<i>Fusarium solani</i>	5 weeks of IV amphotericin B 5 mg/kg daily and voriconazole 4 mg/kg twice daily	Near baseline (residual scarring)	[4, 5, 18, 19, 22]
Patient 2	Systemic scleroderma c/b interstitial pneumonia s/p bilateral lung transplant	Azathioprine 50 mg daily, prednisone 5 mg twice daily	Ulcerative lesion on dor- sum of foot unresponsive to antibiotics	<i>Fusarium solani</i>	Amphotericin B 5 mg/kg daily and voriconazole 4 mg/kg twice daily	Deceased (refractory septic shock)	[4, 5, 18, 19, 23]
Patient 3	Alport Syndrome c/b ESRD s/p kidney transplant	Azathioprine 100 mg daily, tacrolimus 2 mg daily	Numerous violaceous papules on calf and dor- sum of foot unresponsive to antibiotics	<i>Purpureocillium lilacinum</i>	8 weeks of itraconazole 200 mg twice daily	Recurrence after discontinuation of itraconazole, responded to posaconazole	[3, 6, 9, 12, 13, 18, 19]

ALL acute lymphoblastic leukemia, s/p status post, allo-HCT allogeneic hematopoietic cell transplantation, c/b complicated by, cGVHD chronic graft versus host disease, ESRD end-stage renal disease

resulted from *P. lilacinum* cutaneous infection tracking from the toenail.

To aid in diagnosis, differences in the presentation of bacterial and fungal cellulitis may help clinicians to more readily identify atypical and fungal cellulitis. Typical bacterial cellulitis commonly presents with erythema, swelling, and pain, which rapidly improves with antimicrobial therapy. For typical bacterial cellulitis, open wounds often serve as entry portals for bacteria to cause infection, but less commonly do open wounds develop as a result of infection (some exceptions being MRSA, *Streptococcus pyogenes*, and polymicrobial necrotizing fasciitis). For patients 1 and 2, they developed open wounds on the dorsum of the foot. As discussed, this is unusual for typical bacterial cellulitis, and furthermore an unusual location for a traumatic wound. This unusual feature prompted biopsy consistent with fungal infection, which commonly causes ulceration due to invasion of the vascular supply. Additionally, for patient 3, scattered black papules are an unusual feature of bacterial cellulitis, which appropriately prompted further investigation.

Ultimately, tissue is often required for correct identification of fungal pathogens by either standard microbiological or molecular techniques. Identification of fungal pathogens to the species level is essential for selection of appropriate therapy. Although there are no established breakpoints for antifungal susceptibility testing against filamentous fungi, there is often a body of literature to help guide the choice of therapeutics. For example, *E. solani* infections typically respond to combination therapy with amphotericin B and voriconazole, though successful therapy with posaconazole or isavuconazole have also been described [5]. *P. lilacinum* can have some degree of intrinsic *in vitro* resistance to several commonly used mold active agents, such as amphotericin products, itraconazole, and echinocandins, but favorable minimum inhibitory concentrations for voriconazole and posaconazole [15–17]. Given that clinical outcomes have not been well defined for minimal inhibitory or fungicidal concentrations to establish breakpoints, clinicians should closely monitor patients being treated for fungal cellulitis to ensure appropriate clinical response.

Onychomycosis is a common feature in bacterial and fungal cellulitis and has widely been documented as a risk factor for development of cellulitis [18–20]. The exact mechanism behind this association is unclear. However, other reports describe onychomycosis with either paronychia or interdigital intertrigo serving as a suspected cutaneous portal of entry for invasive fungal infection [21]. Given the proximity of all our patients' cellulitis to preexisting onychomycotic infections, we similarly suspect onychomycosis with surrounding skin breakdown led to extension of infection resulting in cellulitis.

These cases highlight the importance of consideration for fungal pathogens when cellulitis displays atypical features in immunocompromised hosts. Prolonged treatment targeted at only bacterial pathogens can potentially lead to significant morbidity or mortality when fungal cellulitis is present. Clinicians should be mindful of these atypical features of cellulitis and investigate possible fungal etiologies in warranted cases.

Abbreviations

HCT: Allogeneic hematopoietic cell transplant; IV: Intravenous; CT: Computed tomography; MRSA: Methicillin-resistant *Staphylococcus aureus*; ESR: Erythrocyte sedimentation rate; CRP: C-reactive protein; PMNs: Polymorphonuclear cells; MRI: Magnetic resonance imaging.

Acknowledgements

Not applicable.

Authors contributions

JP, MMC, RB, and JB all made substantial contributions to the manuscript, approve the submitted version of the manuscript, and agree to be personally accountable for the content of the manuscript. All authors read and approved the final manuscript.

Funding:

No funding was used to prepare this manuscript.

Availability of data and materials

All relevant data is included in the manuscript. A separate data repository is not applicable.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Written consent from one patient was obtained to disclose clinical details and identifying images to participate in the case report. Written consent from next of kin was obtained to disclose clinical details and identifying imaging to participate in the case report for the other two patients, who were deceased at the time of manuscript preparation.

Competing interests

None of the authors have financial or non-financial competing interests.

Author details

¹Department of Medicine, Stanford University School of Medicine, 300 Pasteur Drive, Lane Building, L134 MC:5107, Stanford, CA 94305-5107, USA. ²Division of Infectious Diseases, Stanford University School of Medicine, Stanford, CA, USA. ³Division of Blood and Marrow Transplantation, Stanford University School of Medicine, Stanford, CA, USA.

Received: 19 April 2021 Accepted: 8 April 2022

Published online: 05 May 2022

References

- McNamara DR, Tleyjeh IM, Berbari EF, Lahr BD, Martinez JW, Mirzoyev SA, et al. Incidence of lower-extremity cellulitis: a population-based study in Olmsted county, Minnesota. *Mayo Clin Proc.* 2007;82:817–21.
- Jeng A, Beheshti M, Li J, Nathan R. The role of beta-hemolytic streptococci in causing diffuse, nonculturable cellulitis: a prospective investigation. *Medicine.* 2010;89:217–26.

3. Sotello D, Cappel M, Huff T, Meza D, Alvarez S, Libertin CR. Cutaneous fungal infection in an immunocompromised host. *JMM Case Rep*. 2017;4:e005101.
4. Rekha A, Kindo AJ, Ravi A. *Fusarium solani* in the post-transplant patient: an unusual fungus. *Int J Low Extrem Wounds*. 2008;7:38–40.
5. Muhammed M, Anagnostou T, Desalermos A, Kourkoumpetis TK, Carneiro HA, Glavis-Bloom J, et al. *Fusarium* infection: report of 26 cases and review of 97 cases from the literature. *Medicine*. 2013;92:305–16.
6. Salazar-González MA, Violante-Cumpa JR, Alfaro-Rivera CG, Villanueva-Lozano H, Treviño-Rangel R, de González J. *Purpureocillium lilacinum* as unusual cause of pulmonary infection in immunocompromised hosts. *J Infect Dev Ctries*. 2020;14:415–9.
7. Kaiko GE, Horvat JC, Beagley KW, Hansbro PM. Immunological decision-making: how does the immune system decide to mount a helper T-cell response? *Immunology*. 2008;123:326–38.
8. Bacher P, Kniemeyer O, Schönbrunn A, Sawitzki B, Assenmacher M, Rietschel E, et al. Antigen-specific expansion of human regulatory T cells as a major tolerance mechanism against mucosal fungi. *Mucosal Immunol*. 2014;7:916–28.
9. Luangsa-Ard J, Houbraken J, van Doorn T, Hong S-B, Borman AM, Hywel-Jones NL, et al. *Purpureocillium*, a new genus for the medically important *Paecilomyces lilacinus*. *FEMS Microbiol Lett*. 2011;321:141–9.
10. Li P, Bhattacharjee P, Wang S, Zhang L, Ahmed I, Guo L. Mycoviruses in species: an update. *Front Cell Infect Microbiol*. 2019;9:257.
11. de Sequeira DCM, Menezes RC, Oliveira MME, Antas PRZ, De Luca PM, de Oliveira-Ferreira J, et al. Experimental Hyalohyphomycosis by outcome of the infection in C57BL/6 murine models. *Front Microbiol*. 2017;8:1617.
12. Trinh SA, Angarone MP. *Purpureocillium lilacinum* tattoo-related skin infection in a kidney transplant recipient. *Transpl Infect Dis*. 2017;19. <https://doi.org/10.1111/tid.12689>.
13. Chen W-Y, Lin S-R, Hung S-J. Successful treatment of recurrent cutaneous *Purpureocillium lilacinum* (*Paecilomyces lilacinus*) infection with Posaconazole and Surgical Debridement. *Acta Derm Venereol*. 2019;99:1313–4.
14. Matkoski C, Sharp SE, Kiska DL. Evaluation of the Q score and Q234 systems for cost-effective and clinically relevant interpretation of wound cultures. *J Clin Microbiol*. 2006;44:1869–72.
15. Castelli MV, Alastruey-Izquierdo A, Cuesta I, Monzon A, Mellado E, Rodriguez-Tudela JL, et al. Susceptibility testing and molecular classification of *Paecilomyces* spp. *Antimicrob Agents Chemother*. 2008;52:2926–8.
16. Pastor FJ, Guarro J. Clinical manifestations, treatment and outcome of *Paecilomyces lilacinus* infections. *Clin Microbiol Infect*. 2006;12:948–60.
17. González GM, Fothergill AW, Sutton DA, Rinaldi MG, Loebenberg D. In vitro activities of new and established triazoles against opportunistic filamentous and dimorphic fungi. *Med Mycol*. 2005;43:281–4. <https://doi.org/10.1080/13693780500088416>.
18. Roujeau J-C, Sigurgeirsson B, Korting H-C, Kerl H, Paul C. Chronic dermatomycoses of the foot as risk factors for acute bacterial cellulitis of the leg: a case-control study. *Dermatology*. 2004;209:301–7.
19. Björnsdóttir S, Gottfredsson M, Thórisdóttir AS, Gunnarsson GB, Ríkardsdóttir H, Kristjánsson M, et al. Risk factors for acute cellulitis of the lower limb: a prospective case-control study. *Clin Infect Dis*. 2005;41:1416–22.
20. Zhou D, Zhang X, Du Y, Dong S, Xu Z, Yan L. Insights into the synergistic effect of fungi and bacteria for reactive red decolorization. *J Spectrosc*. 2014;2014:1–4.
21. Nucci M, Varon AG, Garnica M, et al. Increased incidence of invasive fusariosis with cutaneous portal of entry, Brazil. *Emerg Infect Dis*. 2013;19:1567–72. <https://doi.org/10.3201/eid1910.120847>.
22. Mays SR, Bogle MA, Bodey GP. Cutaneous fungal infections in the oncology patient: recognition and management. *Am J Clin Dermatol*. 2006;7:31–43.
23. Bernardeschi C, Foulet F, Ingen-Housz-Oro S, Ortonne N, Sitbon K, Quereux G, et al. French Mycosis Study Group. Cutaneous invasive *Aspergillus*: retrospective multicenter study of the french invasive-aspergillois registry and literature review. *Med (Baltim)*. 2015;94:e1018. <https://doi.org/10.1097/MD.0000000000001018>.

Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

