

Description of *Prevotella rectalis* sp. nov., a new bacterium isolated from human rectum

S. Belkacemi^{1,2}, C. I. Lo^{2,3}, S. Khelaifia^{1,2}, D. Raoult^{1,2}, P.-E. Fournier^{2,3} and F. Fenollar^{2,3}

1) Aix Marseille Université, IRD, AP-HM, MEPHI, 2) IHU-Méditerranée Infection and 3) Aix Marseille Université, IRD, AP-HM, SSA, VITROME, Marseille, France

Abstract

Using a taxonogenomics method, we describe here a Gram-negative bacterium named *Prevotella rectalis* sp. nov., strain Marseille-P4334^T (= CSUR P4334) isolated from the rectum. Strain Marseille-P4334^T has a genome that measure 3.03 Mbp with 43.3 mol% G + C content.

© 2020 The Authors. Published by Elsevier Ltd.

Keywords: Culturomics, Human gut, *Prevotella rectalis* sp. nov., Rectum

Original Submission: 20 December 2019; **Revised Submission:** 19 May 2020; **Accepted:** 22 May 2020

Article published online: 28 May 2020

Corresponding author. F. Fenollar, Institut Hospitalo-Universitaire Méditerranée-Infection, 19–21 Boulevard Jean Moulin, 13385, Marseille, Cedex 05. France.

E-mail: florence.fenollar@univ-amu.fr

Introduction

Studying the diversity of bacteria is fundamental to understanding their involvement in human diseases and normal physiological functions [1]. To unveil human microbial diversity, the culturomics approach, based on diversified culture conditions, was designed to isolate as yet uncultured species and to complement 16S rRNA metagenomics [2–4]. Furthermore, a new taxonomic strategy, named taxonogenomics, was developed to include the analysis of complete genome sequences in combination with phenotypic characteristics [5]. Herein, we report a detailed description of *Prevotella rectalis* sp. nov., for which the type strain Marseille-P4334^T was isolated from the human rectum.

Isolation and growth conditions

In 2017, we isolated an unidentified bacterium strain from a human rectum. The study was approved by the Institut

Federatif de Recherche 48 (agreement number 09-022, Marseille, France), and the patient's consent was obtained. After three tests using matrix-assisted laser desorption/ionization time of flight mass spectrometry (MALDI-TOF MS), the strain was not identified. Screening was carried out on a Microflex LT spectrometer (Bruker Daltonics, Bremen, Germany) as previously described [6]. Spectra obtained from the strain (Fig. 1) were imported and analysed using BIOTYPER 3.0 software against the Bruker database that was permanently updated with the local MEPHI database (<https://www.mediterranee-infection.com/urms-data-base>).

The growth of strain Marseille-P4334^T was possible on 5% sheep's blood-agar (bioMérieux, Marcy l'Étoile, France) under anaerobic atmosphere (anaeroGEN; Oxoid, Dardilly, France) at 37°C after 3 days of pre-incubation and 5 days in incubation.

Strain identification

The 16S rRNA gene was sequenced to classify this bacterium. Amplification was performed using the primer pair fD1 and rP2 (Eurogentec, Angers, France) and sequenced using the Big Dye® Terminator v1.1 Cycle Sequencing Kit and 3500xL Genetic Analyzer capillary sequencer (ThermoFisher, Saint-Aubin, France), as previously described [7]. The 16S rRNA nucleotide sequence was assembled and corrected using CODONCODE ALIGNER software (<http://www.codoncode.com>). Strain Marseille-P4334^T exhibited a 96.4% rRNA sequence identity with




FIG. 1. MALDI-TOF MS reference spectra of *Prevotella rectalis* sp. nov., strain Marseille-P4334^T. Separately the two reference spectra were generated by comparison of spectra from 12 individual colonies from each strain.

Prevotella corporis strain JCM 8529 (GenBank accession number: NR_113099.1), its phylogenetically closest species with standing in nomenclature (Fig. 2). Considering this value less than a threshold value established previously [8], we consequently proposed to classify strain Marseille-P4334^T as a new species within the genus *Prevotella* in the phylum *Bacteroidetes*.

Phenotypic characteristics

Colonies of strain Marseille-P4334^T were black and smooth with a mean diameter of 0.5–1 mm. Bacterial cells were Gram-negative bacilli ranging from 1 to 3 µm in length and 0.4 to 0.6 µm in width. Strain Marseille-P4334^T showed catalase-positive

and oxidase-negative activities. Electron micrograph showing the shapes of this new bacterial strain (Fig. 3) was performed with the Hitachi TM4000 instrument (Hitachi Group, Krefeld, Germany). A comparative study of the biochemical characteristics of strain Marseille-P4334^T with other closely related species in phylum *Bacteroidetes* is presented in Table 1. Results of biochemical tests performed using the API ZYM and 50 CH strips (bioMérieux) are shown in Table 2.

Genome sequencing

Genomic DNA was extracted using the EZ1 biorobot with the EZ1 DNA tissue kit (Qiagen, Hilden, Germany) and then

FIG. 2. Phylogenetic tree highlighting the position of *Prevotella rectalis* sp. nov. strain Marseille-P4334^T with regard to other closely related species. GenBank accession numbers of 16S rRNA are indicated in parentheses. Sequences were aligned using MUSCLE with default parameters, phylogenetic inferences were obtained using the maximum likelihood method and MEGA 7 software. Bootstrap values obtained by repeating the analysis 1000 times to generate a majority consensus tree are indicated at the nodes.





FIG. 3. Scanning electron microscopy (SEM) of the new species. For each strain a colony was collected from agar and immersed into a 2.5% glutaraldehyde fixative solution. Then, a drop of the suspension was directly deposited on a poly-L-lysine-coated microscope slide for 5 minutes and treated with 1% phosphotungstic acid aqueous solution (pH 2.0) for 2 minutes to increase the SEM image contrast. The slide was gently washed in water, air-dried and examined in a tabletop SEM (Hitachi TM4000). Scales and acquisition settings are shown on the figures.

sequenced on a MiSeq sequencer (Illumina Inc., San Diego, CA, USA) with the Nextera Mate Pair sample prep kit and Nextera XT Paired End (Illumina), as previously described [9]. The assembly was performed using a pipeline containing several

softwares (VELVET [10], SPADES [11] and SOAP DENOVO [12]), and trimmed data (MISEQ and TRIMMOMATIC [13] softwares) or untrimmed data (only MISEQ software). GAPCLOSER was used to reduce assembly gaps. Scaffolds <800 bp and scaffolds with a

TABLE 1. Differential characteristics of *Prevotella rectalis* sp. nov., strain Marseille-P4334^T compared with related species within their respective genus

Properties	<i>Prevotella rectalis</i>	<i>Prevotella multiformis</i>	<i>Prevotella bivia</i>	<i>Prevotella melaninogenica</i>	<i>Prevotella aurantiaca</i>
Gram stain	—	—	—	—	—
Motility	—	—	—	—	—
Endospore formation	—	—	—	—	—
Catalase	+	—	NA	NA	—
Glycerol	—	+	—	—	—
D-Cellobiose	—	+	—	—	—
D-Raffinose	—	+	—	+	+
α-Galactosidase	+	—	—	NA	—
β-Galactosidase	—	+	+	+	—
α-Fucosidase	—	NA	+	+	+
α-Glucosidase	—	—	NA	+	NA
β-Glucosidase	+	+	+	+	NA
Arginine aminopeptidase	—	—	—	+	NA
Source	Human rectal	Subgingival plaque	Vagina	Human respiratory tract	Human oral cavity

+, positive result; —, negative result; NA, data not available.

TABLE 2. Phenotypic characterizations of *Prevotella rectalis* sp. nov., based on API tests

API tests	Characteristics	P4334 ^T	
ZYM	Alkaline phosphatase	—	
	Esterase (C4)	+	
	Esterase lipase (C8)	+	
	Lipase (C14)	—	
	Leucine arylamidase	—	
	Valine arylamidase	+	
	Cystine arylamidase	—	
	Trypsin	—	
	α-Chymotrypsin	—	
	Acid phosphatase	—	
	Naphthol-AS-BI-phosphohydrolase	+	
	α-Galactosidase	+	
	β-Galactosidase	—	
	β-Glucuronidase	—	
	α-Glucosidase	—	
	β-Glucosidase	+	
	N-Acetyl-β-glucosaminidase	—	
	α-Mannosidase	—	
	α-Fucosidase	—	
	2-Nitrophenyl βD-galactopyranoside	—	
	50 CH	Glycerol	—
		Erythritol	—
		D-Arabinose	—
L-Arabinose		—	
D-Ribose		—	
D-Xylose		—	
L-Xylose		—	
D-Adonitol		—	
Methyl βD-Xylopyranoside		—	
D-Galactose		+	
D-Glucose		—	
D-Fructose		—	
D-Mannose		—	
L-Sorbose		—	
L-Rhamnose		—	
Dulcitol		—	
Inositol		—	
D-Mannitol		—	
D-Sorbitol		—	
Methyl-αD-Mannopyranoside		—	
Methyl-αD-Glucopyranoside		—	
N-Acetylglucosamine		—	
Amygdalin		—	
Arbutin		—	
Esculin ferric citrate		—	
Salicin		—	
D-Cellobiose		—	
D-Maltose	—		

TABLE 2. Continued

API tests	Characteristics	P4334 ^T
	D-Lactose	—
	D-Melibiose	—
	D-Saccharose	—
	D-Trehalose	—
	Inulin	—
	D-Melezitose	—
	D-Raffinose	—
	Amidon	—
	Glycogen	—
	Xylitol	—
	Gentiobiose	—
	D-Turanose	—
	D-Xylose	—
	D-Tagalose	—
	D-Fucose	—
	L-Fucose	—
	D-Arabitol	—
	L-Arabitol	—
	Potassium gluconate	—
	Potassium 2-ketogluconate	—
	Potassium 5-ketogluconate	—

depth value < 25% of the mean depth were removed. The best assembly was selected using different criteria (number of scaffolds, N50, number of N). Strain Marseille-P4334^T had a genome size of 3 039 397 bp with 43.3 mol% G + C content. The degree of genomic identity of this strain with closely related species was calculated using ORTHOANI software [14]. Results of this analysis are displayed in Fig. 4. For strain Marseille-P4334^T, ORTHOANI values among closely related species ranged from 68.57% between *Prevotella corporis* and *Prevotella bergensis* to 68.70% between *Prevotella bergensis* and *Prevotella salivae*. When *P. rectalis* was compared with these closely related species, values ranged from 69.91% with *P. salivae* to 76.38% with *P. corporis*.




FIG. 4. Heatmap generated with ORTHOANI values calculated using the OAT software for *Prevotella rectalis* sp. nov., strain Marseille-P4334 compared with other closely related species with standing in nomenclature.

Conclusion

On the basis of unique phenotypic features, including the MALDI-TOF spectrum, a 16S rRNA sequence divergence >1.3% and an ORTHOANI value > 95% with the phylogenetically closest species with standing in nomenclature, we formally proposed strain Marseille-P4334^T as the type strain of *Prevotella rectalis* sp. nov., as a new member within the phylum *Bacteroidetes*.

Description of *Prevotella rectalis* sp. nov

Prevotella rectalis (rec.ta'lis. L. gen. n. *rectalis* pertaining to rectum, the chamber that begins at the end of the large intestine from which this bacterium was isolated). It is a Gram-negative bacterium, non-motile and non-spore-forming. Strain Marseille-P4334^T is the type strain of *Prevotella rectalis* sp. nov., and shows catalase-positive and oxidase-negative activities. The strain grows strictly under anaerobic conditions at 37°C. The

potential pathogenicity of the type strain Marseille-P4334^T (= CSUR P4334) is unknown. This strain has a genome size of 3.03 Mbp long with a 43.3 mol% G + C content. The 16S rRNA gene sequence and whole-genome shotgun sequence of strain Marseille-P4334 were deposited in GenBank under accession numbers LS488976 and UWTY00000000, respectively. It has been isolated from the human rectum.

Conflicts of interest

None to declare.

Funding sources

This study was supported by the Institut Hospitalo-Universitaire (IHU) Méditerranée Infection, the National Research Agency under the programme *Investissements d'avenir*,

reference ANR-10-IAHU-03, the Région Provence-Alpes-Côte d'Azur and European funding FEDER PRIMI.

Acknowledgements

The authors thank Hitachi Corporation for providing the TM4000Plus Tabletop microscope and Aurelia Caputo for submitting the genomic sequence to GenBank.

References

- [1] Lagier J-C, Armougom F, Million M, Hugon P, Pagnier I, Robert C, et al. Microbial culturomics: paradigm shift in the human gut microbiome study. *Clin Microbiol Infect* 2012;18:1185–93.
- [2] Lagier J-C, Hugon P, Khelaifia S, Fournier PE, La Scola B, Raoult D. The rebirth of culture in microbiology through the example of culturomics to study human gut microbiota. *Clin Microbiol Rev* 2015;28:237–64.
- [3] Lagier J-C, Khelaifia S, Tidjani Alou M, Ndongo S, Dione N, Hugon P, et al. Culture of previously uncultured members of the human gut microbiota by culturomics. *Nat Microbiol* 2016;1:16203.
- [4] Lagier J-C, Edouard S, Pagnier I, Mediannikov O, Drancourt M, Raoult D. Current and past strategies for bacterial culture in clinical microbiology. *Clin Microbiol Rev* 2015;28:208–36.
- [5] Fournier P-E, Lagier J-C, Dubourg G, Raoult D. From culturomics to taxonomogenomics: a need to change the taxonomy of prokaryotes in clinical microbiology. *Anaerobe* 2015;36:73–8.
- [6] Lo CI, Fall B, Sambe-Ba B, Diawara S, Gueye MW, Mediannikov O, et al. MALDI-TOF mass spectrometry: a powerful tool for clinical microbiology at Hôpital Principal de Dakar, Senegal (West Africa). *PLoS One* 2015;30:10–2.
- [7] Morel AS, Dubourg G, Prudent E, Edouard S, Gouriet F, Casalta JP, et al. Complementarity between targeted real-time specific PCR and conventional broad-range 16S rDNA PCR in the syndrome-driven diagnosis of infectious diseases. *Eur J Clin Microbiol Infect Dis* 2015;34:561–70.
- [8] Stackebrandt E, Ebers J. Taxonomic parameters revisited: tarnished gold standards. *Microbiol Today* 2006;6:152–5.
- [9] Lo CI, Sankar SA, Fall B, Ba B-S, Diawara S, Gueye MW, et al. High-quality draft genome sequence and description of *Haemophilus massiliensis* sp. nov. *Stand Genomic Sci* 2016;11:31.
- [10] Zerbino DR, Birney E. Velvet: algorithms for de novo short read assembly using de Bruijn graphs. *Genome Res* 2008;18:821–9.
- [11] Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 2012;19:455–77.
- [12] Luo R, Liu B, Xie Y, Li Z, Huang W, Yuan J, et al. SOAPdenovo2: an empirically improved memory-efficient short-read de novo assembler. *Gigascience* 2012;1:18.
- [13] Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 2014;30:2114–20.
- [14] Lee I, Ouk Kim Y, Park SC, Chun J. OrthoANI: an improved algorithm and software for calculating average nucleotide identity. *Int J Syst Evol Microbiol* 2016;66:1100–3.