

Effect of triiodothyronine (T_3) augmentation of acute milnacipran administration on monoamine levels: an in vivo microdialysis study in rats

Yuji Kitaichi
Takeshi Inoue
Shin Nakagawa
Shuken Boku
Akiko Kato
Ichiro Kusumi
Tsukasa Koyama

Department of Psychiatry, Hokkaido University Graduate School of Medicine, Sapporo, Japan

Background: Up to 30% of depressed patients are partially or totally resistant to antidepressant therapy. The administration of triiodothyronine (T_3) to antidepressant nonresponders can be an effective augmentation strategy, although the mechanism is not fully understood.

Methods: In vivo microdialysis was used to examine the effect of T_3 augmentation of the antidepressant, milnacipran. Basal extracellular serotonin, norepinephrine, and dopamine levels were measured before and after acute milnacipran administration in the medial prefrontal cortex and amygdala of rats which had received subchronic (7 days) T_3 treatment or control saline.

Results: Subchronic administration of T_3 at 0.1 mg/kg significantly increased basal extracellular levels of serotonin in the medial prefrontal cortex, but not in the amygdala. In contrast, subchronic administration of T_3 at 0.2 mg/kg did not alter basal extracellular serotonin levels in the medial prefrontal cortex. Basal extracellular levels of norepinephrine and dopamine were not modified by either dose of T_3 in either region. Acute administration of milnacipran, a serotonin-norepinephrine reuptake inhibitor, to control animals resulted in a significant increase of extracellular levels of serotonin, norepinephrine, and dopamine. When administered to animals treated subchronically with T_3 at 0.1 mg/kg, milnacipran produced an additional increase in extracellular serotonin levels but not in levels of norepinephrine or dopamine in the medial prefrontal cortex of rats.

Conclusion: These results suggest that the mechanism of the augmentation effect of milnacipran by T_3 administration occurs via enhancement of serotonergic neurotransmission, but not through noradrenergic or dopaminergic neurotransmission.

Keywords: depression, in vivo microdialysis, milnacipran, monoamines, triiodothyronine

Introduction

Major depression is considered by the World Health Organization to be the fourth most disabling disorder worldwide.¹ Despite successive generations of antidepressant drugs, on average, 30%–45% of depressed patients fail to respond adequately to their initial antidepressant treatment and only 25%–35% achieve full symptom remission.² It has been estimated that 10%–30% of depressed patients are partially or totally resistant to antidepressant therapy.³

Among the treatment strategies proposed for nonresponders or partial responders to antidepressants, adjunctive administration of triiodothyronine (T_3) has been shown to be an effective augmentation therapy.^{4–7} STAR*D research has shown that the augmentation effects of T_3 in patients with refractory depression taking selective serotonin reuptake inhibitors, selective serotonin and norepinephrine reuptake inhibitors, or bupropion gave similar remission rates in all cases.^{8,9} However, a recent meta-analysis of four trials where T_3 was initiated simultaneously with a selective

Correspondence: Takeshi Inoue
Department of Psychiatry, Hokkaido University Graduate School of Medicine, North 15, West 7, Kita-ku, Sapporo 060-8638, Japan
Tel +81 11 706 5160
Fax +81 11 706 5081
Email tinoue@med.hokudai.ac.jp

serotonin reuptake inhibitor failed to show any superiority for the augmentation therapy.¹⁰

The mechanism of T_3 augmentation is still unclear. One week of T_3 administration has been shown to result in desensitization of 5-HT_{1A} receptors and an increase in extracellular serotonin levels in the frontal cortex of rats.^{11,12} Coadministration of imipramine and T_3 results in desensitization of 5-HT_{2A} receptors.¹³ Thus, the effect of T_3 augmentation of antidepressant effects has been assumed to involve serotonin neurotransmission.¹⁴ However, to our knowledge, there has been no study investigating extracellular levels of serotonin, norepinephrine, and dopamine simultaneously after T_3 treatment. A role of the norepinephrine or dopamine neurotransmitter cannot thus be ruled out.

Previous mechanistic studies have investigated the augmentation effects of T_3 with tricyclic antidepressants. However, tricyclic antidepressants are not ideal for clarification of the augmentative effects of T_3 because these agents block several receptors in addition to their effects on monoamine transporters.¹⁵ Therefore, we chose to study the effect of the augmentation effects of T_3 on the antidepressant, milnacipran, which inhibits reuptake of serotonin and norepinephrine with similar efficacy but without any receptor interactions.¹⁶⁻¹⁸ A recent review¹⁹ has suggested hyperactivity of the amygdala and hypoactivity of the prefrontal cortex in depressed patients.

In this study we used *in vivo* microdialysis in the medial prefrontal cortex and amygdala to measure the basal extracellular levels of serotonin, norepinephrine, and dopamine and levels after acute administration of milnacipran in rats which had received subchronic treatment (for one week) with T_3 or with control saline.

Materials and methods

Animals

Male Sprague-Dawley rats, obtained from the Shizuoka Laboratory Animal Centre (Shizuoka, Japan), were housed in groups of four and maintained on a 12-hour light-dark cycle (light phase 6.30 am to 6.30 pm) in a temperature-controlled environment ($22^{\circ}\text{C} \pm 1^{\circ}\text{C}$) with free access to food and water. Experiments began after a 10-day period of acclimatization. The Hokkaido University School of Medicine Animal Care and Use Committee approved all procedures, which complied with the Guide for the Care and Use of Laboratory Animals, Hokkaido University School of Medicine.

Drugs

3,3',5-Triiodo-L-thyronine sodium salt (T_3 ; O-(4-hydroxy-3-iodophenyl)-3,5-diiodo-L-tyrosine sodium salt; Sigma,

St Louis, MO) was dissolved in 1 M NaOH with saline, which was used to dilute the solution to the required concentrations (0.1 and 0.2 mg/mL). The final concentration of NaOH was 5 mM for each dose of T_3 ; 5 mM NaOH was used as the vehicle for T_3 . Milnacipran (Asahi Kasei Corporation, Shizuoka, Japan) was dissolved in distilled water, which was used as the vehicle for milnacipran, to achieve a final concentration of 10 mg/mL.

Drug administration

The animals were divided into two groups. Control group rats were given the vehicle subcutaneously once daily in the morning for 7 days and the T_3 group rats were given T_3 0.1 mg/kg or 0.2 mg/kg subcutaneously once daily in the morning for 7 days. On day 7, 2 hours after the last subcutaneous injection, the animals were implanted stereotactically in the medial prefrontal cortex or the amygdala with guide cannulae and microdialysis probes (see below). The rats were administered milnacipran 10 mg/kg intraperitoneally or saline 180 minutes after the first dialysate samples were collected. The 0.1 mg/kg and 0.2 mg/kg doses of T_3 were chosen because administration of these doses for one week has been shown to increase extracellular serotonin levels in the frontal cortex of rats.¹¹

Microdialysis procedures

The experiments were performed according to a procedure described previously.²⁰ Under anesthesia with pentobarbital (30 mg/kg intraperitoneally), AG-4 and AG-8 guide cannulae (Eicom Corporation, Kyoto, Japan) were stereotactically implanted into rats at the surface of the medial prefrontal cortex or amygdala at the following coordinates relative to bregma from the stereotaxic atlas of Paxinos and Watson;²¹ A + 3.2, ML + 0.8, DV + 1.0 mm; A - 2.8, ML + 5.0, + DV 7.4 mm, respectively. Dialysis probes with a 0.22 mm outer diameter (A-I-4-03, A-I-8-02; Eicom Corporation) were then inserted into the guide cannulae so that 3.0 mm and 2.0 mm of the probes were exposed, respectively, in the medial prefrontal cortex and amygdala tissues. Rats were housed individually after the operation. Microdialysis was performed the next day in the freely moving rats. Perfusion was started 20 hours after surgery using artificial cerebrospinal fluid (145 mM NaCl, 3.0 mM KCl, 1.3 mM CaCl₂, 1.0 mM MgCl₂) at a flow rate of 2 μL per minute. Following an initial perfusion for 2 hours, dialysate samples were collected in vials containing 50 μL of 50 mM acetic acid every 20 minutes for 480 minutes. The recovery rates of serotonin, norepinephrine, and dopamine *in vitro* at 30°C were as follows: medial prefrontal cortex, 14.5%, 12.9%,

and 14.1%, respectively; amygdala, 9.4%, 10.4%, and 11.2%, respectively (from Eicom Corporation).

Extracellular serotonin, norepinephrine, and dopamine levels were determined using high-performance liquid chromatography (HPLC) with electrochemical detection. The HPLC system comprised a liquid chromatography pump (EP-300; Eicom Corporation), a degasser (DG-300; Eicom Corporation), a reverse phase ODS column (Eicompak PP-ODS 30 4.6 mm; Eicom Corporation), an electrochemical detector (ECD-300; Eicom Corporation), and a data acquisition system (PowerChrom; AD Instruments Pty Ltd, Sydney, Australia). For serotonin and dopamine analysis, 20 µL of dialysate was injected into the HPLC system, that used a 0.1 M phosphate buffer (pH 6.0) mobile phase containing 1% (v/v) methanol, 50 mg/L Na₂ ethylenediamine tetra-acetic acid and 500 mg/L sodium L-decanesulfonate. Separations were conducted at 25°C with a flow rate of 0.5 mL per minute. In the electrochemical detector, the oxidation potential was set at of 400 mV. Standard solutions for serotonin and dopamine were injected every working day, and the peak heights for the standards were used to determine the amounts of serotonin and dopamine in the samples.

Norepinephrine concentrations were measured using the same equipment with the exception of a different reverse-phase ODS column (Eicompak CA-5ODS 150 2.1 mm; Eicom Corporation). For norepinephrine analysis, 30 µL of dialysate was injected into the HPLC system that used a 0.1 M phosphate buffer (pH 6.0) mobile phase containing 5% (v/v) methanol, 50 mg/L Na₂ ethylenediamine tetra-acetic acid and 500 mg/L L-octanesulfonic acid. Separation was conducted at 25°C with a flow rate of 0.23 mL per minute. The electrochemical detector was set at an oxidation potential of 550 mV. Standard norepinephrine solutions were injected every working day. The peak heights for the standard were used to determine the amount of norepinephrine in the samples.

Statistical analysis

All data are given as the mean values ± standard error of the mean of individual rats from each group. The serotonin, norepinephrine, and dopamine contents of dialysate samples are expressed as absolute values (pg/fraction). The mean absolute values in nine consecutive samples (–160 to 0 minutes) before milnacipran administration were used to determine basal levels. Differences between the two groups for extracellular serotonin, norepinephrine, and dopamine concentrations after milnacipran administration were calculated using a repeated-measures analysis of variance (ANOVA) for absolute values to examine the interaction between pretreatment (T₃) and time

factors (0–300 minutes). When the interaction effects were found to be significant, post hoc comparisons (differences in absolute values at each time point between the two groups, ie, the control and T₃ groups) were analyzed using an unpaired Student's *t*-test (two-tailed). The area under the curve for the 0–180-minute periods was compared between the T₃ 0.1 mg/kg group and the control group using an unpaired Student's *t*-test (two-tailed). Differences were considered to be statistically significant at *P* < 0.05.

Results

Effect of subchronic T₃ treatment on basal extracellular concentrations of serotonin, norepinephrine, and dopamine in the medial prefrontal cortex and amygdala

In the medial prefrontal cortex, basal extracellular serotonin concentrations in the T₃ 0.1 mg/kg group were significantly higher than those of the control group (*P* < 0.05, Table 1). No significant differences in basal extracellular concentrations of norepinephrine or dopamine were found between the control and T₃ 0.1 mg/kg groups. No significant differences were found between the control and T₃ 0.2 mg/kg groups in basal extracellular levels of any of the monoamines. In the amygdala, no significant difference was found in basal extracellular serotonin, norepinephrine or dopamine concentrations between the control and the T₃ 0.1 mg/kg groups.

Table 1 Effect of subchronic treatment with T₃ (for 7 days) on baseline concentrations of serotonin, norepinephrine, and dopamine in the medial prefrontal cortex and the amygdala

	Control n = 7	T ₃ 0.1 mg/kg n = 6
Medial prefrontal cortex T₃ 0.1 mg/kg		
Serotonin	0.793 ± 0.084	1.349 ± 0.234*
Norepinephrine	1.171 ± 0.107	1.060 ± 0.150
Dopamine	0.246 ± 0.027	0.220 ± 0.024
	Control n = 6	T ₃ 0.2 mg/kg n = 6
Medial prefrontal cortex T₃ 0.2 mg/kg		
Serotonin	1.08 ± 0.22	1.12 ± 0.37
Norepinephrine	0.880 ± 0.164	0.827 ± 0.171
Dopamine	0.242 ± 0.039	0.237 ± 0.037
	Control n = 3	T ₃ 0.1 mg/kg n = 4
Amygdala T₃ 0.1 mg/kg		
Serotonin	0.259 ± 0.024	0.342 ± 0.035
Norepinephrine	0.561 ± 0.072	0.760 ± 0.139
Dopamine	0.504 ± 0.039	0.606 ± 0.121

Notes: Values represent mean ± standard error of the mean (pg per 20-minute fraction); n = number of animals in each group; data are calculated from Figures 1–3; **P* < 0.05 compared with control group.

Effect of acute milnacipran administration on extracellular concentrations of serotonin, norepinephrine, and dopamine in the medial prefrontal cortex and amygdala after control and subchronic T₃ treatments

Acute intraperitoneal administration of milnacipran 10 mg/kg to saline-treated (control) animals resulted in an increase in extracellular concentrations of serotonin, norepinephrine, and dopamine in the medial prefrontal cortex (Figures 1 and 2) and in the amygdala (Figure 3).

Following intraperitoneal administration of milnacipran 10 mg/kg to rats treated with subchronic T₃, two-way




Figure 1 Effect of acute intraperitoneal administration of milnacipran 10 mg/kg on extracellular concentrations of serotonin (A), norepinephrine (B), and dopamine (C) in the medial prefrontal cortex after subchronic treatment with T₃ 0.1 mg/kg. **Notes:** Values represent the mean ± standard error of the mean (pg per 20-minute fraction). Black arrows indicate milnacipran administration. [n = 7 (control group), n = 6 (T₃ 0.1 mg/kg group)]. *P < 0.05 versus control group.




Figure 2 Effect of acute intraperitoneal administration of milnacipran 10 mg/kg on extracellular concentrations of serotonin (A), norepinephrine (B), and dopamine (C) in the medial prefrontal cortex after subchronic treatment with T₃ 0.2 mg/kg. **Notes:** Values represent the mean ± standard error of the mean (pg per 20-minute fraction). Black arrows indicate milnacipran administration. [n = 6 (control group), n = 6 (T₃ 0.2 mg/kg group)].

ANOVA with repeated measures (0–300 minutes) of extracellular serotonin concentrations in the medial prefrontal cortex (Figure 1A) showed significant interaction between T₃ treatment and time after milnacipran administration [$F(15,165) = 1.992, P = 0.0185$] and a significant main effect of time [$F(15,165) = 13.111, P < 0.0001$]. However, the main effect of T₃ treatment after milnacipran administration was not significant [$F(1,11) = 4.758, P = 0.0518$]. The T₃ 0.1 mg/kg group showed significantly higher concentrations of extracellular serotonin than those of the control group at two time points, ie, at 40 and 100 minutes ($P < 0.05$, Figure 1).




Figure 3 Effect of acute intraperitoneal administration of milnacipran 10 mg/kg on extracellular concentrations of serotonin (A), norepinephrine (B), and dopamine (C) in the amygdala after subchronic treatment with T_3 0.1 mg/kg.

Notes: Values represent the mean \pm standard error of the mean (pg per 20-minute fraction). Black arrows indicate milnacipran administration. [$n = 3$ (control group), $n = 4$ (T_3 0.1 mg/kg group)].

The area under the curve (0–180 minutes) for extracellular serotonin in the subchronic T_3 0.1 mg/kg group was significantly higher than that of the control group (527.4 ± 101.1 and 282.9 ± 46.5 pg * minutes, respectively, $P < 0.05$). The area under the curve (0–180 minutes) for extracellular norepinephrine and dopamine were not significantly different between the T_3 and control groups.

For the extracellular norepinephrine concentrations in the medial prefrontal cortex (Figure 1B), two-way ANOVA with repeated measures (0–300 minutes) showed a significant

main effect of time [$F(15,165) = 15.234$, $P < 0.0001$]. There was neither a significant main effect of T_3 0.1 mg/kg treatment after milnacipran administration [$F(1,11) = 0.25$, $P = 0.8769$] nor an interaction between subchronic T_3 0.1 mg/kg treatment and time after milnacipran administration [$F(15,165) = 1.107$, $P = 0.3537$].

For the extracellular dopamine concentrations in the medial prefrontal cortex (Figure 1C), two-way ANOVA with repeated measures (0–300 minutes) showed a significant main effect of time [$F(15,165) = 8.449$, $P < 0.0001$]. There was neither a significant main effect of subchronic T_3 0.1 mg/kg treatment after milnacipran administration [$F(1,11) = 0.504$, $P = 0.4926$] nor an interaction between subchronic T_3 0.1 mg/kg treatment and time after milnacipran administration [$F(15,165) = 1.536$, $P = 0.0979$].

In rats treated with subchronic T_3 0.2 mg/kg (Figure 2), two-way ANOVA with repeated measures (0–300 minutes) showed a significant main effect of time for the extracellular serotonin, norepinephrine, and dopamine concentrations [$F(15,150) = 15.068$, $P < 0.0001$; $F(15,150) = 12.058$, $P < 0.0001$; $F(15,150) = 4.241$, $P < 0.0001$, respectively]. However, there was neither a main effect of subchronic T_3 0.2 mg/kg treatment after milnacipran administration [$F(1,10) = 0.036$, $P = 0.853$; $F(1,10) = 0.206$, $P = 0.660$; $F(1,10) = 0.325$, $P = 0.581$, respectively] nor an interaction between subchronic T_3 0.2 mg/kg treatment and time after milnacipran administration [$F(15,150) = 0.046$, $P > 0.999$; $F(15,150) = 0.463$, $P = 0.955$; $F(15,150) = 0.664$, $P = 0.816$, respectively].

In the amygdala (Figure 3), for extracellular serotonin and norepinephrine concentrations, two-way ANOVA with repeated measures (0–300 minutes) showed a significant main effect of time [$F(15,75) = 1.843$, $P = 0.044$; $F(15,75) = 5.197$, $P < 0.0001$, respectively]. There was neither a main effect of subchronic T_3 0.1 mg/kg treatment after milnacipran administration [$F(1,5) = 0.0008$, $P = 0.993$; $F(1,5) = 0.848$, $P = 0.399$, respectively] nor an interaction between subchronic T_3 0.1 mg/kg treatment and time after milnacipran administration [$F(15,75) = 0.547$, $P = 0.905$; $F(15,75) = 0.342$, $P = 0.988$, respectively].

For the extracellular dopamine concentrations in the amygdala (Figure 3C), there was neither a main effect of subchronic T_3 0.1 mg/kg treatment after milnacipran administration [$F(1,5) = 0.165$, $P = 0.701$] nor an interaction between subchronic T_3 0.1 mg/kg treatment and time after milnacipran administration [$F(15,75) = 1.325$, $P = 0.209$]. The main effect of time was not significant [$F(15,75) = 1.217$, $P = 0.279$].

Discussion

Subchronic (7 days) administration of T_3 0.1 mg/kg significantly increased basal extracellular levels of serotonin in the medial prefrontal cortex (Table 1). This finding is consistent with previous studies^{11,12} which have shown that T_3 administered daily for 7 days over the range 0.02–0.2 mg/kg increases basal extracellular levels of serotonin in the rat cortex, in a dose-dependent manner. Subchronic administration of T_3 at 0.1 mg/kg had no effect on basal extracellular levels of norepinephrine or dopamine (Table 1). Although several studies have investigated the role of serotonin in the mechanism of T_3 augmentation,^{11–13} to our knowledge, the present study is the first to examine extracellular serotonin, norepinephrine, and dopamine levels simultaneously after T_3 treatment. It is currently believed that the effect of T_3 on serotonin neurotransmission is integral to the mechanism of T_3 augmentation of antidepressant effects.¹⁴ The results of the present study strongly support this notion, and the absence of effect on the levels of norepinephrine or dopamine makes it unlikely that these neurotransmitters are involved.

The present study found no effects on basal extracellular serotonin levels in the amygdala (Table 1) after subchronic administration of T_3 0.1 mg/kg, in contrast with the increase found in the medial prefrontal cortex (Table 1). A previous study¹¹ has reported that subchronic administration of T_3 at 0.1 mg/kg increased basal extracellular serotonin levels in the frontal cortex of rats, but not in the hippocampus. The effects of T_3 on increases in extracellular serotonin thus appear to be sensitive, not only to dose but to the brain region studied. A more detailed study of brain regions sensitive to the effects of T_3 administration might be helpful in clarifying the mechanism of action of T_3 augmentation.

Results of the present study showed that, in contrast with the effect of subchronic administration of T_3 0.1 mg/kg, similar administration of T_3 0.2 mg/kg did not alter basal extracellular serotonin levels (Table 1). This differs from previous findings.¹² However, higher doses of T_3 0.5 mg/kg have been shown to have no effect on extracellular serotonin levels in the frontal cortex of rats,¹² suggesting a bell-shaped relationship between the dose of T_3 and the increase in serotonin levels. The nature of the bell-shaped curve has not been studied in detail, but it is possible that experimental differences between studies (strain of rats, for example) could result in a leftward shift of the bell-shaped curve, offering a possible explanation of the different results obtained.

As we have reported previously,^{20,22} acute administration of the antidepressant, milnacipran, to rats receiving saline resulted in an increase in basal extracellular concentrations

of serotonin, norepinephrine, and dopamine in the medial prefrontal cortex (Figures 1 and 2) and amygdala (Figure 3). In rats treated subchronically with T_3 0.1 mg/kg, acute administration of milnacipran resulted in an additional increase in extracellular serotonin levels, but not in norepinephrine or dopamine levels. This suggests that the augmentation effect of T_3 administration on a selective serotonin and norepinephrine reuptake inhibitor, such as milnacipran, occurs through enhancement of serotonergic neurotransmission, but not through noradrenergic and dopaminergic neurotransmission.

The present study suffers from several weaknesses. A one-week administration of T_3 was used since this is the duration typically used in similar animal studies.^{11,13,23} However, a recent study reported that coadministration of T_3 and the selective serotonin reuptake inhibitor, fluoxetine, for 3 weeks caused greater increases in neurogenesis in the hippocampus than fluoxetine administration alone.²⁴ The effects of one week of administration of T_3 may thus not be optimal for extrapolating to the effects occurring with long-term treatment as used clinically.

In addition, in the present study, the effect of milnacipran was studied acutely. Subchronic administration of milnacipran to rats has been shown,²⁵ however, to enhance noradrenergic transmission beyond that achieved with acute administration while no such effect is seen on serotonergic or dopaminergic transmission. Thus, it may be interesting to study longer durations of administration of both T_3 and milnacipran in future studies.

Another weakness of the present study is that it is exclusively neurochemical. Simultaneous behavioral studies could possibly give a “functional window” for studying the phenomenon. Unfortunately, the constraints of *in vivo* microdialysis are not compatible with behavioral paradigms.

In conclusion, subchronic administration of T_3 0.1 mg/kg significantly increased basal extracellular levels of serotonin in the medial prefrontal cortex, but not in the amygdala. Subchronic administration of T_3 at 0.2 mg/kg showed no effect on extracellular levels of serotonin in the medial prefrontal cortex, suggesting a bell-shaped dose-effect relationship. Acute administration of milnacipran, which in control animals significantly increased levels of serotonin, norepinephrine, and dopamine, caused an additional increase in extracellular serotonin levels, but not in norepinephrine or dopamine levels, in the medial prefrontal cortex in rats treated subchronically with T_3 0.1 mg/kg. This suggests that the augmentation effect of T_3 administration on selective serotonin and norepinephrine reuptake inhibitors, such as

milnacipran, occurs via enhancement of serotonergic neurotransmission, but not through noradrenergic and dopaminergic neurotransmission.

Acknowledgments

This work was supported in part by a Grant-in-Aid for Scientific Research (21591478 to TI and 17790800 to YK) from the Japanese Ministry of Education, Culture, Sports, Science and Technology. TK has received honoraria from Pfizer, Astellas, and Eli Lilly, and has received a research grant from Dainippon Sumitomo Pharma, Pfizer, Astellas, and GSK. He is an advisory board member for GSK.

Disclosure

The authors report no conflicts of interest in this work.

References

- Murray CJL, Lopez AD. *The Global Burden of Disease: A Comprehensive Assessment of Mortality and Disability from Disease, Injuries, and Risk Factors in 1990 and Projected to 2020*. Cambridge, MA: Harvard University Press; 1996:1.
- Fava M. Diagnosis and definition of treatment-resistant depression. *Biol Psychiatry*. 2003;53:649–659.
- Joffe RT, Levitt AJ, Sokolov ST. Augmentation strategies. *J Clin Psychiatry*. 1996;57:25–31.
- Connolly KR, Thase ME. If at first you don't succeed: a review of the evidence for antidepressant augmentation, combination and switching strategies. *Drugs*. 2011;71:43–64.
- Cooper-Kazaz R, Lerer B. Efficacy and safety of triiodothyronine supplement in patients with major depressive disorder treated with specific serotonin reuptake inhibitors. *Int J Neuropsychopharmacol*. 2008;11:685–699.
- Abraham G, Milev R, Stuart JL. T₃ augmentation of SSRI resistant depression. *J Affect Disord*. 2006;91:211–215.
- Altshuler LL, Bauer M, Frye MA, et al. Does thyroid supplementation accelerate tricyclic antidepressant response? A review and meta-analysis of the literature. *Am J Psychiatry*. 2001;158:1617–1622.
- Nierenberg AA, Fava M, Trivedi MH, et al. A comparison of lithium and T(3) augmentation following two failed medication treatments for depression: a STAR*D report. *Am J Psychiatry*. 2006;163:1519–1530.
- Rush AJ, Trivedi MH, Wisniewski SR, et al. Acute and longer-term outcomes in depressed outpatients requiring one or several treatment steps: a STAR*D report. *Am J Psychiatry*. 2006;163:1905–1917.
- Papakostas GI, Cooper-Kazaz R, Appelhof BC, et al. Simultaneous initiation (coinitiation) of pharmacotherapy with triiodothyronine and a selective serotonin reuptake inhibitor for major depressive disorder: a quantitative synthesis of double-blind studies. *Int Clin Psychopharmacol*. 2009;24:19–25.
- Gur E, Lerer B, Newman ME. Chronic clomipramine and triiodothyronine increases serotonin levels in rat frontal cortex in vivo: relationship to serotonin autoreceptor activity. *J Pharmacol Exp Ther*. 1999;288:81–87.
- Gur E, Lifschytz T, Lerer B, Newman ME. Effects of triiodothyronine and imipramine on basal 5-HT levels and 5-HT(1) autoreceptor activity in rat cortex. *Eur J Pharmacol*. 2002;457:37–43.
- Moreau X, Jeanningros R, Mazzola-Pomietto P. Chronic effects of triiodothyronine in combination with imipramine on 5-HT transporter, 5-HT(1A) and 5-HT(2A) receptors in adult rat brain. *Neuropsychopharmacology*. 2001;24:652–662.
- Bauer M, Heinz A, Whybrow PC. Thyroid hormones, serotonin and mood: of synergy and significance in the adult brain. *Mol Psychiatry*. 2002;7:140–156.
- Richelson E. Interactions of antidepressants with neurotransmitter transporters and receptors and their clinical relevance. *J Clin Psychiatry*. 2003;64 Suppl 13:5–12.
- Moret C, Charveron M, Finberg JP, Couzinier JP, Briley M. Biochemical profile of mitalcipran (F 2207), 1-phenyl-1-diethyl-aminocarbonyl-2-aminomethyl-cyclopropane (Z) hydrochloride, a potential fourth generation antidepressant drug. *Neuropharmacology*. 1985;24:1211–1219.
- Mochizuki D, Tsujita R, Yamada S, et al. Neurochemical and behavioural characterization of milnacipran, a serotonin and noradrenaline reuptake inhibitor in rats. *Psychopharmacology (Berl)*. 2002;162:323–332.
- Vaishnavi SN, Nemeroff CB, Plott SJ, Rao SG, Kranzler J, Owens MJ. Milnacipran: a comparative analysis of human monoamine uptake and transporter binding affinity. *Biol Psychiatry*. 2004;55:320–322.
- Quidé Y, Witterveen AB, El-Hage W, Veltman DJ, Olff M. Differences between effects of psychological versus pharmacological treatments on functional and morphological brain alterations in anxiety disorders and major depressive disorder: a systematic review. *Neurosci Biobehav Rev*. 2012;36:626–644.
- Kitaichi Y, Inoue T, Nakagawa S, Izumi T, Koyama T. Effect of milnacipran on extracellular monoamine concentrations in the medial prefrontal cortex of rats pre-treated with lithium. *Eur J Pharmacol*. 2005;516:219–226.
- Paxinos G, Watson C. *The Rat Brain in Stereotaxic Coordinates*. 3rd ed. San Diego, CA: Academic Press; 1997.
- Kitaichi Y, Inoue T, Izumi T, et al. Effect of co-administration of a serotonin-noradrenaline reuptake inhibitor and a dopamine agonist on extracellular monoamine concentrations in rats. *Eur J Pharmacol*. 2008;584:285–290.
- Lifschytz T, Goltser-Dubner T, Landshut G, Lerer B. Effect of triiodothyronine on 5-HT1A and 5-HT1B receptor expression in rat forebrain and on latency to feed in the novelty suppressed feeding test. *Prog Neuropsychopharmacol Biol Psychiatry*. 2010;34:632–638.
- Eitan R, Landshut G, Lifschytz T, Einstein O, Ben-Hur T, Lerer B. The thyroid hormone, triiodothyronine, enhances fluoxetine-induced neurogenesis in rats: possible role in antidepressant-augmenting properties. *Int J Neuropsychopharmacol*. 2010;13:553–561.
- Kitaichi Y, Inoue T, Izumi T, Nakagawa S, Kato A, Koyama T. Subchronic milnacipran treatment increases basal extracellular noradrenaline concentrations in the medial prefrontal cortex of rats. *Eur J Pharmacol*. 2005;520:37–42.

Neuropsychiatric Disease and Treatment

Publish your work in this journal

Neuropsychiatric Disease and Treatment is an international, peer-reviewed journal of clinical therapeutics and pharmacology focusing on concise rapid reporting of clinical or pre-clinical studies on a range of neuropsychiatric and neurological disorders. This journal is indexed on PubMed Central, the 'PsycINFO' database and CAS.

Submit your manuscript here: <http://www.dovepress.com/neuropsychiatric-disease-and-treatment-journal>

Dovepress

The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.