

# Infectious stentitis after treatment of coarctation of the aorta: a case report

Heleen B. van der Zwaan <sup>1\*</sup>, Gertjan Tj. Sieswerda <sup>1</sup>, Gregor J. Krings<sup>2</sup>, and Michiel Voskuil<sup>1</sup>

<sup>1</sup>Department of Cardiology, University Medical Center Utrecht, Heidelberglaan 100, 3584 CX Utrecht, The Netherlands; and <sup>2</sup>Department of Pediatric Cardiology, Wilhelmina Children's Hospital, University Medical Center Utrecht, Heidelberglaan 100, 3584 CX Utrecht, The Netherlands

Received 17 October 2019; first decision 18 November 2019; accepted 17 March 2020; online publish-ahead-of-print 3 May 2020

## Background

Aortitis is a rare condition that can be caused by inflammatory or infectious aetiologies. The clinical presentation of aortitis includes a heterogeneous range of symptoms and clinical signs.

## Case summary

We present a 53-year-old man whose medical history included presence of a ventricular septal defect, a bicuspid aortic valve, and coarctation of the aorta. The coarctation was treated with percutaneous stent implantation. One and a half years later, he presented to our hospital with complaints of fatigue, night sweats, and shivers. Physical examination revealed a fever, tachycardia, and hypertension. Imaging studies showed no signs of endocarditis. Positron emission tomography–computed tomography (PET–CT) showed an increase in <sup>18</sup>F-fluorodeoxyglucose uptake at the distal end of the stent in the descending aorta. Blood cultures revealed a *Streptococcus gordonii* and antibiotic treatment was adjusted accordingly. The patients' functional status improved quickly, the fever resolved, and the laboratory markers of inflammation returned to normal.

## Discussion

Aortitis is extremely rare after stent implantation. Risk factors for aortitis include congenital vascular malformation and stent implantation. Computed tomography is currently the imaging study of choice for aortitis, while PET–CT seems ideal for identification of stent infection. Mortality associated with infectious aortitis ranges from 21% to 44%, with generally higher mortality if managed with antibiotics alone. The differential diagnosis of stent infection should be taken into account in patients presenting with fever and chills after previous stent procedures.

## Keywords

Infection • Coarctation of the aorta • Stenting • Case report

## Learning points

- The differential diagnosis of stent infection should be taken into account in patients presenting with fever and chills after previous stent procedures.
- Surgery should be considered as treatment in urgent cases, especially when aneurysm formation is present.

## Introduction

Surgical repair or catheter-based stenting is recommended for adult patients with hypertension and significant coarctation of the aorta (CoA).<sup>1,2</sup> Various complications have been described afterwards, including recoarctation, aneurysm formation, and dissection. Long-term outcomes have been provided by various research groups, ranging from 99% at 30 years of age to 65% survival at 70 years of

\* Corresponding author. Tel: +31 (0)88 7574366, Email: [h.b.vanderzwaan-3@umcutrecht.nl](mailto:h.b.vanderzwaan-3@umcutrecht.nl)

Handling Editor: Monika Arzanauskaitė

Peer-reviewers: Christoph Sinning and Julien Adjedj

Compliance Editor: Amir Aziz

Supplementary Material Editor: Vassilios Parisi Memtsas

© The Author(s) 2020. Published by Oxford University Press on behalf of the European Society of Cardiology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact [journals.permissions@oup.com](mailto:journals.permissions@oup.com)

age.<sup>3–5</sup> Event-free survival have been reported from 96.7% 10 years after initial treatment to 53% at 70 years of age.<sup>3–5</sup>

Aortitis is a rare condition that can be caused by inflammatory or infectious aetiologies. The clinical presentation of aortitis includes a heterogeneous range of symptoms and clinical signs, largely determined by the underlying cause, the location of aortic wall thickening and the presence of coexisting arteries at the side of the aortitis.<sup>6</sup> Here, we present a case of infectious aortitis in the presence of a bare-metal stent used for treatment of coarctation of the aorta. Written informed consent by the patient was obtained.

## Timeline

Case	A 53-year-old man with a history of endocarditis, a ventricular septal defect, and coarctation of the aorta treated by percutaneous stent implantation, presented with fatigue, night sweats, and fever. Blood cultures were taken and antibiotic treatment was initiated.
Day 1	Transthoracic echocardiography showed no signs of endocarditis.
Day 3	Transoesophageal echocardiography showed no signs of endocarditis. Blood cultures identified <i>Streptococcus gordonii</i> . Antibiotic treatment was switched accordingly.
Day 7	Positron emission tomography–computed tomography (PET–CT) scan showed an increase in <sup>18</sup> F-fluorodeoxyglucose (FDG) uptake at the distal end of the stent in the descending aorta.
Day 24	Clinical improvement, no more complaints or fever. Markers of inflammation returned towards normal. Repeat PET–CT: normalization of the FDG uptake at the distal stent end. Discharge home.
Day 49	Completion of intravenously administered antibiotic treatment.


## Case presentation

A 53-year-old man was referred to our hospital for a second opinion because of the suspicion of infection of an aortic stent. His medical history included presence of a small, restrictive muscular ventricular septal defect (VSD), a bicuspid aortic valve (BAV) of the right–left coronary cusps type, and coarctation of the aorta. More than 30 years before presentation, he had a sepsis with a *Staphylococcus aureus* and was treated as if it were endocarditis. In 2016, the coarctation was treated percutaneously by uncomplicated implantation of a Cheatham-Platinum/Iridium CP stent (NuMed Inc. Hopkinton, NY, USA) because of persistent hypertension. A non-covered stent was used to allow unrestricted perfusion of the lusoric artery through the stent struts. Approximately one and a half years later, he presented to the referring hospital with fatigue, night sweats, and shivers that persisted for several weeks. Physical examination revealed fever up to 40°C combined with a blood pressure of 153/78 mm of mercury,



**Figure 1** The computed tomography scan showing the CP stent in the aortic arch/descending aorta one day after implantation without any signs of inflammation.

a tachycardia of 106 b.p.m., and a known systolic murmur on the aortic valve. The electrocardiogram showed a sinus tachycardia including a first-degree atrioventricular block. Laboratory results included leucocytes  $11.1 \times 10^9/L$  (reference range  $4.0–10.0 \times 10^9/L$ ), C-reactive protein 66 mg/L (0–10 mg/L), haemoglobin 8.2 mmol/L (8.6–10.7 mmol/L). He was admitted with the working diagnosis of endocarditis and after taking blood cultures antibiotic treatment consisting of ceftriaxone was initiated. Transthoracic echocardiography showed no proof of endocarditis and therefore, transoesophageal echocardiography was performed: the BAV functioned well, the peak velocity was 1.8 m/s, no aortic valve regurgitation was seen, and no vegetation or other characteristics of endocarditis were found. The entry point was extensively searched for but could not be identified. Blood cultures revealed a *Streptococcus gordonii* and antibiotic treatment was adjusted accordingly to penicillin 12 million units per 24 h. A computed tomography (CT) scan was performed and showed no vegetation or thrombus of the BAV, no wall thickening, aneurysm, abscess, or other signs of inflammation (Figure 1). A positron emission tomography (PET)–CT scan with <sup>18</sup>F-fluorodeoxyglucose (FDG) showed an increase in FDG uptake at the distal end of the stent in the descending aorta (Figure 2). Hereafter, antibiotic treatment was switched again to ceftriaxone once daily 2000 mg and continued for 6 weeks, in accordance with treatment of endocarditis with a prosthetic valve.<sup>7</sup> Clinically the patients' functional status improved quickly, the fever resolved and the laboratory markers of inflammation returned to normal. Control PET–CT scan a week after treatment showed normalization of FDG uptake (Figure 3). The patient was sent home where he completed the antibiotic treatment. During outpatient follow-up, he had no fever and control blood cultures



**Figure 2** The positron emission tomography–computed tomography scan with  $^{18}\text{F}$ -fluorodeoxyglucose showing increased  $^{18}\text{F}$ -fluorodeoxyglucose uptake at the distal end of the stent in the proximal descending aorta.

remained negative. At last follow-up, August 2019, he was free of complaints.

## Discussion


We present a rare case of aortitis after stenting of a CoA. Patients with congenital heart disease have an increased risk of developing infective endocarditis and this risk is related to the type of defect encountered.<sup>1,2,8</sup> Beforehand, our patient could have an infection related to the BAV, the VSD, and aortic stent. Transthoracic nor transoesophageal echocardiography showed signs of endocarditis and therefore we chose for a PET–CT scan that is considered an important [supplementary method](#) for patients with suspected infective endocarditis.<sup>7</sup> By doing so, we identified infection of the aortic stent and no abnormalities at the side of the BAV or VSD were seen.

Percutaneous stent implantation is a standard treatment of CoA in adulthood. Aortitis is extremely rare after stent implantation. As we stated earlier, in the current case, a PET–CT scan could demonstrate

situation of the infectious substrate at the stent and its infectious activity dissolved after antibiotic therapy. Under normal conditions, the aorta is resistant to infection, but risk factors such as congenital vascular malformation<sup>9</sup> or stent implantation<sup>6,10</sup> have been identified for the development of infectious aortitis. It seems reasonable that the vessel trauma caused by stent implantation may be a predilection area for aortitis.

Both transoesophageal echocardiography as well as  $^{99\text{m}}\text{Tc}$ -labelled white blood cell scintigraphy have been used to diagnose aortitis and aortic prosthetic stent infection.<sup>11–14</sup> Computed tomography with contrast is currently the imaging study of choice for aortitis,<sup>9</sup> while a PET–CT seems to be an ideal medium for identification of stent infection.<sup>15</sup>

The most common pathogens causing infectious aortitis have been identified, being *Salmonella* species, *S. aureus*, streptococcal species, Gram-negative bacilli other than *Salmonella* and fungi.<sup>9</sup> Mechanisms of infection include haematogenous spread, contiguous seeding from adjacent infection, and traumatic or iatrogenic inoculation.<sup>16</sup> *Streptococcus gordonii*, which was found in our patient, is a normal inhabitant of the



**Figure 3** Control positron emission tomography–computed tomography at follow-up scan showing normalization of  $^{18}\text{F}$ -fluorodeoxyglucose uptake.

human oral cavity. It is regarded as one of the main causative agent in the development of subacute bacterial endocarditis.

The CP stent is widely used for the treatment of large vessel stenosis in various congenital heart diseases. In a recent publication, no stentitis was reported during a 12-month follow-up.<sup>17</sup> In general, intravascular bare metal stent infections are a rare but potentially serious complication as Bosman *et al.*<sup>15</sup> found in their review of literature. Mortality associated with infectious aortitis ranges from 21% to 44%, with generally higher mortality if managed with antibiotics alone.<sup>9</sup> Mlynski *et al.*<sup>18</sup> reported a case of pneumococcal aortitis secondary to endovascular bare-metal stent infection 1 year after stent implantation that was complicated by aortic rupture. Urgent surgical treatment led to a favourable outcome. This underlines the importance of considering surgery in urgent cases, especially when aneurysm formation is present.

In conclusion, infectious aortitis may rarely occur after stent implantation for treatment of coarctation of the aorta. The differential diagnosis of stent infection should be taken into account in patients presenting with fever and chills after previous stent procedures. In our case, it has been treated successfully by medical therapy alone. Surgery has to be considered in urgent cases of aortitis.

## Lead author biography



Heleen B. van der Zwaan currently works as a cardiologist at the University Medical Center in Utrecht where she did specialization in adult congenital heart disease. She completed her thesis on right ventricular function assessment by three-dimensional echocardiography in patients with various congenital heart diseases at the Erasmus Medical

Center in Rotterdam. She is interested in cardiac imaging and treatment of advanced heart failure therapy.

## Supplementary material

Supplementary material is available at *European Heart Journal - Case Reports* online.

**Slide sets:** A fully edited slide set detailing this case and suitable for local presentation is available online as [Supplementary data](#).

**Consent:** The author/s confirm that written consent for submission and publication of this case report including image(s) and associated text has been obtained from the patient in line with COPE guidance.

**Conflict of interest:** none declared.

## References

1. Stout KK, Daniels CJ, Aboulhosn JA, Bozkurt B, Broberg CS, Colman JM, Crumb SR, Dearani JA, Fuller S, Gurvitz M, Khairy P, Landzberg MJ, Saidi A, Valente AM, Van Hare GF. 2018 AHA/ACC Guideline for the management of adults with congenital heart disease: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Circulation* 2019;**139**:e698–ee800.
2. Baumgartner H, Bonhoeffer P, De Groot NM, de Haan F, Deanfield JE, Galie N, Gatzoulis MA, Gohlke-Baerwolf C, Kaemmerer H, Kilner P, Meijboom F, Mulder BJ, Oechslin E, Oliver JM, Serraf A, Szatmari A, Thaulow E, Vouhe PR, Walma E; Task Force on the Management of Grown-up Congenital Heart Disease of the European Society of Cardiology (ESC); Association for European Paediatric Cardiology (AEPIC); ESC Committee for Practice Guidelines (CPG). ESC Guidelines for the management of grown-up congenital heart disease (new version 2010). *Eur Heart J* 2010;**31**:2915–2957.
3. Brown ML, Burkhart HM, Connolly HM, Dearani JA, Cetta F, Li Z, Oliver WC, Warnes CA, Schaff HV. Coarctation of the aorta: lifelong surveillance is mandatory following surgical repair. *J Am Coll Cardiol* 2013;**62**:1020–1025.
4. Lee MGY, Babu-Narayan SV, Kempny A, Uebing A, Montanaro C, Shore DF, d'Udekem Y, Gatzoulis MA. Long-term mortality and cardiovascular burden for adult survivors of coarctation of the aorta. *Heart* 2019;**105**:1190–1196.
5. Choudhary P, Canniffe C, Jackson DJ, Tanous D, Walsh K, Celermajer DS. Late outcomes in adults with coarctation of the aorta. *Heart* 2015;**101**:1190–1195.
6. Gornik HL, Creager MA. Aortitis. *Circulation* 2008;**117**:3039–3051.
7. Habib G, Lancellotti P, Antunes MJ, Bongiorni MG, Casalta JP, Del Zotti F, Dulgheru R, El Khoury G, Erba PA, lung B, Miro JM, Mulder BJ, Plonska-Gosciniak

- E, Price S, Roos-Hesselink J, Snygg-Martin U, Thuny F, Tornos Mas P, Vilacosta I, Zamorano JL; ESC Scientific Document Group. 2015 ESC Guidelines for the management of infective endocarditis: the Task Force for the Management of Infective Endocarditis of the European Society of Cardiology (ESC). Endorsed by: European Association for Cardio-Thoracic Surgery (EACTS), the European Association of Nuclear Medicine (EANM). *Eur Heart J* 2015;**36**:3075–3128.
8. Verheugt CL1, Uiterwaal CS, van der Velde ET, Meijboom FJ, Pieper PG, Veen G, Stappers JL, Grobbee DE, Mulder BJ. Turning 18 with congenital heart disease: prediction of infective endocarditis based on a large population. *Eur Heart J* 2011; **32**:1926–1934.
  9. Foote EA1, Postier RG, Greenfield RA, Bronze MS. Infectious aortitis. *Curr Treat Options Cardiovasc Med* 2005;**7**:89–97.
  10. Aykan AC, Yıldız M, Özkan M. Infective endocarditis, thoracic aortitis, and mycotic aneurysm formation complicating balloon angioplasty of aortic coarctation. *Cardiol Young* 2013;**23**:138–140.
  11. Skinner JR1, Bexton R, Wren C. Aortic coarctation endarteritis and aneurysm: diagnosis by transoesophageal echocardiography. *Int J Cardiol* 1992;**34**: 216–218.
  12. Harris KM, Malenka DJ, Plehn JF. Transesophageal echocardiographic evaluation of aortitis. *Clin Cardiol* 1997;**20**:813–815.
  13. Ryu SW, Allman KC. Native aortic and prosthetic vascular stent infection on <sup>99m</sup>Tc-labeled white blood cell scintigraphy. *J Nucl Med Technol* 2014;**42**:120–121.
  14. Meller J, Strutz F, Siefker U, Scheel A, Sahlmann CO, Lehmann K, Conrad M, Vosshehrich R. Early diagnosis and follow-up of aortitis with [(18)F]FDG PET and MRI. *Eur J Nucl Med Mol Imaging* 2003;**30**:730–736.
  15. Bosman WM, Borger van der Burg BL, Schuttevaer HM, Thoma S, Hedeman Joosten PP. Infections of intravascular bare metal stents: a case report and review of literature. *Eur J Vasc Endovasc Surg* 2014;**47**:87–99.
  16. Restrepo CS, Ocazonez D, Suri R, Vargas D. Aortitis: imaging spectrum of the infectious and inflammatory conditions of the aorta. *Radiographics* 2011;**31**:435–451.
  17. Ewert P, Schubert S, Peters B, Abdul-Khaliq H, Nagdyman N, Lange PE. The CP stent–short, long, covered–for the treatment of aortic coarctation, stenosis of pulmonary arteries and caval veins, and Fontan anastomosis in children and adults: an evaluation of 60 stents in 53 patients. *Heart* 2005;**91**:948–953.
  18. Mlynski A, Mordant P, Dufour G, Augustin P, Lesèche G, Castier Y. Aortic rupture due to pneumococcal infection in aortoiliac stents. *J Vasc Surg* 2011;**53**:1711–1713.