

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

trans-Dibromidotetrakis(pyridine- κ N)-ruthenium(II)

 Xiu-Li Wu,^a Ru-Fei Ye,^b Ai-Quan Jia,^b Qun Chen^a and Qian-Feng Zhang^{b,a*}

^aDepartment of Applied Chemistry, School of Petrochemical Engineering, Changzhou University, Jiangsu 213164, People's Republic of China, and ^bInstitute of Molecular Engineering and Applied Chemistry, Anhui University of Technology, Ma'anshan, Anhui 243002, People's Republic of China

Correspondence e-mail: zhangqf@ahut.edu.cn


Received 25 December 2012; accepted 9 January 2013

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; R factor = 0.025; wR factor = 0.069; data-to-parameter ratio = 19.4.

The title complex, $[\text{RuBr}_2(\text{C}_5\text{H}_5\text{N})_4]$, contains two independent complex molecules in each of which the Ru^{II} atom is located on a site of 222 symmetry and has a distorted octahedral coordination geometry with four pyridine N atoms and two Br atoms. The Br atoms are *trans*-disposed as a result of symmetry.

Related literature

For background to ruthenium complexes: see: Pagliaro *et al.* (2005); van Rijt & Sadler (2009); Wu *et al.* (2009); Zhang *et al.* (2005). For related structures, see: Mirza *et al.* (2003); Wong & Lau (1994); Zhang *et al.* (2006). For a description of the Cambridge Structural Database, see: Allen (2002).



Experimental

Crystal data

 $[\text{RuBr}_2(\text{C}_5\text{H}_5\text{N})_4]$
 $M_r = 577.29$

 Orthorhombic, $Fddd$
 $a = 16.830$ (4) Å

 $b = 22.032$ (5) Å

 $c = 23.221$ (5) Å

 $V = 8610$ (3) Å³
 $Z = 16$

 Mo $K\alpha$ radiation

 $\mu = 4.45$ mm⁻¹
 $T = 296$ K

 $0.22 \times 0.18 \times 0.13$ mm

Data collection

Bruker APEXII CCD diffractometer

Absorption correction: multi-scan (SADABS; Sheldrick, 1996)

 $T_{\text{min}} = 0.441$, $T_{\text{max}} = 0.595$

13382 measured reflections

2430 independent reflections

 1631 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.034$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.025$
 $wR(F^2) = 0.069$
 $S = 1.04$

2430 reflections

125 parameters

H-atom parameters constrained

 $\Delta\rho_{\text{max}} = 0.58$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.34$ e Å⁻³
Table 1

Selected bond lengths (Å).

Ru1—N1	2.086 (2)	Ru2—N2	2.083 (2)
Ru1—Br1	2.5439 (7)	Ru2—Br2	2.5378 (7)

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

This project was supported by the Natural Science Foundation of China (grant Nos. 20771003 and 21201003).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HY2613).

References

- Allen, F. H. (2002). *Acta Cryst.* **B58**, 380–388.
- Bruker (2007). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Mirza, H. A., Farah, A. A., Stynes, D. V. & Lever, A. B. P. (2003). *Acta Cryst.* **E59**, m679–m680.
- Pagliaro, M., Campestrini, S. & Ciriminna, R. (2005). *Chem. Soc. Rev.* **34**, 837–845.
- Rijt, S. H. van & Sadler, P. J. (2009). *Drug Discov. Today*, **14**, 1089–1097.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Wong, W.-T. & Lau, T.-C. (1994). *Acta Cryst.* **C50**, 1406–1407.
- Wu, F. H., Duan, T., Lu, L., Zhang, Q. F. & Leung, W. H. (2009). *J. Organomet. Chem.* **694**, 3844–3851.
- Zhang, Q. F., Cheung, K. M., Williams, I. D. & Leung, W. H. (2005). *Eur. J. Inorg. Chem.* pp. 4780–4787.
- Zhang, L.-Y., Zhu, Y.-M., Shi, L.-X. & Chen, Z.-N. (2006). *Chin. J. Inorg. Chem.* **22**, 1453–1466.

supplementary materials

Acta Cryst. (2013). E69, m105 [doi:10.1107/S1600536813000871]

trans*-Dibromidotetrakis(pyridine- κ N)ruthenium(II)*Xiu-Li Wu, Ru-Fei Ye, Ai-Quan Jia, Qun Chen and Qian-Feng Zhang****Comment**

Coordination chemistry of ruthenium complexes has been studied in last few decades because of their versatile and diverse applications in molecular catalysis (Pagliaro *et al.*, 2005) and bioinorganic chemistry (van Rijt & Sadler, 2009). As part of our long-standing interest in the ruthenium complexes with σ -donor ligands such as thiolate, pyridine and phosphine (Zhang *et al.*, 2005), we have investigated the reactivity of the starting ruthenium compounds such as $\text{RuCl}_2(\text{PPh}_3)_3$, $\text{RuHCl}(\text{CO})(\text{PPh}_3)_3$ and $\text{RuCl}_2(\text{dmsO})_4$ (dmsO = dimethyl sulfoxide) with mono-, bi- and poly-dentate ligands (Wu *et al.*, 2009). Here we report the crystal structure of the mononuclear ruthenium(II) complex.

In the title complex, there are two independent complex molecules with a perpendicular arrangement. Each Ru^{II} atom is located on a 222 symmetry. No significant differences in bonding parameters between these two molecules are found. One of the molecular structures is depicted in Fig. 1. The coordination geometry of the Ru^{II} atom is octahedral with four pyridine N atoms and two Br atoms. The Ru—N bond lengths (Table 1) are in the range of those found in related structures of ruthenium(II) complexes retrieved from the Cambridge Structural Database (Allen, 2002). The Ru—Br bond lengths are comparable to those reported in other ruthenium(II)-bromide complexes such as $[\text{Ru}_2\text{Br}_2(\text{pz})(\text{py})_8][\text{PF}_6]_2 \cdot 2\text{DMF}$ (pz = pyrazine, py = pyridine) [av. 2.5524 (4) Å] (Mirza *et al.*, 2003) and *trans*- $[\text{RuBr}(\text{py})_4\text{C}(\text{CN})_3]$ [2.5453 (12) Å] (Zhang *et al.*, 2006). Two Br atoms are *trans* disposed as indicated by the Br—Ru—Br bond angle of 180° , as a result of symmetry requirements. Similar case was found in analogous complex *trans*- $[\text{RuCl}_2(\text{py})_4]$ (Wong & Lau, 1994).

Experimental


To a THF solution (10 ml) of $\text{RuCl}_2(\text{DMSO})_4$ (97 mg, 0.2 mmol) was added pyridine (63 mg, 0.8 mmol) and Br_2 (32 mg, 0.2 mmol) under a nitrogen atmosphere. The reaction mixture was refluxed for 2 h, developing red. The solvent was evaporated in vacuo and the residue was washed with hexane. Recrystallization from CH_2Cl_2 /hexane afforded red crystals of the title complex within two days (yield: 75 mg, 65 % based on Ru). Analysis, calculated for $\text{C}_{20}\text{H}_{20}\text{Br}_2\text{N}_4\text{Ru}$: C 41.61, H 3.49, N 9.70%; found: C 41.53, H 3.44, N 9.63%.

Refinement


H atoms were placed in geometrically idealized positions and refined as riding atoms, with C—H = 0.93 Å and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.

Computing details

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT* (Bruker, 2007); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008).

**Figure 1**

The molecular structure of the title compound, showing one of the two independent molecules. Displacement ellipsoids are drawn at the 50% probability level. [Symmetry codes: (A) $x, 1/4-y, 1/4-z$; (B) $1/4-x, y, 1/4-z$; (C) $1/4-x, 1/4-y, z$.]

**Figure 2**

Packing diagram of the title compound in a unit cell, viewed along the *c* axis.

***trans*-Dibromidotetrakis(pyridine- κ N)ruthenium(II)**

Crystal data

[RuBr₂(C₅H₅N)₄]

$M_r = 577.29$

Orthorhombic, *Fddd*

Hall symbol: -F 2uv 2vw

$a = 16.830$ (4) Å

$b = 22.032$ (5) Å

$c = 23.221$ (5) Å

$V = 8610$ (3) Å³

$Z = 16$

$F(000) = 4512$

$D_x = 1.781$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 2149 reflections

$\theta = 2.2$ – 26.4°

$\mu = 4.45$ mm⁻¹

$T = 296$ K

Block, red

$0.22 \times 0.18 \times 0.13$ mm

Data collection

Bruker APEXII CCD

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

φ and ω scans

Absorption correction: multi-scan

(*SADABS*; Sheldrick, 1996)

$T_{\min} = 0.441$, $T_{\max} = 0.595$

13382 measured reflections

2430 independent reflections

1631 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.034$

$\theta_{\text{max}} = 27.5^\circ$, $\theta_{\text{min}} = 3.0^\circ$

$h = -21 \rightarrow 21$

$k = -28 \rightarrow 28$

$l = -29 \rightarrow 26$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.025$

$wR(F^2) = 0.069$

$S = 1.04$

2430 reflections

125 parameters

0 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0292P)^2 + 11.6805P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.58 \text{ e } \text{\AA}^{-3}$

$\Delta\rho_{\min} = -0.34 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F , and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Ru1	0.1250	0.1250	0.1250	0.03562 (11)
Ru2	0.6250	0.1250	0.1250	0.03634 (11)
Br1	0.1250	0.009537 (19)	0.1250	0.05758 (14)
Br2	0.6250	0.1250	0.015713 (18)	0.06143 (15)
N1	0.03782 (12)	0.12460 (9)	0.18885 (9)	0.0409 (5)
N2	0.71241 (13)	0.19195 (10)	0.12552 (9)	0.0428 (5)
C1	-0.02375 (16)	0.08600 (14)	0.18763 (12)	0.0513 (7)
H1	-0.0280	0.0594	0.1567	0.062*
C2	-0.08052 (18)	0.08388 (16)	0.22953 (14)	0.0643 (9)
H2	-0.1223	0.0564	0.2268	0.077*
C3	-0.0757 (2)	0.12230 (16)	0.27562 (15)	0.0664 (9)
H3	-0.1137	0.1216	0.3047	0.080*
C4	-0.01284 (18)	0.16198 (15)	0.27768 (13)	0.0565 (8)
H4	-0.0077	0.1887	0.3084	0.068*
C5	0.04201 (16)	0.16198 (13)	0.23442 (11)	0.0463 (6)
H5	0.0842	0.1891	0.2366	0.056*
C6	0.77437 (16)	0.19016 (13)	0.16161 (13)	0.0513 (7)
H6	0.7783	0.1577	0.1870	0.062*
C7	0.83202 (18)	0.23371 (16)	0.16297 (15)	0.0655 (9)
H7	0.8740	0.2304	0.1888	0.079*
C8	0.8280 (2)	0.28178 (15)	0.12661 (16)	0.0721 (10)
H8	0.8670	0.3117	0.1269	0.086*
C9	0.76494 (19)	0.28493 (15)	0.08948 (16)	0.0656 (9)
H9	0.7601	0.3175	0.0642	0.079*
C10	0.70898 (16)	0.23981 (13)	0.08982 (13)	0.0509 (7)
H10	0.6667	0.2424	0.0642	0.061*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Ru1	0.0287 (2)	0.0399 (2)	0.0383 (2)	0.000	0.000	0.000
Ru2	0.0285 (2)	0.0431 (2)	0.0374 (2)	0.000	0.000	0.000
Br1	0.0600 (3)	0.0459 (2)	0.0668 (3)	0.000	-0.0158 (2)	0.000
Br2	0.0616 (3)	0.0806 (3)	0.0421 (2)	-0.0070 (2)	0.000	0.000
N1	0.0331 (11)	0.0465 (12)	0.0430 (12)	-0.0042 (10)	-0.0006 (9)	0.0007 (10)
N2	0.0326 (11)	0.0461 (12)	0.0497 (13)	0.0005 (9)	-0.0006 (10)	0.0022 (11)
C1	0.0398 (15)	0.0614 (18)	0.0526 (17)	-0.0119 (14)	-0.0014 (13)	0.0002 (14)
C2	0.0449 (18)	0.081 (2)	0.067 (2)	-0.0174 (16)	0.0047 (15)	0.0060 (18)
C3	0.0495 (18)	0.091 (2)	0.0590 (19)	-0.0018 (18)	0.0172 (15)	0.0106 (18)
C4	0.0482 (17)	0.072 (2)	0.0490 (17)	0.0020 (15)	0.0079 (13)	-0.0041 (15)
C5	0.0392 (15)	0.0486 (16)	0.0512 (17)	-0.0019 (12)	0.0030 (12)	-0.0028 (13)
C6	0.0382 (15)	0.0530 (17)	0.0627 (19)	0.0023 (13)	-0.0086 (14)	-0.0006 (14)
C7	0.0407 (17)	0.068 (2)	0.088 (2)	-0.0039 (15)	-0.0146 (17)	-0.0092 (18)
C8	0.0504 (19)	0.053 (2)	0.112 (3)	-0.0150 (15)	-0.001 (2)	-0.004 (2)
C9	0.0528 (19)	0.056 (2)	0.088 (2)	-0.0023 (15)	0.0069 (18)	0.0133 (18)
C10	0.0380 (15)	0.0549 (18)	0.0597 (18)	-0.0009 (13)	0.0018 (13)	0.0077 (14)

Geometric parameters (\AA , $^\circ$)

Ru1—N1 ⁱ	2.086 (2)	C1—H1	0.9300
Ru1—N1 ⁱⁱ	2.086 (2)	C2—C3	1.367 (5)
Ru1—N1	2.086 (2)	C2—H2	0.9300
Ru1—N1 ⁱⁱⁱ	2.086 (2)	C3—C4	1.373 (4)
Ru1—Br1	2.5439 (7)	C3—H3	0.9300
Ru1—Br1 ⁱⁱ	2.5439 (7)	C4—C5	1.364 (4)
Ru2—N2 ^{iv}	2.083 (2)	C4—H4	0.9300
Ru2—N2	2.083 (2)	C5—H5	0.9300
Ru2—N2 ⁱ	2.083 (2)	C6—C7	1.365 (4)
Ru2—N2 ^v	2.083 (2)	C6—H6	0.9300
Ru2—Br2	2.5378 (7)	C7—C8	1.356 (5)
Ru2—Br2 ^v	2.5378 (7)	C7—H7	0.9300
N1—C1	1.341 (3)	C8—C9	1.370 (5)
N1—C5	1.343 (3)	C8—H8	0.9300
N2—C6	1.339 (3)	C9—C10	1.369 (4)
N2—C10	1.342 (3)	C9—H9	0.9300
C1—C2	1.365 (4)	C10—H10	0.9300
N1 ⁱ —Ru1—N1 ⁱⁱ	179.52 (12)	C6—N2—C10	116.3 (2)
N1 ⁱ —Ru1—N1	90.60 (12)	C6—N2—Ru2	122.20 (18)
N1 ⁱⁱ —Ru1—N1	89.40 (12)	C10—N2—Ru2	121.48 (18)
N1 ⁱ —Ru1—N1 ⁱⁱⁱ	89.40 (12)	N1—C1—C2	123.2 (3)
N1 ⁱⁱ —Ru1—N1 ⁱⁱⁱ	90.60 (12)	N1—C1—H1	118.4
N1—Ru1—N1 ⁱⁱⁱ	179.52 (12)	C2—C1—H1	118.4
N1 ⁱ —Ru1—Br1	90.24 (6)	C1—C2—C3	119.7 (3)
N1 ⁱⁱ —Ru1—Br1	90.24 (6)	C1—C2—H2	120.2
N1—Ru1—Br1	89.76 (6)	C3—C2—H2	120.2
N1 ⁱⁱⁱ —Ru1—Br1	89.76 (6)	C2—C3—C4	117.9 (3)

N1 ⁱ —Ru1—Br1 ⁱⁱ	89.76 (6)	C2—C3—H3	121.1
N1 ⁱⁱ —Ru1—Br1 ⁱⁱ	89.76 (6)	C4—C3—H3	121.1
N1—Ru1—Br1 ⁱⁱ	90.24 (6)	C5—C4—C3	119.7 (3)
N1 ⁱⁱⁱ —Ru1—Br1 ⁱⁱ	90.24 (6)	C5—C4—H4	120.2
Br1—Ru1—Br1 ⁱⁱ	180.0	C3—C4—H4	120.2
N2 ^{iv} —Ru2—N2	179.34 (11)	N1—C5—C4	123.0 (3)
N2 ^{iv} —Ru2—N2 ⁱ	89.84 (12)	N1—C5—H5	118.5
N2—Ru2—N2 ⁱ	90.16 (12)	C4—C5—H5	118.5
N2 ^{iv} —Ru2—N2 ^v	90.16 (12)	N2—C6—C7	123.2 (3)
N2—Ru2—N2 ^v	89.84 (12)	N2—C6—H6	118.4
N2 ⁱ —Ru2—N2 ^v	179.34 (11)	C7—C6—H6	118.4
N2 ^{iv} —Ru2—Br2	90.33 (6)	C8—C7—C6	120.0 (3)
N2—Ru2—Br2	90.33 (6)	C8—C7—H7	120.0
N2 ⁱ —Ru2—Br2	89.67 (6)	C6—C7—H7	120.0
N2 ^v —Ru2—Br2	89.67 (6)	C7—C8—C9	118.0 (3)
N2 ^{iv} —Ru2—Br2 ^v	89.67 (6)	C7—C8—H8	121.0
N2—Ru2—Br2 ^v	89.67 (6)	C9—C8—H8	121.0
N2 ⁱ —Ru2—Br2 ^v	90.33 (6)	C10—C9—C8	119.5 (3)
N2 ^v —Ru2—Br2 ^v	90.33 (6)	C10—C9—H9	120.2
Br2—Ru2—Br2 ^v	180.0	C8—C9—H9	120.2
C1—N1—C5	116.5 (2)	N2—C10—C9	123.0 (3)
C1—N1—Ru1	122.09 (18)	N2—C10—H10	118.5
C5—N1—Ru1	121.38 (17)	C9—C10—H10	118.5

Symmetry codes: (i) $x, -y+1/4, -z+1/4$; (ii) $-x+1/4, -y+1/4, z$; (iii) $-x+1/4, y, -z+1/4$; (iv) $-x+5/4, -y+1/4, z$; (v) $-x+5/4, y, -z+1/4$.