

CASE REPORT

Airway management and quadratus lumborum block for inguinal hernia repair in context of chromosome 4q deletion: A case report

Katie Lovell¹  | Michael C. Larkins¹  | Melisa Pasli¹ | Adrienne L. Hutton^{1,2}

¹East Carolina University Brody School of Medicine, Greenville, North Carolina, USA

²East Carolina Anesthesia Associates, Greenville, North Carolina, USA

Correspondence

Adrienne L. Hutton, East Carolina Anesthesia Associates, 2430 Emerald Pl #103, Greenville, NC 27834, USA.
Email: asingleton@ecaa.com

Key Clinical Message

Chromosome 4q Deletion Syndrome (4QDS) is a rare chromosomal deletion that presents with various congenital defects that could affect anesthetic management. We present a patient with known 4QDS who underwent successful general anesthetic induction, tracheal intubation, and subsequently, bilateral quadratus lumborum nerve blocks, during laparoscopic inguinal hernia repair surgery.

Abstract

Chromosome 4q deletion syndrome (4QDS) is a rare chromosomal deletion with the potential for congenital defects that could affect the anesthetic management of affected patients, specifically with inconsistent reports of possible laryngeal narrowing which could precipitate intubation difficulties. We present a report of a patient with known 4QDS who underwent successful general anesthetic induction and tracheal intubation, followed by subsequent bilateral quadratus lumborum nerve blocks, in the context of laparoscopic inguinal hernia repair surgery. Given the inconsistent presentation of patients with 4QDS reported in the literature, steps were taken to exercise sufficient caution concerning the overall management of this patient per the most recent guidelines for pediatric patients.

KEYWORDS

challenging pediatric airway, chromosome 4q deletion syndrome, congenital airway malformations, congenital airway malformations, quadratus lumborum block, regional anesthesia

1 | INTRODUCTION

Deletion of the 4q chromosome is a rare chromosomal deletion syndrome estimated to occur in about 1 out of every 100,000 births.¹ The effects of this deletion present with various clinical phenotypes,² such as congenital heart defects, craniofacial dysmorphic features, and

developmental delay.¹ The severity of physical characteristics is dependent on the amount and the site of chromatin that is missing. Most patients pass away before 2 years of age or, if they survive, are confronted with both social and physical barriers.³ With specific reference to anesthesia, patients with this syndrome have inconsistently been reported to have laryngeal narrowing, raising the concern

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2023 The Authors. *Clinical Case Reports* published by John Wiley & Sons Ltd.

for intubation difficulties.^{4,5} Due to the rarity of this syndrome and the myriad of clinical presentations, few guidelines outline the induction of anesthesia in a patient with this syndrome.

In this report, we describe the anesthetic management of a patient with known chromosome abnormality of 46 XY, del(4)q34.1-q35.2, and associated dysmorphic appearance, including micrognathia. Craniofacial abnormalities are of consideration in the selection of tracheal tube sizes and may result in difficult intubation. These same abnormalities can be severe, to the point in which patients require long-term tracheostomy or plastic/orthopedic surgery for jaw reconstruction.⁶ In this report, we discuss the management of anesthesia, utilization of a quadratus lumborum plexus block, and provide images of the characteristic craniofacial abnormalities observed in Chromosome 4q Deletion Syndrome (4QDS). To our knowledge, no case to date has been reported on the anesthetic management of pediatric patients with 4QDS, including nerve block procedures. Additionally, few resources in the literature exist that show clear images of the characteristic appearance of patients with 4QDS or document their intubation.

2 | CASE REPORT

A 13-month-old male, 10.8 kg, presented for anesthesia pre-procedure assessment before a scheduled laparoscopic left inguinal hernia repair under general anesthesia. The patient was born at 36 weeks via spontaneous vaginal delivery. An echocardiogram at 30 hours of life revealed a patent foramen ovale (PFO) and patent ductus arteriosus (PDA) with associated left to right shunting; these defects were later evaluated and confirmed to have closed spontaneously at 1-month follow-up. Past medical history was otherwise significant for genetic testing-confirmed chromosome 4q deletion (q34.1–35.2), constipation, arachnoid cysts, and bilateral hearing loss. This patient had no history of previous general anesthesia but had an uneventful sedation with propofol for MRI at an outside institution. At that time, the patient had a size 4 nasopharyngeal airway successfully placed in one attempt with no complications, and the patient was considered ASA II.

On physical exam, the patient had a moderately dysmorphic appearance, with slightly slanted palpebral fissures, flattened nasal root, low-set ears, left eyelid ptosis, high-arched palate with a triangular mouth and micrognathia (see Figure 1A,B). The patient's cardiovascular exam was unremarkable. The left inguinal hernia defect was easily reducible, and no signs of muscle weakness or lethargy were appreciated.

All other vital signs and electrocardiogram were unremarkable, and their airway was evaluated as not difficult.



FIGURE 1 (A) Frontal view of our patient with Chromosome 4q Deletion Syndrome; note the characteristic slightly slanted palpebral fissures, flattened nasal root, low-set ears, and triangular mouth. (B) Side profile of our patient, demonstrating the micrognathia associated with this syndrome. (C) View of the patient's larynx via C-MAC[®] before tracheal intubation. (D) The same view of our patient's larynx after intubation. No significant laryngeal abnormalities were noted during intubation.

However, given the patient's history of 4QDS and micrognathia on examination, consideration was given for the patient to potentially require a smaller-than-expected tracheal tube, and thus, video laryngoscopy via C-MAC[®] was chosen for intubation. The patient's mother was informed of this plan and written consent was obtained. Shortly after the preoperative assessment, the patient was given 0.5 mg/kg of oral midazolam and 15 mg/kg of oral acetaminophen and then was brought to the operating room for general anesthesia inhalation induction followed by IV placement, which was completed with propofol dosed at 3.2 mg/kg. Tracheal intubation was accomplished in one attempt via C-MAC[®] using a size 1 Miller blade and 3.5 microcuff ETT. An intubating stylet and anterior, back, upward, and right lateral pressure (BURP) were utilized to aid intubation. The laryngeal view was deemed Cormack-Lehane Grade I, with full visualization of the glottis. The

view seen with the C-MAC® can be seen in [Figure 1C,D](#), before and after intubation.

Laparoscopic inguinal hernia repair was completed uneventfully. Following surgical closure, a bilateral quadratus lumborum (QL) block was placed via ultrasound while still under general anesthesia. A total of 0.8 mg/kg of ropivacaine 0.2% with 1:200,000 epinephrine was injected. Additionally, 2 mL of 0.25% bupivacaine was injected by the surgical team into the umbilical incision.

After the QL nerve block, the patient was successfully extubated and transferred to the post-anesthesia care unit. Following surgery, the patient did not require any additional medications for pain control and recovered uneventfully with no pain reported or observed. The patient was discharged later the same day, and both three-day and one-month follow-up appointments were unremarkable; no complaints of pain were noted from the patient's mother beyond mild soreness on Day 3.

3 | DISCUSSION

The deletion of the 4q chromosome is a rare deletion syndrome with only around 200 cases reported in the literature.² Based on the genetic analysis completed by Xu et al., the congenital heart defects associated with this syndrome have been localized to the 4q32.2–q34.3 region.² Our patient demonstrated deletions of the terminal end of this region (q34.1–q35.2), which may have precipitated their PFO and PDA. The heart defects were visualized via echocardiogram on Day Two of life and were expected to likely close spontaneously without intervention. Upon preoperative physical exam by the anesthesia team, the patient's cardiovascular exam was grossly normal. However, close monitoring of their cardiovascular status was implemented due to their history of 4QDS. The patient's status was well-maintained throughout the procedure.

Our patient also presented with dysmorphic features associated with 4QDS such as flattening of the midface and nasal root, upturned nostrils, a high-arched palate, triangular mouth, and posteriorly rotated, low-set ears—all features that could be concerning for intubation. Despite these dysmorphic features, there was no substantial concern for airway complications based on the exam, and the patient was assigned an ASA classification of 2. However, taking into consideration prior reported anomalies for 4QDS, preparation for an unexpectedly difficult airway was planned. This included the availability of smaller endotracheal tubes and video laryngoscopy equipment.^{4,5}

Per the 2022 American Society of Anesthesiologists (ASA), pediatric patients perceived as difficult or potentially difficult can be managed using an airway management algorithm with an emphasis on team-based care.⁶

This care includes the use of video laryngoscopy to maximize visualization of a patient's airway and minimize airway intubation attempts. As such, video laryngoscopy via C-MAC® was chosen for intubation with success in one attempt. Dearlove and Sharples present a case report that highlighted a 4QDS patient whose laryngoscopic imaging demonstrated a small larynx with only the posterior arytenoids visualized.⁴ While our patient presented with normal laryngeal anatomy as seen in [Figures 1C,D](#), this report demonstrates the utility of video laryngoscopy for intubation in patients with 4QDS. There was potential for difficulty intubating our patient, and the video laryngoscope allowed the anesthesiologist to assess this in real-time.

In addition to airway management, the anesthesia team also administered a QL block in the setting of laparoscopic inguinal hernia repair as a component of a multimodal anesthetic technique for pain management. A bilateral block was utilized as surgical exploration for occult hernia was carried out bilaterally. Advances in pediatric regional anesthesia have evolved as the prevalence of ambulatory pediatric surgery has also grown. The utilization of QL blocks has consistently been shown to provide a longer analgesic duration when compared to other techniques for inguinal hernia repairs.⁷ With the association of opioid use and perioperative complications, regional anesthesia has proven to be a safe and effective form of pain control in the ambulatory pediatric surgery setting, while also minimizing the use of narcotics.⁸ When used in conjunction with Tylenol, as in our patient, this constitutes a multimodal anesthetic technique, an approach with proven benefit in ambulatory pediatric patients.⁹ While there is a documented report of pain insensitivity in 4QDS,¹⁰ our patients had previously demonstrated a response to painful stimuli in the setting of incarcerated inguinal hernia, which warranted the adoption of the regional anesthetic technique. The QL nerve block has demonstrated efficacy in multiple randomized control trials, presented by Zhao et al. in a meta-analysis of 346 patients, especially when administered under general anesthesia and ultrasound guidance.¹¹ The uncomplicated nature of our patient's nerve block, in addition to the lack of laryngeal narrowing seen during video-assisted intubation and the lack of persistent cardiac defects, suggest the variance in presentation/phenotype present in patients with 4QDS. This is also in line with the inconsistent reporting of both pain insensitivity and laryngeal narrowing reported in the literature.^{2,4,5,10} Our approach, with emphasis on the ASA 2022 Guidelines, maximized patient safety by emphasizing teamwork, communication, and caution. Some limitations of this approach, though, include the resources and time required to successfully carry out this plan. Smaller institutions may not have access to pediatric video laryngoscopy or providers trained in the use of such equipment.

4 | CONCLUSION

This case demonstrates successful intubation in a pediatric patient with 4QDS despite dysmorphic features and the potential for both laryngeal narrowing and pain insensitivity associated with the patient's genetic condition. While this patient was not expected to be a difficult airway upon preoperative physical examination, the dysmorphic facial features were a cause of concern and extra care was taken in patient intubation. In addition, a bilateral QL nerve block and adjunctive Tylenol, constituting a multimodal anesthesia technique, were utilized in our patient for postoperative pain control. As demonstrated by this case, successful intubation and the use of both general and regional anesthesia can be safely accomplished in patients with 4QDS, though sufficient caution should be taken when assessing and treating this patient population. We suggest other patients with 4QDS be treated with similar caution, given the potential for adverse events related to inconsistently reported characteristics of this genetic syndrome.

AUTHOR CONTRIBUTIONS

Katie Lovell: Investigation; project administration; visualization; writing – original draft; writing – review and editing. **Michael C. Larkins:** Visualization; writing – original draft; writing – review and editing. **Melisa Pasli:** Writing – original draft; writing – review and editing. **Adrienne L. Hutton:** Conceptualization; supervision.

ACKNOWLEDGMENTS

None.

FUNDING INFORMATION

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

CONFLICT OF INTEREST STATEMENT

No conflict of interest is declared by the authors.

DATA AVAILABILITY STATEMENT

Data sharing is not applicable to this article as no new data were created or analyzed in this study.

ETHICS STATEMENT

The authors declare that this manuscript is not published or under consideration in any other journals.

CONSENT

Published with the written consent of the patient's mother.

ORCID

Katie Lovell  <https://orcid.org/0009-0005-0098-8455>

Michael C. Larkins  <https://orcid.org/0000-0002-2322-8345>

REFERENCES

1. Strehle E-M, Yu L, Rosenfeld JA, et al. Genotype-phenotype analysis of 4q deletion syndrome: proposal of a critical region. *Am J Med Genet A*. 2012;158A(9):2139-2151.
2. Xu W, Ahmad A, Dagenais S, Iyer RK, Innis JW. Chromosome 4q deletion syndrome: narrowing the cardiovascular critical region to 4q32.2–q34.3. *Am J Med Genet A*. 2012;158A(3):635-640.
3. Kuldeep CM, Khare AK, Garg A, Mittal A, Gupta L. Terminal 4q deletion syndrome. *Indian J Dermatol*. 2012;57(3):222-224.
4. Dearlove OR, Sharples A. Anaesthesia and partial deletion of chromosome 4. *Anaesthesia*. 1996;51(2):206.
5. Markiewicz MR, Verschuereen D, Assael LA. Chromosome 4q deletion syndrome: craniofacial characteristics associated with monosomy of the long arm of chromosome 4q. *Cleft Palate Craniofac J*. 2010;47(5):518-522.
6. Apfelbaum JL, Hagberg CA, Connis RT, et al. 2022 American Society of Anesthesiologists Practice Guidelines for Management of the Difficult Airway. *Anesthesiology*. 2022;136(1):31-81.
7. Hung TY, Bai GH, Tsai MC, Lin YC. Analgesic effects of regional analgesic techniques in pediatric inguinal surgeries: a systematic review and network meta-analysis of randomized controlled trials. *Anesth Analg*. 2022. doi:10.1213/ANE.0000000000006341. Epub ahead of print.
8. Stein ALS, Baumgard D, Del Rio I, Tutiven JL. Updates in pediatric regional anesthesia and its role in the treatment of acute pain in the ambulatory setting. *Curr Pain Headache Rep*. 2017;21(2):11.
9. Oliver J-A, Oliver L-A, Aggarwal N, et al. Ambulatory pain management in the pediatric patient population. *Curr Pain Headache Rep*. 2022;26:15-23.
10. Cascella M, Muzio MR. Pain insensitivity in a child with a de novo interstitial deletion of the long arm of the chromosome 4: case report. *Rev Chil Pediatr*. 2017;88(3):411-416.
11. Zhao W-L, Li S-D, Wu B, Zhou Z-F. Quadratus lumborum block is an effective postoperative analgesic technique in pediatric patients undergoing lower abdominal surgery: a meta-analysis. *Pain Physician*. 2021;24(5):E555-E563.

How to cite this article: Lovell K, Larkins MC, Pasli M, Hutton AL. Airway management and quadratus lumborum block for inguinal hernia repair in context of chromosome 4q deletion: A case report. *Clin Case Rep*. 2023;11:e7765. doi:10.1002/ccr3.7765