

A Prospective Evaluation of Point of Care Ultrasound Teaching in Switzerland

Connor Byrne¹, Nico Kahl¹, Brian Knight¹, Monica Lee¹, Stephanie Morley¹, Shadi Lahham^{2*}, Roland Bingisser³, Maxwell Thompson², Inna Shniter², Victoria Valdes², John C. Fox²

¹School of Medicine, University of California Irvine, Irvine, California, USA, ²Department of Emergency Medicine, University of California Irvine, Irvine, California, USA, ³University Hospital Basel, Basel, Switzerland

Abstract

Context: As the utility of point-of-care ultrasound (POCUS) continues to expand in the medical field, there is a need for effective educational methods. In Switzerland, medical education follows the European model and lasts 6 years, focusing on preclinical training during the first 2 years. No previous studies have evaluated the optimal time for teaching ultrasound in European medical education. **Aims:** The aim of this study is to provide ultrasound training to medical students in Switzerland at varying times during their clinical training to determine if the level of training plays a role in their ability to comprehend and to apply basic POCUS skills. **Methods:** We performed an observational study utilizing a convenience sample of Swiss medical students between July 11, 2016 and August 6, 2016. They were taught a 2-day POCUS course by five American-trained 1st-year medical students. Following this course, students were evaluated with written and clinical examination. **Results:** 100 Swiss medical students were enrolled in the study. A total of 59 of these students were early clinical students, and 41 students were late clinical students. A two-tailed *t*-test was performed and demonstrated that the late clinical students performed better than the early clinical students on the written assessment; however, no difference was found in clinical skill. **Conclusion:** Our data suggest that Swiss medical students can learn and perform POCUS after a 2-day instructional taught by trained 1st-year American medical students. No difference was found between students in early clinical training and late clinical training for the ability to perform POCUS.

Keywords: Medical education, Switzerland, ultrasound teaching

INTRODUCTION

Point-of-care ultrasound (POCUS) has emerged as a cost-effective, minimally invasive imaging modality. It has a wide array of practical uses across a breadth of medical specialties, and it has been suggested that ultrasound is included in all standard physical examinations to aid physicians in their diagnoses.^[1-3] In addition, including ultrasound in medical education has been shown to later boost physicians' confidence in performing bedside examinations.^[4] As the utility of POCUS continues to expand in the medical field, there is an increasing need for effective educational methods to teach medical students the necessary skill set to master POCUS.

Previous studies have shown variable success and implementation in teaching curriculums.^[5-7] Bahner *et al.*

demonstrated that the integration of ultrasound into curricula is highly variable and is in need of standardization.^[8] A comprehensive study of ultrasound education at Wayne State University has shown that medical students had positive experiences and better outcomes with the inclusion of ultrasound in their medical education.^[9] In addition, a 2010 study conducted by surgeons and medical researchers in the United Kingdom demonstrated that the focused assessment with sonography in trauma (FAST) scan can be effectively taught to inexperienced medical students with significant success.^[10] Although the benefit of including ultrasound education in the medical school curriculum has been well established, there

Address for correspondence: Dr. Shadi Lahham,
333 The City Boulevard West, Suite 640, Rt 128-01, Orange, CA, USA.
E-mail: slahham@uci.edu

Received: 05-05-2018 Accepted: 28-09-2018 Available Online: 26-02-2019

Access this article online

Quick Response Code:



Website:
www.jmuonline.org

DOI:
10.4103/JMU.JMU_57_18

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

How to cite this article: Byrne C, Kahl N, Knight B, Lee M, Morley S, Lahham S, *et al.* A prospective evaluation of point of care ultrasound teaching in Switzerland. *J Med Ultrasound* 2019;27:92-6.

has been limited research as to the most effective time to teach POCUS during medical students' training.^[11] There have been various curriculum models proposed; however, the various cost and benefits of these differing models have not fully been assessed.^[12] Furthermore, the model of peer-to-peer teaching has not been fully evaluated.

In Switzerland, medical education follows the European model and lasts 6 years, focusing on preclinical training during the first 2 years, and focusing on clinical training thereafter. Currently, medical education in Switzerland is devoid of any ultrasound training despite recommendations by the European Federation of Societies for Ultrasound in Medicine and Biology.^[13] This environment allows comparing ultrasound education between Swiss medical students given ultrasound training in the early clinical training period (year 3/4) as compared to during the late clinical training period (year 5/6). The aim of this study is to determine if 1st-year American medical students with minimal ultrasound training can teach ultrasound to medical students in Switzerland at varying times during their clinical training. A secondary objective is to determine if the level of training plays a role in their ability to comprehend and apply basic POCUS skills.

METHODS

Study design

We performed an observational study utilizing a convenience sample of volunteer Swiss medical students between July 11, 2016 and August 6, 2016. All students provided written consent for the study. No financial compensation was provided for participation. The study was approved by the site Institutional Review Board. The study took place at the University of Basel Medical School at the hospital's training site.

Study protocol

We recruited five American-trained 1st-year medical students who consented for study enrollment. All students had completed an entire 1st year of medical education which included human physiology, human anatomy, and basic ultrasound. Before initiation of study enrollment, each student needed to demonstrate the ability to obtain the four required cardiac views including parasternal long-axis view, parasternal short-axis view, apical four-chamber view, and subxiphoid view. Students were required to pass a certification exam given by the institution's POCUS director. Students also needed to be able to identify anatomy for the remainder of the FAST scan including the hepatorenal recess, splenorenal recess, and bladder views.^[14] We asked these instructors to demonstrate an ability to identify all important structures of the FAST scan to be considered qualified to teach POCUS. We performed both instruction and image acquisition using four SonoSite NanoMaxx (SonoSite Inc, Bothell WA) ultrasound machines. Each ultrasound machine was equipped with a phased array probe operating at a frequency of 1–5 MHz.

Intervention

We gave Swiss medical students at various levels of training the opportunity to voluntarily enroll in the study as an elective learning course in addition to their curriculum. All Swiss medical students were offered the opportunity to enroll in the study through an e-mail to the entire student body. The students were separated into two groups. The first group consisted of 3rd and 4th-year Swiss medical students. These students had completed basic science courses as well as basic anatomy but had minimal clinical exposure. The second group of students included 5th- and 6th-year medical students. These students had completed all basic science requirements and were involved in clinical elective rotations. All students in this category had completed internal medicine, general surgery, pediatrics, and emergency medicine.

Students who enrolled in the study were required to take part in a 2-day ultrasound training workshop. We held a total of four ultrasound training sessions. Each training session began with a written pretest, which was then followed by a 1-hr classroom lecture, designed to prepare the students for their hands-on training sessions. A board-certified Swiss Emergency Physician with ultrasound training was present to ensure the accuracy of training and answer any questions. For the hands-on training, we divided the students into groups of four and completed a hands-on pre-test to obtain all components of a FAST examination with a 5-min time limit each.

Ultrasound curriculum

The initial training session began with a 30-min instructional lecture on ultrasound technical handling (termed "knobology"). Next, students underwent a 60-min instructional lecture on the FAST examination, basic cardiac anatomy. Examples of normal anatomy and pathology were shown in lecture. Following didactic instructions, all participants practiced hands-on FAST ultrasound techniques under the direct supervision of one of the medical student instructors for 90-min duration. Instruction began with hepatorenal views followed by splenorenal, cardiac, and bladder views. The remainder of the instructional time was spent practicing POCUS techniques which included image acquisition and image optimization. Techniques included modifications in gain, depth, and image quality. Following instruction, the final hour and a half of the 2-day workshop were spent performing both a written and hands-on posttest. All scans were performed on healthy volunteers. All students were given the same written assessment of general ultrasound use and instructed to perform a practical hands-on demonstration of FAST and cardiac ultrasound examinations before and after the teaching session. Assessment was made in real time, and the students' scores were recorded on standardized data collection sheets. Images were saved on the portable ultrasound machines, and image accuracy was confirmed by the blinded board-certified Swiss Emergency Physician with ultrasound training. Competency was defined as scoring >70% on the examinations.

Data collection and statistics

All POCUS images and videos were recorded to evaluate the students' ultrasound skills. These images were then reviewed and graded based on appropriate probe selection, orientation, image quality, and accuracy of image interpretation. After grading images and compiling scores, images and videos were erased. Researchers stratified the data from the pre- and post-tests based on the years of medical training (Group 1: students in years 3 and 4; Group 2: students in year 5 and 6). Researchers used this information to determine the averages for the pre- and post-test scores of the groups to determine the effectiveness of the workshop. The improvement in mean written and practical examination scores was analyzed with a two-tailed *t*-test for comparison of pre- and post-clinical year students using a *P* = 0.025 to determine significance. The posttest means of all the groups were then compared to determine which group was able to most effectively learn from the workshop. A one-way ANOVA test was used for this portion of the analysis, using a *P* = 0.05 to determine significance. To determine the students' scores during the pre- and post-test assessments, a scoring rubric was used to quantify performance. The scoring rubric is shown in Figures 1 and 2.

RESULTS

A total of 100 Swiss medical students were enrolled in the study. 59 of these students were Group 1 (year 3/4), and 41 students were in Group 2 (year 5/6) [Table 1]. A two-tailed *t*-test was performed and demonstrated that the late clinical students (year 5/6) performed marginally better than the early clinical students (year 3/4) on the written assessment both before and after the teaching sessions (*P* < 0.025). This was the

Table 1: Illustration of written test before 2-day point of care ultrasound course and after 2-day point of care ultrasound course for early and late clinical Swiss students

	Subjects	Written test	
		Preteaching average	Postteaching average
Late clinical	41	11.09	13.68
Early clinical	59	9.07	12.59
<i>P</i>		0.00014	0.00032

Maximum score is 15

only significant difference found in between the groups at any point in the study. Maximum score for the written test was 15.

For the hands-on practical assessment of the cardiac and FAST bedside examination, both early and late clinical groups did not show significant differences in performances before or after the teaching session [Table 2]. After the teaching session, the late clinical students scored an average of 11.74 points

Student ID: _____ Date: _____

University of California, Irvine
School of Medicine

Evaluating Effectiveness of Teaching Methods and Timing in Ultrasound Education at the University of Basel School of Medicine

Practical Exam: Cardiac Ultrasound

Each item in the numbered checklist is worth 1 point.

Step 0: Choose the correct probe.

Parasternal Long

- 1. Identify Right Ventricle
- 2. Identify Left Ventricle
- 3. Identify Mitral Valve
- 4. Identify Aortic Valve
- 5. Identify Interventricular Septum

Parasternal Short

- 6. Identify Aortic Valve
- 7. Identify Mitral Valve
- 8. Identify Papillary Muscles/Chordae Tendinae

Subxiphoid

- 9. Identify 4 chambers of heart and apex.
- 10. Identify pericardium.
- 11. Point out where fluid would be if pericardial effusion.

Apical 4 Chamber

- 12. Identify 4 Chambers
- 13. Identify Pericardium

Student Score: _____ / 13 _____

Figure 1: Practical examination: Cardiac ultrasound

Table 2: Illustration of cardiac and focused assessment with sonography in trauma ultrasound skills before and after 2 day point of care ultrasound course

Year	Subjects	Cardiac skills		FAST skills	
		Preteaching	Postteaching	Preteaching	Postteaching
Late clinical	41	0.5	11.74	1.62	9.5
Early clinical	59	0.98	11.19	1.07	9.09
<i>P</i>		0.056	0.2	0.08	0.1

Please note that maximum score is 13 for cardiac ultrasound and 14 for FAST ultrasound. FAST: Focused assessment with sonography in trauma

Student ID: _____	Date: _____
University of California, Irvine	
School of Medicine	
Evaluating Effectiveness of Teaching Methods and Timing in Ultrasound Education at the University of Basel School of Medicine	
Practical Exam: FAST Ultrasound	
Each item in the numbered checklist is worth 1 point.	
Step 0: Choose the correct probe.	
Right Upper Quadrant:	
<ul style="list-style-type: none"> • 1. Identify Liver • 2. Identify Diaphragm. • 3. Point out mirror image artifact • 4. Identify right kidney • 5. Identify Morrison's pouch/border between liver and kidney. 	
Cardiac (Subxiphoid View):	
<ul style="list-style-type: none"> • 6. Identify 4 chambers of heart and apex. • 7. Identify pericardium. • 8. Point out where fluid would be if pericardial effusion. 	
Left Upper Quadrant:	
<ul style="list-style-type: none"> • 9. Identify spleen. • 10. Identify left kidney. • 11. Point out where fluid would be if hemoperitoneum. 	
Suprapubic (Sagittal/Transverse Views):	
<ul style="list-style-type: none"> • 12. Identify bladder in sagittal view. • 13. Point out where potential free fluid would be in sagittal view. • 14. Repeat steps 12 and 13 in transverse view. 	
Student Score: _____ / 14	

Figure 2: Practical examination: Focused assessment with sonography in trauma ultrasound

(out of a maximum 13) for the cardiac examination and 9.5 (out of a maximum 14) points for the FAST examination. The early clinical students averaged 11.19 points for the cardiac examination and 9.09 points for the FAST examination after the teaching session. While both groups individually improved significantly from their pre-session baseline scores, the final scores were exceedingly similar.

DISCUSSION

The main findings of our study were that POCUS can be taught to medical students in a peer to peer fashion and that there is no difference in knowledge acquisition between

early clinical students and late clinical students. The study evaluated a large sample size of Swiss medical students with no prior formal education with English competency trained by American 1st-year medical students. While it may seem that later clinical students would perform better in the practical assessment of FAST and cardiac bedside ultrasound, we were unable to determine any significant difference between the performance of students in early or late clinical training. Both groups improved significantly, however.

Ultimately, both groups scored well enough to indicate competency in the FAST and cardiac ultrasound examinations (Group 1: FAST = 9.09/14, cardiac = 11.19/13; Group 2: FAST = 9.5/14, cardiac = 11.74/13). This suggests that students in their early clinical training can effectively learn FAST and cardiac bedside POCUS as effectively as their counterparts in late clinical training, indicating that ultrasound can be incorporated into medical education at whichever time is most convenient for each school's curriculum. Their similar increases in pre- to post-test score indicate that they acquired a comparable amount of knowledge over the 2-day workshop, and the similarity of their final scores indicate that both groups were equally able to achieve a high level of competency of the scans.

In the written assessment, we found that students in late clinical training outperformed students in the early clinical years in both the pre-and post-tests. We attributed this finding to the higher probability that students with late clinical training have additional medical education and clinical experience which gives them a larger base of knowledge to draw on for the written test, which was multiple choice. That being said, a written test is not as good an indicator of ultrasound ability as a practical one, but has the advantage of a knowledge focus.

This apparent equality in ability of early and late clinical Swiss medical students to learn POCUS could be very useful for determining at which time point to incorporate ultrasound education into American medical school curricula. The value of integrating ultrasound POCUS into undergraduate medical education is being increasingly recognized, and given the need for national guidelines in American ultrasound education, this study aims to determine when would be the most effective time to learn POCUS.^[5,6,15-17] While there are some studies that illustrate the use of peer-to-peer educational programs in medical school, there are relatively few studies on medical students teaching ultrasound to other medical students.^[18,19] Further investigation is needed into this concept. Potential future studies could include an assessment of long-term retention of the skills and knowledge obtained, to find differences between students of different levels of experience. If there is a difference, the most appropriate time point to start ultrasound education could better further characterized.

Limitations

This study has several limitations. First, there are many differences in curriculum models between Swiss and American medical schools. Swiss medical school contains pre-clinical years, then early and late medical, educational

years. Thus, this difference may affect the performance of Swiss preclinical students and make these conclusions less generalizable to American preclinical students. This model has appeared to be effective with Swiss medical students. It is unclear if similar results would be achieved in other European countries. In addition, it is unclear if there would have been better knowledge acquisition if expert sonographers were used to each ultrasound. Long-term knowledge was not formally tested in this study. Additional studies must be performed to determine the minimum amount of training required to be an instructor and the retention rate of ultrasound knowledge.

CONCLUSION

Our data suggest that Swiss medical students can learn and perform cardiac and FAST ultrasound after a 2-day instructional course taught by trained 1st-year American medical students. In addition, no difference was found between students in their early clinical training and late clinical training for the ability to perform and to interpret FAST and cardiac POCUS.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

- Moore CL, Copel JA. Point-of-care ultrasonography. *N Engl J Med* 2011;364:749-57.
- Frederiksen CA, Juhl-Olsen P, Sloth E. Advances in imaging: Ultrasound in every physician's pocket. *Expert Opin Med Diagn* 2012;6:167-70.
- Siepel T, Clifford DS, James PA, Cowan TM. The ultrasound-assisted physical examination in the periodic health evaluation of the elderly. *J Fam Pract* 2000;49:628-32.
- Fox JC, Schlang JR, Maldonado G, Lotfipour S, Clayman RV. Proactive medicine: The "UCI 30," an ultrasound-based clinical initiative from the University of California, Irvine. *Acad Med* 2014;89:984-9. doi: 10.1097/ACM.0000000000000292.
- Dinh VA, Fu JY, Lu S, Chiem A, Fox JC, Blaivas M, *et al.* Integration of ultrasound in medical education at United States medical schools: A National Survey of Directors' Experiences. *J Ultrasound Med* 2016;35:413-9.
- Dinh VA, Dukes WS, Prigge J, Avila M. Ultrasound integration in undergraduate medical education: Comparison of ultrasound proficiency between trained and untrained medical students. *J Ultrasound Med* 2015;34:1819-24.
- Day J, Davis J, Riesenber LA, Heil D, Berg K, Davis R, *et al.* Integrating sonography training into undergraduate medical education: A study of the previous exposure of one institution's incoming residents. *J Ultrasound Med* 2015;34:1253-7.
- Bahner DP, Goldman E, Way D, Royall NA, Liu YT. The state of ultrasound education in U.S. Medical schools: Results of a national survey. *Acad Med* 2014;89:1681-6.
- Rao S, van Holsbeeck L, Musial JL, Parker A, Bouffard JA, Bridge P, *et al.* A pilot study of comprehensive ultrasound education at the Wayne State University School of Medicine: A pioneer year review. *J Ultrasound Med* 2008;27:745-9.
- Gogalniceanu P, Sheena Y, Kashef E, Purkayastha S, Darzi A, Paraskeva P, *et al.* Is basic emergency ultrasound training feasible as part of standard undergraduate medical education? *J Surg Educ* 2010;67:152-6.
- Bahner DP, Adkins EJ, Hughes D, Barrie M, Boulger CT, Royall NA, *et al.* Integrated medical school ultrasound: Development of an ultrasound vertical curriculum. *Crit Ultrasound J* 2013;5:6.
- Baltarowich OH, Di Salvo DN, Scoutt LM, Brown DL, Cox CW, DiPietro MA, *et al.* National ultrasound curriculum for medical students. *Ultrasound Q* 2014;30:13-9.
- Cantisani V, Dietrich CF, Badea R, Ducea S, Prosch H, Cerezo E, *et al.* EFSUMB statement on medical student education in ultrasound [short version]. *Ultraschall Med* 2016;37:100-2.
- Patel NY, Riherd JM. Focused assessment with sonography for trauma: Methods, accuracy, and indications. *Surg Clin North Am* 2011;91:195-207.
- Hoppmann RA, Rao VV, Poston MB, Howe DB, Hunt PS, Fowler SD, *et al.* An integrated ultrasound curriculum (iUSC) for medical students: 4-year experience. *Crit Ultrasound J* 2011;3:1-2.
- Bell FE 3rd, Wilson LB, Hoppmann RA. Using ultrasound to teach medical students cardiac physiology. *Adv Physiol Educ* 2015;39:392-6.
- Ahn JS, French AJ, Thiessen ME, Kendall JL. Training peer instructors for a combined ultrasound/physical exam curriculum. *Teach Learn Med* 2014;26:292-5.
- Yu TC, Wilson NC, Singh PP, Lemanu DP, Hawken SJ, Hill AG, *et al.* Medical students-as-teachers: A systematic review of peer-assisted teaching during medical school. *Adv Med Educ Pract* 2011;2:157-72.
- Dickerson J, Paul K, Vila P, Whiticar R. The role for peer-assisted ultrasound teaching in medical school. *Clin Teach* 2017;14:170-4.