

ORIGINAL ARTICLE

Up-regulation of T β RIII facilitates the osteogenesis of supraspinous ligament-derived fibroblasts from patients with ankylosing spondylitis

Ying Zhang | Wu-gui Chen | Si-zhen Yang | Hao Qiu | Xu Hu | Yi-yun Qiu |
Xuan Wen | Yue Zhou  | Tong-wei Chu 

Department of Orthopaedics, Xinqiao Hospital, Army Military Medical University, Chongqing, China

Correspondence

Tong-wei Chu, Department of Orthopaedics, Xinqiao Hospital, Army Military Medical University, Xinqiao Main Street of Shapingba District, Chongqing 400037, China.
Email: chtw@sina.com

Funding information

This research was funded by the National Natural Youth Science Fund of China (No. 81501853), the National Natural Science Foundation of China (No. 81570800) and the General project of Chongqing Natural Science Foundation of China (No. cstc2020jcyj-msxmX0688)

Abstract

Spinal supraspinous ligament (SL) osteogenesis is the key risk of ankylosing spondylitis (AS), with an unclear pathogenesis. We previously found that transforming growth factor β 1 (TGF- β 1), bone morphogenetic proteins (eg BMP2) and type III TGF- β 1 receptor (T β RIII) expression were markedly up-regulated in AS-SLs. However, the roles of these closely related molecules in AS are unknown. Here, we showed that BMP2, TGF- β 1, T β RIII and S100A4 (a fibroblast marker) were abundant in active osteogenic AS-SL tissues. In vitro, AS-SL fibroblasts (AS-SLFs) showed high BMP2, TGF- β 1 and T β RIII expression and auto-osteogenic capacity. We further evaluated the role of T β RIII in the osteogenesis of normal SLFs. BMP2 combined with TGF- β 1 induced the osteogenesis of T β RIII-overexpressing SLFs, but the activity was lost in SLFs upon T β RIII knockdown. Moreover, our data suggested that BMP2 combined with TGF- β 1 significantly activated both TGF- β 1/Smad signalling and BMP2/Smad/RUNX2 signalling to induce osteogenesis of SLFs with T β RIII up-regulation. Furthermore, our multi-strategy molecular interaction analysis approach indicated that TGF- β 1 presented BMP2 to T β RIII, sequentially facilitating BMP2 recognition by BMPR1A and promoting the osteogenesis of T β RIII-overexpressing SLFs. Collectively, our results indicate that TGF- β 1 combined with BMP2 may participate in the osteogenic differentiation of AS-SLF by acting on up-regulated T β RIII, resulting in excessive activation of both TGF- β 1/Smad and BMP2/BMPR1A/Smad/RUNX2 signalling.

KEYWORDS

ankylosing spondylitis, bone morphogenetic protein 2, osteogenesis, supraspinous ligament fibroblasts, transforming growth factor β 1, type III transforming growth factor β receptor

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2021 The Authors. *Journal of Cellular and Molecular Medicine* published by Foundation for Cellular and Molecular Medicine and John Wiley & Sons Ltd.

1 | BACKGROUND

Ankylosing spondylitis (AS) is the prototypic phenotype of a class of immune-related spondyloarthropathies that affect the spine, sacroiliac joint and hip joint. AS commonly causes spinal rigidity, which manifests as inflammatory pain, spinal fusion, structural and functional impairment of the sacroiliac joint and hip joint. Moreover, the complications of AS include reactive arthritis, Crohn's disease, ulcerative colitis, psoriatic arthritis and acute anterior uveitis. The late stage of AS may lead to spinal fusion and the development of a 'bamboo-like' spine, ultimately resulting in a high risk of disability. Therefore, investigating the pathogenesis of and therapeutic strategies for AS is valuable.

The pathogenesis of AS, which is complex and incompletely understood, involves changes in spinal joints, cartilage, tendons and ligaments. Heterotopic osteogenesis of supraspinous ligaments (SLs) is the key risk of AS, although the pathogenesis remains unclear. A previous study showed that transforming growth factor β 1 (TGF- β 1) and bone morphogenetic proteins (eg BMP2) are up-regulated in peripheral blood from AS patients.¹ Our previous reports further indicated that the expression of TGF- β 1, BMP2 and the type III TGF- β receptor (T β RIII) was significantly increased in AS-SLs compared with SLs from control patients.^{2,3} However, whether TGF- β 1, BMP2 and T β RIII participate in the osteogenesis of AS-SLs is unclear.

Transforming growth factor β 1 and BMP2 are members of the TGF- β superfamily. Members of this family form a complex with the TGF- β receptor family, including T β RI, T β RII and T β RIII, thereby regulating upstream Smad signalling.⁴ Among them, T β RIII (also known as β -proteoglycan), an abundant membrane-anchoring protein, was originally thought to be an auxiliary receptor of the TGF- β superfamily. Previous reports indicate that T β RIII plays a critical role in TGF- β /Smad signalling.⁵ A report from another group suggests that BMP2, BMP4 and BMP7 can bind to T β RIII using a biosensor method and that loss of T β RIII markedly decreases BMP2 activity.^{6,7} Moreover, a recent report suggests that TGF- β 1 and BMP2 co-regulate T β RIII to activate epithelial-mesenchymal transition (EMT) and invasion in epicardial cells, further suggesting that TGF- β 1 and BMP2 can simultaneously act on T β RIII.⁷ However, whether TGF- β 1 and BMP2 co-regulate T β RIII in AS-SLs is unknown.

Unexpectedly, we found that some fibroblast-like cells isolated from AS-SLs showed auto-osteogenic capacity. These cells expressed high levels of fibroblast-specific protein 1 (Fsp1, also known as S100A4), which is a specific marker of fibroblasts.⁸ Therefore, we hypothesized that TGF- β 1 and BMP2 may participate in the osteogenesis of SL fibroblasts (SLFs) from patients with AS (AS-SLFs) through cooperatively acting on T β RIII. This study aimed to further unveil the role of T β RIII in the osteogenesis of AS-SLFs and provide a new target for developing promising therapies.

2 | EXPERIMENTAL SECTION

2.1 | Sampling

Ankylosing spondylitis-supraspinous ligaments were obtained via spinal orthopaedic surgery from 12 patients (nine males and three females; mean age, 30.2 years) who were in the active stage according to the American Rheumatism Association 1987 revised criteria.⁹ Control supraspinous ligaments (C-SLs) were obtained via reattachment surgery from four patients (two males and two females; mean age, 31.5 years) with spinal fractures. SLs were washed with phosphate-buffered saline (PBS; 0.01 mmol/L; pH 7.2) and were then either stored in liquid nitrogen or fixed with 4% (*m/v*) paraformaldehyde solution and embedded in paraffin. All specimens were obtained with the approval of the ethics committee of Xinqiao Hospital and patient consent.

2.2 | Histological observations

Haematoxylin and eosin (HE) staining was applied to observe the pathologic features of SLs from AS or control patients as described in our previous report.³ To detect protein expression in SLs, immunohistochemistry (IHC) was carried out with antibodies against TGF- β 1, BMP2 (Abcam), S100A4, T β RIII, phospho-Smad2^{Ser465/467}/3^{Ser423/425} (p-Smad2/3), Smad4, phospho-Smad1^{Ser206} (p-Smad1) and runt-related transcription factor 2 (RUNX2; Cell Signaling) as reported previously.³ The histological images were acquired under a light microscope and analysed by ImageJ.

2.3 | Cell cultures

The tissue piece method was applied to isolate supraspinous ligament fibroblasts (SLFs) from AS patients (AS-SLFs) or spinal fracture patients (C-SLFs) according to a previously reported protocol.¹⁰ Briefly, ligaments were washed with Hank's Balanced Salt Solution (HBSS; pH 7.4), minced and cultured in Dulbecco's modified Eagle's medium (DMEM; Gibco) containing 10% (*v/v*) foetal bovine serum (FBS; Gibco), penicillin (100 U/mL) and streptomycin (100 μ g/mL) for 14 days. After SLFs grew, the tissue pieces were removed, and the cells were cultured in complete DMEM.

2.4 | Overexpression and RNA interference

3'-Flag tagged *T β RIII* (GenBank No. NM_001195684) was synthesized and sub-cloned into the vector pcDNA3.1 (Thermo Fisher). The recombinant plasmid used for overexpression was verified by DNA sequencing. Gene knockdown was performed using small

interfering RNA (siRNA; Thermo Fisher). The plasmid was transformed into C-SLFs to obtain C-SLFs with *TβRIII* overexpression using Lipofectamine 3000 (Thermo Fisher) according to the user manuals. Similarly, *TβRIII* siRNA was introduced into C-SLFs with *TβRIII* overexpression to obtain C-SLFs with *TβRIII* knockdown.

2.5 | Osteogenic analysis

Supraspinous ligament fibroblasts cultured in 24-well plates (1×10^4 cells/well) were treated as indicated and differentiated in DMEM containing FBS (10%; *m/v*), β-sodium glycerophosphate (10 mmol/L), dexamethasone (0.1 μmol/L) and vitamin C (30 mmol/L) for 14 days. Then, the cells were further used for mineralization assays or alkaline phosphatase (ALP) activity assays. Mineralization was determined using Alizarin red S staining (Sigma) as described previously.¹¹ In brief, cells were fixed in 95% ethanol, washed with water, incubated with Alizarin red S solution (40 mmol/L pH 4.2) for 15 minutes and washed again with water. The stained cells were imaged using a Leica light microscope (Leica). To detect ALP activity, cells were harvested with cold PBS and gently sonicated. Samples containing equal amounts of protein were used to detect ALP activity using an ALP detection kit (Beyotime).

2.6 | Immunoblotting

Immunoblotting (IB) was performed to detect protein expression as described previously.¹⁰ Briefly, equal amounts of protein extracted from tissues or cells were separated by sodium dodecyl sulphate-polyacrylamide gel electrophoresis (SDS-PAGE). The protein bands were transferred to polyvinylidene fluoride (PVDF) membranes, blocked with 5% (*m/v*) skim milk and probed with the corresponding primary antibodies and horseradish peroxidase (HRP)-conjugated secondary antibodies (Abcam). Chemiluminescence imaging was carried out with Clarity Western ECL substrate (Bio-Rad) in a ChemiDoc™ Touch Imaging System (Bio-Rad). ImageJ was used to calculate protein expression ratios normalized to the expression of GAPDH and that in the control group.

2.7 | Immunoprecipitation

Immunoprecipitation (IP) was performed using a Protein A/G Magnetic Beads kit (Thermo Fisher) according to the user manual. C-SLFs grown in dishes were transfected with the flag-*TβRIII* construct using Lipofectamine 3000. After treatment with TGF-β1 and/or BMP2 as indicated, the cell lysate was harvested using the cell lysis-wash buffer supplied in the kit, and 1 mg of protein was incubated with primary antibodies or anti-mouse IgG (negative control) overnight at 4°C. Immune complexes were captured by the Protein A/G beads and detected by IB.

2.8 | Immunofluorescence

Supraspinous ligament fibroblasts grown on glass slides in a 24-well plate (1×10^4 cells/well) were treated as indicated, fixed with 4% (*m/v*) paraformaldehyde solution and blocked with PBS containing 1% (*m/v*) skim milk and 0.1% (*m/v*) Triton X-100. The cell cytoskeleton was stained with rhodamine-phalloidin (Sigma). TβRIII was probed with a rabbit anti-TβRIII antibody (Abcam) followed by an Alexa Fluor 555-conjugated secondary antibody (Sigma; red fluorescence). TGF-β1 and BMP2 were probed with mouse primary antibodies (Abcam) followed by an Alexa Fluor 488-conjugated secondary antibody (Sigma; green fluorescence). Cell nuclei were stained with DAPI. Fluorescence images were acquired with a laser confocal microscope (Zeiss). ImageJ was used to calculate the cell area, mean fluorescence intensity and the Pearson correlation coefficient (*Rr*) to indicate colocalization.

2.9 | Statistics

Data in histograms are shown as the means ± SDs. Student's *t* test was used for pairwise comparisons. Differences with a *P* value of <.05 were considered statistically significant, <.01 and highly statistically significant.

3 | RESULTS

3.1 | S100A4, TβRIII, TGF-β1 and BMP2 are up-regulated in AS-SLs

The HE staining results showed that the fibres of AS-SLs were broken and disordered, with collagen deposition, vascular proliferation and ectopic bone formation (as shown in dotted line), indicating an osteogenic morphology (Figure 1A). The immunohistochemical results suggested that S100A4 (a fibroblast marker), TβRIII, TGF-β1 and BMP2 were up-regulated in the tissue near the ectopic bone in the contiguous sections, and the expression region was very consistent (Figure 1A,B). To further assess the expression of these proteins, total protein was extracted from AS-SLs and C-SLs. As expected, the IB results also showed that the expression of S100A4, TβRIII, TGF-β1 and BMP2 was markedly increased in AS-SLs (Figure 1C). Together, these results suggest that TβRIII, TGF-β1 and BMP2 are up-regulated in AS-SLs with abundant fibroblasts. However, further studies are needed to determine whether these proteins are expressed in the additional fibroblasts.

3.2 | AS-SLFs exhibit auto-osteogenic capacity and up-regulation of TβRIII, TGF-β1 and BMP2

Fibroblasts play crucial roles in extracellular matrix (ECM) alteration, organ remodelling and tissue repair. To further investigate the role of fibroblasts in AS, we isolated and obtained C-SLFs and AS-SLFs. By measuring the mineralization capacity and ALP activity of both




FIGURE 1 Supraspinous ligaments of AS patients exhibit up-regulation of S100A4, T β RIII, TGF- β 1 and BMP2. A, Sections of supraspinous ligaments from patients with spinal fracture (control) or AS were stained with haematoxylin and eosin (HE) or subjected to immunohistochemistry (IHC) using antibodies against S100A4, T β RIII, TGF- β 1 and BMP2. The histological changes were evaluated under a light microscope (bar = 100 μ m). The dotted lines indicate the edge of the ectopic bone. B, Expression levels as indicated by IHC. Expression levels were detected by measuring the positive staining area in ImageJ and were calculated by normalization to a control. C, Total tissue lysates of samples as described in (A) were examined by immunoblotting using the corresponding antibodies. The histograms indicate the protein expression ratios normalized to GAPDH expression. The dotted line indicates the edge of the ectopic bone. ** P < .01, *** P < .001 and **** P < .0001 (control: n = 4; AS: n = 12)

C-SLFs and AS-SLFs, we found that compared to C-SLFs, AS-SLFs showed strong auto-osteogenic capacity (Figure 2A,B). Moreover, AS-SLFs had a larger cell area than C-SLFs (Figure 2C). Furthermore, the expression of T β RIII, TGF- β 1 and BMP2 was markedly up-regulated in AS-SLFs (Figure 2D). These results strongly suggest that up-regulation of T β RIII, TGF- β 1 and BMP2 may mediate osteogenic differentiation of AS-SLFs; however, more evidence is needed to confirm this hypothesis.

3.3 | T β RIII up-regulation facilitates the osteogenesis of C-SLFs

To further confirm that up-regulation of T β RIII, TGF- β 1 and BMP2 may induce the osteogenesis of SLFs, we constructed a C-SLF cell

line with T β RIII up-regulation (Figure 3A). The cytoskeleton of C-SLFs with T β RIII up-regulation was stained with rhodamine-phalloidin, and the effect of TGF- β 1 and BMP2 on the cell area was determined. The cell morphology of the medium group was spindle or polygonal, consistent with that of normal C-SLFs, as shown in Figure 2C. Unsurprisingly, culture with either TGF- β 1 or BMP2 alone markedly induced cell hypertrophy compared with that in the medium group. Notably, the cell size in the group cultured with TGF- β 1 combined with BMP2 was further increased compared with that in the first three groups mentioned above (Figure 3B). Furthermore, TGF- β 1 combined with BMP2 also resulted in more mineralization nodules than medium, TGF- β 1 or BMP2 alone (Figure 3C). Consistent with these results, the combination group exhibited a synergistic effect on increasing ALP activity compared with that in the medium and single treatment groups (Figure 3D). These results indicate that the




FIGURE 2 AS-SLFs exhibit auto-osteogenic capacity and up-regulation of T β RIII, TGF- β 1 and BMP2. A, B, C-SLFs and AS-SLFs were plated in 24-well plates (1×10^4 cells/well). Then, the cells were differentiated in DMEM containing FBS (10%; m/v), β -glycerophosphate sodium (10 mmol/L), dexamethasone (0.1 $\mu\text{mol/L}$) and vitamin C (30 mmol/L) for 14 days and further used for mineralization assays (A) or ALP activity assays (B). Bar = 50 μm . C, C-SLFs and AS-SLFs were seeded on glass slides in a 24-well plate (1×10^4 cells/well) for 48 h, stained with rhodamine-phalloidin and observed via laser confocal microscopy. The cell area was measured in ImageJ ($n = 50$; bar = 50 μm). D, Cells cultured as described in (C) were stained for T β RIII, TGF- β 1 and BMP2 (Alexa Fluor 555; red). The mean fluorescence intensity was measured in ImageJ ($n = 50$; bar = 50 μm). ** $P < .01$; *** $P < .001$

combination of TGF- β 1 and BMP2 induces the hypertrophy and osteogenic differentiation of C-SLFs with T β RIII up-regulation and further enhances the osteogenic capacity of AS-SLFs.

3.4 | T β RIII knockdown inhibits the osteogenesis of C-SLFs

The above data suggest that TGF- β 1 combined with BMP2 can induce osteogenic differentiation of C-SLFs with T β RIII up-regulation. To further confirm these findings, we constructed C-SLFs with T β RIII knockdown using specific siRNA targeting T β RIII (Figure 4A). Consistent with the results shown in Figure 3B, TGF- β 1 combined with BMP2 showed a synergistic effect on increasing the size of C-SLFs with T β RIII up-regulation (transfected with control siRNA). However, neither TGF- β 1 nor BMP2 alone or in combination could change the size of C-SLFs with T β RIII knockdown (transfected with T β RIII siRNA; Figure 4B). Consistent with this pattern, the combination treatment showed no synergistic effect on inducing mineralization nodule formation or ALP activity in C-SLFs with T β RIII knockdown (Figure 4C,D). Collectively, these results indicate that T β RIII knockdown inhibits the osteogenic differentiation of C-SLFs induced by TGF- β 1 plus with BMP2.

3.5 | T β RIII up-regulation facilitates osteogenesis via TGF- β 1-Smad2/3 and BMP2-Smad1-RUNX2 signalling

The above results indicate that T β RIII up-regulation is critical for the osteogenic differentiation of C-SLFs, but the detailed mechanism remains unclear. Using C-SLFs with T β RIII up-regulation, we showed that compared to TGF- β 1 or BMP2 alone, TGF- β 1 in combination with BMP2 up-regulated the expression of T β RIII but not T β RI and T β RII (Figure 5A). However, these effects were lost in C-SLFs with T β RIII knockdown (Figure 5B).

In addition, we further observed the effect of TGF- β 1 in combination with BMP2 on Smad signalling downstream of TGF- β 1 and BMP2 in these two cell lines. TGF- β 1 in combination with BMP2 markedly increased the phosphorylation of Smad2/3 (TGF- β 1 signalling), Smad1 (BMP2 signalling) and Smad4 (the co-Smad of both TGF- β 1 and BMP2 signalling) as well as the expression of RUNX2 (BMP2 signalling) (Figure 5C,D). However, the synergistic effects were lost in C-SLFs with T β RIII knockdown (Figure 5E,F). These results suggest that TGF- β 1 combined with BMP2 further up-regulates T β RIII expression in SLFs with T β RIII up-regulation and then sequentially activates both TGF- β 1-Smad2/3 and BMP2-Smad1-RUNX2 signalling, leading to cell osteogenic differentiation.




FIGURE 3 TβRIII up-regulation facilitates osteogenic differentiation of C-SLFs. A, C-SLFs were plated in 6-well plates (1×10^5 cells/well) and transfected with a 3'-Flag tagged TβRIII construct using a liposome method. TβRIII expression was assessed by IB. B, C-SLFs with TβRIII up-regulation were cultured as shown in Figure 2C and treated with TGF-β1 (10 μmol/L) and/or BMP2 (50 μg/mL) for 24 h. Cells were then stained, observed and analysed as shown in Figure 2C ($n = 50$; bar = 20 μm). C, D, C-SLFs with TβRIII up-regulation were plated in 24-well plates (1×10^4 cells/well), treated as shown in (B), differentiated as shown in Figure 2A, and then further used for mineralization assays (C) or ALP activity assays (D). Bar = 50 μm. ** $P < .01$; *** $P < .001$

3.6 | TGF-β1 presents BMP2 to TβRIII, thereby increasing BMP2-BMPR1A binding

The above data indicate that TGF-β1 combined with BMP2 up-regulates TβRIII-mediated TGF-β1 and BMP2 signalling. In addition to the expression levels, whether and how TβRIII up-regulation contributes to the binding of TGF-β1 or BMP2 to their own receptors are unclear. Therefore, colocalization and co-IP assays were used to evaluate whether TβRIII up-regulation affects TGF-β1 and BMP2 ligand-receptor interactions.

In the colocalization experiments, TGF-β1 or BMP2 was labelled green, TβRIII was labelled red, and colocalization of TβRIII-TGF-β1 and TβRIII-BMP2 was observed as yellow fluorescence. Neither TGF-β1 nor BMP2 alone or in combination increased TβRIII-TGF-β1 colocalization (Figure 6A,B). In addition, neither TGF-β1 nor BMP2 alone or in combination increased TβRIII-BMP2 colocalization; however, TGF-β1 combined with BMP2 markedly increased TGFβR3-BMP2 colocalization (Figure 6C,D). These results suggest that TGF-β1 promotes the binding of BMP2 to TβRIII, but BMP2 does not affect the TβRIII-TGF-β1 interaction.

To further confirm the colocalization assay results, antibodies against TGF-β1, BMP2 and Flag-TβRIII were separately used to pull down the associated proteins in total cell lysate extracted from C-SLFs with TβRIII up-regulation. The results of IB on the Flag-TβRIII immunoprecipitate showed that BMP2 combined with TGF-β1 decreased the TβRIII-TGF-β1 interaction while significantly increasing the TβRIII-BMP2 interaction, indicating that TGF-β1 presents BMP2 to TβRIII (Figure 6E,F). Unsurprisingly, in the TGF-β1 immunoprecipitate, BMP2 combined with TGF-β1 did not increase the binding of TGF-β1 to TβRIII or TβRI (Figure 6G,H). Remarkably, in the BMP2 immunoprecipitate, BMP2 combined with TGF-β1 increased both the TβRIII-BMP2 and BMPR1A-BMP2 interactions, suggesting that TGF-β1 presents BMP2 to TβRIII, thereby facilitating the recognition of BMP2 by its own receptor, BMPR1A (Figure 6I,J). Not surprisingly, no TGF-β1, BMP2 or TβRIII band was detected in pull-down products of anti-mouse IgG used as a negative control (data not shown). These results suggest that TβRIII binds TGF-β1 and BMP2 simultaneously and further indicate that TGF-β1 presents BMP2 to TβRIII and facilitates the recognition of BMP2 by BMPR1A.




FIGURE 4 TβRIII siRNA decreases osteogenic differentiation of C-SLFs with TβRIII up-regulation. A, C-SLFs with TβRIII up-regulation were plated in 6-well plates (1×10^5 cells/well) and transfected with TβRIII siRNA or control siRNA using a liposome method. TβRIII expression was assessed by IB. B, C-SLFs with TβRIII knockdown were treated and assessed as shown in Figure 3B (n = 50). Bar = 20 μm. C, D, C-SLFs with TβRIII knockdown were treated as shown in Figure 3B and further used for mineralization assays (C) or ALP activity assays (D). Bar = 50 μm. **P < .01; #P > .05




FIGURE 5 TβRIII up-regulation facilitates the osteogenesis of C-SLFs via TGF-β1-Smad2/3 and BMP2-Smad1-RUNX2 signalling. A, C, D, C-SLFs with TβRIII up-regulation were plated in 6-well plates (1×10^5 cells/well) and treated with TGF-β1 (10 μmol/L) and/or BMP2 (50 μg/mL) for 24 h. The total lysate was extracted and used for IB with the indicated antibodies. B, E, F, C-SLFs with TβRIII knockdown were cultured, treated and assessed as shown in (A). The histograms indicate the protein expression ratios (n = 3). **P < .01; #P > .05




FIGURE 6 TGF- β 1 presents BMP2 to T β RIII, thereby increasing BMP2-BMPR1A binding. A, B, C-SLFs with T β RIII up-regulation were seeded on glass slides in a 24-well plate (1×10^4 cells/well) and treated with TGF- β 1 (10 μ mol/L) and/or BMP2 (50 μ g/mL) for 24 h. Cells were then stained for TGF- β 1 (Alexa Fluor 488; green) and T β RIII (Alexa Fluor 555; red) and observed under a laser confocal microscope (bar = 5 μ m). Colocalization of TGF- β 1 with T β RIII was assessed using the Pearson correlation coefficient (Rr) via ImageJ (n = 50). C, D, Experiments were performed as described in (A), and cells were stained for BMP2 (Alexa Fluor 488; green) and T β RIII (Alexa Fluor 555; red) (bar = 5 μ m). E–J, C-SLFs with T β RIII up-regulation were seeded in 10-cm dishes (1×10^6 cells/dish) and treated as in (A). The total lysate was used for immunoprecipitation (IP) using antibodies against (E, F) Flag, (G, H) TGF- β 1 or (I, J) BMP2, and the associated proteins, as indicated, were detected via immunoblotting (IB). Histograms represent the protein expression ratios (n = 3). #*P* > .05; ***P* < .01

3.7 | p-Smad2/3, Smad4, p-Smad1 and RUNX2 levels are increased in AS-SLs

Paraffin sections of C-SLs and AS-SLs (the same sections as shown in Figure 1) were used for IHC to assess the activation of

TGF- β 1-Smad2/3 and BMP2-Smad1-RUNX2 signalling. No large areas of brown coloration were observed in C-SLs, suggesting that the levels of p-Smad2/3, p-Smad1, Smad4 and RUNX2 were decreased. However, the immunohistochemical results showed that the levels of p-Smad2/3, p-Smad1, Smad4 and RUNX2 were

increased in the tissue near the ectopic bone in the contiguous sections (Figure 7). The results further indicate that TGF- β 1-Smad2/3 and BMP2-Smad1-RUNX2 signalling are activated in AS-SLs.

4 | DISCUSSION

Numerous reports indicate that inflammation, bone erosion and cartilage formation are the basic characteristics of AS pathology. Ectopic osteogenesis within the SL is the key risk of AS and is the main cause of joint fusion/stiffness and mobility loss. As shown in Figure 1, significant fibre breakage and disorganization, with collagen deposition, angiogenesis and ectopic bone formation, were observed in AS ligaments. The data presented here suggest that AS-SLFs have a larger cell area and exhibit obvious self-osteogenic capacity in vitro, consistent with other reports.¹² The understanding of the molecular mechanisms underlying new bone formation in AS has significantly improved but remains incomplete. Here, we demonstrate that TGF- β 1 combined with BMP2 can induce the osteogenesis of AS-SLFs by acting on up-regulated T β RIII, thus resulting in excessive activation of both TGF- β 1/Smad and BMP2/BMPRI1A/Smad/RUNX2 signalling. These findings establish a

novel pathway downstream of T β RIII in AS progression that contributes to bone formation by regulating the function of TGF- β 1 and BMP2.

The key role of inflammation in regulating bone remodelling has been extensively reported, and spatiotemporally controlled release of inflammatory signals and factors is essential for regulating bone remodelling.¹³ Dysregulated inflammation leads to increased bone resorption and suppressed bone formation in bone diseases such as arthritis and fracture, but its detailed mechanism in promoting ectopic osteogenesis in AS is unclear. Typically, proinflammatory cytokines, including tumour necrosis factor- α (TNF- α), IL-1, IL-11, IL-6 and IL-17, promote bone resorption by enhancing osteoclast differentiation and activity and/or inhibiting osteoblast differentiation, collagen synthesis and bone formation, while anti-inflammatory cytokines, such as TGF- β , IL-10 and IL-13, show the opposite effects. Recent evidence suggests that a series of inflammatory mediators, such as TGF- β , interleukin and TNF- α , not only play a key role in AS but also affect osteoclast and osteoblast activity and lead to bone remodelling. Therefore, some clinical studies have shown that inflammation inhibitors, such as monoclonal antibodies against TNF- α and new biological inhibitors targeting IL-17/IL-23, could slow radiological progression in AS patients.¹⁴ However, their impact on slowing bone structural damage is still not clearly established.^{15,16} Inflammatory cytokines such as




FIGURE 7 AS-SLs from patients exhibit up-regulation of p-Smad2/3, Smad4, p-Smad1 and RUNX2. A, The sections from the same specimens as described in Figure 1A were subjected to immunohistochemistry (IHC) using antibodies against phospho-Smad2Ser465/467/3Ser423/425 (p-Smad2/3), Smad4, phospho-Smad1Ser206 (p-Smad1) and RUNX2. The sections were examined under a light microscope (bar = 100 μ m). The dotted lines indicate the edge of the ectopic bone. B, Expression levels were detected by measuring positive staining area in ImageJ and were calculated by normalization to the control. *** P < .001 and **** P < .0001 (n = 3)

TNF- α and IL-1 β have also been reported to play a coordinated role in regulating osteoblast differentiation *in vitro* and *in vivo*.¹⁷ In addition to the proinflammatory factors TNF- α and IL-6, the following factors play a significant role in osteoblast differentiation: Wnt/wingless protein family members, BMP, TGF- β , fibroblast growth factor (FGF) and insulin-like growth factor (IGF).

The TGF- β superfamily comprises a group of polypeptide factors with similar structures, including TGF- β , activin/inhibitors, BMPs and growth differentiation factor (GDF), that play key roles in inflammation, bone reconstruction and cancer progression.¹⁸ Both TGF- β 1 and BMP2 belong to the TGF- β superfamily and are involved in inflammation and bone formation. They are important osteogenic mediators and contribute to the progression of multiple diseases,¹⁹ including AS.^{2,3,20} Previous studies indicate that TGF- β and BMP2 have synergistic effects on promoting osteogenic differentiation,²¹⁻²³ but the detailed mechanism remains unclear. Wang and He noted that TGF- β 3 and BMP2 have a synergistic effect on regulating the osteogenic differentiation of bone marrow mesenchymal stem cells but did not propose a specific mechanism.^{23,24} As our results in Figure 2 show, TGF- β 1 combined with BMP2 further induces the hypertrophy and osteogenic differentiation of C-SLFs and may participate in the osteogenic differentiation of AS-SLFs. Furthermore, our study further unveils the mechanism by which TGF- β 1 synergizes with BMP-2 to promote osteogenic differentiation in AS.

T β RIII (also known as β -proteoglycan) was originally thought to be an auxiliary receptor for members of the TGF- β superfamily, including TGF- β and BMP2, and to enhance their cellular effects.^{6,7} In endocardial cells, T β RIII is required to activate the Par6/Smurf1/RhoA pathway via TGF- β and BMP-2 to induce EMT and invasion.^{7,25,26} T β RIII binds all three TGF β ligands and BMP2, but its contribution to AS is unknown. Our research further indicated that the expression of TGF- β 1, BMP2 and T β RIII was significantly increased in AS-SLFs compared with C-SLFs. As shown in Figure 5A, TGF- β 1 combined with BMP2 significantly up-regulated the expression of T β RIII but not T β RI or T β RII. Subsequent findings indicated that after up-regulation of T β RIII expression in C-SLFs, TGF- β 1 combined with BMP2 further induced cell hypertrophy and mineralization nodule formation compared with these effects in the single treatment groups. However, the synergistic effects were lost in C-SLFs with T β RIII knockdown, further suggesting that T β RIII up-regulation may participate in the osteogenic differentiation of AS-SLFs, which has rarely been reported.^{7,19}

Additionally, our experiments further confirmed that TGF- β 1 promotes the binding of BMP2 to T β RIII, but BMP2 does not affect the T β RIII-TGF- β 1 interaction. Moreover, in the presence of TGF- β 1, up-regulated T β RIII presents BMP2 for binding to T β RIII, which facilitates the recognition of BMP2 by BMPR1A and then may have a role in enhancing the osteogenesis of AS-SLFs. These data further demonstrated another possible important mechanism underlying the T β RIII-mediated function of TGF- β 1 and BMP2 that contributes to osteogenic differentiation in AS that has not been reported.

Multiple signalling pathways, such as the BMP-Smads, Wnt/ β -catenin, Notch, Hedgehog and FGF signalling pathways, have been indicated to play critical roles in the regulation of osteogenic

differentiation. These signalling pathways not only have complex regulatory mechanisms but are also interrelated and influence each other, thus forming a more complex and sophisticated regulatory network to regulate osteogenic differentiation. To our knowledge, both TGF- β 1 and BMP-2 activate Smad signalling, and phosphorylated Smads translocate into the nucleus to promote the expression of genes involved in osteoblast differentiation.^{27,28} Typically, BMP-2 primarily utilizes Smad1, whereas TGF- β 1 utilizes Smad2/3.^{27,29} Herein, we demonstrate that TGF- β 1 combined with BMP2 up-regulates T β RIII expression in SLFs and sequentially activates both TGF- β 1-Smad2/3 and BMP2-Smad1-RUNX2 signalling, leading to cell osteogenic differentiation. In addition, the immunohistochemical results indicated TGF- β 1-Smad2/3 and BMP2-Smad1-RUNX2 signalling activation in paraffin sections of human AS-SLFs but not C-SLFs. However, further animal experiments are needed.

5 | CONCLUSIONS

In conclusion, we demonstrated that TGF- β 1 combined with BMP2 may participate in the osteogenic differentiation of AS-SLFs by acting on up-regulated T β RIII, thus resulting in excessive activation of both TGF- β 1/Smad and BMP2/BMPR1A/Smad/RUNX2 signalling. These data force us to reconsider the interactive roles and contributions of these ligands in the activation of canonical signalling pathways through T β RIII in AS.

ACKNOWLEDGEMENTS

The authors thank the Central Laboratory of Xinqiao Hospital for their technical support during the experiment and Dr Min Ma, Dr Rujie Wang and Assistant Ling Yang for their help and support during the experiment.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

AUTHOR CONTRIBUTION

Ying Zhang: Conceptualization (lead); Data curation (lead); Formal analysis (lead); Funding acquisition (lead); Project administration (lead); Validation (equal); Writing-original draft (lead); Writing-review & editing (lead). **Wu-gui Chen:** Data curation (equal); Formal analysis (equal); Investigation (equal); Methodology (lead); Software (lead); Writing-original draft (equal); Writing-review & editing (equal). **Si-zhen Yang:** Data curation (equal); Formal analysis (equal); Investigation (equal); Writing-original draft (equal). **Hao Qiu:** Formal analysis (supporting); Software (supporting); Supervision (supporting); Writing-review & editing (equal). **Xu Hu:** Data curation (supporting); Formal analysis (supporting); Writing-review & editing (supporting). **Yi-yun Qiu:** Data curation (supporting); Formal analysis (supporting); Writing-review & editing (supporting). **Xuan Wen:** Data curation (supporting); Formal analysis (supporting); Writing-review & editing (supporting). **Yue Zhou:** Project administration (equal); Resources (supporting); Writing-review & editing (supporting). **Tong-wei Chu:** Conceptualization (lead);

Project administration (lead); Resources (equal); Writing-original draft (equal); Writing-review & editing (equal).

ETHICAL APPROVAL AND CONSENT TO PARTICIPATE

This study was approved by the Research Ethics Committee of Xinqiao Hospital, and informed consent was obtained from all participants.

CONSENT FOR PUBLICATION

Written informed consent for publication was obtained from all participants.

DATA AVAILABILITY STATEMENT

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

ORCID

Yue Zhou  <https://orcid.org/0000-0001-7102-6484>

Tong-wei Chu  <https://orcid.org/0000-0003-0309-7082>

REFERENCES

1. Reveille JD. The genetic basis of ankylosing spondylitis. *Curr Opin Rheumatol*. 2006;18:332-341.
2. Zhang Y, Hu X, Zhang C, Zhou Y, Chu TW. Suppression subtractive hybridization reveals differentially expressed genes in supraspinous ligaments of patients with ankylosing spondylitis. *Mol Med Rep*. 2015;11:4482-4488.
3. Zhang Y, Xu H, Hu X, Zhang C, Chu T, Zhou Y. Histopathological changes in supraspinous ligaments, ligamentum flava and paraspinous muscle tissues of patients with ankylosing spondylitis. *Int J Rheum Dis*. 2014;19:420-429.
4. Kawaguchi Y, Furushima K, Sugimori K, Inoue I, Kimura T. Association between polymorphism of the transforming growth factor-beta1 gene with the radiologic characteristic and ossification of the posterior longitudinal ligament. *Spine*. 2003;28:1424-1426.
5. Kamiya M, Harada A, Mizuno M, Iwata H, Yamada Y. Association between a polymorphism of the transforming growth factor-beta1 gene and genetic susceptibility to ossification of the posterior longitudinal ligament in Japanese patients. *Spine*. 2001;26:1264-1266; discussion 6-7.
6. Kirkbride KC, Townsend TA, Bruinsma MW, Barnett JV, Globe GC. Bone morphogenetic proteins signal through the transforming growth factor-beta type III receptor. *J Biol Chem*. 2008;283:7628-7637.
7. Sanchez NS, Barnett JV. TGFbeta and BMP-2 regulate epicardial cell invasion via TGFbetaR3 activation of the Par6/Smurf1/RhoA pathway. *Cell Signal*. 2012;24:539-548.
8. Boye K, Maelandsmo GM. S100A4 and metastasis: a small actor playing many roles. *Am J Pathol*. 2010;176:528-535.
9. Arnett FC, Edworthy SM, Bloch DA, et al. The American Rheumatism Association 1987 revised criteria for the classification of rheumatoid arthritis. *Arthritis Rheum*. 1988;31:315-324.
10. Pan XC, Liu Y, Cen YY, et al. Dual role of triptolide in interrupting the NLRP3 inflammasome pathway to attenuate cardiac fibrosis. *Int J Mol Sci*. 2019;20:360.
11. Wang X, Wang G, Liu L, Zhang D. The mechanism of a chitosan-collagen composite film used as biomaterial support for MC3T3-E1 cell differentiation. *Sci Rep*. 2016;6:39322.
12. Yang M, Yuan H, Miao M, Xu W. The osteogenic potential of ligament fibroblasts is greater in ankylosing spondylitis patients than in patients with osteoarthritis. *Z Rheumatol*. 2015;74:340-345.
13. Adamopoulos IE. Inflammation in bone physiology and pathology. *Curr Opin Rheumatol*. 2018;30:59-64.
14. Sieper J, Poddubnyy D. Axial spondyloarthritis. *Lancet*. 2017;390:73-84.
15. Maas F, Spoorenberg A, Brouwer E, et al. Spinal radiographic progression in patients with ankylosing spondylitis treated with TNF- α blocking therapy: a prospective longitudinal observational cohort study. *PLoS One*. 2015;10:e0122693-e.
16. Braun J, Baraliakos X, Deodhar A, et al. Effect of secukinumab on clinical and radiographic outcomes in ankylosing spondylitis: 2-year results from the randomised phase III MEASURE 1 study. *Ann Rheum Dis*. 2017;76:1070-1077.
17. Ding J, Ghali O, Lencel P, et al. TNF- α and IL-1 β inhibit RUNX2 and collagen expression but increase alkaline phosphatase activity and mineralization in human mesenchymal stem cells. *Life Sci*. 2009;84:499-504.
18. Chen W, Ten Dijke P. Immunoregulation by members of the TGFbeta superfamily. *Nat Rev Immunol*. 2016;16:723-740.
19. Song R, Fullerton DA, Ao L, Zheng D, Zhao KS, Meng X. BMP-2 and TGF- β 1 mediate biglycan-induced pro-osteogenic reprogramming in aortic valve interstitial cells. *J Mol Med*. 2015;93:403-412.
20. Xie Z, Wang P, Li Y, et al. Imbalance between bone morphogenetic protein 2 and noggin induces abnormal osteogenic differentiation of mesenchymal stem cells in ankylosing spondylitis. *Arthritis Rheumatol*. 2016;68:430-440.
21. Fromigué O, Marie PJ, Lomri A. Bone morphogenetic protein-2 and transforming growth factor-beta2 interact to modulate human bone marrow stromal cell proliferation and differentiation. *J Cell Biochem*. 1998;68:411-426.
22. Asparuhova MB, Caballé-Serrano J, Buser D, Chappuis V. Bone-conditioned medium contributes to initiation and progression of osteogenesis by exhibiting synergistic TGF- β 1/BMP-2 activity. *Int J Oral Sci*. 2018;10:20.
23. Wang Y, He T, Liu J, et al. Synergistic effects of overexpression of BMP-2 and TGF- β 3 on osteogenic differentiation of bone marrow mesenchymal stem cells. *Mol Med Rep*. 2016;14:5514-5520.
24. He W, Chen L, Huang Y, et al. Synergistic effects of recombinant Lentiviral-mediated BMP2 and TGF-beta3 on the osteogenic differentiation of rat bone marrow mesenchymal stem cells in vitro. *Cytokine*. 2019;120:1-8.
25. Townsend TA, Wrana JL, Davis GE, Barnett JV. Transforming growth factor-beta-stimulated endocardial cell transformation is dependent on Par6c regulation of RhoA. *J Biol Chem*. 2008;283:13834-13841.
26. Townsend TA, Robinson JY, Deig CR, et al. BMP-2 and TGF β 2 shared pathways regulate endocardial cell transformation. *Cells Tissues Organs*. 2011;194:1-12.
27. Chen G, Deng C, Li Y-P. TGF- β and BMP signaling in osteoblast differentiation and bone formation. *Int J Biol Sci*. 2012;8:272-288.
28. Wu M, Chen G, Li Y-P. TGF- β and BMP signaling in osteoblast, skeletal development, and bone formation, homeostasis and disease. *Bone Res*. 2016;4:16009.
29. Ding L-Z, Teng X, Zhang Z-B, Zheng C-J, Chen S-H. Mangiferin inhibits apoptosis and oxidative stress via BMP2/Smad-1 signaling in dexamethasone-induced MC3T3-E1 cells. *Int J Mol Med*. 2018;41:2517-2526.

How to cite this article: Zhang Y, Chen W-G, Yang S-Z, et al. Up-regulation of T β RIII facilitates the osteogenesis of supraspinous ligament-derived fibroblasts from patients with ankylosing spondylitis. *J Cell Mol Med*. 2021;25:1613-1623. <https://doi.org/10.1111/jcmm.16262>