

POSTER PRESENTATION

Open Access

P72. Transgenic expression of a chimeric signaling receptor to facilitate T cell costimulation in the tumour environment

R Schlenker^{1*}, M Leisegang², W Uckert², E Noessner¹

From 1st Immunotherapy of Cancer Conference (ITOC1)
Munich, Germany. 12-14 March 2014

Tumour therapy with T cell receptor (TCR) engineered T cells is reported to induce clinical responses but shortcomings regarding poor *in vivo* persistence and loss of function in the tumour milieu have been observed. Providing costimulation to adoptively transferred T cells may improve these shortcomings. However, human T effector cells are largely CD28 negative and epithelial tumours do not express CD80 or CD86. Therefore, costimulation of human CD8 T effector cells cannot be triggered via the classical way of CD28 ligation. We propose to facilitate costimulation of CD8 T effector cells in the tumour milieu through retroviral engineering of T cells with a chimeric signaling molecule (CSM). This CSM is consisted of an intracellular costimulatory domain fused to an extracellular domain with binding capacity for a ligand expressed by a great variety of tumours.

Human activated PBL retrovirally transduced to express the CSM exhibited a survival advantage during *in vitro* expansion according to clinical protocol. The effect of the chimeric molecule on T cell function was analyzed using T cells expressing a tumour antigen specific TCR alone or in combination with the CSM. Transduced T cells were stimulated with target cells positive or negative for the CSM ligand (CSM-L). CSM expressing T cells responded better to CSM-L⁺ target cells showing higher phosphorylation of ERK and RPS6 compared to stimulation with CSM-L⁻ target cells. CSM⁻ T cells responded equally to both target cells. Accordingly, CSM⁺ but not CSM⁻ T cells secreted more IL-2 and IFN- γ upon co-culture with CSM-L⁺ target cells. In summary, transduction of PBL with the chimeric signaling

molecule supported T cell survival and TCR induced signaling leading to enhanced T cell function.

Authors' details

¹Helmholtz Zentrum Muenchen, Institute of Molecular Immunology, Munich, Germany. ²Max Delbrueck Center for Molecular Medicine, Molecular Cell Biology and Gene Therapy, Berlin, Germany.

Published: 12 March 2014

doi:10.1186/2051-1426-2-S2-P46

Cite this article as: Schlenker et al.: P72. Transgenic expression of a chimeric signaling receptor to facilitate T cell costimulation in the tumour environment. *Journal for ImmunoTherapy of Cancer* 2014 2(Suppl 2):P46.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



¹Helmholtz Zentrum Muenchen, Institute of Molecular Immunology, Munich, Germany

Full list of author information is available at the end of the article