

Revision of the xyleborine ambrosia beetle genus *Microperus* Wood, 1980 (Curculionidae, Scolytinae, Xyleborini) of Thailand with four new species and four newly recorded species

Wisut Sittichaya¹, Sarah M. Smith², Roger A. Beaver³, Narit Thaochan¹

1 Agricultural Innovation and Management Division, Faculty of Natural Resources, Prince of Songkla University, Songkhla, 90110, Thailand **2** Department of Entomology, Michigan State University, East Lansing, MI 48824, USA **3** 161/2 Mu 5, Soi Wat Pranon, T. Donkaew, A. Maerim, Chiangmai 50180, Thailand

Corresponding author: Wisut Sittichaya (wisut.s@psu.ac.th)

Academic editor: M. Alonso-Zarazaga | Received 7 October 2021 | Accepted 14 November 2021 | Published 3 December 2021

<http://zoobank.org/91828A73-84D9-49ED-8D99-5E94BDE3C1B7>

Citation: Sittichaya W, Smith SM, Beaver RA, Thaochan N (2021) Revision of the xyleborine ambrosia beetle genus *Microperus* Wood, 1980 (Curculionidae, Scolytinae, Xyleborini) of Thailand; with four new species and four newly recorded species. ZooKeys 1074: 191–214. <https://doi.org/10.3897/zookeys.1074.76235>

Abstract

Microperus Wood, 1980 ambrosia beetles in Thailand are reviewed. Four species, *M. bidentatus* **sp. nov.**, *M. bucolicus* **sp. nov.**, *M. globodeclivis* **sp. nov.**, and *M. serratus* **sp. nov.** are described. Four new combinations are given: *Microperus armaticeps* (Schedl, 1942) **comb. nov.**, *Microperus exsculptus* (Eggers, 1927) **comb. nov.**, *Microperus pedellus* (Schedl, 1969) **comb. nov.**, and *Microperus spicatus* (Browne, 1986) **comb. nov., stat. res.**, all from *Xyleborus*. Two new synonyms are proposed: *Microperus cruralis* (Schedl, 1975) (= *Xyleborus myllus* Browne, 1986 **syn. nov.**), *Microperus exsculptus* (Eggers, 1927) (= *Xyleborus dentipennis* Browne, 1983 **syn. nov.**). Four species are reported from Thailand for the first time: *Microperus chrysophylli* (Eggers, 1930), *Microperus exsculptus*, *Microperus nanus* (Browne, 1949) and *Microperus quercicola* (Eggers, 1926). With the inclusion of the *Microperus* species described and recorded herein, the diversity of *Microperus* is increased to 35 species, of which 18 are recorded in Thailand. An updated key to the *Microperus* of the Indochinese Peninsula and China is provided. The taxonomy, diagnostic characters, and distribution of species are briefly discussed.

Keywords

Key, *Microperus*, new combinations, new records, new species, new synonymy, Oriental region, Thailand

Introduction

The xyleborine ambrosia beetle genus *Microperus* was first erected by Wood (1980) for members of the *Xyleborus theae* Eggers, 1940 species group, but the genus name was initially proposed in an unpublished manuscript by F. G. Browne (Wood 1980). Wood diagnosed *Microperus* species as slender, at least 2× as long as wide, posterior face of the antennal club with at least one suture visible and apical margin of corneous area never costate, scutellum invisible and strial punctures usually seriate. Later Wood (1986) synonymized *Microperus* with the similar genus *Coptodryas* Hopkins, 1915 without comment. *Microperus* was resurrected and separated from *Coptodryas* Hopkins, 1915 as a result of a cladistic review of xyleborine taxonomic characters (Hulcr et al. 2007). The genera are distinguished by multiple differences in the antennal characters, and body shape. Subsequent molecular phylogenetic studies including both *Coptodryas* and *Microperus* have confirmed that the genera are closely related (Cognato et al. 2011; Mandelshtam et al. 2019). However, Hulcr et al.'s (2007) diagnosis and description of both *Coptodryas* and *Microperus* were based on a limited taxon sampling of five species per genus and as a result, did not reflect the full morphological variation exhibited by species within these genera. Smith et al. (2020) recently expanded Hulcr et al.'s (2007) work in a revision of the fauna of Indochina and China and provided revised descriptions and diagnoses. Despite these recent advances, additional revisionary work is required to identify other *Microperus* species still erroneously placed in *Coptodryas*.

Microperus currently contains 27 species distributed in Asia from Far East Russia, Korea and Japan, west to Sri Lanka and southeast to Papua New Guinea and Australia. Eighteen species are distributed in Indochina and China, and Thailand has the greatest diversity (Smith et al. 2020). The first *Microperus* species recorded in Thailand was *M. pomelianus* (Schedl 1939), reported as *Xyleborus pomelianus* (Nunberg and Chùjò 1961). Several researchers have subsequently recorded additional species from Thailand (Browne 1980; Beaver 1999; Hutacharern et al. 2007; Sittichaya et al. 2012). The first synoptic list of Thai *Microperus* included nine species (Beaver et al. 2014). Smith et al. (2020) added three additional species, increasing the *Microperus* diversity in Thailand to 12. In this paper, we describe four new species of the genus from Thailand, transfer four species from *Coptodryas* and *Cyclorhipidion* Hagedorn, 1912 to *Microperus* and place two species into synonymy, increasing the total number of *Microperus* species to 35. We also provide a key to the species of *Microperus* in the Indochinese peninsula and China updated from that in Smith et al. (2020).

Materials and methods

Specimens were collected from 27 study sites in 24 conservation areas across all regions of Thailand. Southern Thailand was sampled between 2014–2015 and the North, Northeast and East were sampled between 2019–2020 (Fig. 1). In Southern Thailand 20 ethanol-baited flight intercept traps were deployed in each study site, whereas five




Figure 1. Insect survey localities in Thailand, red stars indicate surveyed areas, blue stars indicate locations where new species and newly recorded *Microperus* species were captured. Source: the map was modified from a Royal Forest Department of Thailand map.

ethanol-baited cross-vane panel traps were used at each site in the other parts of the country (N, NE, E). *Microperus* specimens were collected from Hala-Bala Wildlife Sanctuary, Narathiwat Province; Ton Nga Chang Wildlife Sanctuary, Songkhla Province; Khao Ban Tad Wildlife Sanctuary, Trang province; Khao Nan National Park, Nakhon Sri Thammarat Province, Songkhla Zoo: Songkhla Province, Khao Lak-Lam Ru National Park, Phang Nga Province; Doi Inthanon National Park, Chiang Mai Province and Doi Phu Nang National Park, Phayao Province.

Photographs were taken with a Canon 6D digital Camera with a Canon MP-E 65 mm Macro Photo Lens (Canon, Tokyo, Japan) and StackShot-Macrorail (Cognisys Inc, Michigan, USA). The photos were then combined with Helicon Focus 6.8.0. (Helicon Soft, Ukraine); all photos were improved with Adobe Photoshop CS6 (Adobe Systems, California, USA). The antennal and pronotum types and characters follow those proposed by Hulcr et al. (2007) and subsequently elaborated upon by Smith et al. (2020). Length was measured from pronotum apex to the apex of the declivity and width was measured at the widest part of specimen.

Abbreviations used for entomological collections

MSUC	Albert J. Cook Arthropod Research Collection, Michigan State University, East Lansing, USA;
NHML	Natural History Museum, London, U.K.;
NHMW	Museum of Natural History of Vienna, Austria;
NMNH	National Museum of Natural History, Washington D.C., USA;
RABC	Roger A. Beaver collection, Chiang Mai, Thailand;
THNHM	Natural History Museum of the National Science Museum, Thailand;
WSTC	Private collection of Wisut Sittichaya, Songkhla, Thailand.

Taxonomic treatment

Microperus Wood, 1980

Microperus Wood, 1980: 94.

Type species. *Xyleborus theae* Eggers, 1940 (= *Xyleborus myristicae* Schedl, 1939); original designation.

Diagnosis. This genus is distinguished by the following combination of characters: small to minute size, 1.2–3.1 mm long and cylindrical and elongate form, 1.9–3.2× as long as wide; antennal club truncate or flattened, segment 1 costate, segment 2 or segments 2 and 3 visible on anterior side small and appearing soft, segment 2 or segments 2 and 3 visible on posterior side (types 2, 3, 4 of Hulcr et al. 2007 and Smith et al. 2020); elytral base sinuate, rarely transverse, with a dense tuft of setae present along

elytral base associated with an elytral mycangium; scutellum minute or not apparent; striae and interstriae punctures arranged in parallel rows, striae punctate, prominent; pronotum from lateral view taller than basic (type 2 of Hulcr et al. 2007) or with pronotal disc longer than anterior slope (type 7); pronotum from dorsal view basic and parallel-sided (type 2), or subquadrate (type 3), and anterior margin of pronotum without a row of serrations.

Similar genera. *Coptodryas*, *Xyleborinus*

New species

***Microperus bidentatus* Sittichaya, Smith & Beaver, sp. nov.**

<http://zoobank.org/2F84550F-FF90-4485-955E-C3B5DE6783EE>

Figure 2

Type material. *Holotype*, female, THAILAND, Trang Province, Khao Banthat Wildlife Sanctuary, 7°24'54.6"N, 99°49'47.5"E, tropical rainforest, ethanol baited trap, 01.xii.2013, W. Sittichaya, (NHMW). *Paratype*, female, Nakhon Sri Thammarat Province, Khao Nan National Park, 8°48'31.1"N, 99°33'28.4"E, tropical rainforest, ethanol baited trap, 01.xii.2013, (W. Sittichaya), (1 WSTC).

Similar species. *M. recidens*.

Diagnosis. 1.68–1.76 mm (mean = 1.72 mm; n = 2) long, 2.80–2.89 (mean 2.85× as long as wide). This species is distinguished by its cylindrical appearance, elytral disc shining and convex, declivity steep, declivital interstriae 1 and 2 flat, slightly impressed, declivital interstriae 3–6 convex on basal half, with a pair of prominent tubercles on interstriae 3, apical half flattened. This species differs from *M. recidens* by the elytral disc more flattened, declivital summit less steep, declivital face more flattened and broader, and declivity armed with a pair of prominent tubercles on interstriae 3 (those much smaller in *M. recidens*).

Description (female). Body greenish brown to brown. **Head:** epistoma entire, transverse, with a row of hair-like setae. Frons weakly convex shagreened, covered with sparse vestiture, punctate, punctures shallow with a short erect hair-like setae. Eye shallowly emarginate just above antennal insertion, upper portion slightly smaller than lower part. Submentum large, distinctly triangular, moderately impressed. Antennal scape short and thick, subequal in length with club. Pedicel slightly broader than scape, shorter than funicle. Funicle 4-segmented, segment 1 shorter than pedicel. Club longer than wide (5:4), type 3, segment 1 corneous, feebly convex on anterior face, occupying basal 1/3, covering 2/3 of posterior face; segment 2 short, soft; segment 2 present on posterior face, soft. **Pronotum:** 1.20× as long as wide, elongate and subquadrate, type 7 in dorsal view, lateral sides parallel to anterior 3/4, rounded anteriorly; anterior margin without serrations; base weakly concave, posterior angles rounded. In lateral view disc slightly longer than anterior slope, type 7, summit at apical 2/5, anterior slope densely asperate; disc slightly




Figure 2. *Microperus bidentatus* sp. nov. holotype **A** lateral view, **B** dorsal view, **C** posterolateral view, **D** front, **E** head ventral view, **F** declivity.

convex, subshiny, impunctate; lateral margins obliquely costate. **Elytra:** 1.55× as long as wide, 1.33× as long as pronotum. Scutellum minute, convex, slightly raised above elytral surface. Elytral mycangium indicated by a disperse median setal tuft along elytral base. Base sinuate, edge carinate, humeral angles rounded, parallel-sided in basal 7/8, then narrowly rounded to apex. Disc shiny, moderately convex, striae not impressed, with large shallow punctures, setose, setae very short, semi-erect, hair-like; interstriae flat, impunctate, setose, setae long erect, hair-like. Declivity abruptly commencing, declivital face on striae and interstriae 1 and 2 shallowly impressed, upper portion on 3–6 convex with a pair of prominent tubercles on interstriae 3, lower portion flattened; declivital striae 1 and 2 slightly impressed, punctate, punctures round, prominent, moderately impressed, striae 3–6 flat, punctate, punctures small; interstriae flat with a row of long erect hair-like setae. Posterolateral margin carinate to interstriae 7. **Legs:** procoxae contiguous; prosternal coxal piece short, inconspicuous. Protibiae slender, broadest at the middle; posterior face smooth; margin armed with six large socketed denticles. Meso- and metatibiae rounded, armed with seven large socketed denticles.

Etymology. *L. bi* = two, *dentatus* = toothed, the name refers to the two prominent denticles on each declivital interstriae 3. An adjective.

Distribution. Southern Thailand (Trang Province, Nakhon Sri Thammarat Province).

Host plants. Unknown.

***Microperus bucolicus* Sittichaya, Smith & Beaver, sp. nov.**

<http://zoobank.org/8AD72B58-7FF5-4CD0-97C1-EC006009329A>

Figure 3

Type material. *Holotype*, female, THAILAND: Nakhon Sri Thammarat Province, 8°21'40.8"N, 99°39'17.3"E, durian orchard, ethanol baited trap, 1.vi.2011, W. Sittichaya, (MSUC).

Diagnosis. 1.6 mm (n = 1), 2.67× as long as wide. This species is distinguished by the elytral disc flat with short, steep declivity, declivital posterolateral margin costate and denticulate, declivity with sparse minor denticles, less abundant than striae punctures, and denticles uniform in size.

Similar species. *M. alpha*.

Description (female). Head, pronotum and elytra light brown, antennae and legs yellow brown. **Head:** epistoma entire, transverse, with a row of hair-like setae. Frons weakly convex to upper level of eyes, subshiny, punctate; punctures large, shallow, sparse, each bearing an erect hair-like seta. Eyes moderately emarginate just above antennal insertion, upper part smaller than lower part. Submentum large, distinctly triangular, slightly impressed. Antennal scape short and thick, shorter than club. Pedicel as wide as scape, shorter than funicle. Funicle 4-segmented, segment 1 shorter than pedicel. Club longer than wide, flattened, type 3; segment 1 corneous, transverse on anterior face, occupying basal 1/4 of club; segment 2 narrow, soft; segments 1 and 2 present on posterior face. **Pronotum:** 1.09× as long as wide. In dorsal view basic and parallel-sided (type 2), sides parallel in basal 2/3, rounded anteriorly; anterior margin without serrations. Base weakly bisinuate, posterior angles obliquely rounded. In lateral view elongate with disc much longer than anterior slope, type 8, summit low, at apical 2/5. Anterior slope with densely spaced, broad asperities, becoming lower and more strongly transverse towards summit. Disc shagreened, alutaceous, finely punctate, glabrous, some moderately long hair-like setae at margins. Lateral margins obliquely costate. **Elytra:** 1.53× as long as wide, 1.4× as long as pronotum. Scutellum minute, convex, slightly raised above elytral surface. Elytral mycangium indicated by a disperse median setal tuft along elytral base. Base transverse, edge oblique, humeral angles rounded, parallel-sided in basal 4/5, then narrowly rounded to apex. Disc flat, shiny, striae clearly impressed, with large deep punctures separated by 2–3 diameters of a puncture, glabrous; interstriae flat, impunctate, setose, setae short, sparse, erect hair-like. Declivity occupying 1/3 of elytral length, steeply rounded, its margins denticulate, weakly shagreened, subshiny; striae flat, punctate, each puncture bearing a seta as long as a puncture; interstriae regularly denticulate along their lengths, denticles uniformly sized, each bearing a long erect hair-like seta. Posterolateral margin costate, denticulate to interstriae 7. **Legs:** procoxae contiguous; prosternal coxal piece tall, pointed. Protibiae slender, broadest at the middle; posterior face smooth; margin armed with at least six large socketed denticles (broken).




Figure 3. *Microperus bucolicus* sp. nov. holotype **A** lateral view **B** dorsal view **C** posterolateral view **D** head ventral view **E** front **F** declivity.

Etymology. *L. bucolicus* = rural, rustic. In reference to the agrarian habitat in which the species was collected. An adjective.

Distribution. Southern Thailand (Nakhon Sri Thammarat Province).

Host plants. Unknown.

***Microperus globodeclivis* Sittichaya, Smith & Beaver, sp. nov.**

<http://zoobank.org/CAF6E1EF-5095-4B32-A17A-3C0ABCAAA609>

Figure 4

Type material. *Holotype*, female, THAILAND: Payao Province, Doi Phu Nang National Park, 18°51'57.7"N, 100°10'51.0"E, dry deciduous dipterocarp forest, ethanol baited trap, 01.i.2019, W. Sittichaya (NHMW).

Diagnosis. 1.6 mm long ($n = 1$), 2.66× as long as wide. This species is distinguished by its cylindrical appearance, convex elytral disc and declivity, elytral apex globose and posterolateral margin broadly rounded without a carina. This species is differentiated from the morphologically similar *M. pedellus* (Schedl, 1969) by the following combination of characters (*M. globodeclivis* given first): less elongate form, 2.66× as long as wide vs. 2.8× as long as wide), stouter pronotum 1.08× as long as wide vs. 1.17× as long as wide, elytra 1.5× as long as pronotum vs. 1.4× as long as pronotum, and elytral base sinuate vs. broadly V-shaped.

Similar species. *M. pedellus*.




Figure 4. *Microperus globodeclivis* sp. nov. holotype **A** lateral view **B** dorsal view **C** posterolateral view **D** head ventral view **E** front **F** declivity.

Description (female). Appearing bicolored, moderately setose: head, anterior slope of pronotum and elytra dark brown, remainder of pronotum, antennae, and legs light brown. **Head:** epistoma entire, transverse, with a row of dense hair-like setae. Frons weakly convex to upper level of eyes, shagreened lower part densely covered with long erect hair-like setae, punctate, punctures shallow and bearing a long, erect hair-like seta. Eye shallowly emarginated just above antennal insertion, upper and lower parts subequal. Submentum medium in size, distinctly triangular, shallowly impressed. Antennal scape short and thick, as long as club. Pedicel as wide as scape, shorter than funicle. Funicle 4-segmented, segment 1 shorter than pedicel. Club circular, broader than tall; type 3; segment 1 corneous covering about half of posterior face, anterior face costate concave, narrow; segment 2 narrowly corneous on anterior face, posterior face fully visible, segment 3 soft, visible on posterior face. **Pronotum:** 1.08× as long as wide. In dorsal view basic type 2; lateral side parallel to anterior two thirds; anterior margin round without medial serrations. Base weakly bisinuate, posterior angles acutely rounded. In lateral view type 7, disc slightly longer than anterior slope; disc convex, summit at apical 2/5; anterior slope densely covered with asperities; disc shagreened, impunctate; pronotum sparsely covered with moderately long hair-like setae, lateral margins obliquely costate. **Elytra:** 1.62× as long as wide, 1.5× as long as pronotum. Scutellum minute, convex, slightly raised above elytral surface. Elytral mycangium indicated by a disperse median setal tuft along elytral base. Base shallowly sinuate, edge oblique, humeral angles rounded. Lateral margins parallel beyond basal half,

apex broadly attenuated at apical 4/5, apex broadly rounded. Declivity gradually commencing, broadly convex without posterolateral carina; elytra setose, elytral disc shining, flat near scutellum broadly convex beyond, striae punctate, parallel laterally, puncture small, round moderately deep, punctures with shorter hair-like setae, interstriae flat, impunctate bearing a row of long erect hair-like setae, setae on disc and declivity equal in length. **Legs:** procoxae contiguous, prosternal coxal piece short, inconspicuously covered with a tuft of long hair-like setae. Protibiae slender, broadest at the middle; posterior face smooth; margin armed with six small socketed denticles. Meso- and metatibiae inflated; outer margin evenly rounded with seven and six small socketed denticles, respectively.

Etymology. *L. globus* = globe, round; *declivis* = downhill sloping (declivity). In reference to its round declivity which lacks a posterolateral carina. An adjective.

Distribution. Northern Thailand (Payao Province).

Host plants. Unknown.

***Microperus serratus* Sittichaya, Smith & Beaver, sp. nov.**

<http://zoobank.org/B8E6B1EB-89F9-466C-88B9-F4785B2FF2CE>

Figure 5

Type material. **Holotype**, female, THAILAND: Songkhla Province, Ton Nga Chang Wildlife Sanctuary, 6°59'32.1"N, 100°08'57.8"E, tropical rainforest, ethanol baited trap, 01.iii.2015, (W. Sittichaya) (NHMW). **Paratypes**, female, same data as holotype, 01.iv.2014 (1) (THNHM), 01.vii.2014 (1) (WSTC); Songkhla Province, Songkhla Zoo, 7°08'53.7"N, 100°36'29.4"E, secondary tropical rainforest, ethanol baited trap, 01.iii.2019, (1) (WSTC); Narathiwat Province, Hala-Bala Wildlife Sanctuary, 5°47'44"N, 101°50'07"E, lowland tropical rainforest, ethanol baited trap, 01.ii.2015 (1) (RABC) (1) (MSUC). MALAYSIA: Sabah, Sipitang, Mendolong, A1L, 5.iv.1988, S. Adebratt (1) (RABC); as previous except: 6.iv.1988 (1) (RABC).

Similar species. *M. nugax*, *M. sagmatus*, *M. undulatus* (Sampson, 1919).

Diagnosis. 1.66–1.72 mm long (mean = 1.70 mm; n = 6) 2.59–2.69× as long as wide (mean 2.64× as long as wide). This species is distinguished by elytral disc convex, posterior half of elytra striae impressed with large round punctures, impression increased posteriorly, interstriae costate with strong interstitial spines posterior to declivital summit, declivity steep, shallowly impressed on the middle to interstriae 2, interstriae 3 with small but prominent tubercles.

This species is related to *M. undulatus* and similar species, but elytral disc is convex, lacking a transverse or saddle-like impression.

Description. **Head:** epistoma entire, transverse, with a row of hair-like setae. Frons subshiny, finely punctate, lower portion flat sparsely covered with very fine hair-like setae, upper level of eyes slightly convex; sparsely setose, setae fine. Eye shallowly emarginate just above antennal insertion, upper part smaller than lower part. Submentum large, distinctly triangular, slightly impressed. Antennal scape short and




Figure 5. *Microperus serratus* sp. nov. holotype **A** lateral view **B** dorsal view **C** posterolateral view **D** head ventral view **E** front **F** declivity.

thick, as long as club. Pedicel as wide as scape, shorter than funicle. Funicle 4-segmented, segment 1 shorter than pedicel. Club longer than wide, type 3, anterior face segment 1 corneous, concave, segment 2 narrowly visible, posterior face segment 1 covering basal 2/3, segment 2 visible, appearing soft. **Pronotum:** 1.05× as long as wide. In dorsal view subquadrate and parallel-sided, type 3, sides parallel in basal 2/3, weakly rounded anteriorly with prominent anterolateral corners; anterior margin without a row of serrations. Base weakly bisinuate, posterior angles acutely rounded, almost subquadrate. In lateral view tall, type 2, disc flat, summit at midpoint. Anterior slope with densely spaced, broad asperities, becoming lower and more strongly transverse towards summit. Disc shagreened, alutaceous, impunctate, lateral margins obliquely costate. **Elytra:** 1.53× as long as wide, 1.53× as long as pronotum, Scutellum minute, convex, slightly raised above elytral surface. Elytral mycangium indicated by a disperse median setal tuft along elytral base. Base shallowly sinuate, edge oblique, humeral angles rounded, parallel-sided in basal 4/5, then narrowly rounded to apex. Disc shiny, basal half convex, striae and interstriae flat, posterior half rugose with deeply impressed striae and costate interstriae bearing strong incurved spines, becoming more prominent toward declivital summit which bears the largest incurved spine. Declivity steep, declivital face dull, striae and interstriae 1 and 2 shallowly impressed, striae 1 and 2 with prominent round punctures, interstriae 3 with two small acute denticles. Striae with short erect hair-like setae and interstriae with long

setae. Posterolateral margin acutely carinate to interstriae 7. **Legs:** procoxae contiguous; prosternal coxal piece tall, conical. Protibiae obliquely triangular, broadest at apical 1/3; posterior face smooth; apical 1/3 of outer margin with six moderate socketed denticles. Meso- and metatibiae flattened; outer margin evenly rounded with eight moderate to large socketed denticles.

Etymology. *L. serratus* = toothed like a saw. In reference to the serrate margin of the elytral declivity. An adjective.

Distribution. Thailand (Narathiwat Province, Songkhla Province), East Malaysia (Sabah).

Host plants. Unknown.

New combinations and new synonymy

Microperus armaticeps (Schedl, 1942), comb. nov.

Xyleborus armaticeps Schedl, 1942: 198.

Cyclorhipidion armaticeps (Schedl, 1942): Wood and Bright 1992: 698.

Remarks. A dorsal habitus image of the lectotype (NMHW) was examined by all authors and the species is transferred from *Cyclorhipidion* to *Microperus* because of the following characters: dense tuft of setae present along elytral base associated with an elytral mycangium; elytral bases broadly V-shaped, costate; scutellum minute, convex and slightly raised above elytra; pronotum from dorsal view basic and parallel sided (type 2), from lateral view disc longer than anterior slope (type 7), and pronotal base weakly sinuate.

This species is a probable synonym of *M. exsculptus* (Eggers, 1927) but has larger denticles along the declivital margin. Additional investigations are needed to assess intraspecific variation and clarify species limits.

Distribution. West Malaysia (Pahang).

Microperus cruralis (Schedl, 1975)

Xyleborus cruralis Schedl, 1975b: 456.

Cnestus cruralis (Schedl): Wood & Bright, 1992: 802.

Coptodryas cruralis (Schedl): Beaver 1995: 201.

Microperus cruralis (Schedl): Smith et al. 2020: 287.

Xyleborus myllus Browne, 1986a: 92. syn. nov.

Remarks. The holotypes of *Xyleborus myllus* (NHML) and *M. cruralis* (NHMW), and a further specimen in RABC have been compared by RAB, and found to be conspecific. *X. myllus* is accordingly here placed in synonymy with *M. cruralis*.

Distribution. Cambodia, Laos, Thailand.

***Microperus exsculptus* (Eggers, 1927), comb. nov.**

Figure 6

Xyleborus exsculptus Eggers, 1927: 101.*Coptodryas exsculptus* (Eggers, 1927): Wood and Bright 1992: 824.*Xyleborus dentipennis* Browne, 1983: 558. syn. nov.**Similar species.** *M. armaticeps*, *M. cruralis* (Schedl, 1975).

Diagnosis. 2.12 mm long [1.9–2.0 mm Browne (1983)], 2.4× as long as wide. Robust and globular in appearance. This species is distinguished by its large size, stout and robust elytral disc broadly, deeply transversely impressed with a saddle-like depression from scutellum to declivital base; elytral declivity deeply sulcate, its margins begin at interstriae 2 and margined by large tubercles on interstriae 2–6, tubercles on interstriae 1 absent; declivity dull, punctures on declivity small and indistinct; elytral bases slightly emarginated from sutural margin to interstriae 4 to accommodate mycangial tuft, mycangial tuft setae long, very dense; and posterolateral costa absent.

This species is distinguished from *M. cruralis* by its smaller size, elytral bases slightly emarginated from sutural margin to interstriae 4 to accommodate mycangial tuft, elytral disc more deeply depressed, deeper posteriorly, declivital summit without a pair of large




Figure 6. *Microperus exsculptus* (Eggers, 1927) comb. nov. **A** dorsal view **B** lateral view.

tubercles on interstriae 1, margins of declivital sulcus begin at interstriae 2, tubercles on declivital margin slender and acute, and punctures on declivity small and indistinct.

New records. Brunei: Kuala Belalong FSC, 116.7°E, 4.34°N, 270 m alt, dipterocarp forest, aerial FIT 4, 17.vi.91, N. Mawdsley (1) (RABC). Thailand: Narathiwat Province, Hala-Bala Wildlife Sanctuary, 5°47'44"N, 101°50'07"E, lowland tropical rainforest, ethanol baited trap, 01.viii.2015, (1) (W. Sittichaya) (WSTC).

Distribution. Brunei, East Malaysia (Sarawak), Philippines, Thailand.

Host plants. This species was collected from a seraya log (*Shorea* sp.) (Dipterocarpaceae) imported to Japan (Browne 1983).

Remarks. The holotype of *Xyleborus dentipennis* (NHML) and images of the *M. exsculptus* lectotype (NMNH) have been examined and compared by RAB and determined to be conspecific. Accordingly, *X. dentipennis* is here placed in synonymy. The species is transferred to *Microperus* based on the characters given for *M. armaticeps*.

***Microperus pedellus* (Schedl, 1969), comb. nov.**

Xyleborus pedellus Schedl, 1969: 231.

Coptodryas pedellus (Schedl, 1969): Wood and Bright 1992: 826.

Remarks. The paratype deposited in NHMW was examined by the senior author. This species is included in *Microperus* with the following combination of characters: antennal club truncate, type 2, segment 2 visible on posterior side, dense tuft of setae present along elytral base associated with an elytral mycangium; elytral bases broadly V-shaped, costate; scutellum minute, convex and slightly raised above elytra; pronotum from dorsal view basic and parallel sided (type 2), from lateral view disc longer than anterior slope (type 7), and pronotal base weakly sinuate.

Distribution. Philippines.

***Microperus spicatus* (Browne, 1986), comb. nov., stat. res.**

Xyleborus spicatus Browne, 1986b: 667.

Remarks. This species was previously considered a synonym of *Coptodryas dentipennis* (= *M. exsculptus*) by Bright and Skidmore (1997: 165). The holotypes of *Xyleborus spicatus* and *X. dentipennis* (NHML) and further specimens in NHML and RAB, and images of the *M. exsculptus* lectotype (NMNH), have been compared by RAB, and *X. spicatus* was found to be distinct. It is here recognized as a distinct species and transferred to *Microperus* based on the generic characters given for *M. armaticeps*. *Microperus spicatus* can be distinguished from *M. exsculptus* by the following combination of characters (*M. spicatus* given first): body elongate 3.0–3.1× as long as wide, vs. 2.1–2.2×; larger size (2.2–2.4 mm vs. 1.85–2.0 mm); elytral disc without a discal depression vs. with a discal depression; the number of teeth on each declivital margin 3–5 vs. 7–8.

New records. [Indonesia]: Sumatra, ex *Dipterocarpus* s. [no further data] [1 specimen in NHML determined by F. G. Browne as *Xyleborus dentipennis* Browne, but not that species.] Laos: Sekong prov. ca. 12 km S Seking, Tao Faek waterfalls, 15°14.7'N, 106°45.1'E, 118 m, at light, 8–12.v.2010, J. Hájek (1) (RABC).

Distribution. Indonesia (Sumatra), Laos, East Malaysia (Sabah).

Newly recorded species for Thailand

Microperus chrysophylli (Eggers, 1930)

Figure 7

Xyleborus chrysophylli Eggers, 1930: 205.

Coptodryas chrysophylli (Eggers): Wood and Bright 1992: 823.

Microperus chrysophylli (Eggers): Saha and Maiti 1996: 824.

Diagnosis. 2.52–2.6 mm long (2.56 mm; n = 2) 2.83–2.86× as long as wide [(2.6–2.7 mm long (mean = 2.68 mm; n = 5); 2.6–2.7× as long as wide (Smith et al. 2020)]; robust elongate in appearance. This species is distinguished by the elytral disc flat; declivity long, gradual; large size; declivital interstriae 2 lacking granules on declivital face; declivital face strongly shagreened, weakly impressed along striae 2 and interstriae



Figure 7. *Microperus chrysophylli* (Eggers, 1930) **A** dorsal view **B** lateral view.

2; declivital stria punctures small, indistinct; posterolateral costa granulate; interstriae densely covered with long erect hair-like setae, setae longer than two interstitial widths; and striae setose, setae short, semi-recumbent, as long as stria width.

This species strongly resembles *M. corporaali* and is distinguished by the less strongly sulcate declivity, and declivital stria punctures very large and distinct.

New record. Thailand: Chiang Mai Province, Doi Inthanon National Park, Chomthong District, 18°31'39"N, 98°30'00"E, 1650 msl, Hill evergreen forest, ethanol-baited trap, 01.v.2019 (2) (W. Sittichaya) (WSTC).

Distribution. Bangladesh, China (Yunnan), India (West Bengal) (Smith et al. 2020).

Host plants. This species is recorded from *Cinnamomum* (Lauraceae), *Chrysophyllum* (Sapotaceae), (Maiti and Saha 2004), and *Hevea brasiliensis* (Euphorbiaceae) (Smith et al. 2020).

Microperus nanus (Browne, 1949)

Figure 8

Cryptoxyleborus nanus Browne, 1949: 903.

Xyleborus caelator Browne, 1955: 354 (unnecessary replacement name).

Microperus nanus (Browne): Beaver and Hulcr 2008: 151.



Figure 8. *Microperus nanus* (Browne, 1949) **A** dorsal view **B** lateral view.

Diagnosis. 1.12–1.24 mm (mean = 1.19 mm; n = 6), 2.8–3.0× as long as wide; minute species. This species is distinguished by antennal club type 2, segment 2 visible on anterior side, elytra tapering, declivity obliquely attenuate without posterolateral carina, and elytral posterior margin acute.

New record. Thailand: Narathiwat Province, Hala-Bala Wildlife Sanctuary, 5°47'44"N, 101°50'07"E, lowland tropical rainforest, ethanol-baited trap, 01.xii.2014 (3), 01.iii.2015 (2), (W. Sittichaya) (WSTC).

Distribution. Brunei Darussalam, East Malaysia (Sabah, Sarawak), West Malaysia.

Host plants. This species is only known from Dipterocarp hosts including *Shorea* sp., *Hopea ferrea*, and *Parashorea malaanonan* (Browne 1961; Beaver and Browne 1979). It is unusual in breeding between the bark and wood, and not making galleries in the xylem (Browne 1961).

Microperus quercicola (Eggers, 1926)

Figure 9

Xyleborus quercicola Eggers, 1926: 146.

Microperus quercicola (Eggers): Smith et al. 2018: 396.

Xyleborus izuensis Murayama, 1952: 16. Synonymy: Smith et al. 2018: 396.



Figure 9. *Microperus quercicola* (Eggers, 1926) **A** dorsal view **B** lateral view.

Diagnosis. 1.8–1.92 (mean = 1.85 mm; n = 4) 2.50–2.53× as long as wide [1.8–2.0 mm long (mean = 1.96 mm; n = 5); 2.38–2.86× as long as wide (Smith et al. 2020)]. This species is distinguished by the elytral disc flat; declivity short, steep; declivity granulate from base to apex, granules small, as abundant as stria punctures; granules dense, separated by the width of one granule; declivital surface shiny; posterolateral costa strongly carinate; interstriae densely setose, setae fine, hair-like as long as the width of an interstria; and stria punctures setose, setae recumbent, hair-like, less than a stria width.

New records. Thailand: Narathiwat Province, Hala-Bala Wildlife Sanctuary, 5°47'44"N, 101°50'07"E, lowland tropical rainforest, 01.xii.2017 (1 WSTC) ethanol-baited trap; Songkhla Province, Ton Nga Chang Wildlife Sanctuary, tropical rainforest, 6°59'32.1"N, 100°08'57.8"E, 01.ii.2015 (2), ethanol-baited trap, (all W. Sittichaya) (WSTC).

Distribution. China (Guizhou, Hong Kong, Jiangxi, Sichuan, Zhejiang), Japan, Russia (Far East), South Korea, Taiwan.

Host plants. This species is polyphagous and has been recorded from *Cinnamomum* (Lauraceae) (Murayama 1952), *Diospyros* (Ebenaceae), *Fraxinus* (Oleaceae), *Carpinus* (Betulaceae) (Mandelshtam et al. 2018) and “oak trees” (Fagaceae) (Eggers 1926).

Discussion

Key to *Microperus* species of Indochinese Peninsula and China including species found in Thailand (females only) [modified from Smith et al. (2020)]

- 1 Elytral disc broadly, deeply transversely impressed with a saddle-like depression from scutellum to declivital base; declivity deeply sulcate, its margins costate; elytral bases slightly emarginated to accommodate a prominent and dense mycangial tuft **2**
- Elytral disc either medially impressed and appearing humped, or flat, or broadly convex; declivity round or flat or convex; elytral bases not emarginated and mycangial tuft minute, either not readily apparent or lightly setose **3**
- 2 Elytral base emargination broadly recurved; elytral disc shallowly depressed; declivital summit armed with a pair of large tubercles at interstriae 1; punctures on declivity distinct and broad; larger species, 3.0–3.1 mm ***cruralis***
- Elytral base emargination broadly V-shaped; elytral disc deeply depressed, more deeper posteriorly; interstriae 1 on declivital summit without a pair of large tubercles; punctures on declivital face small and indistinct; smaller species, 2.12 mm ***exsculptus***
- 3 Posterolateral margin of elytra smoothly rounded, without any indication of a costa or carina **4**
- Posterolateral margin of elytra marked by an elevated costa or carina **5**

- 4 Elytral lateral margin tapering posteriorly from the middle to apex; elytral apex acute; minute species, 1.12–1.24 mm *nanus*
- Elytral lateral margin subparallel, broadly tapering posteriorly; apex broadly rounded, not acute; larger species, 1.60 mm *globodeclivis* sp. nov.
- 5 Declivity obliquely truncate; posterolateral declivital margin rounded and denticulate **6**
- Declivity rounded; posterolateral declivital margin costate or carinate, with or without granules **9**
- 6 Declivital interstriae 2 and 3 strongly laterally broadened from base to declivital midpoint and then narrowing towards apex *latesalebrinus*
- Declivital interstriae parallel from base to apex, never laterally broadened **7**
- 7 Denticles on declivital summit and margins larger and more sharply acute than those on declivital face *kirishimanus*
- Denticles on declivital summit of equal size and shape as those on declivital face **8**
- 8 Denticles on declivital summit as dense as those on declivital face; declivital face opalescent, subshiny *nudibrevis*
- Denticles on declivital summit denser than those on declivital face; declivital face shagreened, dull *perparvus*
- 9 Larger, 2.55–2.95 mm **10**
- Smaller, 1.2–2.1 mm **12**
- 10 Stout, 1.93–2.19× as long as wide; elytral posterolateral margin strongly carinate and unarmed *fulvulus*
- Elongate, 2.5–2.9× as long as wide; elytral posterolateral margin costate and granulate **11**
- 11 Declivital strial punctures very large, distinct *chrysophylli*
- Declivital strial punctures small, indistinct *corporaali*
- 12 Declivity with granules, denticles or tubercles distinctly less abundant than strial punctures **13**
- Declivity with abundant granules or denticles, at least as abundant as strial punctures **19**
- 13 Elytral disc shallowly transversely impressed with a saddle-like impression ... **14**
- Elytral disc without a depression **15**
- 14 Discal impression deeper, antero-posteriorly narrower, with steeper anterior and posterior slopes, strial punctures on impression with rounded granules; interstitial spines on disc behind impression stronger and backwardly hooked *sagmatus*
- Discal impression shallower, antero-posteriorly broader, with gentler anterior and posterior slopes, strial punctures on impression without granules; interstitial tubercles on disc behind impression moderate with rounded apices pointing dorsally *undulatus*

- 15 Declivital tubercles uniformly sized..... **16**
 – Declivital tubercles not uniformly sized, 1–2 pairs of larger tubercles on declivital interstriae 3 **17**
 16 Posterolateral declivital margin carinate, unarmed..... *alpha*
 – Posterolateral declivital margin costate and denticulate *bucolicus* **sp. nov.**
 17 Declivity steeply rounded, declivital interstriae 1 bearing 1–2 small tubercles on declivital face; larger, 1.9–2.0 mm *recidens*
 – Declivity gradually rounded, declivital interstriae 1 unarmed on declivital face; smaller, 1.66–1.76 mm **18**
 18 Declivital interstriae 3 with two pairs of prominent denticles; posterior half of elytral disc rugose with deeply impressed striae and costate interstriae, interstriae bearing strong incurved spines that become more prominent toward declivital summit *serratus* **sp. nov.**
 – Declivital interstriae 3 with one pair of prominent tubercles; posterior half of elytral disc smooth, striae and interstriae flush, interstriae with a few small tubercles at summit..... *bidentatus* **sp. nov.**
 19 Elytral disc convex on basal 1/3, appearing humped in lateral view..... **20**
 – Elytral disc flat, never appearing humped..... **22**
 20 Declivital interstriae densely covered with short semi-erect scales
 *kadoyamaensis*
 – Declivital interstriae densely covered with long fine, erect hair-like setae... **21**
 21 Antennal club flat, type 3 with two sutures visible on apical 1/3 of posterior face; larger, declivity smooth, shiny; larger, 1.95–2.0 mm and more elongate, 2.79–2.86× as long as wide *minax*
 – Antennal club obliquely truncate, type 2 with segment 1 almost covering posterior face (Fig. 2); one suture visible on posterior face near apex; declivity shagreened, dull; smaller, 1.8–1.9 mm and less elongate, 2.57–2.71× as long as wide..... *nugax*
 22 Antennal club flat, type 3 with two sutures visible on apical 1/3 of posterior face (Fig. 3 *diversicolor*
 – Antennal club obliquely truncate, type 2 with segment 1 almost covering posterior face; one suture visible on posterior face near apex **23**
 23 Declivital interstitial granules dispersed, separated by the width of at least three granules; posterolateral margin of declivity weakly carinate and granulate; interstitial vestiture consisting of short semi-erect bristles, shorter in length than the width of an interstria; smaller, 1.2–1.7 mm..... *pometianus*
 – Declivital interstitial granules dense, separated by the width of one granule; posterolateral margin of declivity strongly carinate; interstitial vestiture consisting of long semi-erect hair-like setae, longer in length than the width of an interstria (easily abraded); larger, 1.8–2.0 mm *quercicola*

Our study increased the diversity of *Microperus* to 35 species of which 18 are recorded in Thailand. Since 2010, *Microperus* has received considerable attention with

Table 1. Synoptic list of the *Microperus* fauna of Thailand.

Species	First Record	Thai distribution
<i>Microperus alpha</i> (Beeson, 1929)	Hutachareern et al. (2007)	C: Nakhon Nayok; N: Chiang Mai, Phetchabun; N-E: Chaiyaphum, Nakhon Ratchasima, Sakhon Nakhon
<i>Microperus bidentatus</i> Sittichaya, Smith and Beaver sp. nov.	This publication	S: Songkhla, Trang
<i>Microperus bucolicus</i> Sittichaya, Smith and Beaver sp. nov.	This publication	S: Nakhon Sri Thammarat
<i>Microperus chrysophylli</i> (Eggers, 1930)	New record	N: Chiangmai
<i>Microperus corporaali</i> (Eggers, 1923)	Sittichaya et al. (2012)	S: Chumphon, Nakhon Sri Thammarat
<i>Microperus diversicolor</i> (Eggers, 1923)	Browne (1980)	S: Nakhon Sri Thammarat
<i>Microperus exculptus</i> (Eggers, 1927)	New record	S: Narathiwat
<i>Microperus globodeclivis</i> Sittichaya, Smith and Beaver sp. nov.	This publication	N: Phayao
<i>Microperus nanus</i> (Browne, 1949)	New record	S: Narathawat
<i>Microperus nudibrevis</i> (Schedl, 1942)	Beaver et al. (2014)	C: Suphanburi; N: Chiang Mai; N-E: Chaiyaphum; S: Nakhon Sri Thammarat, Surat Thani
<i>Microperus nugax</i> (Schedl, 1939)	Smith et al. (2020)	C: Chanthaburi; S: Chumphon, Nakhon Sri Thammarat, Surat Thani
<i>Microperus perparvus</i> (Sampson, 1922)	Beaver et al. (2014)	C: Nakhon Nayok; N: Chiang Mai, Mae Hong Son, Nan, Phetchabun, Phitsanulok; N-E: Chaiyaphum, Loei; S: Nakhon Sri Thammarat, Surat Thani
<i>Microperus pometianus</i> (Schedl, 1939)	Smith et al. (2020)	N: Chiang Mai
<i>Microperus quercicola</i> (Eggers, 1926)	New record	S: Songkhla, Narathiwat
<i>Microperus recidens</i> (Sampson, 1923)	Beaver et al. (2014)	C: Phetchaburi; N: Chiang Mai, Nan; S: Chumphon, Nakhon Sri Thammarat
<i>Microperus sagmatus</i> Smith, Beaver & Cognato (2020)	Smith et al. (2020)	S: Chumphon, Nakhon Sri Thammarat, Songkhla, Surat Thani
<i>Microperus serratus</i> Sittichaya, Smith and Beaver sp. nov.	This publication	S: Songkhla, Narathiwat
<i>Microperus undulatus</i> (Sampson, 1919)	Beaver et al. (2014)	C: Prachuab Khiri Khan, Uthaithani; N: Chiang Mai; N-E: Chaiyaphum; S: Chumphon, Nakhon Sri Thammarat, Songkhla, Surat Thani, Trang

eight species described and 12 species transferred from a diversity of genera including *Xyleborus*, *Coptodryas* and *Cyclorhipidion* (Beaver and Liu 2010; Hulcr and Cognato 2013; Beaver et al. 2014; Smith et al. 2018; Park et al. 2020; Smith et al. 2020). However, the faunal emphasis in these studies has largely focused on the faunas of Papua New Guinea, Indochina and the Palearctic. Three of the four new species described herein, and 14 of the 18 Thai species are recorded from the Southern Region (Table 1) which is characterized by species with affinities to the Indo-Malayan and Malayan faunas (Beaver et al. 2014). Based on our findings the *Microperus* faunas of these regions are in need of considerable revision and are undoubtedly more diverse than currently known.

Acknowledgements

We are most grateful to Dr. H. Schillhammer (NHMW) for access to specimens. Special thanks also go to the members of National Parks, Wildlife Sanctuaries and Songkhla Zoo for their facilitation of insect collecting. This work was supported by the Prince of Songkla University annual government statement of expenditure under the Plant Genetic Conservation Project under the Royal initiative of Her Royal Highness

Princess Maha Chakri Sirindhorn Year 2019 and Thailand Research Fund (TRF), project number DBG- 6180023 to WS and SMS was funded in part, by a Cooperative Agreement (AP21PPQS&T00C034) from the United States Department of Agriculture's Animal and Plant Health Inspection Service (APHIS). It may not necessarily express APHIS' views.

References

- Beaver RA, Browne FG (1979) The Scolytidae and Platypodidae (Coleoptera) of Penang, Malaysia. *Oriental Insects* 12: 575–624. <https://doi.org/10.1080/00305316.1978.10432538>
- Beaver RA (1995) New synonymy and taxonomic changes in Oriental and Australasian Scolytidae and Platypodidae (Coleoptera). *Annalen des Naturhistorischen Museums in Wien*, 97B: 197–204.
- Beaver RA (1999) New records of bark and ambrosia beetles from Thailand (Coleoptera: Scolytidae). *Serangga* 4: 175–183.
- Beaver RA, Hulcr J (2008) A review of the ambrosia beetle genus *Cryptoxyleborus* Schedl (Coleoptera: Curculionidae: Scolytinae). *The Coleopterists Bulletin* 62: 133–153. <https://doi.org/10.1649/1026.1>
- Beaver RA, Liu L-Y (2010) An annotated synopsis of Taiwanese bark and ambrosia beetles, with new synonymy, new combinations and new records (Coleoptera: Curculionidae: Scolytinae). *Zootaxa* 2602: 1–47. <https://doi.org/10.11646/zootaxa.2602.1.1>
- Beaver RA, Sittichaya W, Liu L-Y (2014) A synopsis of the scolytine ambrosia beetles of Thailand (Coleoptera: Curculionidae: Scolytinae). *Zootaxa* 3875: 001–082. <http://dx.doi.org/10.11646/zootaxa.3875.1.1>
- Bright DE, Skidmore RE (1997) A Catalog of Scolytidae and Platypodidae (Coleoptera), Supplement 1 (1990–1994). NRC Research Press, Ottawa, 368 pp.
- Browne FG (1949) Notes on Malayan Scolytoidea (Coleoptera) with descriptions of new species. *The Annals and Magazine of Natural History*, series 12(1): 892–912. <https://doi.org/10.1080/00222934808653953>
- Browne FG (1955) Synonymy and descriptions of some oriental Scolytidae and Platypodidae (Coleoptera). *Sarawak Museum, Journal* 6: 343–373.
- Browne FG (1961) The biology of Malayan Scolytidae and Platypodidae. *Malayan Forest Records* 22: 1–255.
- Browne FG (1980) Bark beetles and ambrosia beetles (Coleoptera, Scolytidae and Platypodidae) intercepted at Japanese ports, with descriptions of new species, III. *Kontyû* 48: 482–489.
- Browne FG (1983) Bark beetles and ambrosia beetles (Coleoptera, Scolytidae and Platypodidae) intercepted at Japanese ports, with descriptions of new species, VII. *Kontyû* 51: 554–572.
- Browne FG (1986a) Bark beetles and ambrosia beetles (Coleoptera, Scolytidae and Platypodidae) intercepted at Japanese ports, with descriptions of new species, XII. *Kontyû* 54: 89–99.
- Browne FG (1986b) Bark beetles and ambrosia beetles (Coleoptera, Scolytidae and Platypodidae) intercepted at Japanese ports, with descriptions of new species, XV. *Kontyû* 54: 661–671.

- Cognato AI, Hulcr J, Dole SA, Jordal BH (2011) Phylogeny of haplo-diploid, fungus-growing ambrosia beetles (Curculionidae: Scolytinae: Xyleborini) inferred from molecular and morphological data. *Zoologica Scripta* 40: 174–186. <https://doi.org/10.1111/j.1463-6409.2010.00466.x>
- Eggers H (1926) Japanische Borkenkäfer, I. *Entomologische Blätter* 22: 145–148.
- Eggers H (1927) Neue Indo-Malayische Borkenkäfer (Ipidae). II. Nachtrag. *The Philippine Journal of Science* 33: 67–108.
- Eggers H (1930) Neue *Xyleborus*-Arten (Col. Scolytidae) aus Indien. *Indian Forest Records, Entomology Series* 14: 177–208.
- Eggers H (1940) Neue indomalayische Borkenkäfer (Ipidae) III Nachtrag (Forstsetzung). *Tijdschrift voor Entomologie* 83: 132–154.
- Hulcr J, Dole SA, Beaver RA, Cognato AI (2007) Cladistic review of generic taxonomic characters in Xyleborina (Coleoptera: Curculionidae: Scolytinae). *Systematic Entomology* 32: 568–584. <https://doi.org/10.1111/j.1365-3113.2007.00386.x>
- Hulcr J, Cognato AI (2013) Xyleborini of New Guinea: A Taxonomic Monograph. Thomas Say Publications in Entomology, Entomological Society of America, Lanham, 176 pp.
- Hutacharern C, Tubtim N, Dokmai C (2007) Checklist of insects and mites in Thailand. Department of National Parks, Wildlife and Plant Conservation, Bangkok, 319 pp.
- Maiti PK, Saha N (2004) Fauna of India and the Adjacent Countries. Scolytidae: Coleoptera (Bark- and Ambrosia-beetles) (Vol. 1). Part-1. Introduction and Tribe Xyleborini. *Zoological Survey of India, Kolkata*, 268 pp.
- Mandelsham MYu, Yakushkin EA, Petrov AV (2018) Oriental ambrosia beetles (Coleoptera: Curculionidae: Scolytinae): new inhabitants of Primorsky krai in Russia. *Russian Journal of Biological Invasions* 9: 355–365. <https://doi.org/10.1134/S2075111718040082>
- Mandelsham MYu, Petrov AV, Smith SM, Cognato AI (2019) Resurrection of *Heteroborips* Reitter, 1913 (Coleoptera: Curculionidae: Scolytinae) from synonymy with *Xyleborus* Eichhoff, 1864. *The Coleopterists Bulletin* 73: 387–394. <https://doi.org/10.1649/0010-065X-73.2.387>
- Murayama J (1952) Notes on the scolytid-beetles (Coleoptera) from southern and western parts of Izu Peninsula, Shizuoka Prefecture. *Bulletin of the Faculty of Agriculture, Yamaguchi University* 3: 15–23.
- Nobuchi A (1983) The Scolytidae (Coleoptera) of the Philippines. *JARQ* Vol. 16, No. 4: 296–305.
- Nunberg M, Chûjō M (1961) Coleoptera from South-East Asia. 25. Scolytidae, 26. Platypodidae. *Nature and Life in South-East Asia* 1: e358.
- Park S, Smith SM, Cognato AI, Beaver RA (2020) Catalogue of Korean Xyleborine ambrosia beetles (Coleoptera: Curculionidae) with seven new species. *Journal of Asia-Pacific Biodiversity* 13: 210–228. <https://doi.org/10.1016/j.japb.2020.01.002>
- Saha N, Maiti PK (1996) Insecta: Coleoptera: Scolytidae. *Fauna of West Bengal* 6B: 775–866.
- Schedl KE (1939) Scolytidae and Platypodidae 47 Beitrag. *Tijdschrift voor Entomologie* 82: 30–53.
- Schedl KE (1942) Forschungsberichte zur Scolytoiden-Fauna der Malayischen Halbinsel, V. 80. Beitrag zur Morphologie und Systematik der Scolytoidea. *Kolonialforstliche Mitteilungen* 5: 169–218.
- Schedl KE (1969) Bark beetles and pinhole borers (Scolytidae and Platypodidae) intercepted from imported logs in Japanese ports. III. *Kontyû* 37: 202–219.

- Schedl KE (1975) Indian bark and timber beetles VI. 312. Contribution to the morphology and taxonomy of the Scolytoidea. *Revue Suisse de Zoologie* 82: 445–458. <https://doi.org/10.5962/bhl.part.78268>
- Sittichaya W (2012) Bark and ambrosia beetles (Coleoptera: Curculionidae: Scolytinae and Platypodinae) infesting mango trees (*Mangifera indica* L.) in Southern Thailand, with new species recorded for Thailand. *Songklanakarin Journal of Science and Technology* 34: 153–155.
- Smith SM, Beaver RA, Cognato AI (2018) New synonymy, new combinations and taxonomic changes in Japanese xyleborine ambrosia beetles (Coleoptera: Curculionidae: Scolytinae). *Zootaxa* 4521: 391–403. <https://doi.org/10.11646/zootaxa.4521.3.5>
- Smith SM, Beaver RA, Cognato AI (2020) A monograph of the Xyleborini (Coleoptera, Curculionidae, Scolytinae) of the Indochinese Peninsula (except Malaysia) and China. *ZooKeys* 983: 1–442. <https://doi.org/10.3897/zookeys.983.52630>
- Wood SL (1980) New genera and new generic synonymy in Scolytidae (Coleoptera). *Great Basin Naturalist* 40: 89–97. <https://doi.org/10.5962/bhl.part.20006>
- Wood SL (1992) Nomenclatural changes and new species of Platypodidae and Scolytidae (Coleoptera), part II. *The Great Basin Naturalist* 52: 78–88.
- Wood SL, Bright DE (1992) A catalog of Scolytidae and Platypodidae (Coleoptera), Part 2: Taxonomic index. *The Great Basin Naturalist Memoirs* 13: 1–1553.