


# Self-management strategies to consider to combat endometriosis symptoms during the COVID-19 pandemic

Mathew Leonardi <sup>1,2,3,\*</sup>, Andrew W. Horne <sup>4</sup>, Katy Vincent <sup>5</sup>, Justin Sinclair <sup>6</sup>, Kerry A. Sherman <sup>7</sup>, Donna Ciccia<sup>8</sup>, George Condous <sup>1,2</sup>, Neil P. Johnson<sup>9,10,11</sup>, and Mike Armour <sup>6,12</sup>

<sup>1</sup>Acute Gynaecology, Early Pregnancy, and Advanced Endoscopy Surgery Unit, Nepean Hospital, Kingswood, Australia <sup>2</sup>Sydney Medical School Nepean, University of Sydney, Sydney, Australia <sup>3</sup>Department of Obstetrics and Gynecology, McMaster University, Hamilton, Canada <sup>4</sup>MRC Centre for Reproductive Health, University of Edinburgh, Edinburgh, UK <sup>5</sup>Nuffield Department of Women's and Reproductive Health, University of Oxford, John Radcliffe Hospital, Oxford, UK <sup>6</sup>NICM Health Research Institute, Western Sydney University, Penrith, Australia <sup>7</sup>Department of Psychology, Centre for Emotional Health, Macquarie University, North Ryde, Australia <sup>8</sup>Endometriosis Australia, Sydney South, Australia <sup>9</sup>Robinson Research Institute, University of Adelaide, Adelaide, Australia <sup>10</sup>Department of Obstetrics and Gynaecology, University of Auckland, Auckland, New Zealand <sup>11</sup>Auckland Gynaecology Group and Repromed Auckland, Auckland, New Zealand <sup>12</sup>Translational Health Research Institute (THRI); Western Sydney University, Penrith, Australia

\*Correspondence address. Acute Gynaecology, Early Pregnancy, and Advanced Endoscopy Surgery Unit, Sydney Medical School Nepean, Nepean Hospital, University of Sydney, Sydney 2747 NSW, Australia. Tel: +61 2 4734 4777; Fax: +61 2 4734 4887; E-mail: mathew.leonardi@sydney.edu.au  <https://orcid.org/0000-0001-5538-6906>

Submitted on April 18, 2020; resubmitted on April 30, 2020; editorial decision on May 4, 2020

**ABSTRACT:** The care of patients with endometriosis has been complicated by the coronavirus disease 2019 (COVID-19) pandemic. Medical and allied healthcare appointments and surgeries are being temporarily postponed. Mandatory self-isolation has created new obstacles for individuals with endometriosis seeking pain relief and improvement in their quality of life. Anxieties may be heightened by concerns over whether endometriosis may be an underlying condition that could predispose to severe COVID-19 infection and what constitutes an appropriate indication for presentation for urgent treatment in the epidemic. Furthermore, the restrictions imposed due to COVID-19 can impose negative psychological effects, which patients with endometriosis may be more prone to already. In combination with medical therapies, or as an alternative, we encourage patients to consider self-management strategies to combat endometriosis symptoms during the COVID-19 pandemic. These self-management strategies are divided into problem-focused and emotion-focused strategies, with the former aiming to change the environment to alleviate pain, and the latter address the psychology of living with endometriosis. We put forward this guidance, which is based on evidence and expert opinion, for healthcare providers to utilize during their consultations with patients via telephone or video. Patients may also independently use this article as an educational resource. The strategies discussed are not exclusively restricted to consideration during the COVID-19 pandemic. Most have been researched before this period of time and all will continue to be a part of the biopsychological approach to endometriosis long after COVID-19 restrictions are lifted.

**Key words:** endometriosis / pelvic pain / COVID-19 / coronavirus / self-management / self-care / mind-body therapies / counseling / lifestyle management / psychology

## Introduction

The coronavirus disease 2019 (COVID-19) pandemic, and the resultant required self-isolation, is having a huge impact on the care of patients with endometriosis worldwide. With the support of

several gynecological associations (AAGL – Elevating Gynecologic Surgery, 2020), many medical centers have temporarily stopped offering surgical management for endometriosis. Similarly, many outpatient appointments for endometriosis with family physicians, gynecologists, fertility specialists and/or other allied healthcare

## WHAT DOES THIS MEAN FOR PATIENTS?

This is an article that aims to help patients with endometriosis-related pain symptoms through the difficult obstacles posed by coronavirus disease 2019 (COVID-19). Across the world, care for patients with endometriosis has changed dramatically: patients may be unable to visit their doctors or allied healthcare providers in person or obtain imaging tests, surgery or fertility treatments. Patients with endometriosis may be experiencing increased stress during this time.

In addition to the more typical treatments for endometriosis, such as drug treatments or surgery, it is widely accepted that complementary or alternative treatments should be used. We encourage patients and healthcare providers to more strongly consider using these strategies at this time. Some examples of those that aim to directly target the problem of endometriosis-related pain are improving sleep hygiene, low-intensity physical activity (including pelvic exercises, yoga), dietary changes, application of heat and medical cannabis. Other examples target the emotional aspect of endometriosis-related pain and these include relaxation/mindfulness, acceptance of chronic illness (e.g. via Acceptance and Commitment Therapy (ACT)) and improving a balance toward positive attitude.

Some patients may already be using these strategies during normal times, but many will not have been counseled on them or advised to use them. Some of these strategies could be implemented directly by patients whilst others may require the guidance of healthcare providers through telephone and online avenues. Initiating any of these strategies now to combat endometriosis symptoms may prove beneficial for patients and may encourage ongoing use once life gets back to normal.

professionals (e.g. physiotherapists, psychologists, dieticians) have been canceled or postponed. In addition, the International Society of Ultrasound in Obstetrics and Gynecology has recommended postponing ultrasound evaluation of non-acute pelvic pain (Bourne et al., 2020). In addition to the effects of restrictions on healthcare services, patients with endometriosis, like everyone else, are at risk of negative psychological effects from mandated self-isolation (Brooks et al., 2020). To offset the direct healthcare obstacles, we hope that as much as possible, patients are still able to connect with their healthcare providers by telephone or virtually online. We encourage healthcare providers and patients alike to consider medical management options that they have not yet trialed and that are within reach during this period of self-isolation despite the closure of services they may otherwise access. In combination with medical therapies or as an alternative, we encourage patients to consider self-management strategies that may be effective in combating the negative physical or psychological effects triggered by COVID-19 restrictions. The importance of integrating self-management strategies into the broad biopsychological approach to endometriosis is not novel (Buggio et al., 2017) nor will it be exclusively relevant during this time. The necessity of long-term and multimodal treatments for endometriosis, including the self-management strategies described here, means the usefulness of the content of this article will extend beyond the current situation. We have therefore prepared the following expert opinion-based guidance on self-management strategies for patients to consider using during self-isolation and beyond (Table I).

### Problem-focused strategies

Research in psychology provides a useful framework for understanding how an individual with endometriosis will respond to a serious health threat such as COVID-19. The transactional model of stress and coping (Lazarus and Folkman, 1987; Folkman and Moskowitz, 2004) indicates that the way any person adapts to this kind of stressor varies according to how they think about the stressor and the actions and behaviors they initiate to manage the demands of the situation; that is,

the pain and challenges of living with endometriosis (Folkman and Moskowitz, 2004). If one adopts active, engagement coping strategies (Kvillemo and Bränström, 2014), such as problem-solving, information-seeking, acceptance, seeking support and planning, one is more likely to effectively cope with the ongoing stresses of living with endometriosis. On the other hand, disengagement coping strategies that focus attention away from thoughts and feelings about endometriosis, such as denial, substance use and self-blame, are likely to heighten feelings of being out of control and being distressed (Dijkstra and Homan, 2016). Women with endometriosis often use both problem- and emotion-based coping strategies (Roomaney and Kagee, 2016). Women with endometriosis who engage in positive, problem-focused coping strategies report lower stress levels and less depressive symptoms, which results in less non-cyclical pelvic pain and dysmenorrhea compared to those with maladaptive coping strategies (Donatti et al., 2017). The key here is for individuals with endometriosis to adopt self-management and self-care strategies that will enhance feelings of being in control and minimize feelings of worry in the face of the additional serious health threat of COVID-19. While medical and/or surgical intervention in endometriosis may be effective (Leonardi et al., 2020), they are not a 'magic bullet' and do not stand alone. Encouragement from healthcare providers to implement other self-management strategies should improve the confidence patients have in those strategies. Below, we review a series of active and self-directed strategies that may assist in coping with endometriosis in the face of COVID-19.

### Education about endometriosis

Having a coherent understanding of endometriosis is an important step in being able to effectively self-manage this condition, as is the case with other chronic diseases such as diabetes and heart disease (Griva et al., 2000; Weinman et al., 2000). While awareness of endometriosis is building amongst the general population thanks to efforts by advocacy organizations and academics working in the field, endometriosis remains an enigmatic disease. Many patients learn of the word 'endometriosis' at the time of consultation with their healthcare provider as a possible explanation for their

**Table 1** Self-management strategies for patients with endometriosis to consider using during self-isolation and beyond, and recommendation statements.

Strategy	Recommendation
Problem-focused strategies	
Education about endometriosis	Patients could access information from reputable organizations that have demonstrated clear dedication to advancing the care of those with endometriosis.
Managing work and study	Patients could be honest with employers/educators about anticipated and possibly unanticipated episodes of pain. Patients could discuss modifications of work/school obligations during times of pain exacerbation, if appropriate.
Social activities	Patients should make a strong effort to maintain social activities (in the context of current isolation rules). Patients could use video conferencing applications, social media portals or websites to best replace their typical social activities.
Social support	Patients should consider connecting with their social supports by video link instead of messaging or the telephone. Patients could consider joining online endometriosis support groups.
Sleep	Patients should talk to their family physician/general practitioner about dysfunctional sleep or improving sleep before initiating over-the-counter treatments.
Physical exercise	Patients should initiate or maintain exercise based on their personal experiences and take a graduated approach to increase physical activity. Patients could access a large selection of online material to support exercise activities.
Yoga	Patients could initiate or maintain an at-home yoga program, guided by online material.
Pelvic exercise and physiotherapy	Patients could consider at-home pelvic exercises but should do so with caution and most ideally with the past/current guidance of a physiotherapist.
Diet	Patients could consider trialing modifications in their diet. Possible considerations include: anti-inflammatory diet; gluten-free diet; low FODMAP diet.
Heat	Patients could consider trialing stick-on heat patches that provide a stable temperature for long durations (8–12 h at a time).
Cannabis	Patients could consider medicinal cannabis as a potential option if they live in a location with legal access.
Emotion-focused strategies	
Relaxation and mindfulness	Patients should consider trialing relaxation, breathing exercises and mindfulness. Patients could access online material to support relaxation through websites or smartphone applications.
Acceptance and commitment therapy	Patients could consider self-initiating aspects of Acceptance and Commitment Therapy using online resources. Alternative methods to learn how to 'accept' endometriosis could be utilized.
Adopting a positive attitude	Patients could consider self-initiating the process of learning to adopt a positive attitude using online resources and social support (e.g. online endometriosis groups).

FODMAP, fermentable oligosaccharides, disaccharides, monosaccharides, and polyols.

symptoms. Patients may be given a diagnosis following clinical assessment (Agarwal *et al.*, 2019), imaging (Nisenblat *et al.*, 2016) or surgery (Wykes *et al.*, 2004). Depending on how they receive a diagnosis and by whom they are diagnosed, the information they receive may be inconsistent and ambiguous. Endometriosis education in schools has recently been introduced in some countries to raise awareness early and standardize messaging (Bush *et al.*, 2017), but this has not been successfully attempted with adults. The tangible nature of endometriosis means that education is often focused on what can be seen (i.e. endometriosis deposits, endometriomas) rather than on the invisible downstream effects like chronic pelvic pain and central sensitization.

Learning more about endometriosis independently has been useful for some individuals (Roomaney and Kagee, 2016). Education on the role of the nervous system in endometriosis would be helpful for patients to understand how the various non-medical and non-surgical treatments may be useful. Knowing how and where to access patient-friendly information can be challenging.

### Recommendation

Beyond the education delivered to patients directly by healthcare providers, we recommend guiding patients to reputable organizations that have demonstrated clear dedication to advancing the care of individuals with endometriosis. For some during self-isolation due to COVID-19, there may be more opportunities to educate themselves and so this should be especially encouraged. Sources of education may include university websites, where 'Information for patients' can be found (EXPPECT, 2020a) or other reputable sources, such as regional or national advocacy organizations, which often involve collaboration between healthcare providers and patient representatives (Endometriosis Australia, 2020; Endometriosis New Zealand, 2020; Endometriosis UK, 2020; The Endometriosis Network Canada, 2020).

### Work and study

During normal circumstances, episodes of escalated pain (frequently often associated with the menstrual cycle) lead to an inability to complete work/school obligations resulting in loss of productivity (Soliman

*et al.*, 2016). Working from home (WFH) or attending school virtually has become a new normal for many individuals. This may create unfamiliar challenges for reporting sickness that limits one's ability to complete work/school obligations. People may be sensing a change in attitude on what warrants a 'sick day' since WFH no longer requires commuting or the same degree of human interaction.

#### *Recommendation*

During this unprecedented time, we should endeavor to be empathetic of new obstacles (e.g. WFH with children, exacerbations of mental health issues, the necessity to devote extra time to ensuring financial stability) that may lead to difficulty fulfilling typical obligations. Honesty with employers and educators about anticipated and possibly unanticipated episodes of endometriosis-related pain may help to set expectations and decrease anxiety when it is necessary to request a sick day (Roomaney and Kagee, 2016). This period of time could also be an opportunity to discuss modifications of work/school obligations during times of pain exacerbation, if appropriate, and deemed possible by the person experiencing the troublesome symptoms. As a positive, it may demonstrate that removing the commute and having a comfortable environment allows the woman to be adequately productive, meaning a sick day is not required.

### Social activities

During normal circumstances, episodes of escalated pain (or associated menstrual symptoms) or the development of chronic pelvic pain may necessitate social isolation (Mellado *et al.*, 2016). In the past, social activities, such as going to the theatre, having coffee with friends or hosting a games night, might have been canceled or postponed due to pain. Now, during mandatory social isolation, these activities and others are no longer viable. While social isolation is being mandated broadly globally due to COVID-19 and this might seem advantageous to those with endometriosis, isolation is recognized as a contributor to psychosocial issues, especially amongst those who are vulnerable (Usher *et al.*, 2020). A frequent strategy for those suffering from endometriosis is to schedule social events during more reliable periods of decreased or no pain.

#### *Recommendation*

More than ever, making a concerted effort to maintain social activities (in the context of current isolation rules) is vital. Using video conferencing applications and social media portals should be considered as a replacement for in-person gatherings. Many arts and culture organizations are providing access to material online. For those living with others, honesty about when and why they may be able or unable to socialize could be helpful. As a positive, this experience may teach us ways to connect that do not require leaving the house, which may facilitate a greater sense of social connectivity for women unable to leave the house in the future due to endometriosis-associated symptoms.

### Social support

During the pandemic, social support networks will undoubtedly change in appearance, but their function and critical importance to help with coping with the dual stressors of endometriosis and COVID-19 should not be underestimated. The World Endometriosis Society has long

espoused the value of endometriosis support groups in providing a valuable forum for mutual sharing of experiences and quality of life improvement through teaching coping mechanisms (Johnson and Hummelshoj, 2013). As noted above, utilizing technology to fill the void of in-person social interaction will be essential. However, there is an increasingly common belief that social media use is negatively associated with health-related outcomes (Shakya and Christakis, 2017). While there is certainly evidence for a concerning association, there is a suggestion that how one uses social media matters more than the quantity of use (Bekalu *et al.*, 2019; Killam, 2020).

#### *Recommendation*

If possible, patients should consider connecting with their social supports by video instead of messaging or the telephone. This method of communication allows for the next best thing to in-person interaction. There is also a plethora of established online support through social media websites such as Twitter and Facebook. An example is the 'Endometriosis Australia' Facebook page, which is, in part, a social network for those living with endometriosis where they may get the emotional support they need from others who similarly suffer from endometriosis (Endometriosis Australia, 2020). Similar groups may exist locally and patients can be pointed in the direction of their local or national endometriosis advocacy organization.

### Sleep

Sleep plays an important role in regulating an individual's quality of life and social, emotional and physical well-being. Dysfunction in the physiology of sleep is common in those suffering from chronic pain (Smith and Haythornthwaite, 2004). To face the COVID-19 pandemic, ensuring good sleep practices is even more important because sleep can modulate physical and mental health. Thinking and planning about sleep are relevant as our routines have changed, whereby some individuals may be WFH or studying from home, possibly even in their bedroom. Individuals with endometriosis are prone to issues with sleep (Nunes *et al.*, 2015; Leone Roberti Maggiore *et al.*, 2017). Arion *et al.* (2020) found that overall quality of sleep was poorer in those also reporting more severe pain, poorer functional quality of life on the Endometriosis Health Profile-30 questionnaire, and more depression symptoms. Thus far, the literature on sleep dysfunction in women with endometriosis is limited. However, one randomized controlled trial (RCT) implemented an administration of 10 mg of daily melatonin, which was found to improve the quality of sleep, reduce pain scores and reduce analgesia use (Schwertner *et al.*, 2013).

#### *Recommendation*

Adult patients should attempt to obtain 7–9 h of sleep each night (Hirshkowitz *et al.*, 2015). Mindfulness activities may be helpful in falling asleep. Since sleep can be affected by many factors, patients should try to assess what factors are immediately modifiable (e.g. surrounding noise, light, distractions in bed). For extra support, the National Sleep Foundation has produced ready-to-use patient guidelines, which can be found here: <https://www.sleepfoundation.org/sleep-guidelines-covid-19-isolation#guidelines> (National Sleep Foundation, 2020). Talking to one's family physician about dysfunctional sleep or improving sleep is recommended before initiating over-the-counter agents such as melatonin. Online sleep programs based on sound scientific approaches (cognitive behavioral therapy (CBT) for insomnia (Horsch

*et al.*, 2017)) are also available and may be of benefit to those women with significant sleep disruption.

## Physical exercise

Exercise is increasingly being recommended to manage a range of chronic pain conditions (Geneen *et al.*, 2017), including primary dysmenorrhea (Armour *et al.*, 2019a) and endometriosis (Awad *et al.*, 2017; Mira *et al.*, 2018). A recent systematic review found that regular exercise in women with primary dysmenorrhea, irrespective of intensity, reduced menstrual pain severity (Armour *et al.*, 2019a). Conversely, a focused systematic review on endometriosis and physical exercise found inconclusive results from only six studies (Bonoche *et al.*, 2014). Armour *et al.* (2019b) have found in a survey of Australian women that exercise led to episodes of exacerbated pelvic pain, more 'flare-ups', and increased fatigue in approximately one-third of respondents.

### Recommendation

The limited research on endometriosis and general physical activity makes formulating a recommendation difficult. However, during the COVID-19 pandemic, outdoor exercise remains one of the few reasons people can leave their homes, which undoubtedly will have an overall positive impact on well-being, for everyone (Fallon, 2020). For those with endometriosis, we suggest using previous and current personal experiences to judge what exercise is appropriate and to take a graduated approach to advance this exercise, setting achievable goals. There is a wide selection of exercise types that could be considered.

## Yoga

In a survey of Australian women, Armour *et al.* (2019b) reported 35% of 372 participants used yoga/Pilates to self-manage their endometriosis symptoms. An RCT by Gonçalves *et al.* (2017) found that yoga was associated with a reduction in levels of chronic pelvic pain and an improvement in the quality of life in women with endometriosis. Similarly, yoga therapy in the forms of asanas, pranayama and relaxation has been shown to decrease pain intensity and improve the quality of life for women with chronic pelvic pain.

### Recommendation

Patients should be aware that at-home yoga led by an online instructor may not replicate the formal 8-week program utilized by Gonçalves and colleagues, but may still be useful. A simple search of YouTube for 'yoga for endometriosis' identifies many videos and patients may wish to engage in a telehealth style consultation with a trained yoga therapist if they wish to have more personalized information.

## Pelvic exercises and physiotherapy

Pelvic physiotherapy is increasingly being recognized as an important component of the multidisciplinary treatment approach to endometriosis (Ball and Khan, 2020; Finklee *et al.*, 2020), especially when there is coexisting chronic pelvic pain. The British Pain Society (2013) considers a physiotherapist an essential member of the multidisciplinary team for patients with chronic pain. While evidence on the efficacy for endometriosis specifically is scarce (Fuentes-Márquez *et al.*, 2019; Klotz *et al.*, 2019), there is evidence that patients are seeking this intervention (Schwartz *et al.*, 2019).

### Recommendation

Pelvic physiotherapy is not something a patient can initiate on their own at home while in isolation. However, if patients have established relationships with pelvic physiotherapists, they could consider connecting with that individual to receive advice on what they may achieve at home while in isolation. In some cases, often due to funding limitations for the services of physiotherapy, patients are given instructions or lessons to perform at home, minimizing the number of in-person appointments necessary. Patients should consider implementing these strategies whilst at home. Patients can be directed to this simple pelvic exercise program recommended by the patient organization 'Endometriosis UK' (Supplementary Data S1). Besides this document, which has been approved by experts in physiotherapy and contains important instructions on safety, patients should be cautious about using videos without previous consultation with a physiotherapist. In some cases, the pelvic floor muscles can already be hypertonic or non-relaxed, requiring relaxation and coordination of pelvic floor muscles rather than strengthening (Faubion *et al.*, 2012). Hence, Kegel-based programs should be avoided unless otherwise specified by one's physiotherapist.

## Diet

A recent review of the complex relationship between diet and endometriosis highlights many of the studies that have evaluated this concept (Saguyod *et al.*, 2018). Though often limited by bias intrinsic in the data collected from observational studies and the complexity of the subject being evaluated, there is evidence that there may be an increase in endometriosis risk with increased red meat consumption (Yamamoto *et al.*, 2018) and a decrease in risk with increased consumption of fish oils, green vegetables, fruits (especially citrus fruits) (Harris *et al.*, 2018) and dairy products (Missmer *et al.*, 2010; Harris *et al.*, 2013; Nodler *et al.*, 2020). However, we do not currently have evidence that altering the consumption of these potential risk factors once endometriosis has occurred leads to any symptomatic improvements. There is some evidence for the following dietary modifications that may improve symptoms:

- dietary factors (e.g. green tea, resveratrol, fish oil, soy isoflavones) with known anti-inflammatory effects have been shown to mitigate disease progression, assessed as lesion size in mouse models;
- there may be a decreased risk for advanced endometriosis stage and/or reduction in pain symptoms in afflicted patients who use an anti-inflammatory diet;
- a gluten-free diet showed a significant positive effect in reducing pain symptoms in endometriosis patients (Marziali *et al.*, 2012).

In addition, a low fermentable oligosaccharides, disaccharides, monosaccharides and polyols (known as FODMAP) diet may reduce the gastrointestinal symptoms, such as bloating, that many women with endometriosis report (Moore *et al.*, 2017). Armour *et al.* (2019b) found that 18% of survey respondents who used a specific diet reported being able to reduce their endometriosis-related medication by 50% or more. In some cases, it is unclear if there are confounding reasons for improvements (e.g. coexisting celiac disease in the case of a gluten-free diet). There is increasing evidence that visceral pain comorbidities (e.g. bowel or bladder conditions) can exacerbate chronic pain conditions and vice-versa (Costantini *et al.*, 2017). As

such, mitigating visceral pain triggers may have a positive downstream effect on endometriosis-associated pain. While these findings are interesting, and in some cases promising, there is no strong evidence for dietary interventions improving endometriosis symptoms.

There is some preliminary evidence that dietary supplements may be beneficial in the treatment of endometriosis. A small observational cohort study found that, compared to no treatment, daily consumption of 1800 mg of N-acetylcysteine for 3 months was associated with a greater likelihood of canceling planned laparoscopic surgery compared to the no treatment group (Porpora et al., 2013). Similarly, a small RCT of melatonin (10 mg per day) for 8 weeks improved non-cyclical pelvic pain and dysmenorrhea while reducing analgesic consumption (Schwertner et al., 2013). Resveratrol, commonly found in significant quantities in grapes, wine and berries, may also assist in the reduction of endometriosis-related pain, however, the results from clinical trials in humans are mixed (Dull et al., 2019). Finally, palmitoylethanolamide, in combination with a fatty acid and myrrh, taken twice daily for 6 months resulted in a reduction of non-cyclical pelvic pain and dysmenorrhea, however, there was no placebo group so caution is advised (De Leo et al., 2019).

#### Recommendation

It is too early to make formal recommendations on dietary changes that will improve the lives of those living with endometriosis. Nevertheless, simple advice like ensuring adequate fiber and fluid intake to avoid constipation and avoiding known triggers of bladder symptoms (personal to the individual but may include alcohol, acidic fruits, etc.) may be useful. The potential positive impact of dietary improvements extends far beyond endometriosis. Therefore, patients may consider experimenting as the risk of adverse events is low and there may be a potential benefit, especially during COVID-19 where spending more time eating at home may allow more opportunities to modify dietary intake. New practices that one may initiate during COVID-19 self-isolation to combat endometriosis-pain could lead to longer-lasting positive health effects.

## Heat

There are no RCTs directly looking at the effectiveness of heat in endometriosis. Despite this, heat is commonly used and rated as effective for endometriosis-related pain (Armour et al., 2019b). Heat therapy has been shown to be effective in primary dysmenorrhea, where heat therapy using adhesive heat patches on the lower abdomen showed a moderate improvement in pain intensity compared to placebo patches, placebo pills or no treatment and a small to moderate improvement compared to ibuprofen (Armour et al., 2019c). While these are promising findings, several caveats may reduce heat's effectiveness in real-world use compared to clinical trials. Most women in the community are unlikely to be able to maintain constant heat at 38–40°C for 8–12 h using heat packs or hot water bottles, which are the most common forms used (O'Connell et al., 2006). Stick-on heat patches that provide stable temperatures for long periods of time may be the safest and most effective form of sustained heat.

#### Recommendation

More evidence is necessary at this time, but patients can consider trialing adhesive heat patches that provide a stable temperature for long durations (8–12 h at a time).

## Medicinal cannabis

Similar to heat, there are currently no RCTs that have investigated the efficacy of medicinal cannabis for endometriosis; however, emergent evidence for its use in managing chronic pain has been reported (Klimkiewicz and Jasinska, 2018). There exists a plausible rationale for the use of medicinal cannabis in endometriosis-associated pain and concurrent symptoms via modulation of the endocannabinoid system, which can impact neuropathic and inflammatory pain, along with the psychological experience of the pain (Bouaziz et al., 2017). A recent survey of Australian women with confirmed endometriosis noted that 1 in 10 respondents was utilizing cannabis, for symptom management, with cannabis rated as the highest self-management strategy for the pain of all those assessed (Armour et al., 2019b). Furthermore, significant improvements in sleep, nausea, gastrointestinal upset, anxiety and depression were self-reported, along with 56% of women in the cannabis-using cohort being able to reduce their pharmaceutical usage by 50% or more (Sinclair et al., 2020). Medicinal cannabis is not risk-free but the majority of reported adverse events tend to be mild and self-limiting (Lintzeris et al., 2018). Potential cannabis dependence (reported in 4–9% of users) (Anthony et al., 1994; Volkow et al., 2014) and possible psychosis must also be considered in heavy, chronic users, although whether this applies to medicinal products, as it is in illicit use, is unclear at this time.

#### Recommendation

Whilst more evidence is required for its use in the endometriosis population, women who are not able to access their normal medical or surgical treatment during this time may wish to consider medicinal cannabis as a potential option if they live in a location with legal access. Ideally, the use of medicinal cannabis should be discussed with one's family physician or pain specialist through telephone or video consultation before initiation.

## Emotion-focused strategies

Health anxiety that interferes with everyday life can be heightened in individuals with pre-existing chronic conditions, such as endometriosis. Along with the problem-based self-management strategies, individuals with endometriosis must adopt strategies to manage negative emotional responses to these threats posed by COVID-19. These strategies are always important, but there is a heightened need for adopting these strategies during this time. There are many self-help options to assist with managing these concerns, along with readily available support from psychologists and counselors. Consequently, several strategies that address the negative psychological effects of living with endometriosis exist and may be helpful for patients (Roomaney and Kagee, 2016). It is the responsibility of healthcare providers to guide patients through these emotion-focused, non-medical and non-surgical strategies. Some of these approaches are outlined below.

### Relaxation and mindfulness

A systematic review of psychological and mind–body interventions for endometriosis highlighted several studies that identified a significant reduction in pain when patients were exposed to physical therapy with CBT, yoga, biofeedback, mindfulness and psychotherapy,

psychotherapy combined with acupuncture and progressive muscle relaxation (Evans *et al.*, 2019). Not all of these interventions are relevant to self-management at home during isolation, but some are within reach of patients. Moreover, this category of treatment is included and encouraged by The British Pain Society (2013) guidelines on pain management for adults. In a survey of Australian women, 47% of 372 participants used meditation/breathing to self-manage their endometriosis symptoms (Armour *et al.*, 2019b).

Performing mindfulness meditation is not a skill most people naturally have so it must be learned and practiced. From YouTube (San Bruno, CA, USA) to smartphone applications, to innumerable websites, the resources necessary are accessible and sometimes free. A Danish research group has published two studies on mindfulness, the second being a 6-year follow-up study (Kold *et al.*, 2012; Hansen *et al.*, 2017). While only 10 participants were recruited, they found significant improvements in participants' pain level, well-being and ability to function in daily life after the first study (Kold *et al.*, 2012). Six years later, 9 out of the 10 women were still using the mindfulness-techniques and other mental techniques learned during the original study, and seven of these nine women experienced benefits from these techniques to a great or very great extent (Hansen *et al.*, 2017).

### Recommendation

During self-isolation, relaxation, breathing exercises and mindfulness seem to be helpful to patients with endometriosis. Progressive muscle relaxation and breathing exercises can be undertaken at home using a variety of websites and/or audio downloads. One example can be found at (Beyond Blue, 2020): <https://www.beyondblue.org.au/get-support/staying-well/relaxation-exercises>.

More evidence is needed to support this as an effective intervention in the context of endometriosis, however, there is a body of evidence in chronic pain more generally (Ball *et al.*, 2017; Hilton *et al.*, 2017), but in the absence of harmful side effects, these strategies could be safely trialed by patients. Patients can access online material through websites or smartphone applications. If health-related anxiety and worries are interfering with a patient's everyday life, then it is critical that they are referred for professional support from a health professional trained in appropriate psychotherapy (i.e. psychologist). There are telehealth options in many jurisdictions providing high-quality psychological support, using therapeutic techniques such as CBT (focusing on identifying and changing unhelpful thought patterns, and thereby changing behaviors) and ACT (discussed below) that can be effective at managing health-related negative emotions.

## Acceptance

The concept of accepting chronic illness goes well beyond the sphere of endometriosis. ACT is a form of evidence-based psychotherapy that aims to increase psychological flexibility (Wetherell *et al.*, 2011). In other words, the goal is to give patients the knowledge to recognize internal and external triggering events or experiences (e.g. pain) and equipping them with appropriate and useful thoughts and behaviors (Kuba and Weißflog, 2017). ACT aims to provide the following skills (Hayes *et al.*, 2012):

- accept automatic thoughts, sensations and urges;
- defuse from thinking (i.e. observe thoughts without believing them or following their directions);

- experience self as context (i.e. a continuous, stable sense of self as an observer of psychological experiences);
- attend to the present moment with self-awareness;
- clearly articulate values (i.e. self-chosen, desirable ways of behaving);
- engage in committed action (i.e. participating in values-consistent activities, even when psychologically challenging).

### Recommendation

Telephone and video consultation options will enable individuals with endometriosis to access support from psychologists and counselors trained in these techniques, and basic information about this therapeutic approach is freely available in the first two chapters of 'ACT made simple' by Dr Harris (2009). As with CBT, it is important to stress that these therapies are not designed as self-help approaches, and should be utilized only in consultation with a trained health professional (e.g. psychologist). Moreover, freely available online therapy delivered and supported by psychologists for managing general anxiety and depression may be available. For example, in Australia support can be accessed online through the MindSpot Clinic (2020). While the literature on ACT for endometriosis is currently scarce (Hällstam *et al.*, 2018), there is an ongoing RCT that includes therapy based on ACT as intervention (Forman *et al.*, 2016), and some evidence that ACT may be of some help in managing chronic pain, generally (Öst, 2014).

## Adopting a positive attitude

Some patients report needing to make a conscious effort to positively adjusting their attitude, almost as a last resort (Roomaney and Kagee, 2016). The concept of attitude in chronic illness is not new: correlation between optimism and well-being in patients with chronic illness has been investigated with findings suggestive that optimism improves well-being (Gustavsson-Lilius *et al.*, 2012; Hurt *et al.*, 2014). Though no studies have been completed assessing how a change in attitude in patients with endometriosis impacts outcomes, there has been a study that found a lower level of optimism in women with endometriosis compared to those without (Morán-Sánchez *et al.*, 2020). Recently, the Survey of Pain Attitudes (SOPA) was tested and demonstrated reliability and validity in a sample of women with chronic pain due to endometriosis (Ferreira-Valente *et al.*, 2019). This is important as the SOPA tool could be used to measure pain-related beliefs in future studies. It would be ideal to have scientific evidence to support or refute the suggestion to adopt a positive attitude to improve well-being. However, in the absence of this evidence, the default advice will surely be 'adopt a positive attitude'. Similar to the concept of *accepting the disease*, trying to adopt a positive attitude may not be so straightforward without guidance. Maintaining a remote relationship with established psychological healthcare providers may allow this to be a focus, but establishing new connections and building a rapport may not be possible.

### Recommendation

Like the other self-management strategies, it may be within reach of patients to self-initiate this process by finding and reading resources online. Utilizing social supports, particularly online endometriosis community groups, may also aid in motivating patients to achieve this goal (EXPPECT, 2020b).

## Discussion

Our review article briefly touches on the various problem-focused and emotions-focused strategies that are non-medication and non-surgical, which patients with endometriosis and their healthcare providers could discuss and consider implementing during the pandemic of COVID-19. These strategies could be attempted in combination with medical management strategies that may still be prescribed via telephone or video consultations. In addition, regardless of whether exacerbated by endometriosis or not, the COVID-19 restrictions may trigger negative psychological effects (Brooks et al., 2020) in patients with endometriosis, which require diverse and multimodal strategies. The strategies discussed here may have merit not necessarily because they specifically target endometriosis, but because they target overall well-being. In relation to endometriosis, the literature on these self-management strategies is scarce and/or limited by lower methodological quality. In many cases, no RCTs have been completed within endometriosis populations to test the efficacy or safety of the strategies. For some (e.g. diet, sleep, mindfulness), the likelihood and severity of possible adverse events are low enough that the potential benefits would outweigh the risk. These may be more reasonable to attempt as novel strategies and patients could use their judgment and independent research to guide their decisions. For other strategies (e.g. pelvic floor exercises, cannabis), the risks are greater or they require guidance by healthcare providers. As such, greater caution should be used in initiating these strategies, though maintenance may be easier if the patient had initiated them before COVID-19 and had a period of adequate supervision or training.

For many of the problem-focused strategies, access to electricity, technological devices and the internet is required. Not all individuals may have access or the knowledge to navigate online resources. Some individuals may also experience difficulty accessing other necessities such as food to fulfill new diet trials or exercise equipment to complete at-home exercises. Unfortunately, providing solutions to this larger-scale disparity is beyond the aim of the article, which hopes to guide those who may read this article themselves or via their healthcare providers whom they may still be seeing through the telephone/virtual consultations.

While the postponement of surgeries and the limits to visiting healthcare providers for diagnosis and management of endometriosis are tragic, this is a time to seek alternative solutions. While they may be initiated as 'temporary bandages' until the traditional options resurface, some individuals may find the lessons learned and outcomes experienced valuable, resulting in long-lasting changes to how their endometriosis is managed. Besides, in many ways, the strategies laid out here have possible benefits for general well-being and non-endometriosis conditions that may be useful during the present global scenario of COVID-19, which is undoubtedly eliciting new stressors.

It should be noted that this article is not meant to prescribe specific self-management strategies to patients, but rather provide education so patients may discuss options with their healthcare providers or make independent decisions that might be right for them. Healthcare providers may also direct their patients to the following article lay summary on 'Medical and self-care advice for women with endometriosis during the COVID-19 pandemic', which has been recently published by Endometriosis UK (Supplementary Data S2).

## Supplementary data

Supplementary data are available at *Human Reproduction Open* online.

## Acknowledgments

We would like to acknowledge the input of Mrs Tiffany Marlow, pelvic floor physiotherapist of Bluewater Pelvic Health Centre in Ontario, Canada.

## Authors' roles

All authors meet justification criteria of authorship as per ICMJE that is they all provided substantial contributions to conception and design, or acquisition of data, or analysis and interpretation of data, were involved in drafting the article or revising it critically for important intellectual content, gave final approval of the version to be published and agreed to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

## Funding

No external funding was either sought or obtained for this study.

## Conflict of interest

Dr M.L. reports grants from Australian Women and Children's Research Foundation, outside the submitted work. Prof. A.W.H. reports grants from Chief Scientist Office, grants from NIHR EME, grants from MRC, grants from Wellbeing of Women, grants from Ferring, other from Roche Diagnostics, during the conduct of the study; other from Honoraria for consultancy for Ferring, Roche and AbbVie, outside the submitted work. Prof. K.V. reports grants from IMI2, NIHR Bayer Healthcare outside the submitted work; other from Bayer Healthcare, Grunenthal GmbH, Eli Lilly and AbbVie outside the submitted work. Mr J.S. reports other from BioCeuticals, personal fees from Australian Medicinal Cannabis Research and Education Collaboration (AMCREC), grants from BioCeuticals/Blackmores, other from United in Compassion, outside the submitted work. Prof. G.C. reports personal fees from Roche, personal fees from GE Healthcare, grants from Australian Women and Children's Research Foundation, outside the submitted work. Prof. N.P.J. reports personal fees from Guerbet, personal fees from Vifor Pharma, grants and personal fees from Myovant Sciences, grants from AbbVie, personal fees from Roche, outside the submitted work. Dr M.A. reports grants from Metagenics and Spectrum outside the submitted work. Other authors report nothing to disclose.

## References

AAGL – Elevating Gynecologic Surgery. *COVID-19: Joint Statement on Elective Surgeries*. 2020. <https://www.aagl.org/news/covid-19-joint-statement-on-elective-surgeries/> (5 April 2020, date last accessed).



- Agarwal SK, Chapron C, Giudice LC, Laufer MR, Leyland N, Missmer SA, Singh SS, Taylor HS. Clinical diagnosis of endometriosis: a call to action. *Am J Obstet Gynecol* 2019;**220**:354.e1–354.e12.
- Anthony JC, Warner LA, Kessler RC. Comparative epidemiology of dependence on tobacco, alcohol, controlled substances, and inhalants: basic findings from the National Comorbidity Survey. *Exp Clin Psychopharmacol* 1994;**2**:244–268.
- Arion K, Orr NL, Noga H, Allaire C, Williams C, Bedaiwy MA, Yong PJ. A quantitative analysis of sleep quality in women with endometriosis. *J Women's Heal* 2020. doi:10.1089/jwh.2019.8008.
- Armour M, Ee CC, Naidoo D, Ayati Z, Chalmers KJ, Steel KA, de Manincor MJ, Delshad E. Exercise for dysmenorrhoea. *Cochrane Database Syst Rev* 2019a;**19**:CD004142. doi:10.1002/14651858.CD004142.
- Armour M, Sinclair J, Chalmers KJ, Smith CA. Self-management strategies amongst Australian women with endometriosis: a national online survey. *BMC Complement Altern Med* 2019b;**19**:1–16. doi:10.1186/s12906-019-2431-x.
- Armour M, Smith CA, Steel KA, Macmillan F. The effectiveness of self-care and lifestyle interventions in primary dysmenorrhoea: a systematic review and meta-analysis. *BMC Complement Altern Med* 2019c;**19**:22.
- Awad E, Ahmed HAH, Yousef A, Abbas R. Efficacy of exercise on pelvic pain and posture associated with endometriosis: within subject design. *J Phys Ther Sci* 2017;**29**:2112–2115.
- Ball E, Khan KS. Recent advances in understanding and managing chronic pelvic pain in women with special consideration to endometriosis. *F1000Research* 2020;**9**:1–8.
- Ball EF, Nur Shafina Muhammad Sharizan E, Franklin G, Rogozińska, E. Does mindfulness meditation improve chronic pain? A systematic review. *Curr Opin Obstet Gynecol* 2017;**29**:359–366.
- Bekalu MA, McCloud RF, Viswanath K. Association of social media use with social well-being, positive mental health, and self-rated health: disentangling routine use from emotional connection to use. *Heal Educ Behav* 2019;**46**:69S–80S.
- Beyond Blue. *Relaxation Exercises*. 2020. <https://www.beyondblue.org.au/get-support/staying-well/relaxation-exercises> (16 April 2020, date last accessed).
- Bonoche CM, Montenegro ML, Rosa e Silva JC, Ferriani RA, Meola J. Endometriosis and physical exercises: a systematic review. *Reprod Biol Endocrinol* 2014;**12**:1–5. doi:10.1186/1477-7827-12-4.
- Bouaziz J, Bar On A, Seidman DS, Soriano D. The clinical significance of endocannabinoids in endometriosis pain management. *Cannabis Cannabinoid Res* 2017;**2**:72–80.
- Bourne T, Leonardi M, Kyriacou C, Al-Memar M, Landolfo C, Cibula D, Condous G, Metzger U, Fischerova D, Timmerman D et al. ISUOG Consensus Statement on rationalization of gynecological ultrasound services in context of SARS-CoV-2. *Ultrasound Obstet Gynecol* 2020. doi:10.1002/uog.22047.
- Brooks SK, Webster RK, Smith LE, Woodland L, Wessely S, Greenberg N, Rubin GJ. The psychological impact of quarantine and how to reduce it: rapid review of the evidence. *Lancet* 2020;**395**:912–920.
- Buggio L, Barbara G, Facchin F, Frattaruolo MP, Aimi G, Berlanda N. Self-management and psychological-sexological interventions in patients with endometriosis: strategies, outcomes, and integration into clinical care. *Int J Womens Health* 2017;**9**:281–293.
- Bush D, Brick E, East MC, Johnson N. Endometriosis education in schools: a New Zealand model examining the impact of an education program in schools on early recognition of symptoms suggesting endometriosis. *Aust New Zeal J Obstet Gynaecol* 2017;**57**:452–457.
- Costantini R, Affaitati G, Wesselmann U, Czakanski P, Giamberardino MA. Visceral pain as a triggering factor for fibromyalgia symptoms in comorbid patients. *Pain* 2017;**158**:1925–1937.
- De Leo V, Cagnacci A, Cappelli V, Biasioli A, Leonardi D, Seracchioli R. Role of a natural integrator based on lipoic acid, palmitoiletanolamide and myrrh in the treatment of chronic pelvic pain and endometriosis. *Minerva Ginecol* 2019;**71**:191–195.
- Dijkstra MTM, Homan AC. Engaging in rather than disengaging from stress: Effective coping and perceived control. *Front Psychol* 2016;**7**:1–12.
- Donatti L, Ramos DG, Andres M de P, Passman LJ, Podgaec S. Patients with endometriosis using positive coping strategies have less depression, stress and pelvic pain. *Einstein (Sao Paulo)* 2017;**15**:65–70.
- Dull AM, Moga MA, Dimienescu OG, Sechel G, Burtea V, Anastasiu CV. Therapeutic approaches of resveratrol on endometriosis via anti-inflammatory and anti-angiogenic pathways. *Molecules* 2019;**24**:1–21.
- Endometriosis Australia. 2020. <https://www.endometriosisaustralia.org/> (9 April 2020, date last accessed).
- Endometriosis New Zealand. 2020. <https://nzendo.org.nz/> (9 April 2020, date last accessed).
- Endometriosis UK. 2020. <https://www.endometriosis-uk.org/> (9 April 2020, date last accessed).
- Evans S, Fernandez S, Olive L, Payne LA, Mikocka-Walus A. Psychological and mind-body interventions for endometriosis: a systematic review. *J Psychosom Res* 2019;**124**:1–11. doi:10.1016/j.jpsychores.2019.109756.
- EXPECT. Information for Patients. 2020a. <https://www.ed.ac.uk/centre-reproductive-health/expect-endometriosis/information-for-patients> (9 April 2020, date last accessed).
- EXPECT. Alternative Ways of Managing Your Pain. 2020b. <https://www.ed.ac.uk/centre-reproductive-health/expect-endometriosis/covid19-advice-nhs-lothian-endometriosis/covid19-alternative-ways-managing-pain> (13 April 2020, date last accessed).
- Fallon K. Exercise in the time of COVID-19. *Aust J Gen Pract* 2020;**49**.
- Faubion SS, Shuster LT, Bharucha AE. Recognition and management of nonrelaxing pelvic floor dysfunction. *Mayo Clin Proc* 2012;**87**:187–193.
- Ferreira-Valente A, Garcia IQ, Rosa AM, Pereira A, Pais-Ribeiro JL, Jensen MP. The Portuguese 35-item survey of pain attitudes applied to Portuguese women with endometriosis. *Scand J Pain* 2019;**19**:553–563.
- Findeklee S, Radosa JC, Hamza A, Haj Hamoud B, Iordache I, Sklavounos P, Takacs ZF, Solomayer EF, Radosa M. Treatment algorithm for women with endometriosis in a certified Endometriosis Unit. *Minerva Ginecol* 2020;**72**:43–49.
- Folkman S, Moskowitz JT. Coping: pitfalls and promise. *Annu Rev Psychol* 2004;**55**:745–774.
- Forman A, Vase L, Hansen KE. *Psychological Pain Treatment in Endometriosis*. Identifier: NCT02761382. *ClinicalTrials.gov*. 2016.

- <https://clinicaltrials.gov/ct2/show/NCT02761382> (11 April 2020, date last accessed).
- Fuentes-Márquez P, Cabrera-Martos I, Valenza MC. Physiotherapy interventions for patients with chronic pelvic pain: a systematic review of the literature. *Physiother Theory Pract* 2019;**35**:1131–1138.
- Geneen LJ, More RA, Clarke C, Martin D, Colvin LA, Smith BH. Physical activity and exercise for chronic pain in adults. *J Social* 2017;**1**:135–139.
- Gonçalves AV, Barros NF, Bahamondes L. The practice of hatha yoga for the treatment of pain associated with endometriosis. *J Altern Complement Med* 2017;**23**:45–52.
- Griva K, Myers LB, Newman S. Illness perceptions and self efficacy beliefs in adolescents and young adults with insulin dependent diabetes mellitus. *Psychol Health* 2000;**15**:733–750.
- Gustavsson-Lilius M, Julkunen J, Keskivaara P, Lipsanen J, Hietanen P. Predictors of distress in cancer patients and their partners: the role of optimism in the sense of coherence construct. *Psychol Health* 2012;**27**:178–195.
- Hällstam A, Stålnacke BM, Svensén C, Löfgren M. Living with painful endometriosis—a struggle for coherence. A qualitative study. *Sex Reprod Healthc* 2018;**17**:97–102.
- Hansen KE, Kesmodel US, Kold M, Forman A. Long-term effects of mindfulness-based psychological intervention for coping with pain in endometriosis: a six-year follow-up on a pilot study. *Nord Psychol* 2017;**69**:100–109.
- Harris HR, Chavarro JE, Malspeis S, Willett WC, Missmer SA. Dairy-food, calcium, magnesium, and vitamin D intake and endometriosis: a prospective cohort study. *Am J Epidemiol* 2013;**177**:420–430.
- Harris HR, Eke AC, Chavarro JE, Missmer SA. Fruit and vegetable consumption and risk of endometriosis. *Hum Reprod* 2018;**33**:715–727.
- Harris R. In: C Honeychurch (ed). *ACT Made Simple: An Easy-to-Read Primer on Acceptance and Commitment Therapy*. 2009, CA: New Harbinger Publications, Inc.
- Hayes SC, Strosahl KD, Wilson KG. In: Guilford Press (ed). *Acceptance and Commitment Therapy: The Process and Practice of Mindful Change*. 2nd edn. 2012, New York, NY: Guilford Press.
- Hilton L, Hempel S, Ewing BA, Apaydin E, Xenakis L, Newberry S, Colaiaco B, Maher AR, Shanman RM, Sorbero ME et al. Mindfulness meditation for chronic pain: systematic review and meta-analysis. *Ann Behav Med* 2017;**51**:199–213.
- Hirshkowitz M, Whiton K, Albert SM, Alessi C, Bruni O, DonCarlos L, Hazen N, Herman J, Katz ES, Kheirandish-Gozal L et al. National sleep foundation's sleep time duration recommendations: methodology and results summary. *Sleep Heal* 2015;**1**:40–43.
- Horsch CHG, Lancee J, Griffioen-Both F, Spruit S, Fitrianie S, Neerincx MA, Beun RJ, Brinkman W-P. Mobile phone-delivered cognitive behavioral therapy for insomnia: a randomized waitlist controlled trial. *J Med Internet Res* 2017;**19**:e70.
- Hurt CS, Burn DJ, Hindle J, Samuel M, Wilson K, Brown RG. Thinking positively about chronic illness: an exploration of optimism, illness perceptions and well-being in patients with Parkinson's disease. *Br J Health Psychol* 2014;**19**:363–379.
- Johnson NP, Hummelshoj L; World Endometriosis Society Montpellier Consortium. Consensus on current management of endometriosis. *Hum Reprod* 2013;**28**:1552–1568.
- Killam K. How to Prevent Loneliness in a Time of Social Distancing - Scientific American. *Scientific American*. 2020. <https://www.scientificamerican.com/article/how-to-prevent-loneliness-in-a-time-of-social-distancing/> (10 April 2020, date last accessed).
- Klimkiewicz A, Jasinska A. The health effects of cannabis and cannabinoids. *Psychiatra* 2018;**15**:88–92.
- Klotz SGR, Schön M, Ketels G, Löwe B, Brünahl CA. Physiotherapy management of patients with chronic pelvic pain (CPP): a systematic review. *Physiother Theory Pract* 2019;**35**:516–532.
- Kold M, Hansen T, Vedsted-Hansen H, Forman A. Mindfulness-based psychological intervention for coping with pain in endometriosis. *Nord Psychol* 2012;**64**:2–16.
- Kuba K, Weißflog G. Acceptance and commitment therapy in the treatment of chronic disease. *Psychother Psychosom Medizinische Psychol* 2017;**67**:525–536.
- Kvillemo P, Bränström R. Coping with breast cancer: a meta-analysis. *PLoS One* 2014;**9**:1–26.
- Lazarus RS, Folkman S. Transactional theory and research on emotions and coping. *Eur J Pers* 1987;**1**:141–169.
- Leonardi M, Gibbons T, Armour M, Wang R, Glanville E, Hodgson R, Cave AE, Ong J, Tong YF, Jacobson TZ et al. When to do surgery and when not to do surgery for endometriosis: a systematic review and meta-analysis. *J Minim Invasive Gynecol* 2020;**27**:390–407.e3.
- Leone Roberti Maggiore U, Bizzarri N, Scala C, Tafi E, Siesto G, Alessandri F, Ferrero S. Symptomatic endometriosis of the posterior cul-de-sac is associated with impaired sleep quality, excessive daytime sleepiness and insomnia: a case-control study. *Eur J Obstet Gynecol Reprod Biol* 2017;**209**:39–43.
- Lintzeris N, Driels J, Elias N, Arnold JC, McGregor IS, Allsop DJ. Medicinal cannabis in Australia, 2016: the Cannabis as Medicine Survey (CAMS-16). *Med J Aust* 2018;**209**:211–216.
- Marziali M, Venza M, Lazzaro S, Lazzaro A, Micossi C, Stolfi VM. Gluten-free diet: a new strategy for management of painful endometriosis related symptoms? *Minerva Chir* 2012;**67**:499–504.
- Mellado BH, Falcone ACM, Poli-Neto OB, Rosa E Silva JC, Nogueira AA, Candido-Dos-Reis FJ. Social isolation in women with endometriosis and chronic pelvic pain. *Int J Gynecol Obstet* 2016;**133**:199–201.
- MindSpot Clinic. MindSpot. 2020. <https://mindspot.org.au/> (16 April 2020, date last accessed)
- Mira TAA, Buen MM, Borges MG, Yela DA, Benetti-Pinto CL. Systematic review and meta-analysis of complementary treatments for women with symptomatic endometriosis. *Int J Gynecol Obstet* 2018;**143**:2–9.
- Missmer SA, Chavarro JE, Malspeis S, Bertone-Johnson ER, Hornstein MD, Spiegelman D, Barbieri RL, Willett WC, Hankinson SE. A prospective study of dietary fat consumption and endometriosis risk. *Hum Reprod* 2010;**25**:1528–1535.
- Moore JS, Gibson PR, Perry RE, Burgell RE. Endometriosis in patients with irritable bowel syndrome: specific symptomatic and demographic profile, and response to the low FODMAP diet. *Aust New Zeal J Obstet Gynaecol* 2017;**57**:201–205.
- Morán-Sánchez I, Adoamnei E, Sánchez-Ferrer ML, Prieto-Sánchez MT, Areñse-Gonzalo JJ, Casanova-Mompeán V, Carmona-Barnosi A, Mendiola J, Torres-Cantero AM. Is dispositional optimism associated

- with endometriomas and deep infiltrating endometriosis? *J Psychosom Obstet Gynecol* 2020. doi:10.1080/0167482X.2020.1729732.
- National Sleep Foundation. *Sleep Guidelines and Help during the COVID-19 Pandemic*. 2020. <https://www.sleepfoundation.org/sleep-guidelines-covid-19-isolation#guidelines> (30 April 2020, date last accessed).
- Nisenblat V, Bossuyt PMM, Farquhar C, Johnson N, Hull ML. Imaging modalities for the non-invasive diagnosis of endometriosis. *Cochrane Database Syst Rev* 2016;2:1-293; CD009591. doi:10.1002/14651858.CD009591.
- Nodler JL, Harris HR, Chavarro JE, Frazier AL, Missmer SA. Dairy consumption during adolescence and endometriosis risk. *Am J Obstet Gynecol* 2020;**222**:257.e1–257.e16.
- Nunes FR, Ferreira JM, Bahamondes L. Pain threshold and sleep quality in women with endometriosis. *Eur J Pain* 2015;**19**:15–20.
- O'Connell K, Davis AR, Westhoff C. Self-treatment patterns among adolescent girls with dysmenorrhea. *J Pediatr Adolesc Gynecol* 2006;**19**:285–289.
- Öst L-G. The efficacy of acceptance and commitment therapy: an updated systematic review and meta-analysis. *Behav Res Ther* 2014;**61**:105–121.
- Porpora MG, Brunelli R, Costa G, Imperiale L, Krasnowska EK, Lundeberg T, Nofroni I, Piccioni MG, Pittaluga E, Ticino A et al. A promise in the treatment of endometriosis: an observational cohort study on ovarian endometrioma reduction by N-acetylcysteine. *Evidence-based Complement Altern Med* 2013;**2013**:1–7.
- Roomaney R, Kagee A. Coping strategies employed by women with endometriosis in a public health-care setting. *J Health Psychol* 2016;**21**:2259–2268.
- Saguyod SJU, Kelley AS, Velarde MC, Simmen RCM. Diet and endometriosis-revisiting the linkages to inflammation. *J Endometr Pelvic Pain Disord* 2018;**10**:51–58.
- Schwartz ASK, Gross E, Geraedts K, Rauchfuss M, Wölfler MM, Häberlin F, Orelli S von, Eberhard M, Imesch P, Imthurn B et al. The use of home remedies and complementary health approaches in endometriosis. *Reprod Biomed Online* 2019;**38**:260–271.
- Schwertner A, Conceição Dos Santos CC, Costa GD, Deitos A, De Souza A, De Souza ICC, Torres ILS, Da Cunha Filho JSL, Caumo W. Efficacy of melatonin in the treatment of endometriosis: a phase II, randomized, double-blind, placebo-controlled trial. *Pain* 2013;**154**:874–881.
- Shakya HB, Christakis NA. Association of facebook use with compromised well-being: a longitudinal study. *Am J Epidemiol* 2017;**185**:203–211.
- Sinclair J, Smith CA, Abbott J, Chalmers KJ, Pate DW, Armour M. Cannabis use, a self-management strategy among Australian women with endometriosis: results from a national online survey. *J Obstet Gynaecol Canada* 2020;**42**:256–261.
- Smith MT, Haythornthwaite JA. How do sleep disturbance and chronic pain inter-relate? Insights from the longitudinal and cognitive-behavioral clinical trials literature. *Sleep Med Rev* 2004;**8**:119–132.
- Soliman AM, Yang H, Du EX, Kelley C, Winkel C. The direct and indirect costs associated with endometriosis: a systematic literature review. *Hum Reprod* 2016;**31**:712–722.
- The British Pain Society. *Guidelines for Pain Management Programmes for Adults*. 2013, London: The British Pain Society.
- The Endometriosis Network Canada. 2020. <https://endometriosisnetwork.com/> (16 April 2020, date last accessed)
- Usher K, Bhullar N, Jackson D. Life in the pandemic: social isolation and mental health. *J Clin Nurs* 2020. doi:10.1111/jocn.15290.
- Volkow ND, Baler RD, Compton WM, Weiss SRB. Adverse health effects of marijuana use. *N Engl J Med* 2014;**370**:2219–2227.
- Weinman J, Petrie KJ, Sharpe N, Walker S. Causal attributions in patients and spouses following first-time myocardial infarction and subsequent lifestyle changes. *Br J Health Psychol* 2000;**5**:263–273.
- Wetherell JL, Afari N, Rutledge T, Sorrell JT, Stoddard JA, Petkus AJ, Solomon BC, Lehman DH, Liu L, Lang AJ et al. A randomized, controlled trial of acceptance and commitment therapy and cognitive-behavioral therapy for chronic pain. *Pain* 2011;**152**:2098–2107.
- Wykes CB, Clark TJ, Khan KS. Accuracy of laparoscopy in the diagnosis of endometriosis: a systematic quantitative review. *Br J Obstet Gynaecol* 2004;**111**:1204–1212.
- Yamamoto A, Harris HR, Vitonis AF, Chavarro JE, Missmer SA. A prospective cohort study of meat and fish consumption and endometriosis risk. *Am J Obstet Gynecol* 2018;**219**:178.e1–178.e10.