

## Comparison of salty taste acuity and salty taste preference with sodium intake and blood pressure based on zinc nutritional status in two rural populations in Korea

Jeong-Sook Choe<sup>1§</sup>, Eun-Kyung Kim<sup>1</sup> and Eun-Kyung Kim<sup>2</sup>

<sup>1</sup>National Academy of Agricultural Science, Rural Development Administration, 30 Seoho-ro, Suwon 441-707, Korea

<sup>2</sup>Department of Food and Nutrition, Gangneung-Wonju National University, Gangneung 210-702, Korea

### Abstract

This study examined salty taste acuity and salty taste preference and sodium intake in relation to zinc nutritional status in 2 rural populations in Korea. And we also examined the main food contributors of their sodium intakes. We enrolled 218 adults (66 men and 152 women) from the Kangneung and Samcheok regions in Korea's Kangwon province in our study conducted from December 2011 to February 2012. Participants from each region were divided into 3 groups based on their serum zinc level (T1: lowest, T2: intermediate, T3: highest). We compared the salty taste acuity and preference, Na index (Dish Frequency Questionnaire for estimation of habitual sodium intake), blood pressure, and intakes of nutrients including sodium by 3 groups of serum zinc level. The results were as follows: a higher serum zinc level indicated a lower sodium intake and Na index ( $P < 0.05$ ). The salty taste acuity was considerably higher for participants from the Kangneung region than those from the Samcheok region ( $P < 0.05$ ). And the serum zinc level was significantly higher in participants from the Kangneung region than those from the Samcheok region ( $P < 0.05$ ). We further divided the participants into 2 groups: those who consumed more zinc than the recommended intake (RI) and the others. We compared salty taste acuity and salty taste preference in the 2 groups. The salty taste threshold and palatable salty taste concentrations were lower for the group with a zinc intake above RI than for the group with zinc intake below the RI. However, the difference was not significant. This study confirms that taste function differs depending on zinc nutritional status. In future, it is required to a large-scale, long-term, prospective study on the correlation between zinc intake, serum zinc levels, and taste perception function and blood pressure.

**Key Words:** Zinc nutritional status, salty taste acuity, salty taste preference, sodium intake, blood pressure

### Introduction

According to the 2010 Korean National Health and Nutrition Examination Survey, the prevalence of high blood pressure in the Korean population of above the age of 30 years is 28.9%. Since 1998, the prevalence of high blood pressure has declined, and the awareness, then treatment rate, the control rate have increased [1]. However, statistics on the causes of death from the Korea National Statistical Office show that high blood pressure-related diseases rank 10th among all causes of death in Korea. This proves that these diseases remain among the main detrimental diseases [2].

Korean meals are traditionally rice-based. Kimchi, vegetables pickled in soy sauce (jangajji), soy bean pastes (jang), and other products that contain a large amount of salt are consumed with rice; this shows that salt preference is high in the Korea population, and the Korea population has one of the highest salt intakes in the world [3,4]. In addition, the sodium intake is almost 15.5% higher in rural areas as compared to the urban areas. This is because the food culture in these rural areas is yet to be modernized,

unlike that in Korean cities. Rural residents still consume a large quantity of grains, kimchi, soy bean pastes, and vegetables pickled in soy sauce, which increases the risk of high blood pressure [5]. Such excessive intake of sodium not only leads to increased blood pressure, but also influences the occurrence of stroke, brain hemorrhage, chronic heart failure, and other cardiovascular diseases caused by high blood pressure [6-8]. In March 2012, the Korean Food and Drug Administration established 'the Sodium Intake Reduction Movement Headquarter'; this center is now making efforts towards improving awareness among citizens regarding the need to reduce sodium intake and change the food culture. The center is of the view that the most important task in this regard is to change societal attitudes, in general, rather than to change the private habits of individuals. Currently, 'Healthy Restaurant' by Korean Food and Drug Administration are involved in efforts to reduce the salt contents of seasoning or meat broth [9].

According to epidemiology studies, there is generally a quantitative correlation between sodium intake and blood pressure. It is reported that for populations with a lower intake of table salt, blood pressure does not increase with age; in

This research was funded by Rural Development Administration (the number of agreement; PJ008244).

§ **Corresponding Author:** Jeong-Sook Choe, Tel. 82-31-299-0483, Fax. 82-31-299-0454, Email. swany@korea.kr

Received: July 26, 2012, Revised: December 12, 2012, Accepted: December 15, 2012

©2012 The Korean Nutrition Society and the Korean Society of Community Nutrition

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

addition, in groups with low blood pressure, salt intake is generally low [10].

The microelement zinc is a key constituent of enzymes that is required for various metabolic processes and is therefore necessary for tissue and skeleton formation, skin protection, cell-mediated immunity, DNA and RNA synthesis, and metabolism of carbohydrates, etc [11]. Zinc is found not only in the salivary glands, but also in saliva and is involved in the proper functioning of the gustatory nerve [12]. Zinc deficiency can lead to a worsening of the sense of taste [13], which can, in turn, damage the taste acuity [14]. It is known that salt taste perception and salty taste preference are influenced by the zinc nutritional status; this not only influences sodium intake, but is also related to blood pressure [15-20]. Previous research conducted outside of Korea has shown that the consumption of zinc in food and via supplements can increase taste acuity [21]. Korean research on zinc is mainly focused on the current status of zinc intake, including the zinc nutritional status [22,23]. There have also been studies on the zinc nutritional status in pregnant and breastfeeding women [24,25] and in diabetes patients [26]. However, there is a lack of Korean research on salty taste acuity and salty taste preference. Thus, the objective of this study is to investigate differences in zinc nutritional status and their relation with salty taste acuity and salty taste preference, blood pressure. And we also determined which foods majorly contributed to the sodium intake.

## Subjects and Methods

### *Study participants*

We enrolled male and female adult participants from the Kangneung and Samcheok regions of Kangwon province from November 2011 to March 2012. The total number of participants was 252, and in our analysis, we used data from 218 participants (66 men, 152 women) who completed all tests of the study; these were blood tests, a dietary intake survey, and a salty taste acuity and salty taste preference test. All subjects provided informed consent, and this study was approved by the National Academy of Agricultural science under Rural Development Administration

### *Salty taste acuity and salty taste preference tests*

A sensory test was conducted to assess the salty taste acuity and salty taste preference of participants. To estimate salty taste acuity, we used a salt-water solution comprising water and refined salt. This was a 2-stage experiment. First, we offered the participants 10 different salt-water solutions, starting with a solution with a low salt concentration and increasing the concentration as follows: salt concentration of 0.03, 0.06, 0.09, 0.12, 0.15, 0.18, 0.21, 0.24, 0.27, and 0.3% at room temperature. Participants had to note when they began perceiving saltiness.

After this, a solution with 0.33% salinity was added to the series, and 11 forms of salt water were offered, from the highest concentration to the lowest concentration. The participants then had to note when they stopped perceiving the saltiness. The average value between the 2 results was considered the threshold saltiness concentration.

To test salty taste preference, we used bean sprout soup made only of bean sprouts and water. Salt concentrations of 0.15% and 1.0% were reached with the addition of refined salt. The participants first tried the soup with the less salty 0.15% concentration and then added the salty 1.0% soup to suit their preference. After this, salinity of the soup was measured twice with the help of a salinometer, and the average value between the 2 measurements was considered to be salty taste preference. The bean sprout soup was given at 50-60°C, and the participants rinsed their mouths with water each time after they tried the soup.

### *Dietary intake survey*

The dietary intake of study participants was surveyed using the 24-h recall method. Information was collected using one-on-one interviews on 1 day in the middle of the week, and additional phone surveys were carried out on 1 day during the weekend. Trained interviewers with expertise in food and nutrition asked the participants about the food they had eaten the day before and recorded the ingredients and the intake amount for each meal. To increase the accuracy of the dietary intake survey, models of real food and picture materials were used [27]. With the help of CAN pro 3.0 Software (The Korean Nutrition Society, Seoul, Republic of Korea), we calculated the nutrient intake from the dietary intake data collected through our survey.

### *Dish frequency questionnaire for estimation of habitual sodium intake (Na index)*

To estimate habitual sodium intakes, we used an amended version of the dish frequency questionnaire (DFQ 15) developed by Son *et al.* [28]. The survey data was recorded by trained interviewers in one-on-one interviews. The DFQ was designed to consider the contribution to intake frequency, sodium intake per one portion, and total sodium intake. Based on the sodium content per one portion of each type of food, extra points were given (more than 700 mg of sodium per one portion: 3 points; 500-699 mg: 2 points; and 300-499 mg: 1 point). The points were then multiplied by the intake frequency (every day: 5 points; more than 3 times a week: 4 points; 1-2 times a week: 3 points; 2-3 times a month: 2 points; once a month: 1 point; and almost never eat: 0 points) to yield the Na index.

### *Blood collection and blood pressure measurement*

To analyze subjects' serum zinc concentrations, we collected 3.5 ml of venous blood gathered after subjects had fasted for

12 h. This was lightly mixed and left at room temperature for 30 min. The serum was then separated by centrifugation. The Neodin medical laboratory performed the blood analysis.

### Statistical analysis

All data were analyzed using the SAS 9.2 statistical program (Cary, USA); data appears as mean and standard deviation. The participants from both regions, Kangneung and Samcheok, were further divided into 3 groups according to serum zinc level (T1: lowest; T2: intermediate; T3: highest). We adjusted the values for sex and age and compared the Na index, blood pressure, nutrient intake, and salty taste acuity and salty taste preference. We also divided the participants into 2 groups: those whose zinc intake was higher than the dietary reference intakes Koreans (2010) and the others. We calculated food intake of each food group for each of these groups. We identified the main sodium sources and calculated their rate and then analyzed the characteristics of the sodium sources for each region. All the adjusted averages were calculated using a general linear model.

## Results

### Smoking and health nutritional supplement use and dish frequency questionnaire

Table 1 shows the division of participants into smokers and non-smokers and users and non-users of health nutritional supplements based on their serum zinc level. Among the men, 14 were smokers and 52 were non-smokers, and 21 were taking health nutritional supplements, while 45 were not. Among the women, 2 were smokers and 150 were non-smokers, while 70 were taking health nutritional supplements and 82 were not. There was no significant difference in participants' zinc serum level on the basis of smoking or consumption of health nutritional supplements

The Na index used for estimation of habitual sodium intake are shown in Table 2. The Na index based on the serum zinc level was 68.6 (T1), 71.4 (T2), and 63.9 (T3). This index was

**Table 1.** Smoking and health nutritional supplement use of the subjects

		Total (n = 218)	T1 <sup>1)</sup> (n = 72)	T2 (n = 72)	T3 (n = 74)	X <sup>2</sup>
Sex	Male	66	20	23	23	0.8477 <sup>2)</sup>
	Female	152	52	49	51	
Male (n = 66)						
Smoking	Yes	14	3	7	4	1.8333
	No	52	17	16	19	
Health nutritional supplement	Yes	21	7	8	6	0.5347
	No	45	13	15	17	
Female (n = 152)						
Health nutritional supplement	Yes	70	20	25	25	2.0819
	No	82	33	23	26	

<sup>1)</sup> Tertile of serum zinc level (T1: lowest, T2: intermediate, T3: highest)

<sup>2)</sup> Values were tested by chi-square test.

**Table 2.** Eating behavior score related to salty food intake according to the serum zinc level

	T1 <sup>1)</sup> (n = 72)	T2 (n = 72)	T3 (n = 74)	P-value <sup>2)</sup>
Na index	68.6 ± 2.3 <sup>3)</sup>	71.4 ± 2.3	63.9 ± 2.3	0.0001

<sup>1)</sup> Tertile of serum zinc level (T1: lowest, T2: intermediate, T3: highest)

<sup>2)</sup> All the P-value were calculated by ANCOVA (sex, age, region adjusted).

<sup>3)</sup> Mean ± SD

significantly lower in the group (T3) with the highest serum zinc level; this indicates that participants in this group consumed fewer food with high sodium content.

### Serum zinc level and blood pressure

Data on the blood pressure of participants depending on their zinc serum level are presented in Table 3. This serum zinc level was significantly higher for participants from Kangneung (90.8 µg/dl) than those from Samcheok (81.2 µg/dl). Systolic blood pressure was higher in Kangneung participants (125.7 mmHg) than in Samcheok participants (121.8 mmHg), but this difference was not significant. However, for participants from Kangneung, the group with high serum zinc level had a significantly lower systolic blood pressure than the group with low serum zinc level (Table 3). The higher blood pressure in the Kangneung population, despite a high serum zinc level, than the Samcheok can be explained by other environmental factors than zinc intake

**Table 3.** Serum zinc and blood pressure according to the serum zinc level

	Gangneung					Samcheok					P-value <sup>3)</sup>
	Total (n = 133)	T1 <sup>1)</sup> (n = 44)	T2 (n = 44)	T3 (n = 45)	P-value <sup>2)</sup>	Total (n = 85)	T1 (n = 28)	T2 (n = 28)	T3 (n = 29)	P-value <sup>2)</sup>	
Serum zinc (µg/dl)	90.8 ± 1.6 <sup>4)</sup>	73.7 ± 1.2	89.6 ± 1.1	109.0 ± 1.1	0.0001	81.2 ± 2.0	61.3 ± 1.7	80.8 ± 1.7	98.9 ± 1.7	0.0001	0.0003
Age (yrs)	53.1 ± 0.7	53.3 ± 1.1	53.8 ± 1.1	52.2 ± 1.1	0.4329	63.0 ± 0.9	63.4 ± 1.8	61.4 ± 1.7	64.3 ± 1.7	0.6847	0.0001
Systolic blood pressure (mmHg)	125.7 ± 1.4	124.0 ± 2.3	126.2 ± 2.3	122.6 ± 2.2	0.0238	121.8 ± 2.0	123.9 ± 2.8	124.0 ± 2.7	125.4 ± 2.4	0.8882	0.0566
Diastolic blood pressure (mmHg)	78.9 ± 0.9	77.9 ± 1.6	80.0 ± 1.6	79.9 ± 1.6	0.8335	77.8 ± 1.2	77.1 ± 1.5	76.7 ± 1.5	77.7 ± 1.5	0.0906	0.2597

<sup>1)</sup> Tertile of serum zinc level (T1: lowest, T2: intermediate, T3: highest)

<sup>2)</sup> The P-value of difference in the serum zinc level were calculated by ANCOVA (sex, age adjusted)

<sup>3)</sup> Different between two groups by t-test (sex, age adjusted)

<sup>4)</sup> Mean ± SD

**Table 4.** Daily nutrient intake according to the serum zinc level



	Gangneung					Samcheok					
	Total (n = 133)	T1 <sup>1)</sup> (n = 44)	T2 (n = 44)	T3 (n = 45)	P-value <sup>2)</sup>	Total (n = 85)	T1 (n = 28)	T2 (n = 28)	T3 (n = 29)	P-value <sup>2)</sup>	P-value <sup>3)</sup>
Energy (kcal)	1,688.1 ± 35.9 <sup>4)</sup>	1,682.1 ± 60.6	1,763.5 ± 60.0	1,642.8 ± 59.3	0.0001	1,632.4 ± 46.6	1,618.3 ± 70.5	1,606.9 ± 69.7	1,634.8 ± 68.3	0.7265	0.0001
Carbohydrate (g)	254.3 ± 5.5	250.4 ± 9.2	263.9 ± 9.1	248.0 ± 9.0	0.0027	264.6 ± 7.1	254.8 ± 10.6	263.0 ± 10.5	276.2 ± 10.3	0.5344	0.0017
Protein (g)	69.2 ± 2.3	69.2 ± 3.6	73.6 ± 3.6	65.7 ± 3.6	0.1956	69.9 ± 3.0	73.4 ± 5.2	69.9 ± 5.2	65.4 ± 5.1	0.6292	0.3135
Fat (g)	39.7 ± 1.8	42.6 ± 3.3	42.2 ± 3.3	37.5 ± 3.3	0.2077	31.6 ± 2.4	32.5 ± 3.1	30.6 ± 3.1	27.2 ± 3.0	0.5932	0.0002
Fe (mg)	11.3 ± 0.6	11.0 ± 0.6	12.2 ± 0.6	11.2 ± 0.5	0.3713	13.9 ± 0.7	13.2 ± 1.7	14.0 ± 1.6	13.5 ± 1.4	0.8951	0.0424
Na (mg)	4,246.4 ± 133.5	4,125.2 ± 181.0	4,751.3 ± 179.3	3,944.2 ± 177.2	0.0016	4,856.2 ± 173.3	5,248.9 ± 327.5	4,802.1 ± 323.6	4,425.5 ± 317.0	0.0927	0.0008
Zn (mg)	8.11 ± 0.42	7.7 ± 0.6	8.2 ± 0.6	8.4 ± 0.6	0.9328	8.04 ± 0.54	7.5 ± 1.0	8.2 ± 1.0	8.3 ± 1.0	0.9495	0.9994
Vitamin A (μgRE)	716.6 ± 31.9	573.7 ± 58.6	829.2 ± 58.1	739.0 ± 57.4	0.0460	427.2 ± 41.3	475.0 ± 46.1	425.3 ± 45.6	390.7 ± 44.7	0.7771	0.0001

<sup>1)</sup> Tertile of serum zinc level (T1: lowest, T2: intermediate, T3: highest)

<sup>2)</sup> The P-value of difference in the serum zinc level were calculated by ANCOVA (sex, age adjusted).

<sup>3)</sup> Different between two groups by t-test (sex, age adjusted)

<sup>4)</sup> Mean ± SD

**Fig 1.** Distribution of salty taste acuity in study subjects**Fig 2.** Distribution of salty taste preference in study subjects

status.

### Nutrient intake status of participants

Data on the nutrient intake of the study participants are presented in Table 4. The energy intake of Kangneung participants (1688.1 kcal/day) was significantly higher than that of Samcheok participants (1632.4 kcal/day). The intake of carbohydrates, iron, and sodium was significantly higher for participants from Samcheok compared with those from Kangneung; conversely, the intake of fat, vitamin A, and niacin was significantly higher in Kangneung participants than in Samcheok participants. A comparison of the nutrient intake for the 2 regions based on serum zinc level shows that for Kangneung, the group with the low serum zinc level (T1) had a higher intake of energy, carbohydrates, and sodium and a lower intake of vitamin A than the group with a high serum zinc level (T3).

### Salty taste acuity and salty taste preference of participants

The distribution of salty taste acuity and salty taste preference of the study participants are shown in Fig. 1 and 2. The lowest threshold concentration for salt was 0.045%, and the highest

**Table 5.** Salt perception and preference by age

	< 50 (n = 43)	50-64 (n = 120)	≥ 65 (n = 55)	P-value <sup>1)</sup>
Acuity (%)	0.15 ± 0.01 <sup>2)</sup>	0.13 ± 0.01	0.16 ± 0.01	0.0001
Preference (%)	0.42 ± 0.02	0.41 ± 0.01	0.43 ± 0.02	0.0462

<sup>1)</sup> The P-value of difference by the serum zinc level were calculated by ANCOVA (sex, age adjusted).

<sup>2)</sup> Mean ± SD

threshold was 0.285%. The minimum salty taste preference level was 0.11%, while the maximum level was 0.77%. Most participants perceived saltiness at a concentration of 0.09%-0.12%, and most had a salty taste preference of 0.4%-0.5%. In a study by Ahn *et al.* [29] that looked at salty taste acuity and salty taste preference among adults in their twenties, most participants perceived saltiness at 0.06%-0.09% concentration and most preferred the 0.5%-0.6% salt concentration; this is a slightly lower value for acuity and slightly higher value for preference than that reported in the current study.

Values for salty taste acuity and salty taste preference by age are shown in Table 5. There were significant differences depending on age, with the saltiness threshold concentration at 0.15% for participants younger than 50 years of age, 0.13% for participants between 50 and 64 years of age, and 0.16% for participants above 65 years of age. The salty taste preference

**Table 6.** Salt acuity and preference by zinc nutritional status

	Gangneung				Samcheok				<i>P</i> -value <sup>2)</sup>
	Total (n = 133)	< RI (n = 57)	≥ RI (n = 76)	<i>P</i> -value <sup>1)</sup>	Total (n = 85)	< RI (n = 38)	≥ RI (n = 47)	<i>P</i> -value <sup>1)</sup>	
Acuity (%)	0.13 ± 0.01 <sup>3)</sup>	0.13 ± 0.01	0.12 ± 0.01	0.5327	0.17 ± 0.01	0.18 ± 0.01	0.16 ± 0.01	0.1082	0.0001
Preference (%)	0.40 ± 0.01	0.40 ± 0.02	0.40 ± 0.01	0.6378	0.45 ± 0.02	0.46 ± 0.03	0.44 ± 0.02	0.4340	0.0255
Serum zinc	90.26 ± 1.62	91.8 ± 2.2	90.5 ± 1.9	0.9553	81.65 ± 2.14	79.4 ± 3.0	81.5 ± 2.6	0.6310	0.0002

<sup>1)</sup> The *P*-value of difference by the serum zinc level were calculated by ANCOVA (sex, age adjusted)

<sup>2)</sup> Different between two groups by t-test (sex, age adjusted)

<sup>3)</sup> Mean ± SD

**Table 7.** Salt acuity and preference by serum zinc level

	Gangneung				Samcheok			
	T1 <sup>1)</sup> (n = 44)	T2 (n = 44)	T3 (n = 45)	<i>P</i> -value <sup>2)</sup>	T1 (n = 28)	T2 (n = 28)	T3 (n = 29)	<i>P</i> -value <sup>2)</sup>
Acuity (%)	0.13 ± 0.01 <sup>3)</sup>	0.11 ± 0.01	0.14 ± 0.01	0.1372	0.17 ± 0.01	0.17 ± 0.01	0.16 ± 0.01	0.2300
Preference (%)	0.38 ± 0.02	0.39 ± 0.02	0.41 ± 0.02	0.5429	0.43 ± 0.03	0.43 ± 0.03	0.50 ± 0.03	0.1367

<sup>1)</sup> Tertile of serum zinc level (T1: lowest, T2: intermediate, T3: highest)

<sup>2)</sup> The *P*-value of difference by the serum zinc level were calculated by ANCOVA (sex, age adjusted)

<sup>3)</sup> Mean ± SD

**Table 8.** Food intake of subjects by zinc intake

Food group	Gangneung(n = 133)			Samcheok(n = 85)		
	< RI <sup>1)</sup> (n = 57)	≥ RI (n = 76)	<i>P</i> -value <sup>2)</sup>	< RI (n = 38)	≥ RI (n = 47)	<i>P</i> -value <sup>2)</sup>
Grains	493.1 ± 17.8 <sup>3)</sup>	547.3 ± 26.1	0.0001	526.2 ± 18.1	598.7 ± 28.0	0.0648
Potatoes	151.7 ± 18.9	107.2 ± 27.4	0.1646	156.2 ± 28.1	113.7 ± 51.4	0.2911
Sugar	7.1 ± 0.8	9.2 ± 1.2	0.5017	8.6 ± 1.1	8.5 ± 1.6	0.0699
Beans	60.3 ± 7.8	55.1 ± 10.0	0.1728	49.5 ± 7.0	55.3 ± 10.9	0.5991
Seeds	9.1 ± 2.9	4.4 ± 4.2	0.2752	3.9 ± 1.7	10.6 ± 2.0	0.0041
Vegetables	340.0 ± 19.7	423.1 ± 28.9	0.1046	259.2 ± 13.4	300.4 ± 20.8	0.3312
Mushrooms	27.2 ± 4.0	26.1 ± 5.0	0.6390	26.1 ± 5.3	13.7 ± 8.1	0.3696
Fruits	236.4 ± 27.0	294.1 ± 34.1	0.1394	249.9 ± 34.5	393.0 ± 50.5	0.0673
Meats	73.1 ± 12.0	200.9 ± 15.6	0.0001	53.0 ± 18.8	185.5 ± 25.8	0.0015
Eggs	32.2 ± 3.9	47.1 ± 5.2	0.0066	27.3 ± 3.5	28.6 ± 6.1	0.1139
Fish & Shellfish	66.5 ± 7.4	84.2 ± 10.7	0.3415	76.5 ± 10.8	124.0 ± 16.9	0.0537
Seaweeds	5.4 ± 2.7	10.8 ± 3.7	0.6338	19.5 ± 5.0	23.3 ± 7.8	0.4765
Milk & Dairy products	172.3 ± 19.9	197.5 ± 27.1	0.5910	211.9 ± 42.2	234.2 ± 45.2	0.8184
Oils	7.4 ± 0.7	9.2 ± 0.9	0.3087	4.4 ± 0.5	7.0 ± 0.8	0.0205
Seasoning & Spices	37.8 ± 2.2	44.2 ± 3.2	0.3114	35.1 ± 2.1	40.5 ± 3.3	0.4698
Others	7.8 ± 2.1	8.3 ± 1.8	0.9590	-	-	-
Total	1,291.4 ± 35.9	1,685.9 ± 52.6	0.0001	1,218.0 ± 41.7	1,595.5 ± 64.7	0.0001

<sup>1)</sup> Dietary reference intakes for Koreans 2010 : 9 mg/day (male ≥ 30 y), 8 mg/day (30 y ≤ female < 65 y), 7 mg/day (female ≥ 65 y)

<sup>2)</sup> Different between two groups by t-test (sex, age adjusted)

<sup>3)</sup> Mean ± SD

concentration was 0.42% for participants below 50 years of age, 0.41% for those between 50 and 64 years of age, and 0.43% for those above 65 years of age.

#### *Salty taste acuity and salty taste preference and zinc nutritional status*

Results for salty taste acuity and salty taste preference and their relationship with zinc nutritional status for participants from Kangneung and Samcheok are presented in Table 6. Salty taste threshold and salty taste preference concentrations were both significantly higher in Samcheok participants (0.17% and 0.45%, respectively) than in Kangneung participants (0.13% and 0.40%,

respectively). Conversely, serum zinc level was 90.26 µg/dL in Kangneung participants and 81.65 µg/dL in Samcheok participants. However, comparison of salty taste acuity and salty taste preference for the groups of participants who had a zinc intake lower than the recommended nutrient intake(RI) of dietary reference intakes for Koreans and those who had a zinc intake higher than the Korean RI showed no significant differences. It should be noted, though, that in Kangneung, the salty taste threshold were higher for the group that had a zinc intake lower than the RI (threshold concentration, 0.13% [*<* RI] and 0.12% [*≥* RI], respectively). For both groups, preferred concentrations of salty taste was 0.40%, and there was no difference.

For Samcheok, the threshold concentration was 0.18% (*<* RI)

**Table 9.** Intake and major food sources of Na

Rank	Gangneung (n = 133)				Samcheok (n = 85)			
	Dish name	Na (mg)	Contribution rate (%)	Accumulated percent (%)	Dish name	Na (mg)	Contribution rate (%)	Accumulated percent (%)
1	Baechukimchi	1344.01	17.29	17.29	Baechukimchi	1815.74	19.26	19.26
2	Doenjang-jjigae	499.60	6.43	23.72	Gajami-sikhae	757.82	8.04	27.30
3	Doenjangguk	257.68	3.32	27.04	Doenjangguk	456.09	4.84	32.14
4	Ramyeon	223.73	2.88	29.92	Dorumuk-jorim	422.84	4.49	36.63
5	Kimchi-jjigae	182.27	2.35	32.26	Cheonggukjang-jjigae	386.83	4.10	40.73
6	Mixed soy paste (ssamjang)	180.47	2.32	34.59	Doenjang-jjigae	354.33	3.76	44.49
7	Chueotang	169.00	2.17	36.76	Kimchi-jjigae	321.10	3.41	47.90
8	Miyeokguk	168.77	2.17	38.93	Mandutguk	243.59	2.58	50.48
9	Yeolmu-kimchi	141.83	1.83	40.76	Miyeokguk	188.17	2.00	52.48
10	Guksu-jangguk	131.31	1.69	42.45	Ramyeon	184.11	1.95	54.43
11	Godeungeo-jorim	114.45	1.47	43.92	Guksu-jangguk	182.57	1.94	56.37
12	Stir-fried pumpkin	113.38	1.46	45.38	Dongchimi	145.44	1.54	57.91
13	Makguksu	110.27	1.42	46.80	Miyeok-muchim	136.73	1.45	59.36
14	Nabak kimchi	101.05	1.30	48.10	Kimchitguk	128.44	1.36	60.73
15	Mackerel	89.86	1.16	49.25	Dongtae-jjigae	124.11	1.32	62.04
16	Kalguksu	83.83	1.08	50.33	Crisp laver	121.29	1.29	63.33
17	Stir-fried Anchovy	80.30	1.03	51.37	Hobakjuk	107.19	1.14	64.47
18	Crisp laver	76.46	0.98	52.35	Stir-fried anchovies	102.90	1.09	65.56
19	Gamja-jorim	73.75	0.95	53.30	Myeongnanjeot	93.26	0.99	66.55
20	Oi-saengchae	68.78	0.89	54.18	Myeongtae-sikhae	91.83	0.97	67.52

and 0.16% ( $\geq$  RI) and preference was 0.46% ( $<$  RI) and 0.44% ( $\geq$  RI). There was no significant difference between the 2 groups, although the group that consumed less than the RI showed higher values; the serum zinc level was 79.4  $\mu$ g/dl ( $<$  RI) and 81.5  $\mu$ g/dl ( $\geq$  RI), and this was higher for the group that consumed more than the RI, but the difference was not significant.

Data on salty taste acuity and salty taste preference and their relation to serum zinc levels for Kangneung and Samcheok are presented in Table 7. For both regions, there was almost no difference in the salty taste threshold concentration based on the serum zinc level. Salty taste preference concentrations were 0.38% (T1), 0.39% (T2), 0.41% (T3) for Kangneung participants and 0.43% (T1), 0.43% (T2), 0.50% (T3) for Samcheok participants. Thus, there was a tendency for an increased salty taste preference concentration with increased serum zinc level, but there was no significant difference among the 3 groups.

#### *Food intake status of study participants*

The food intakes of the study participants and its relation with zinc intakes is presented in Table 8. For Kangneung, the groups with a zinc intake higher than RI showed a significant larger intake of grains, meat, and eggs than the group with an intake lower than RI. In Samcheok, there was a significantly higher intake of seeds and nuts, meat, fat, and oils. The grain intake was 493.1 g ( $<$  RI) and 547.3 g ( $\geq$  RI) for Kangneung participants and 526.2 g ( $<$  RI) and 598.7 g ( $\geq$  RI) for Samcheok participants; the group that had zinc intake higher than RI had a significantly higher grain intakes in Gangneung. The results

of the 2010 Korean National Health and Nutrition Examination Survey showed that the grain intake for people in small towns or townships was 321.7 g/day [1].

Meat intake was significantly higher in the groups with a zinc intake higher than RI in both regions, which confirms that meat is a good source of zinc.

The main sodium sources are shown in Table 9. The most important product for increasing the intake of sodium in both regions was Baechukimchi. The amount of sodium consumed through Baechukimchi was 1,344.0 mg in Kangneung and 1,815.7 mg in Samcheok. In Kangneung, other sodium sources included Doenjang-jjigae, Doenjangguk, Ramyeon, Kimchi-jjigae in order of importance. In Samcheok, this order was Gajami-sikhae, (salted flatfish with grains), Doenjangguk, Dorumuk-jorim, Cheonggukjang-jjigae. There is clear difference between the 2 regions in terms of the diet. The presence of Gajami-sikhae in the list for Samcheok reflects food culture in the East Sea regions. It is the typical local dish in these regions. Apart from the foods already listed, among the top 20 foods that increase sodium intake are Ramyeon, Godeungeo-jorim, Kalguksu, etc.

#### **Discussion**

This study was conducted to explore differences in zinc nutritional status and their relation with salty taste acuity and salty taste preference, blood pressure, and nutrient and food intake for residents of Kangneung and Samcheok regions in Korea's Kangwon province.

We found that the population Kangneung has significantly lower salty taste threshold concentrations and salty taste preference concentrations than the population from Samcheok, while the serum zinc level of Kangneung population was significantly higher. In this study, we also explored salty taste acuity and salty taste preference depending on zinc intake status in both regions. In Samcheok, the group with higher zinc intake showed a lower salty taste threshold and lower salty taste preference concentrations. However, the difference groups was not significant.

In our study, the results of a comparison of sodium intake and the Na index (acquired using food frequency questionnaire for estimation of sodium intakes) showed that a higher serum zinc level indicated a lower sodium intake and Na index score. A previous study of zinc nutritional status and salty taste acuity and salty taste preference by Ahn *et al.* [29] showed that only participants with a serum zinc level lower than the standard had higher salty taste preference concentration as the concentration of serum zinc level lowered. Although zinc deficiency influences salty taste preference, it has been reported that differences within the normal score of zinc concentration variation do not have a large influence on salty taste preference. Conversely, Stewart-Knox *et al.* [31] found no correlation between serum zinc level and salty taste acuity, but they found that a higher zinc concentration in red blood cells was associated with a higher taste acuity.

Studies on zinc intake and salty taste acuity and preference also show differences based on sex. In the study by McDaid *et al.* [20], women with high zinc intake had better salty taste acuity, but there was no significant finding for men. Ahn *et al.* [29] also analyzed the connection between zinc intake and salty taste acuity and preference and found that the only significant correlation was found in women. In addition, zinc intake has been shown to be related to taste function, which shows differences between men and women in terms of the zinc intake, and salty taste acuity and preference. A previous study on the food intake of rural adults according to saline sensitivity showed that the participants less sensitive to salt had a high intake rate of salt-preserved foods such as salted fish or vegetables pickled in soy sauce, as well as poultry and meat and eggs [30]. With increased age, salty taste acuity decreases, and people tend to eat very salty food, thus increasing their salt intake. Accordingly, there is a need to raise awareness about low salt intake among older people.

Other studies have demonstrated that better zinc nutritional status leads to improve cognitive performance and the recognition threshold for salt. Tupe and Chiplonkar [21] investigated the influence of zinc supplements on taste acuity and found that a significant fall in recognition thresholds for salt was noted after both diet and zinc supplementation for 10 weeks ( $P < 0.01$ ): however, it remained the same in the group that did not receive the supplements for 10 weeks.

The findings of this study, together with the results of previous

studies, indicate that zinc deficiency is linked with a reduction in the senses, including eyesight, taste, and smell [32,33]. One of the symptoms of zinc deficiency is damage to the sense of taste [34,35], and this symptom can be successfully treated with zinc supplements [36,37]. However, studies have shown conflicting results in terms of the relation between zinc nutritional status (serum zinc levels or zinc plasma levels) and taste acuity as well as zinc intake and taste perception function. These results differ in terms of sex and the level of zinc deficiency. Therefore, to establish the relation between zinc nutritional status, salty taste acuity, and salty taste preference and, in turn, blood pressure, there is the need for a large scale longitudinal study that includes people from different strata of society. The higher blood pressure in the Kangneung population, despite a high serum zinc level, can be explained by other factors than zinc Intake status.

There is a need for further research on the relationship between intake of different minerals and high blood pressure. Potassium plays an important role in terms of limiting sodium consumption to reduce blood pressure; it helps in excreting sodium from the body. In addition, calcium increases sensitivity to salt, and when renin activity is low, it acts to reduce blood pressure. Calcium facilitates the excretion of sodium with urine, which causes a decrease in the amount of intracellular fluid. Subsequently, this causes a decrease in blood volume, in turn lowering the blood pressure. Thus, potassium, calcium, and magnesium influence the control of blood pressure.

In addition, salty taste acuity appears to decrease with age. Thus, there is a need to explore what factors influence this change.

Finally, future studies should focus on determining how to preserve the taste characteristics and safety of low-salt Korean traditional foods. For example, a low-salt Gajami-sikhae recipe could be developed for the residents of the Samcheok region.

## References

1. Korea Centers for Disease Control and Prevention. The Fourth Korea National Health and Nutrition Examination Survey (KNHANES IV). Cheongwon; 2010.
2. Statistics Korea. 2010 Annual Report on the Cause of Death Statistics. Daejeon; 2010.
3. Song MR, Lee KJ. Salinity and consumption patterns of Kimchi and soup, stew in Jeonju area. *Korean J Food Cookery Sci* 2008;24:84-91.
4. Korea Centers for Disease Control and Prevention. Korea National Health and Nutrition Examination Survey (KNHANES). Cheongwon; 2002.
5. Choe M, Kim JD, Kim SS. A study on correlation between blood pressure and Na, K intakes pattern in the family members of normal and hypertension patients. *J Korean Soc Food Sci Nutr* 1996;25:1045-9.
6. He J, Gu D, Chen J, Jaquish CE, Rao DC, Hixson JE, Chen JC, Duan X, Huang JF, Chen CS, Kelly TN, Bazzano LA, Whelton PK; GenSalt Collaborative Research Group. Gender difference in

- blood pressure responses to dietary sodium intervention in the GenSalt study. *J Hypertens* 2009;27:48-54.
7. de Wardener HE, MacGregor GA. Harmful effects of dietary salt in addition to hypertension. *J Hum Hypertens* 2002;16:213-23.
  8. Sanders PW. Dietary salt intake, salt sensitivity, and cardiovascular health. *Hypertension* 2009;53:442-5.
  9. Korea Food & Drug Administration [Internet]. Cheongwon: Korea Food & Drug Administration; [cited 2012 July 16]. Available from: <http://www.kfda.go.kr/index.kfda?mid=56&pageNo=11&seq=17623&cmd=v>.
  10. U.S. Department of Health and Human Services. The Surgeon General's Report on Nutrition and Health. Washington, D.C.: U.S. Government Printing Office; 1988. p.149-52.
  11. Johnson PE, Hunt CD, Milne DB, Mullen LK. Homeostatic control of zinc metabolism in men: zinc excretion and balance in men fed diets low in zinc. *Am J Clin Nutr* 1993;57:557-65.
  12. Tanaka M. Secretory function of the salivary gland in patients with taste disorders or xerostomia: correlation with zinc deficiency. *Acta Otolaryngol Suppl* 2002;(546):134-41.
  13. Pluhator MM, Thomson ABR, Fedorak RN. Clinical aspects of trace elements: zinc in human nutrition - assessment of zinc status. *Can J Gastroenterol* 1996;10:37-42.
  14. Bales CW, Steinman LC, Freeland-Graves JH, Stone JM, Young RK. The effect of age on plasma zinc uptake and taste acuity. *Am J Clin Nutr* 1986;44:664-9.
  15. Bertino M, Beauchamp GK, Engelman K. Long-term reduction in dietary sodium alters the taste of salt. *Am J Clin Nutr* 1982;36:1134-44.
  16. Kim JY, Kang YR, Lee MY, Paik HY. Sodium intake and preference for salty taste in elementary school children residing in rural and urban areas of Korea. *Korean J Nutr* 1990;23:248-56.
  17. Isezuo SA, Saidu Y, Anas S, Tambuwal BU, Bilbis LS. Salt taste perception and relationship with blood pressure in type 2 diabetics. *J Hum Hypertens* 2008;22:432-4.
  18. Rabin M, Poli de Figueiredo CE, Wagner MB, Antonello IC. Salt taste sensitivity threshold and exercise-induced hypertension. *Appetite* 2009;52:609-13.
  19. Ueda C, Takaoka T, Sarukura N, Matsuda K, Kitamura Y, Toda N, Tanaka T, Yamamoto S, Takeda N. Zinc nutrition in healthy subjects and patients with taste impairment from the view point of zinc ingestion, serum zinc concentration and angiotensin converting enzyme activity. *Auris Nasus Larynx* 2006;33:283-8.
  20. McDaid O, Stewart-Knox B, Parr H, Simpson E. Dietary zinc intake and sex differences in taste acuity in healthy young adults. *J Hum Nutr Diet* 2007;20:103-10.
  21. Tupe RP, Chiplonkar SA. Zinc supplementation improved cognitive performance and taste acuity in Indian adolescent girls. *J Am Coll Nutr* 2009;28:388-96.
  22. Park JS, Chyun JH. Dietary zinc analysis and changes of zinc nutriture with zinc supplementation in Korean adults. *Korean J Nutr* 1993;26:1110-7.
  23. Kim CH, Paik HY, Joung H. Evaluation of zinc and copper status in Korean college women. *Korean J Nutr* 1999;32:277-86.
  24. Lee HS. Zinc status of South Korean women in the second trimester and the effect of those on pregnancy outcome. *Korean J Nutr* 2005;38:578-85.
  25. Yu KH. The study of zinc nutritional status of pregnant women visiting in public health centers in Ulsan. *Korean J Nutr* 2000;33:848-56.
  26. Lee JH, Lee HJ, Lee IK, Yoon JS. Zinc and copper status of middle- and old-aged women in type 2 diabetes. *Korean J Nutr* 2005;38:56-66.
  27. Rural Development Administration. Standard Tables of Food Composition in Korea. Suwon; 2009.
  28. Son SM, Huh GY, Lee HS. Development and evaluation of validity of dish frequency questionnaire (DFQ) and short DFQ using Na index for estimation of habitual sodium intake. *Korean J Community Nutr* 2005;10:677-92.
  29. Ahn EJ, Noh HY, Chung J, Paik HY. The effect of zinc status on salty taste acuity, salty taste preference, sodium intake and blood pressure in Korean young adults. *Korean J Nutr* 2010;43:132-40.
  30. Kim MK, Han JI, Chung YJ. Dietary behavior related to salty food intake of adults living in a rural area according to saline sensitivity. *Korean J Nutr* 2011;44:537-50.
  31. Stewart-Knox BJ, Simpson EE, Parr H, Rae G, Polito A, Intorre F, Meunier N, Andriollo-Sanchez M, O'Connor JM, Coudray C, Strain JJ. Zinc status and taste acuity in older Europeans: the ZENITH study. *Eur J Clin Nutr* 2005;59 Suppl 2:S31-6.
  32. Russell RM, Cox ME, Solomons N. Zinc and the special senses. *Ann Intern Med* 1983;99:227-39.
  33. Van Wouwe JP. Clinical and laboratory assessment of zinc deficiency in Dutch children. A review. *Biol Trace Elem Res* 1995;49:211-25.
  34. Henkin RI, Bradley DF. Hypogeusia corrected by Ni<sup>++</sup> and Zn<sup>++</sup>. *Life Sci* 1970;9:701-9.
  35. Mahajan SK, Prasad AS, Rabbani P, Briggs WA, McDonald FD. Zinc deficiency: a reversible complication of uremia. *Am J Clin Nutr* 1982;36:1177-83.
  36. Henkin RI, Schechter PJ, Friedewald WT, Demets DL, Raff M. A double blind study of the effects of zinc sulfate on taste and smell dysfunction. *Am J Med Sci* 1976;272:285-99.
  37. Sakai F, Yoshida S, Endo S, Tomita H. Double-blind, placebo-controlled trial of zinc picolinate for taste disorders. *Acta Otolaryngol Suppl* 2002;(546):129-33.