

Gut as viral reservoir: lessons from gut viromes, HIV and COVID-19

Markus F Neurath ¹, Klaus Überla,² Siew C Ng ³

BACKGROUND: VIRUSES IN THE HEALTHY GUT

The gut microbiota is composed of bacteria, fungi and viruses.^{1–3} While the presence and composition of the bacteriome have been extensively studied in recent years, the ‘dark matter’ of microbiomes including viruses (viral microbiome or virome) remains incompletely understood.⁴ Analysis of microbiotic communities in the gut showed that bacteria rapidly colonise the gut after birth, followed by the appearance of prophages by 1 month which are harboured by these bacteria. Subsequently, a gut virome dominated by bacteriophages develops and human viruses can be detected 4 months after birth.^{5–6} In adults, recent studies have analysed the gut virome in healthy individuals by using metagenomic approaches.^{7–8} For instance, a metagenome analysis in 1986 individuals identified 33 242 unique viral populations, demonstrating that the human gut contains a vast array of bacteriophages and eukaryotic RNA and DNA viruses with a predominance of bacteriophages. Specifically, several groups of prokaryotic crAss-like, Caudovirales and Microviridae bacteriophages and eukaryotic adenoviruses and herpesviruses were characterised as stable colonisers of the human gut.

Further studies described the existence of individual-specific persistent personal viromes. These viromes displayed marked interindividual heterogeneity with age-dependent changes and were critically dependent on host factors such as geography and ethnicity-distinct diets.^{9–10} Moreover, the duration of urban residence was associated with the presence of multiple bacteriophages, including *Lactobacillus* and *Lactococcus* phages.⁹ Although the functions of most viruses in the gut viromes have not been fully characterised, it has been

suggested that viromes play a crucial role in shaping the gut microbiome and intestinal health under homeostatic conditions.

This concept was underlined by experimental studies in murine dextran sulfate sodium (DSS) colitis.^{11–12} Treatment of DSS colitis with a cocktail of antiviral nucleoside analogues led to augmented mucosal inflammation, while administration of agonists for Toll-like receptor 3 (TLR3: sensor for double-stranded RNA) or TLR7 (sensor for single-stranded RNA) suppressed colitis activity. Triggering of TLR3 and TLR7 responses was associated with production of interferon- β by mucosal, plasmacytoid dendritic cells. Collectively, these results indicated that recognition of resident luminal viruses regulates protective mucosal immunity during experimental colitis.¹¹ This protective role of the gut virome in the intestinal microbiota is reminiscent of the well-established protective role of commensal bacteria in intestinal inflammation.¹²

ALTERATIONS OF THE GUT VIROME IN INTESTINAL INFLAMMATORY AND NEOPLASTIC DISORDERS

In spite of the relative stability of personal gut viromes, marked changes may occur in patients with chronic disorders of the intestine. A recent study in human UC used deep metagenomic sequencing in 91 patients and 76 healthy controls.¹³ An expansion of mucosal viruses was noted that correlated with the presence of inflammation. In particular, Caudovirales, *Escherichia* and Enterobacteria phages were more frequently detected in patients with UC as compared with controls. Functions of bacteriophages with relationship to host bacteria fitness and pathogenicity were increased in UC mucosa, suggesting that structural and functional changes in the gut virome may play an important role in disease pathogenesis.¹³ A second metagenomic study compared the gut virome using colonic/ileal tissues in treatment-naive UC with Crohn’s disease (CD) and controls¹⁴ and found an increase of Hepadnaviridae in UC, while Hepeviridae were more frequently detected in CD as compared with controls. In contrast, Virgaviridae and Polydnviridae were reduced in CD and UC, respectively. Finally, two additional

studies reported expansion of Caudovirales or Adenoviridae and Herpesviridae in both CD and UC, respectively.^{15–16}

There is evidence that enteric viruses can modulate disease activity in IBD. For instance, rotavirus infections have been shown to induce epithelial damage and proinflammatory intestinal immune responses through increased secretion of cytokines such as interferon- β and thus may augment gut inflammation in IBD.¹⁷ Moreover, reduction of mucosal blood type antigens in non-secretor patients carrying polymorphisms of the α 1,2-fucosyltransferase (FUT2) and FUT3 genes has been linked to altered host–microbiome interactions in IBD.^{18–20} Specifically, the FUT2 W143X mutation may cause asymptomatic norovirus infection and favour the development of ileitis in CD.²¹

In addition to UC and CD, changes in the virome of faecal samples were noted in IBS. A study in 55 patients with IBS and 51 controls showed lower alpha diversity of viral clusters in the former as compared with the latter patients and abundant bacteriophage clusters (Caudovirales) in IBS. Additional analysis of CRISPR (clustered regularly interspaced short palindromic repeats) spacers indicated that changes in the faecal virome were related in part to the alterations in the bacteriome, suggesting that their cross-sectional relationship may affect IBS pathogenesis.²² After allogeneic haematopoietic stem cell transplantation (HSCT), a significant increase in the viral sequences of faecal samples was found in a recent study in 44 patients. Moreover, enteric graft-versus-host disease (GVHD) following HSCT was characterised by reduced phage richness and expansion of persistent DNA viruses such as anelloviruses, herpesviruses, papillomaviruses and polyomaviruses. In particular, the presence of picobirnaviruses was predictive of severe enteric GVHD.²³ In addition, after repeated faecal microbiota transplantation (FMT) in severe colitis due to GVHD, the gut virome showed a stable rise in diversity. While the levels of Torque teno viruses decreased after FMTs, an increase in the relative abundance of Caudovirales bacteriophages in faecal samples was observed.²⁴ Collectively, these findings demonstrated marked alterations of the gut virome in GVHD and GVHD-associated enteritis that may affect the course of the disease.

In addition to intestinal inflammation, it should be noted that viruses have also been implicated in the pathogenesis of neoplastic GI disorders such as colorectal cancer.²⁵ Several studies supported the concept that the gut virome is involved in cancerogenesis and tumour growth in this disease by viral

¹First Department of Medicine & Deutsches Zentrum Immuntherapie DZI, Friedrich-Alexander-Universität Erlangen-Nürnberg, Erlangen, Germany

²Institute of Clinical and Molecular Virology & Deutsches Zentrum für Immuntherapie (DZI), University of Erlangen-Nürnberg, Erlangen, Germany

³Medicine and Therapeutics, Chinese University of Hong Kong, Hong Kong, Hong Kong

Correspondence to Professor Markus F Neurath, First Department of Medicine & Deutsches Zentrum Immuntherapie DZI, Friedrich-Alexander-Universität Erlangen-Nürnberg, Erlangen, Bayern, Germany; markus.neurath@uk-erlangen.de

infection of epithelial cells or modulation of the composition of bacterial communities.^{25,26} For instance, the large T-antigen of polyomavirus was suggested to drive activation of the β -catenin/Wnt pathway, which favours cell proliferation and apoptosis resistance in colorectal cancer.^{27–29} Moreover, the diversity of the gut bacteriophage community was significantly increased in patients with colorectal cancer as compared with control patients without cancer,³⁰ and certain bacteriophages (Caudovirales, *Inovirus*, *Orthobunyavirus*) were associated with dysbiosis as potential driver of cancer development.^{30,31} Finally, virus-based therapeutic concepts (engineered oncolytic viruses and bacteriophages) have been suggested as potential future therapy for colorectal cancer.²⁵

THE GUT AS RESERVOIR FOR HIV AND SARS-COV-2

The mucosal surface of the gut plays a key role during infection with HIV.^{32,33} In fact, the gut contains a large number of infected cells, even in individuals receiving antiretroviral therapy (ART). This process is facilitated by the increased susceptibility of mucosal CD4+ T cells for HIV infection due to greater C-C chemokine receptor (CCR)-5 expression and local T cell activation.^{34,35} On infection, there is a rapid loss of mucosal barrier function with translocation of microbial products³⁶ and loss of mucosal CD4+ T cells.³⁷ These changes favour local and systemic inflammation in HIV-infected individuals. Probiotic

treatment of HIV-infected patients has been shown to suppress microbial translocation and serum inflammatory markers in ART-naïve patients,^{38,39} highlighting a leaky barrier as key component of HIV-induced gut inflammation. Moreover, it has been shown that the mucosa, as the largest lymphoid organ of the body, is an important reservoir for HIV-infected cells. Consistently, studies reported that HIV-DNA levels in CD4+ T cells were on average 5–6 times higher in the ileum compared with blood in patients on ART,⁴⁰ and 2-fold to 12-fold higher in the duodenum, ileum, right colon and rectum as compared with peripheral blood.⁴¹ Based on these findings it was estimated that the gut harbours 83%–95% of all HIV-infected cells in the body and thus provides a key reservoir for disease persistence. Local, infected CD13+ myeloid cells and CD4+ T cells may reach regional lymph nodes and the bloodstream via immune cell trafficking, thereby favouring viral dissemination throughout the body.^{35,42,43} This concept has important implications for clinical therapy. In fact, ART does not cure HIV infection due to the persistence of HIV reservoirs in long-lived memory CD4 T cells present in the blood, lymph nodes and intestinal tract.⁴⁴

The COVID-19 pandemic is caused by SARS-CoV-2. Infection of more than 144 million cases has been reported and has led to 3 million confirmed deaths.⁴⁵ As intestinal epithelial cells express high amounts of the SARS-CoV-2 receptor ACE2 and transmembrane protease serine subtype 2

(TMPRSS2), a cellular protease important for viral entry, GI infection is frequently seen in COVID-19 and GI symptoms have been reported in 30%–70% of patients.^{46–48} Remarkably, GI infection results in limited or absent signs of local inflammation as well as low mortality in COVID-19, suggesting a potential role of the GI tract in attenuating SARS-CoV-2 infection.⁴⁹ Stool analyses confirmed the presence of SARS-CoV-2 genomic and subgenomic RNA in affected patients,⁵⁰ but isolation of the virus from stool samples has rarely been successful,⁵¹ indicating that GI infection may be self-limiting. However, local infection of intestinal epithelial cells in COVID-19 may affect the gut virome. In patients with COVID-19, a significantly reduced viral Shannon diversity was noted as compared with the gut viromes of healthy controls.⁵² Specifically, viromes were composed of DNA/RNA viruses (mainly Herelleviridae and Virgaviridae families) as well as bacteriophages (Caudovirales, crAss-like phage, Inoviridae, Microviridae, Myoviridae, Podoviridae and Siphoviridae families).⁵² A recent metagenomic study profiled the faecal RNA and DNA viromes of patients with COVID-19 and found that the faecal virome in SARS-CoV-2 infection harboured more stress-associated, inflammation-associated and virulence-associated gene-encoding capacities, including those pertaining to bacteriophage integration, DNA repair, and metabolism and virulence associated with their bacterial host. Several gut viruses also correlated with COVID-19 disease severity, suggesting that the gut virome may calibrate host immunity and regulate severity to SARS-CoV-2 infection.⁵³

Cellular and humoral immunity mediated by T and B cells plays a key role in COVID-19.^{54–57} In particular, B cell-derived antibodies against the spike protein and its receptor-binding domain are relevant in this context, as they prevent virus binding to epithelial cells.^{54,55} Moreover, an expansion of T follicular helper cells is indicative of a matured humoral immune response allowing memory B cells to prevent possible reinfection.⁵⁴ However, the reasons for persistent antigen-specific memory responses in COVID-19 remained mysterious, despite the above findings, suggesting the presence of reservoirs for viral proteins that might trigger prolonged immune responses. Interestingly, there is a high rate of positive PCR findings in stool even weeks or months after respiratory samples became negative, indicating persistence of mucosal SARS-CoV-2 mRNA in patients with COVID-19.⁵⁸ This could be due to infection of intestinal epithelial cells, as SARS-CoV-2 could infect and




Figure 1 The gut virome and microbiome are displayed in the lumen. SARS-CoV-2 may infect intestinal epithelial cells, leading to persistent activation of memory B and T cell responses. Moreover, in HIV-infected patients, the gut serves as a reservoir for infected cells, which may reach the bloodstream via lymph nodes and immune cell trafficking. DC, dendritic cell; IEC, intestinal epithelial cell; ILC, innate lymphoid cell; M, macrophage; TRM, tissue resident memory T cell.

productively replicate in human intestinal tissues *ex vivo* with subsequent release of infectious virus particles, suggesting that the GI tract serves as a potential route of virus dissemination within an infected host.⁵⁹ Consistently, a recent study showed that SARS-CoV-2 nucleoprotein (N) is present in intestinal epithelial cells of approximately 35% of patients with COVID-19 even several weeks or months after initial diagnosis, indicating antigen persistence in the intestine after resolution of clinical illness.⁶⁰ Given the high turnover of intestinal epithelial cells persistence of the viral antigen is also suggestive of continuous virus replication. As even small amounts of persistent viral antigen have been suggested to fuel antibody evolution, the observation on persistent N protein expression over prolonged periods of time is consistent with the relative persistence of SARS-CoV-2 IgA antibodies and continued antibody evolution, as well as persistent polyfunctional SARS-CoV-2 antigen-specific B and T cell memory responses.⁶¹ These immune responses in turn are likely to support a rapid and effective adaptive immune response to the virus on re-exposure and thus provide a critical cornerstone in immune protection against COVID-19. However, one cannot exclude the possibility that long-term shedding of viruses might contribute to long-term COVID-19 and we feel that this point requires further investigation.

IMPLICATIONS AND CONCLUSION

Recent studies have started to explore the composition and function of the gut virome, which provides a rich reservoir for bacteriophages and viruses. These studies have highlighted the concept that bacteriophages and viruses play a seminal role in shaping the bacterial microbiome and are important in controlling gut homeostasis.^{7–8} Moreover, transfer of bacteriophages and viruses during FMT may contribute to treatment efficacy in inflammatory disorders. This concept was underlined by a recent study demonstrating that transfer and effective colonisation of donor Caudovirales bacteriophages correlates with treatment efficacy of FMT in *Clostridioides difficile* infection (CDI).⁶² Others demonstrated that sterile faecal filtrates from donor faeces were effective in treating recurrent CDI in five patients and the gut virome at several months post-FMT related to the donor virome.^{63–64} These findings indicate that apart from live bacteria, other components of the microbiota such as bacteriophages, antimicrobial compounds or metabolites contribute to re-establishment of the intestinal microbiota in FMT. These findings

have important implications for future design of FMT trials in patients with inflammatory disorders and suggest that careful characterisation of the gut virome should be considered for FMT donors. In addition, the potential of selective transfer of viral communities should be explored. For instance, a recent study⁶⁵ described that selective transfer of caecal viral communities from mice with a lean phenotype into mice with an obese phenotype leads to reduced weight gain and normalised blood glucose parameters, suggesting that faecal virome transfer might be used in the future for treatment of diseases.

Another important concept arising from studies in patients with viral infections relates to the gut as potential reservoir for recurrent disease or continuous immune cell stimulation (figure 1). For instance, the intestine has been identified as a crucial reservoir for HIV-infected immune cells that may reach the bloodstream via immune cell trafficking, thereby potentially impairing the efficacy of ART in affected patients. Furthermore, more recent studies of patients with COVID-19 on SARS-CoV-2 infection uncovered a potentially important role of persistent infection of intestinal epithelial cells for prolonged memory immune responses and prevention of reinfection. Collectively, these data shed new light on the role of the intestine as reservoir for viral infections and provide fascinating insights into the function of intestinal epithelial and immune cells in controlling host responses during infections.

Contributors All authors drafted the manuscript and discussed the content.

Funding This work was supported by Deutsche Forschungsgemeinschaft (DFG) with grant numbers SFB1181, TRR241 and FOR2438, as well as by IZKF Erlangen.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not required.

Provenance and peer review Commissioned; externally peer reviewed.

This article is made freely available for use in accordance with BMJ's website terms and conditions for the duration of the covid-19 pandemic or until otherwise determined by BMJ. You may use, download and print the article for any lawful, non-commercial purpose (including text and data mining) provided that all copyright notices and trade marks are retained.

© Author(s) (or their employer(s)) 2021. No commercial re-use. See rights and permissions. Published by BMJ.



To cite Neurath MF, Überla K, Ng SC. *Gut* Epub ahead of print: [please include Day Month Year]. doi:10.1136/gutjnl-2021-324622

Received 22 March 2021

Revised 13 April 2021

Accepted 15 April 2021

Gut 2021;0:1–4.

doi:10.1136/gutjnl-2021-324622

ORCID iDs

Markus F Neurath <http://orcid.org/0000-0003-4344-1474>

Siew C Ng <http://orcid.org/0000-0002-6850-4454>

REFERENCES

- 1 Franzosa EA, Sirota-Madi A, Avila-Pacheco J, *et al*. Gut microbiome structure and metabolic activity in inflammatory bowel disease. *Nat Microbiol* 2019;4:293–305.
- 2 Ananthakrishnan AN, Luo C, Yajnik V, *et al*. Gut microbiome function predicts response to Anti-integrin biologic therapy in inflammatory bowel diseases. *Cell Host Microbe* 2017;21:603–10.
- 3 Chiriac MT, Mahapatro M, Neurath MF, *et al*. The microbiome in visceral medicine: inflammatory bowel disease, obesity and beyond. *Visc Med* 2017;33:153–62.
- 4 Vemuri R, Shankar EM, Chieppa M, *et al*. Beyond just bacteria: functional biomes in the gut ecosystem including Virome, Mycobiome, Archaeome and helminths. *Microorganisms* 2020;8. doi:10.3390/microorganisms8040483. [Epub ahead of print: 28 Mar 2020].
- 5 Liang G, Zhao C, Zhang H, *et al*. The stepwise assembly of the neonatal virome is modulated by breastfeeding. *Nature* 2020;581:470–4.
- 6 York A. Prophages are gut virome pioneers. *Nat Rev Microbiol* 2020;18:317.
- 7 Shkoporov AN, Clooney AG, Sutton TDS, *et al*. The human gut Virome is highly diverse, stable, and individual specific. *Cell Host Microbe* 2019;26:e525:527–41.
- 8 Gregory AC, Zablocki O, Zayed AA, *et al*. The gut Virome database reveals age-dependent patterns of Virome diversity in the human gut. *Cell Host Microbe* 2020;28:e728:724–40.
- 9 Zuo T, Sun Y, Wan Y, *et al*. Human-Gut-DNA Virome variations across geography, ethnicity, and urbanization. *Cell Host Microbe* 2020;28:e744:741–51.
- 10 Lim ES, Zhou Y, Zhao G, *et al*. Early life dynamics of the human gut virome and bacterial microbiome in infants. *Nat Med* 2015;21:1228–34.
- 11 Yang J-Y, Kim M-S, Kim E, *et al*. Enteric viruses ameliorate gut inflammation via Toll-like receptor 3 and Toll-like receptor 7-mediated interferon- β production. *Immunity* 2016;44:889–900.
- 12 Karst SM. Viral safeguard: the enteric Virome protects against gut inflammation. *Immunity* 2016;44:715–8.
- 13 Zuo T, Lu X-J, Zhang Y, *et al*. Gut mucosal virome alterations in ulcerative colitis. *Gut* 2019;68:1169–79.
- 14 Ungaro F, Massimino L, Furfaro F, *et al*. Metagenomic analysis of intestinal mucosa revealed a specific eukaryotic gut virome signature in early-diagnosed inflammatory bowel disease. *Gut Microbes* 2019;10:149–58.
- 15 Norman JM, Handley SA, Baldrige MT, *et al*. Disease-Specific alterations in the enteric virome in inflammatory bowel disease. *Cell* 2015;160:447–60.
- 16 Wang W, Jovel J, Halloran B, *et al*. Metagenomic analysis of microbiome in colon tissue from subjects with inflammatory bowel diseases reveals interplay of viruses and bacteria. *Inflamm Bowel Dis* 2015;53:1–1427.
- 17 Frias AH, Jones RM, Fifadara NH, *et al*. Rotavirus-Induced IFN- β promotes anti-viral signaling and apoptosis that modulate viral replication in intestinal epithelial cells. *Innate Immun* 2012;18:294–306.

- 18 Hu D, Zhang D, Zheng S, *et al.* Association of ulcerative colitis with FUT2 and FUT3 polymorphisms in patients from Southeast China. *PLoS One* 2016;11:e0146557.
- 19 McGovern DPB, Jones MR, Taylor KD, *et al.* Fucosyltransferase 2 (FUT2) non-secretor status is associated with Crohn's disease. *Hum Mol Genet* 2010;19:3468–76.
- 20 Miyoshi J, Yajima T, Okamoto S, *et al.* Ectopic expression of blood type antigens in inflamed mucosa with higher incidence of FUT2 secretor status in colonic Crohn's disease. *J Gastroenterol* 2011;46:1056–63.
- 21 Chamailard M, Cesaro A, Lober P-E, *et al.* Decoding norovirus infection in Crohn's disease. *Inflamm Bowel Dis* 2014;20:767–70.
- 22 Coughlan S, Das A, O'Herlihy E, *et al.* The gut virome in irritable bowel syndrome differs from that of controls. *Gut Microbes* 2021;13:1–15.
- 23 Legoff J, Resche-Rigon M, Bouquet J, *et al.* The eukaryotic gut virome in hematopoietic stem cell transplantation: new clues in enteric graft-versus-host disease. *Nat Med* 2017;23:1080–5.
- 24 Zhang F, Zuo T, Yeoh YK, *et al.* Longitudinal dynamics of gut bacteriome, mycobiome and virome after fecal microbiota transplantation in graft-versus-host disease. *Nat Commun* 2021;12:65.
- 25 Turkington CJR, Varadan AC, Grenier SF, *et al.* The viral Janus: viruses as aetiological agents and treatment options in colorectal cancer. *Front Cell Infect Microbiol* 2020;10:601573.
- 26 Giuliani L, Ronci C, Bonifacio D, *et al.* Detection of oncogenic DNA viruses in colorectal cancer. *Anticancer Res* 2008;28:1405–10.
- 27 Cho Y-H, Ro EJ, Yoon J-S, *et al.* 5-Fu promotes stemness of colorectal cancer via p53-mediated Wnt/ β -catenin pathway activation. *Nat Commun* 2020;11:5321.
- 28 Boltin D, Vilkin A, Levi Z, *et al.* JC virus T-antigen DNA in gastrointestinal mucosa of immunosuppressed patients: a prospective, controlled study. *Dig Dis Sci* 2010;55:1975–81.
- 29 Uleri E, Piu C, Caocci M, *et al.* Multiple signatures of the JC polyomavirus in paired normal and altered colorectal mucosa indicate a link with human colorectal cancer, but not with cancer progression. *Int J Mol Sci* 2019;20:5965.
- 30 Nakatsu G, Zhou H, Wu WKK, *et al.* Alterations in enteric Virome are associated with colorectal cancer and survival outcomes. *Gastroenterology* 2018;155:529–41.
- 31 Hannigan GD, Duhaime MB, Ruffin MT, *et al.* Diagnostic potential and interactive dynamics of the colorectal cancer Virome. *mBio* 2018;9. doi:10.1128/mBio.02248-18. [Epub ahead of print: 20 Nov 2018].
- 32 Smith PD, Quinn TC, Strober W, *et al.* Nih conference. gastrointestinal infections in AIDS. *Ann Intern Med* 1992;116:63–77.
- 33 Sim JH, Mukerji SS, Russo SC, *et al.* Gastrointestinal dysfunction and HIV comorbidities. *Curr HIV/AIDS Rep* 2021;18:57–62.
- 34 Lapenta C, Boirivant M, Marini M, *et al.* Human intestinal lamina propria lymphocytes are naturally permissive to HIV-1 infection. *Eur J Immunol* 1999;29:1202–8.
- 35 Wong JK, Yukl SA. Tissue reservoirs of HIV. *Curr Opin HIV AIDS* 2016;11:362–70.
- 36 Godfrey C, Bremer A, Alba D, *et al.* Obesity and fat metabolism in human immunodeficiency virus-infected individuals: immunopathogenic mechanisms and clinical implications. *J Infect Dis* 2019;220:420–31.
- 37 Lim SG, Condez A, Lee CA, *et al.* Loss of mucosal CD4 lymphocytes is an early feature of HIV infection. *Clin Exp Immunol* 1993;92:448–54.
- 38 Gori A, Rizzardini G, Van't Land B, *et al.* Specific prebiotics modulate gut microbiota and immune activation in HAART-naïve HIV-infected adults: results of the "COPA" pilot randomized trial. *Mucosal Immunol* 2011;4:554–63.
- 39 Villar-García J, Hernández JJ, Güerri-Fernández R, *et al.* Effect of probiotics (*Saccharomyces boulardii*) on microbial translocation and inflammation in HIV-treated patients: a double-blind, randomized, placebo-controlled trial. *J Acquir Immune Defic Syndr* 2015;68:256–63.
- 40 Chun T-W, Nickle DC, Justement JS, *et al.* Persistence of HIV in gut-associated lymphoid tissue despite long-term antiretroviral therapy. *J Infect Dis* 2008;197:714–20.
- 41 Yukl SA, Gianella S, Sinclair E, *et al.* Differences in HIV burden and immune activation within the gut of HIV-positive patients receiving suppressive antiretroviral therapy. *J Infect Dis* 2010;202:1553–61.
- 42 Yukl SA, Sinclair E, Somsouk M, *et al.* A comparison of methods for measuring rectal HIV levels suggests that HIV DNA resides in cells other than CD4+ T cells, including myeloid cells. *AIDS* 2014;28:439–42.
- 43 Neurath MF. Targeting immune cell circuits and trafficking in inflammatory bowel disease. *Nat Immunol* 2019;20:970–9.
- 44 Alzahrani J, Hussain T, Simar D, *et al.* Inflammatory and immunometabolic consequences of gut dysfunction in HIV: parallels with IBD and implications for reservoir persistence and non-AIDS comorbidities. *EBioMedicine* 2019;46:522–31.
- 45 Jhu. COVID-19 Dashboard by the center for systems science and engineering (CSSE), 2021. Available: <https://coronavirus.jhu.edu/map.html>
- 46 Hoffmann M, Kleine-Weber H, Schroeder S, *et al.* SARS-CoV-2 cell entry depends on ACE2 and TMPRSS2 and is blocked by a clinically proven protease inhibitor. *Cell* 2020;181:271–80.
- 47 Patankar JV, Chiriac M, Lehmann M. The SARS-CoV-2 attachment receptor ACE2 is decreased in Crohn's disease and regulated by microbial and inflammatory signaling. *Gastroenterology* 2020:S0016-5085(0020)35278-35271.
- 48 Lamers MM, Beumer J, van der Vaart J, *et al.* SARS-CoV-2 productively infects human gut enterocytes. *Science* 2020;369:50–4.
- 49 Livanos AE, Jha D, Cossarini F, *et al.* Intestinal host response to SARS-CoV-2 infection and COVID-19 outcomes in patients with gastrointestinal symptoms. *Gastroenterology* 2021;gastro.2021.02.056. doi:10.1053/j.gastro.2021.02.056
- 50 Wölfel R, Corman VM, Guggemos W, *et al.* Virological assessment of hospitalized patients with COVID-2019. *Nature* 2020;581:465–9.
- 51 Amirian ES. Potential fecal transmission of SARS-CoV-2: current evidence and implications for public health. *Int J Infect Dis* 2020;95:363–70.
- 52 Cao J, Wang C, Zhang Y, *et al.* Integrated gut virome and bacteriome dynamics in COVID-19 patients. *Gut Microbes* 2021;13:1–21.
- 53 Zuo T, Liu Q, Zhang F, *et al.* Temporal landscape of human gut RNA and DNA virome in SARS-CoV-2 infection and severity. *Microbiome* 2021;9:91.
- 54 Cox RJ, Brokstad KA. Not just antibodies: B cells and T cells mediate immunity to COVID-19. *Nat Rev Immunol* 2020;20:581–2.
- 55 Grifoni A, Weiskopf D, Ramirez SI, *et al.* Targets of T cell responses to SARS-CoV-2 coronavirus in humans with COVID-19 disease and unexposed individuals. *Cell* 2020;181:e1415:1489–501.
- 56 Sekine T, Perez-Potti A, Rivera-Ballesteros O, *et al.* Robust T cell immunity in convalescent individuals with asymptomatic or mild COVID-19. *Cell* 2020;183:e114:158–68.
- 57 Swadling L, Maini MK. T cells in COVID-19 - united in diversity. *Nat Immunol* 2020;21:1307–8.
- 58 Wu Y, Guo C, Tang L, *et al.* Prolonged presence of SARS-CoV-2 viral RNA in faecal samples. *Lancet Gastroenterol Hepatol* 2020;5:434–5.
- 59 Chu H, Chan JF-W, Wang Y, *et al.* SARS-CoV-2 Induces a More Robust Innate Immune Response and Replicates Less Efficiently Than SARS-CoV in the Human Intestines: An Ex Vivo Study With Implications on Pathogenesis of COVID-19. *Cell Mol Gastroenterol Hepatol* 2021;11:771–81.
- 60 Gaebler C, Wang Z, Lorenzi JCC, *et al.* Evolution of antibody immunity to SARS-CoV-2. *Nature* 2021;591:639–44.
- 61 Breton G, Mendoza P, Hägglöf T, *et al.* Persistent cellular immunity to SARS-CoV-2 infection. *J Exp Med* 2021;218. doi:10.1084/jem.20202515. [Epub ahead of print: 05 Apr 2021].
- 62 Zuo T, Wong SH, Lam K. Bacteriophage transfer during faecal microbiota transplantation in *Clostridium difficile* infection is associated with treatment outcome. *Gut* 2018;67:634–43.
- 63 Ott SJ, Waetzig GH, Rehman A, *et al.* Efficacy of sterile fecal filtrate transfer for treating patients with *Clostridium difficile* infection. *Gastroenterology* 2017;152:e797:799–811.
- 64 Broecker F, Russo G, Klumpp J, *et al.* Stable core virome despite variable microbiome after fecal transfer. *Gut Microbes* 2017;8:214–20.
- 65 Rasmussen TS, Mentzel CMJ, Kot W, *et al.* Faecal virome transplantation decreases symptoms of type 2 diabetes and obesity in a murine model. *Gut* 2020;69:2122–30.