

# *Brachy bacterium muris* Detected in a Hepatocellular Carcinoma Patient with Pleural Effusion: A Case Report

Yujing Wu<sup>1,\*</sup>, Sumei Wang<sup>1,\*</sup>, Guangye Wu<sup>2</sup>, Jiandong Zhang<sup>1</sup>, Shuye Liu<sup>1</sup>

<sup>1</sup>The Third Central Hospital of Tianjin, Tianjin Key Laboratory of Extracorporeal Life Support for Critical Diseases, Artificial Cell Engineering Technology Research Center, Tianjin Institute of Hepatobiliary Disease, Tianjin, People's Republic of China; <sup>2</sup>Second Teaching Hospital of Tianjin University of Traditional Chinese Medicine, Tianjin Traditional Chinese Medicine Rehabilitation Center, Tianjin, People's Republic of China

\*These authors contributed equally to this work

Correspondence: Shuye Liu, Tel +86-022-84112332, Fax +86 022-24384350, Email liushuye@tmu.edu.cn

**Background:** *Brachy bacterium muris* is a species of Gram positive and strictly aerobic bacterium. It was first reported in 2003 after being isolated from the liver of a laboratory mouse strain. It was also found on human skin and nasal cavity. Herein, we present the first case pleural effusion infection in humans caused by *Brachy bacterium muris*.

**Case Presentation:** A 65-year-old man was admitted to our hospital for a 4-week history of fever, accompanied by chills, occasional abdominal pain, occasional chest tightness and shortness of breath. On the day of hospitalization, thoracentesis was performed and 1000mL of yellow cloudy fluid was released. Result of pleural fluid culture was positive and *B. muris* was identified using 16S rDNA amplification and sequence comparisons.

**Conclusion:** To our knowledge, this is the first report of pleural effusion infection caused by *B. muris*. *B. muris* can be pathogenic in humans.

**Keywords:** *Brachy bacterium muris*, pleural effusion, infection

## Background

Genus *Brachy bacterium* was first discovered and isolated in 1988.<sup>1</sup> Up to now, there were 25 child taxa with a validly published and correct name (<https://lpsn.dsmz.de/genus/brachybacterium>). Genus *Brachy bacterium* can be isolated from various environmental samples, such as seawater,<sup>2</sup> plant roots and branches,<sup>3,4</sup> river,<sup>5</sup> surface of cheeses,<sup>6</sup> coastal sand<sup>7</sup> and soil.<sup>8</sup> A few studies have also reported the isolation of this genus from human samples.<sup>9–11</sup>

*Brachy bacterium muris* was first reported in 2003 after being isolated from the liver of a laboratory mouse strain.<sup>12</sup> In 2017, *Brachy bacterium muris* was also isolated from Chaka Salt Lake (Qinghai, China).<sup>13</sup> *Brachy bacterium muris* was also found in human samples, such as human nasal cavity<sup>14</sup> and skin.<sup>15</sup> However, to our knowledge, there are no reports of the species *Brachy bacterium muris* infecting humans and causing disease. We herein report the first case of *Brachy bacterium muris* isolated from pleural effusion.

## Case Presentation

A man with a medical history of hepatitis B for more than ten years was diagnosed with hepatocellular carcinoma in June 2021. The patient accepted treatment with radiofrequency on July 1, 2021. But in March 2022, the hepatocellular carcinoma recurred and he underwent radiofrequency treatment again. In November 2022, this 65-year-old man presented to Tianjin Third Central Hospital (Tianjin), with a 4-week history of fever, accompanied by chills, occasional abdominal pain, chest tightness and shortness of breath. On admission, this patient presented chronic liver disease face, liver palm and loss of breath sounds in the right lung. The patient had decompensated cirrhosis of the liver with a Child-Pugh classification of B and a small amount of ascites.

On the day of hospitalization, thoracentesis was performed and 1000mL of yellow cloudy fluid was released. Laboratory tests of whole blood showed the following results: white blood cell count  $2.17 \times 10^9/L$  (neutrophil 77.1%), red blood cell count  $3.06 \times 10^{12}/L$ , hemoglobin 90 g/L and platelet count  $46 \times 10^9/L$ . His serum hepatitis B surface antigen and hepatitis B core antibody were 13.41 IU/mL and 8.78 IU/mL, respectively. His procalcitonin level in serum was 0.33 ng/mL. The result of G test was 114.30pg/mL. The G test detects (1,3)-b-D-glucan, a component of the cell wall of fungi. Pleural effusion tests revealed a white blood cell count of  $2.3 \times 10^9/L$  with 20% neutrophils and 80% lymphocytes. The specific gravity of the pleural fluid was 1.015 and the Rivalta test result was positive. The above results suggest that the pleural fluid was exudative and the patient had a pleural fluid infection. Empirical anti-infective treatment with piperacillin/tazobactam was initiated after collection of pleural fluid for culture. A 4.5g of piperacillin/tazobactam was dissolved in 100mL of saline and administered intravenously every 8 hours, three times a day. The next day the patient underwent thoracentesis drainage and released 700 mL of pleural fluid. Two days after the patient received anti-infection and intermittent pleural fluid release treatment, the patient's temperature returned to normal.

The infecting agent was isolated from pleural fluid by performing in vitro culture and then the infecting agent was purified in blood agar culture. Next, the organism was subjected to matrix-assisted laser desorption/ionization time-of-flight mass spectrometry (MALDI-TOF MS) analysis. We used a Vitek MS system. Unfortunately, our database did not include this microorganism. We outsourced microbial sequencing to Beijing RuiBio BioTech Co. The microorganism was identified using 16S rDNA amplification and sequence comparisons. Two primers were used. Their sequences were "AGAGTTTGATCATGGCTC" and "TAGGGTTACCTTGTTACGACTT". Comparison with sequences in the BLAST database (<http://www.ncbi.nlm.nih.gov/BLAST>) revealed that the sequence has 99.131% similarity to the *Brachybacterium muris* sequence (GenBank accession number NR\_024571.1). The sequence was attached in [Supplementary 1](#). As shown in [Figure 1](#), culture on blood agar plate grew white, moist, and smooth colonies after 24 h of incubation at 35 °C in a 5% CO<sub>2</sub> atmosphere.



**Figure 1** Culture on blood agar plate.

## Discussion and Conclusion

*Brachybacterium muris* is a species of Gram positive, strictly aerobic, yellow-pigmented bacterium. The cells are coccoid during the stationary phase, and irregular rods during the exponential phase. It was first isolated from the liver of a laboratory mouse. The species was first described in 2003, and the name is derived from the Latin *muris* (mouse). The optimum growth temperature for *B. muris* is 25–37 °C. It can grow in the 15–42 °C range and in pH 6.0–9.0.<sup>12</sup> *B. muris* have been reported in the literature to be found in human skin/faeces.<sup>15,16</sup> However, there are few reports of human diseases caused by this bacterium.

In this case, the patient had a history of hepatitis B-related liver cancer and was treated with radiofrequency. The infecting organism may have come from the environment or from the patient's skin. It was not entirely clear that how he got the infection. Empirical treatment with piperacillin/tazobactam seemed to be effective, as he experienced significant relief of the infection two days after the administration of the drug. We identified the isolated strains by MALDI-TOF and 16S rRNA sequencing.

In summary, we present the first case of pleural effusion infection caused by *B. muris*. The organism was identified by 16S ribosomal RNA gene-sequencing analysis. Although human infection is rare, *B. muris* can be pathogenic in humans.

## Consent for Publication

Written informed consent was provided by the patient for the publication of the case details and images. Institutional approval was obtained to publish the case details.

## Funding

This work is supported by Tianjin Key Medical Discipline (Specialty) Construction Project (Grants No. TJYXZDXK-047A).

## Disclosure

The authors report no conflicts of interest in this work.

## References

- Collins MD, Brown J, Jones D. *Brachybacterium faecium* gen. nov., sp. nov., a coryneform bacterium from poultry deep litter. *Int J Syst Bacteriol.* 1988;38:45–48. doi:10.1099/00207713-38-1-45
- Kaur G, Kumar N, Mual P, et al. *Brachybacterium aquaticum* sp. nov., a novel actinobacterium isolated from seawater. *Int J Syst Evol Microbiol.* 2016;66(11):4705–4710. doi:10.1099/ijsem.0.001414
- Gontia I, Kavita K, Schmid M, et al. *Brachybacterium saurashtrense* sp. nov., a halotolerant root-associated bacterium with plant growth-promoting potential. *Int J Syst Evol Microbiol.* 2011;61(Pt 12):2799–2804. doi:10.1099/ijms.0.023176-0
- Liu Y, Zhai L, Yao S, et al. *Brachybacterium hainanense* sp. nov., isolated from noni (*Morinda citrifolia* L.) branch. *Int J Syst Evol Microbiol.* 2015;65(11):4196–4201. doi:10.1099/ijsem.0.000559
- Alam SA, Saha P. Biodegradation of p-nitrophenol by a member of the genus *Brachybacterium*, isolated from the river Ganges. *Biotech.* 2022;12(9):213. doi:10.1007/s13205-022-03263-7
- Schubert K, Ludwig W, Springer N, et al. Two coryneform bacteria isolated from the surface of French Gruyère and Beaufort cheeses are new species of the genus *Brachybacterium*: *brachybacterium alimentarium* sp. nov. and *Brachybacterium tyrofermentans* sp. nov. *Int J Syst Bacteriol.* 1996;46(1):81–87. doi:10.1099/00207713-46-1-81
- Chou JH, Lin KY, Lin MC, et al. *Brachybacterium phenoliresistens* sp. nov., isolated from oil-contaminated coastal sand. *Int J Syst Evol Microbiol.* 2007;57(Pt 11):2674–2679. doi:10.1099/ijms.0.65019-0
- Singh H, Du J, Yang JE, et al. *Brachybacterium horti* sp. nov., isolated from garden soil. *Int J Syst Evol Microbiol.* 2016;66(1):189–195. doi:10.1099/ijsem.0.000696
- Renvoise A, Aldrovandi N, Raoult D, et al. *Helcobacillus massiliensis* gen. nov., sp. nov., a novel representative of the family Dermabacteraceae isolated from a patient with a cutaneous discharge. *Int J Syst Evol Microbiol.* 2009;59(Pt 9):2346–2351. doi:10.1099/ijms.0.003319-0
- Kuete E, Mbogning Fonkou MD, Mekhalif F, et al. *Brachybacterium timonense* sp. nov., a new bacterium isolated from human sputum. *New Microbes New Infect.* 2019;29(31):100568. doi:10.1016/j.nmni.2019.100568
- Tamai K, Akashi Y, Yoshimoto Y, et al. First case of a bloodstream infection caused by the genus *Brachybacterium*. *J Infect Chemother.* 2018;24(12):998–1003. doi:10.1016/j.jiac.2018.06.005
- Buczolits S, Schumann P, Weidler G, et al. *Brachybacterium muris* sp. nov., isolated from the liver of a laboratory mouse strain. *Int J Syst Evol Microbiol.* 2003;53(Pt 6):1955–1960. doi:10.1099/ijms.0.02728-0
- Gu D, Jiao Y, Wu J, et al. Optimization of EPS Production and Characterization by a Halophilic Bacterium, *Kocuria rosea* ZJUQH from Chaka Salt Lake with Response Surface Methodology. *Molecules.* 2017;22(5):814. doi:10.3390/molecules22050814
- Ruokolainen L, Paalanen L, Karkman A, et al. Significant disparities in allergy prevalence and microbiota between the young people in Finnish and Russian Karelia. *Clin Exp Allergy.* 2017;47(5):665–674. doi:10.1111/cea.12895

15. Leung MHY, Tong X, Bastien P, et al. Changes of the human skin microbiota upon chronic exposure to polycyclic aromatic hydrocarbon pollutants. *Microbiome*. 2020;8(1):100. doi:10.1186/s40168-020-00874-1
16. Paramsothy S, Kamm MA, Kaakoush NO, et al. Multidonor intensive faecal microbiota transplantation for active ulcerative colitis: a randomised placebo-controlled trial. *Lancet*. 2017;389(10075):1218–1228. doi:10.1016/S0140-6736(17)30182-4

Infection and Drug Resistance

Dovepress

## Publish your work in this journal

Infection and Drug Resistance is an international, peer-reviewed open-access journal that focuses on the optimal treatment of infection (bacterial, fungal and viral) and the development and institution of preventive strategies to minimize the development and spread of resistance. The journal is specifically concerned with the epidemiology of antibiotic resistance and the mechanisms of resistance development and diffusion in both hospitals and the community. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/infection-and-drug-resistance-journal>