

Can biowarfare agents be defeated with light?

Fatma Vatansever^{1,2}, Cleber Ferraresi^{1,3,4,5}, Marcelo Victor Pires de Sousa^{1,6}, Rui Yin^{1,2,7}, Ardeshir Rineh^{1,8}, Sulbha K Sharma^{1,9}, and Michael R Hamblin^{1,2,10,*}

¹Wellman Center for Photomedicine; Massachusetts General Hospital; Boston MA USA; ²Harvard Medical School; Department of Dermatology; Boston, MA USA;

³Laboratory of Electro-thermo-phototherapy; Department of Physical Therapy; Federal University of São Carlos; São Paulo, Brazil; ⁴Post-Graduation Program in Biotechnology; Federal University of São Carlos; São Paulo, Brazil; ⁵Optics Group; Physics Institute of Sao Carlos; University of São Paulo; São Carlos, Brazil; ⁶Laboratory of Radiation Dosimetry and Medical Physics; Institute of Physics, São Paulo University, São Paulo, Brazil; ⁷Department of Dermatology; Southwest Hospital; Third Military Medical University; Chongqing, PR China; ⁸School of Chemistry; University of Wollongong; Wollongong, NSW Australia; ⁹Raja Ramanna Centre for Advanced Technology; Indore, India;

¹⁰Harvard-MIT Division of Health Sciences and Technology; Cambridge, MA USA

Keywords: biowarfare, bioterrorism, photocatalysis, photocatalytic inactivation, microbial cells, ultraviolet light, germicidal ultraviolet, photo inactivation, UV dosimeters, titanium dioxide, psorales, photodynamic therapy, blue light inactivation

Biological warfare and bioterrorism is an unpleasant fact of 21st century life. Highly infectious and profoundly virulent diseases may be caused in combat personnel or in civilian populations by the appropriate dissemination of viruses, bacteria, spores, fungi, or toxins. Dissemination may be airborne, waterborne, or by contamination of food or surfaces. Countermeasures may be directed toward destroying or neutralizing the agents outside the body before infection has taken place, by destroying the agents once they have entered the body before the disease has fully developed, or by immunizing susceptible populations against the effects. A range of light-based technologies may have a role to play in biodefense countermeasures. Germicidal UV (UVC) is exceptionally active in destroying a wide range of viruses and microbial cells, and recent data suggests that UVC has high selectivity over host mammalian cells and tissues. Two UVA mediated approaches may also have roles to play; one where UVA is combined with titanium dioxide nanoparticles in a process called photocatalysis, and a second where UVA is combined with psoralens (PUVA) to produce “killed but metabolically active” microbial cells that may be particularly suitable for vaccines. Many microbial cells are surprisingly sensitive to blue light alone, and blue light can effectively destroy bacteria, fungi, and *Bacillus* spores and can treat wound infections. The combination of photosensitizing dyes such as porphyrins or phenothiaziniums and red light is called photodynamic therapy (PDT) or photoinactivation, and this approach cannot only kill bacteria, spores, and fungi, but also inactivate viruses and toxins. Many reports have highlighted the ability of PDT to treat infections and stimulate the host immune system. Finally pulsed (femtosecond) high power lasers have been used to inactivate pathogens with some degree of selectivity. We have pointed to some of the ways light-based technology may be used to defeat biological warfare in the future.

Introduction: Biological Warfare and Bioterrorism Agents

In recent years, the possibility of biological warfare and bioterrorism has become of increasing concern to both military planners and civil defense authorities. The mailing of anthrax spore containing letters to destinations within the United States in 2001 brought the sudden realization that bioterrorism is not merely a theoretical threat but a real and present danger. Since then, much thought and planning has gone into defining possible biowarfare and bioterrorism agents. There are six requirements for these agents that are relevant here:

- 1) A high degree of morbidity and lethality.
- 2) Highly infectious microbes or highly toxic substances.
- 3) Easy to distribute widely in an active form.
- 4) Easy to produce in bulk and store until delivered.
- 5) Reasonably hardy in the environment after distribution.
- 6) Bacteria should be genetically engineered to be resistant to known antibiotic drugs.

The 2001 bioterrorist attacks in the US using anthrax spores and the US Postal Service as the spreading medium have once more emphasized the need of early detection and decontamination of critical facilities in the shortest possible time. During the recent decade there has been a remarkable progress in the detection, protection, and decontamination of biological warfare agents since various and sophisticated detection/decontamination methods have been developed and implemented. Nevertheless the threat of biological warfare agents and their possible use in bioterrorist attacks still remains a leading cause of concern in the global community. Furthermore, in the past decade there have been threats to the global society due to the emergence of new infectious diseases and/or re-emergence of old infectious diseases that were considered eliminated. Adding to the milieu the observed global rise in the antimicrobial resistance, the preparedness of societies against these agents becomes obvious. Under these circumstances it becomes obvious that the field requires better knowledge about the disease agents, more research, better training and diagnostic facilities, and improved public health system¹ (see Table 1).

The emergence of bacterial strains that are resistant to all known antibiotics represents a major challenge to human health.

*Correspondence to: Michael R Hamblin; Email:

hamblin@helix.mgh.harvard.edu

Submitted: 06/03/2013; Revised: 09/10/2013; Accepted: 09/12/2013

<http://dx.doi.org/10.4161/viru.26475>

Table 1. Common biological warfare agent characteristics

Disease	Etiologic agent	Organism persistence	Symptoms	Person to person?	Infective dose (aerosol)	Incubation period	Mortality	Treatment
	Bacterial agents							
Anthrax (inhalation)	Spores of <i>Bacillus anthracis</i> (encapsulated gram-positive bacillus); reservoir: the soil	Spores can be viable for >40 years	Fever, malaise, fatigue, cough, mild chest discomfort, respiratory distress, shock	No	8000–50 000 spores	1–6 d	High once symptoms appear	Ciprofloxacin or doxycycline
Brucellosis	Genus <i>Brucella</i> (<i>B. melitensis</i> , <i>B. abortus</i> , <i>B. suis</i> , <i>B. canis</i>)	6 wks in dust and 10 wks in soil or water	Irregular fever, headache, malaise, chills, sweating, myalgia, joint pain, depression	No	10–100 organisms	5–60 d	5% untreated	Doxycycline + rifampin
Pneumonic plague	<i>Yersinia pestis</i> ; reservoir: rodents	Up to 1 year in soil, 270 d in live tissue	High fever, chills, headache, productive cough-water then bloody	Yes, highly	<100 organisms	2–3 d	High unless treated in 12–24 h	Gentamycin or doxycycline
Q fever	<i>Coxiella burnetii</i> ; reservoir: animals	Withstands heat and drying; persists in environment for weeks to months	Fever, chills, headache, diaphoresis, malaise, fatigue, anorexia, weight loss	Rarely	1–10 organisms	2–14 d	Very low	Tetracycline or doxycycline
Tularemia	<i>Francisella tularensis</i> ; reservoir: rabbits, rodents	Months in moist soil or other media	Fever, headache, malaise, weight loss, nonproductive cough	No	10–50 organisms	1–21 d	Moderate if untreated	Ciprofloxacin, doxycycline, or gentamycin
Glanders	<i>B. mallei</i> ; reservoir horses, mules, donkeys	Stable	Fever, rigors, sweating, myalgia, headache, pleuritis, chest pain, splenomegaly, pustular eruptions	Yes	?	10–14 d	Varies	Amoxicillin, tetracycline, or trimethoprim/sulfa
Melioidosis	<i>Burkholderia pseudomallei</i> ; reservoir: soil and water	Stable	Fever, aching chest pain, cough-productive and nonproductive, severe dyspnea, diarrhea, flushing of the skin, cyanosis, rash that can progress to pustular exanthem	No	?	10–14 d	Moderate if untreated	Amoxicillin, tetracycline, trimethoprim/sulfa, ceftazidime

^aLD₅₀ lethal dose µg/kg; ^bMay be effective; ^cRicin and botulinum toxin are lethal at all levels. The mortality levels terminology is as defined by the Centers for Disease Control. Compiled and modified from reference 246.

Table 1. Common biological warfare agent characteristics (continued)

Disease	Etiologic agent	Organism persistence	Symptoms	Person to person?	Infective dose (aerosol)	Incubation period	Mortality	Treatment
	Viral agents							
Smallpox	Variola, poxvirus family; reservoir: humans	Very stable	Fever, rigors, severe headache, backache, malaise, vomiting, delirium, acute papular dermatitis on the face, hands, and forearms which is spreading to the lower extremities	Yes, highly	Assumed low (10–100 organisms)	7–17 d	High to moderate	Cidofovir ^b
Venezuelan viral encephalitis	VEE virus, an arthropodborne alphavirus; reservoir: rodent-mosquito cycles; transmission through mosquitos	Relatively unstable in the environment	Fever, rigors, severe headache, photophobia, malaise, nausea, vomiting, diarrhea	Low	10–100 organisms	1–5 d	Varies	Supportive care
Viral hemorrhagic fevers	VHF virus, lipid-enveloped viruses with single-stranded RNA families	Relatively unstable in the environment	Fever, malaise, myalgia, prostration, vascular permeability may present as conjunctival injection and petechial hemorrhage and progress to mucous membrane hemorrhage and shock	Moderate	1–10 organisms; All VHF transmitted via aerosols, exception dengue	4–21 d	5–90% case fatality rate depending on the virus	Ribavirin or supportive care
Ebola	Four viruses: Bundibugyo virus, Ebola virus, Sudan virus, and Tai Forest virus of the genus <i>Ebolavirus</i> , family <i>Filoviridae</i> ; reservoir: fruit bats <i>Pteropodidae</i> family, plants, arthropods, birds	Stable	Intense weakness, muscle pain, headache, soar throat, vomiting, diarrhea, rash, impaired kidney and liver functions	Yes	?	1–21 d	90% fatality	
Lassa	Lassa virus, a member of <i>Arenaviridae</i> virus family, single-stranded RNA virus; reservoir: rodents	Stable	Fever, retrosternal pain, sore throat, back pain, cough, abdominal pain, vomiting, diarrhea, facial swelling, proteinuria, mucosal bleeding, hearing loss, tremors	Yes	?	1–3 wk	Moderate	Ribavirin or supportive care

^aLD₅₀ lethal dose µg/kg; ^bMay be effective; ^cRicin and botulinum toxin are lethal at all levels. The mortality levels terminology is as defined by the Centers for Disease Control. Compiled and modified from reference 246.

Table 1. Common biological warfare agent characteristics (continued)

Disease	Etiologic agent	Organism persistence	Symptoms	Person to person?	Infective dose (aerosol)	Incubation period	Mortality	Treatment
	Toxins							
Botulism clostridium^c	Group of seven toxins produced by <i>Clostridium botulinum</i> ; reservoir: soil, animals, fish	Weeks in non-moving water and soil	Drooping eyelids, general weakness, dizziness, dry mouth and throat, blurred and double vision, progressive descending symmetrical paralysis	No	0.001 mg/kg LD50 ^a	12–36 h up to several days	High without respiratory support	Antitoxin, supportive care
Ricin^c	Derived from the beans of the castor plant <i>Ricinus communis</i> ; reservoir: castor beans	Stable	Aerosol route: fever, chest tightness, cough, hypothermia; Oral route: gastro-intestinal hemorrhage	No	3–5 ul/kg LD50	18–24 h	High	Inhalation: supportive; care; GI: lavage, charcoal, cathartics
Staphylococcal Enterotoxin B	Produced by <i>S. aureus</i>	Resistant to freezing; heat-stable	Sudden onset of fever, chills, headache, myalgias, non-productive cough	No	30 ug/person incapacitation	3–12 h after inhalation	<1%	Supportive care
Saxitoxin	Marine dinoflagellates of the genus <i>Gonyaulax</i> ; reservoir: shellfish	Stable	Severe to life-threatening paralytic neuromuscular condition, respiratory paralysis and failure	No	?	10 min to several hours after ingestion	Low	Superactivated charcoal
T-2 Mycotoxins trichothecene	A group of 40 compounds produced by molds of the genus <i>Fusarium</i>	Stable for years at room temp	Skin pain, redness, necrosis, sloughing of epidermis, wheezing, chest pain, hemoptysis	No	Moderate	Minutes to hours	Moderate	Supportive care

^aLD₅₀ lethal dose µg/kg; ^bMay be effective; ^cRicin and botulinum toxin are lethal at all levels. The mortality levels terminology is as defined by the Centers for Disease Control. Compiled and modified from reference 246.

One of the most common bacteria, *Staphylococcus aureus* has developed resistance to β-lactams (known as methicillin-resistant *S. aureus* or MRSA) and its vancomycin-resistant counterpart (VRSA) have been isolated from infected patients in various parts of the world. Other species, such as *Streptococcus pyogenes*, are highly virulent and systemic infection can result in death in times as short as 48 h. As a consequence, antibiotic-resistant microorganisms are potentially near-ideal biological weapons that could be used either by enemy combatants on foreign battlefields or by terrorists who have infiltrated the country. Antibiotic-resistant, virulent strains of common microorganisms are particularly attractive as terrorist weapons because no security screening is in effect for common species. Even if detected, the antibiotic-resistant nature of the microorganism would initially remain hidden and no alarms would be raised until large-scale contamination and infection had occurred. These issues make it imperative that broadly-based alternative strategies be developed for the neutralization of drug-resistant biological pathogens.

The deliberate creation of pan-resistant bacterial strains is forbidden in laboratories in most Western countries, but the techniques of genetic engineering are relatively well understood and could easily be replicated in countries that are rumored to sponsor terrorism. Therefore effective countermeasures against biological weapons should be able to deal with multiple classes of biological agents regardless of whether they have been engineered to be resistant to all known antibiotics.

There are many potential bioterrorism agents such as bacteria, viruses, fungi and toxins that can be spread by air, water or food. In this context, we emphasize some of these microorganisms due their elevated capabilities of being lethally dangerous or easily dispersible:

1) In gram-negative bacteria, *Francisella tularensis* causes tularemia or rabbit fever, which is debilitating or

even fatal.² *Brucella melitensis* is also gram-negative and responsible for the contagious disease of brucellosis in sheep, goats, cattle, and in humans causing fever, sweats, anorexia, fatigue, malaise, weight loss, and depression.³ A third gram-negative bacterium is *Yersinia pestis*, which infects humans and other animals causing plague or “the black death”. This bacterium is primarily a disease of rodents or other wild mammals that usually is transmitted by fleas and often is fatal. Human *Yersinia* infection takes three main forms: pneumonic, septicemia, and bubonic plagues.⁴ A fourth gram-negative species is *Burkholderia pseudomallei*, which causes glanders in animals and melioidosis in humans with a mortality rate of 20–50%.⁵

2) Among the gram-positive bacteria, *S. aureus* is the most well-known bacterium and is frequently found in the human respiratory tract and on the skin causing skin infections and respiratory diseases beyond promote infections through potent protein toxins produced by it. In addition, MRSA is a widespread antibiotic-resistant strain and has become a major problem in hospitals in the United States.⁶ *S. pyogenes* is also a gram-positive bacterium that causes invasive and severe infection including sepsis and osteomyelitis partly due to its ability to carry out hemolysis releasing hemoglobin.⁷

3) *Bacillus anthracis*, *Bacillus cereus*, and *Bacillus thuringiensis* are gram-positive bacteria that produce hardy endospores that can be easily disseminated. *B. cereus* is endemic and can be transmitted through food while *B. thuringiensis* produces intracellular protein crystals toxic to a wide number of insect larvae. *B. anthracis* is a rod-shaped bacterium that causes anthrax disease and often is lethal. In addition, these bacteria are similar because they can produce spores and thus infect larger areas in bioterrorism actions.⁸

4) Viruses such the etiologic agents of Variola, Ebola, and Lassa are very dangerous. Variola virus is the etiological agent of smallpox, causes 20–30% mortality, and persists in an infectious state for many days in dried crusts from skin lesions as well as in fluid from vesicles.⁹ Ebola virus causes severe hemorrhagic fever in humans and primates resulting in mortality rates between 80–90%.⁹ Lassa virus causes Lassa fever that is endemic in West Africa, infecting 2 million people per year and resulting in 5000–10 000 fatalities annually.⁹

5) *Clostridium botulinum* is a gram-positive anaerobic bacterium and produces the most potent known neurotoxin responsible for botulism, which promotes neuromuscular weakness or paralysis.¹⁰

Historical evidence of the use of biological warfare is somewhat sketchy. In April and May 1979, an unusual anthrax epidemic occurred in Sverdlovsk, Union of Soviet Socialist Republics. Soviet officials attributed it to consumption of contaminated meat but US agencies attributed it to inhalation of spores accidentally released at a military microbiology facility in the city. Epidemiological data show that most victims worked or lived in a narrow zone extending from the military facility to the southern city limit. Further south, livestock died of anthrax along the extended axis of the zone. The zone paralleled the northerly wind that prevailed shortly before the outbreak. It was concluded

that the escape of an aerosol of anthrax pathogen at the military facility caused the outbreak.¹¹

The difficulty faced in decontaminating the environment from biological weapons agents can be illustrated by the historical story of Gruinard Island. British military scientists from Porton Down in 1942, during the Second World War, had tested methods to disseminate anthrax spores on a remote and uninhabited island off the Scottish coast. Military scientists exploded a series of anthrax-spore laden bombs, testing their killing efficiency using sheep.¹² Initial efforts to decontaminate the island after the biological warfare trials failed due to the high durability of anthrax spores. After 44 years of complete quarantine, Gruinard Island was finally decontaminated in 1986 with 280 tons of formaldehyde diluted in seawater being sprayed over all 196 hectares of the island and the worst-contaminated topsoil around the dispersal site being physically removed.¹³ A flock of sheep was then placed on the island and remained healthy.¹⁴

In Kosovo, rural villagers reported an unusual massive die-off of mice and rats in the summer of 1999 in war-devastated areas. Clusters and small outbreaks of a human disease with fever, lymphadenopathy, and ulcerations of skin and mucosa occurred, which were initially diagnosed as tonsillitis until tularemia was suspected clinically. Rumors started to circulate in some villages that wells had been deliberately contaminated with the pathogen. The Albanian authorities asked World Health Organization to send in a Global Outbreak Alert and Response Network (GOARN) team in order to help in the diagnostics and to investigate the origin and cause of this “unusual” tularemia epidemic. Since the strain was thought to be Biovar B (the endemic European strain) rather than the more virulent Biovar A, the epidemic was attributed to war-related destruction of the ecosystem and infrastructure leading to an increased population density of rodents and producing human cases of tularemia.¹⁵

There have been some documented occurrences of bioterrorism. In 1984, two large cohorts of salmonellosis cases (a total of 751 individuals) occurred in The Dalles, Oregon. The size and nature of this outbreak initiated a criminal investigation. The cause only became known when the Federal Bureau of Investigation (FBI) investigated a nearby cult (Rajneeshees) for other criminal violations. In October 1985, a vial containing a culture of *Salmonella* Typhimurium was discovered by authorities in the Rajneeshee clinic laboratory.¹⁶ As gastroenteritis cases occurred in increasing numbers, health authorities closed all salad bars in The Dalles.

In 1996 between 29 October and 1 November 1996, 12 clinical laboratory workers at the St. Paul Medical Center in Dallas, TX developed severe acute diarrheal illness as a result of eating muffins and doughnuts left in their break room on 29 October. *Shigella dysenteriae* type 2 was cultured from 8 patients that was identical to the laboratory stock strain (some of which was missing) by pulsed field gel electrophoresis and it was concluded the pastries had been deliberately contaminated.¹⁷

On 4 October 2001, a case of inhalational anthrax was reported in a 63-year-old male in Florida. Authorities initially announced this individual had probably contracted the illness

by hunting. There were two further cases in Florida, and a fourth case of cutaneous anthrax was identified in a female employee at NBC news in New York City (NYC). Investigators then realized that exposures had occurred from anthrax-containing letters sent in the mail. On 15 October, the Senate Majority Leader received an anthrax-containing letter, which led to the closure of the Hart Senate Office Building in Washington, DC.¹⁸ By the end of the year, anthrax-laden letters had caused 22 cases of anthrax (10 confirmed inhalational and 12 cutaneous, of which 7 were confirmed and 5 suspected) and 5 deaths, mostly among postal workers and mail handlers.¹⁹ A twelfth case of cutaneous anthrax occurred in March 2002 in a Texas laboratory where the anthrax samples were processed.²⁰

The mode of dispersal of a biological weapons agent may to some extent depend on whether the biological agent is being used as a form of biological warfare or as bioterrorism. In warfare it is more likely that the agent will be dispersed from an aircraft, loaded into a bomb or an explosive shell that can be directed toward enemy forces, while in bioterrorism it is more likely to be surreptitiously released into a subway tunnel or other enclosed space, or introduced into the water or food supply, or even sent through the mail. Therefore the countermeasures chosen may have to take into account widely differing environments that the agent may be in.

Countermeasures against biological weapons agents can be divided into three broad classes. The first broad class is what can be loosely described as disinfectants, or in other words, treatments that can destroy or neutralize the agent in a wide range of inorganic, organic, or living environments before the agent has had a chance to come into contact with human beings in a sufficiently large dose to cause infection of harm. The second broad class consists of treatments that can kill or neutralize the agent after it has come into contact with human beings, either before or after infection or intoxication has become established, and this class may include some drugs that can reduce symptoms without destroying the agent. The third broad class consists of strategies to vaccinate or immunize people who have been exposed to the agent, or who are at risk of exposure, in order to avoid infection or to reduce the severity of the consequences of exposure.

It is the hypothesis of the present review that light-based approaches can be effective in all three of these broad classes of countermeasures, and moreover that many of these light-based approaches can be effective against all known classes of biological weapons agents. We have divided our coverage into sections depending on which part of the electromagnetic spectrum is being employed (see Fig. 1). These wavelength ranges are: UV C (UVC, 220–280 nm); photocatalysis (UVA 320–400 + titanium dioxide); psoralens + UVA (PUVA); blue light (400–470




Figure 1. Electromagnetic spectrum and its physiological effects on various microorganisms.

nm); photodynamic inactivation (PDI, visible light 400–700-nm + photosensitizers); and near infrared short-pulsed lasers (700–1400 nm femtosecond). All of the techniques that are listed above act as disinfectants to some degree, and can kill or inactivate bacteria, fungi, viruses, and toxins in more or less challenging environments. Some of them (UVC, blue light, PDI) have been shown to be effective in inactivating pathogens without harming host tissue, after they have come into contact with a subject that would otherwise develop an infection, or who already has developed an infection. PUVA in particular has been shown to be highly effective in inactivating pathogens in such a manner to make them good vaccine preparations.

Light has several advantages over alternative disinfectants, biocides, and anti-infectives.

- Light is environmentally friendly and non-polluting.
- Light is relatively safe and non-toxic.
- Light does not cause excessive damage to the material surrounding the biological agent, whether inorganic, organic, or living.
- Light is relatively cheap to produce.
- Light acts rapidly, usually within seconds.
- Light can be applied to human skin, wounds, mucosa, and other sites of exposure without causing undue injury.
- There have been no reports of microbial cells developing resistance to light-based anti-infectives.

UV Light and Its Effects over Living Organisms

Light can be classified according to its wavelength and its interaction with matter, ionizing or non-ionizing effects. For instance, gamma rays (3×10^{-3} nm) have higher energy than radio waves (3×10^{13} nm) and as such can promote ionizing effects^{21,22} (see Fig. 1).

Due to its electromagnetic properties, the interaction of the light (at all regions of the electromagnetic spectrum) with matter leads to triggering of various phenomena. For instance, wavelengths less than 100 nm result in changes in the atomic charge (ionization) of atoms of the material interacting with the photon. However, as the wavelengths increase, the energy is not sufficient to produce ionization but can excite electrons

of this material and elevate them to higher-energy states as well as inducing conformational changes in the molecular structures^{21,22}

The UV (UV) wavelength region is set between the X-ray (≤ 100 nm) and the visual (>400 nm) bands of the electromagnetic spectrum. As such, UV light can be classified into four wavelengths according to its interaction with molecules: vacuum UV (VUV) at 100–200 nm; UV C (UVC) at 200–280 nm; UV B (UVB) at 280–315 nm; and UV A (UVA) at 315–400 nm.^{21–24} The main physiologic effects, stemming from the photonic energy, can be described as:

- VUV light: including wavelengths <200 nm; is harmful due to its capability of immediate reaction with oxygen atoms and organic molecules even at low doses.

- UVC light: wavelength range lies between 200 and 280 nm; this electromagnetic spectrum has biocidal effects and generally is reported as “germicidal” or more usually “ultraviolet germicidal irradiation” (UVGI).

- UVB light: comprises wavelengths between 280 to 315 nm; these photons are known for “sun burning” of the skin and have been implicated in photocarcinogenesis and photoaging.

- UVA light: comprises wavelengths between 315 to 400 nm; it is becoming realized that the shorter UVA wavelengths (called UVA1, 315–340 nm) can have also have detrimental effects on the skin due to production of reactive oxygen species.

Energetically UVC is very important in the context of inactivation of microorganisms, since UVC directly affects deoxyribonucleic acid (DNA) and ribonucleic acid (RNA) by inducing molecular transformation (i.e., producing photoproducts in the genetic material). The pyrimidines and purines can absorb UV light and this way DNA and RNA can be inactivated by UV light, especially UVC at 254 nm by oxidation of these bases or through base dimerization and formation of *cis-syn* cyclobutane pyrimidine dimers in the DNA molecules.^{3,21,22} When DNA is damaged it becomes very difficult for the nucleic acids to replicate, and if replication does occur, it often produces a defect that prevents the bacterium from being viable.^{23,25} There are six possible photoproduct “defects” in the DNA induced by UV light: thymine–thymine dimer; cytosine–cytosine dimer; cytosine–thymine dimer; uracil–uracil dimer; uracil–thymine dimer; and uracil–cytosine dimer.²¹

What are the factors governing the effective photonic interaction with living organisms? The Grotthus–Draper law (first law of photochemistry) states that photons must be absorbed for the photochemical reaction to occur and the Stark–Einstein law (second law of photochemistry) states that, if a photon is absorbed, then only one photon should be enough for a photoproduct formation.²⁶ On the other hand, it is well known that microbial inactivation is a dose-dependent process (Bunsen–Roscoe reciprocity law) based on the UV intensity in the irradiation area.²¹ UV light (as also applies to all wavelengths) has energy measured in joules (J), power in watts (W), area irradiated in cm^2 or m^2 , time of irradiation in seconds (s), irradiance (W/area), and fluence or dose (J/area) for calculation of dose-response. In addition, environmental condition such as humidity, temperature, and particle size also affect the dose-response and need to be

considered, although the duration of exposure required for lethal effect of UVC is short.²⁴

UVC light (200 to 280 nm) is the most used light for inactivation of microorganisms.^{3,9,25,27–35} This inactivation can use monochromatic or polychromatic light sources. Indeed, the main difference between these UVC lamps is that monochromatic lamps such as mercury lamp emitting at 254 nm cause genetic damage to microorganism, whereas polychromatic sources with other UV regions also affect aromatic proteins (i.e., can also affect function and structure of microbial proteins which depends on primary, secondary, and tertiary structures).²¹

UV light sources

The main source of UV light used to kill microorganisms has been produced by mercury vapor arc lamps for a long time, predominately at a wavelength of 253.7 nm (UVC electromagnetic spectrum)²⁴ (see Table 2). This kind of lamp is low-pressure mercury (Hg) and are 30% efficient at converting input power to UVC at 253.7 nm.³⁶ Currently, and owing to its wider application ranges, there is a need for UVC light to be emitted from lamps or devices containing non-toxic materials with better efficiency and lower costs to make them more affordable, owing to the potential risks of mercury lamps being broken and exposing its hazardous material to the environment. In this context, light-emitting diode (LED) and xenon lamps have gained prominence.³⁶

A UVC LED has been tested in a single-pass flow-through device. Unfortunately, the LED is very inefficient at producing UVC radiation (0.3%). However, arrays of LEDs can be more efficient and produce the expected inactivation.³⁶ The xenon lamp emits a peak wavelength at 240 nm. This lamp can have a total emission of 10 W of which approximately 1.4 W is UVC radiation. This lamp is a non-toxic alternative to mercury but it produces ozone, which is a strong oxidant and toxic air pollutant.³⁶ Thus, more research needs to be done in order to improve LED efficiency and/or discover others sources of UV light.³⁶

UV light as a viable decontamination technique for potential biological warfare agents

The first observation how microorganisms respond to light was in nineteenth century with experiments using sunlight and inactivation or disinfection of test tubes containing Pasteur solution. At this time it was already known that inactivation or disinfection of surfaces was dependent on intensity, duration, and wavelength of the light, starting the concept of dose-response. Especially in this context, it was observed differences of sensitivity between different bacteria.²³

Since the last century the source of light used to kill microorganisms have been the low-pressure mercury (Hg) lamps emitting primarily a short wave (254 nm) of UVC electromagnetic spectrum.²³ UVC light affects pyrimidines, purines, and flavins promoting the formation of dimers in RNA (uracil and cytosine) and DNA (thymine and cytosine), which promotes inactivation of many microorganisms. Thus, UVC is an established means of disinfection and can be used to kill agents causing many infectious diseases.^{21,23} There have been some studies to determine which wavelength in the UVC region is actually best to inactivate microorganisms. Lakretz et al.³⁷ compared UV wavelengths between 220 and 280 nm and concluded that 254 and 270 nm

Table 2. UV irradiation effect on microorganisms

Microorganism	Description	UV light	Light source	Irradiance	Dose and/or time of irradiation	Reference
<i>Francisella tularensis</i>	Petri dishes	UVC 254 nm	Mercury lamp	—	1.4 mJ/cm ²	35
<i>Brucella melitensis/abortus</i>	Petri dishes	UVC 254 nm	Mercury lamp (5 × 8W)	18.7 mW/cm ² and 19.5 mW/cm ²	120 to 240 s	34
<i>Staphylococcus aureus</i>	Infected wounds	UVC 254 nm	Mercury lamp	2.7 mW/cm ²	2.59 J/cm ² (16 min)	25
Methicillin-resistant <i>Staphylococcus aureus</i> (MRSA)	Petri dishes	UV continuous (peak at 245 and 261 nm)	Xenon flash lamp (6 W)	250 μW/cm ²	1 to 10 s	33
<i>Streptococcus pyogenes</i>	Petri dishes	UVC 265 nm	Light-emitting diode	1.93 mW/cm ²	1.93 mJ/cm ² (1 s) to 57.95 mJ/cm ² (30 s)	32
<i>Yersinia pestis</i>	Petri dishes	UVC 254 nm	Mercury lamp	—	~1.4 mJ/cm ²	35
<i>Bacillus anthracis</i>	Petri dishes	UVC 254 nm	Mercury lamp	—	~25 mJ/cm ²	35
<i>Bacillus cereus</i>	Petri dishes	UV continuous (peak at 245 and 261 nm)	Xenon flash lamp (6 W)	250 μW/cm ²	5 to 20 s	33
		UV pulsed (53 Hz) (peak at 245 and 261 nm)	Xenon flash lamp (6 W)	250 μW/cm ²	5 s	
<i>Bacillus thuringiensis</i>	Air circulation system	UVC 254 nm	Mercury lamp	1870 μW/cm ² to 3720 μW/cm ²	30 min to 48 h	31
<i>Bacillus subtilis</i>	Petri dishes	White light pulsed (250 μs) (200 to 1100 nm)	Xenon flash lamp	—	0.17 J/cm ² and 5.28 J/cm ² (1 to 2 min)	30
	Petri dishes	UVC 254 nm	Mercury lamp	~1.34 × 10 ⁻³ W/cm ²	25 to 1000 mJ/cm ² (19 s to 12 min)	
<i>Bacillus atrophaeus</i>	Air circulation system	UVC 254 nm	Mercury lamp	1870 μW/cm ² to 3720 μW/cm ²	30 min to 48 h	31
<i>Bacillus megaterium</i>	Air circulation system	UVC 254 nm	Mercury lamp	1870 μW/cm ² to 3720 μW/cm ²	30 min to 48 h	31
Variola virus	Airborne disinfection (hospital)	UVC 254 nm	Mercury lamp: 5 W		17 J/m ² to 68 J/m ² (10 min)	3
Ebola	Petri dishes	UVC 254 nm	Mercury lamp: 15 W	—	4 J/m ² (0 to 30 s)	9
Lassa	Petri dishes	UVC 254 nm	Mercury lamp: 15 W		16 J/m ² (0 to 30 s)	9
<i>Saccharomyces cerevisiae</i>	Fresh nectar from fruits	UVC 254 nm	Two mercury lamps	25 mW/cm ²	75 to 450 kJ/m ² (15 min at 0.073 and 1.02 L/min)	29
<i>Trichophyton rubrum</i>	Petri dishes	UVC 254 nm	Germicidal lamp	—	120 mJ/cm ²	245
<i>Trichophyton mentagrophytes</i>					36–864 J/cm ²	

^aInactivation of 90% (1log₁₀)

Table 2. UV irradiation effect on microorganisms (continued)

Microorganism	Description	UV light	Light source	Irradiance	Dose and/or time of irradiation	Reference
<i>Aspergillus fumigatus</i>	Water disinfection	UVC 254 nm	Mercury lamp	0.83 mW/cm ²	12.45 mJ/cm ²	28
<i>Aspergillus flavus</i>				0.83 mW/cm ²	16.6 mJ/cm ²	
<i>Aspergillus niger</i>				0.83 mW/cm ²	20.75 mJ/cm ²	
<i>Clostridium botulinum</i> toxin	Petri dishes	UVC 254 nm	Germicidal lamp	15 ergs/mm ² s or 1.5 μW/mm ²	675 to 900 ergs/mm ² or 67.5 to 90 μJ/mm ²	27

^aInactivation of 90% (1log₁₀)

were better at carrying out bacterial inactivation and biofilm disruption than 239 and 280 nm. Medium pressure mercury lamps emit a wider range of wavelengths than low pressure lamps including lines between 365 nm and 578 nm and it has been claimed that they are actually better than low pressure lamps at inactivating pathogens.³⁸ There have also been studies aimed at comparing pulsed with continuous wave (CW) UV light. Using 365-nm LEDs, Li et al. showed³⁹ that pulsing at 100 Hz was superior to CW for inactivating *E. coli* and *C. albicans* biofilms. Moreover pulsed xenon light technology (broad spectrum including both UV and visible) has also been much studied⁴⁰ for microbial inactivation.

Due to its killing effects on microorganisms, other applications of the UVC have been extended into the food processing industry, disinfecting heating-cooling coils, ventilating and air-conditioning systems, whole room/surface disinfection, and into killing of all human pathogens (bacterial, viral, and protozoan) transmitted via water.^{21,23,24,36}

Considering food-processing, UVC has shown a great potential for surface disinfection of fresh-cut fruit and vegetables, reducing deterioration, prolonging storage life, and becoming a viable alternative to chemical sanitizers as titanium dioxide (TiO₂) and chlorine.²¹ It is important highlight that UV treatment is increasingly common because the process is effective against a wide range of microorganisms, overdose is not possible, chemical residues or byproducts are avoided, and water quality is unaffected and therefore UV treatment has also been an important tool for water and wastewater treatments.²¹

Another significant use of UV light is air disinfection because a wide variety of fungal, bacterial, and viral pathogens may be transmitted by airborne droplets as e.g., *Mycobacterium tuberculosis*, influenza viruses, SARS corona virus, *Aspergillus* spp., and *Legionella* spp.²¹ UV has successfully reduced the concentration of airborne microorganisms in operating rooms during surgery. The installation of UV light in air handling units and ventilation systems reduced the concentration of airborne bacteria and fungi in indoor air as well as the total amount of bacteria collected at the edge of the surgical site was significantly reduced.²¹ These

results foreshadowed the use of UV light in 1935, specifically UVC in the ducts of ventilation systems.²³

The initial success of air disinfection by UVC in surgical rooms stimulated an expansion of UVC application in hospitals. For instance, UVC light sources were arranged such that to provide a kind of “light curtain” and prevent respiratory cross-infections in infant wards²³ and in neonatal intensive care units; UVC was used successfully for coil cleaning and promoting significantly the reduction of tracheal microbial colonization, as well as ventilator-associated pneumonia and the use of antibiotics.³⁶

UVC can be used for whole room disinfection, cleaning the air and surfaces under this light. Generally, air disinfection by UVC is accomplished through: irradiation on the upper-room air; irradiation of the entire room; or irradiation of the air that passes through enclosed air-circulation and heating, ventilation, and air-conditioning systems.²³ For faster results, high-powered lamps that generate high fluence levels can be used for whole room disinfection, but in unoccupied spaces in order to prevent erythema to the skin and photokeratitis in humans³⁶ or when people wear specific clothes for their protection.²³ Currently in the United States, UVC has been installed in air-handling units in heating, ventilating, and air conditioning systems to irradiate the surfaces of the coil and disinfect system components.³⁶

Although biosafety is a public health concern, most of the attention is cornered to hospital environments and microbiology laboratories, and bioterrorism concerns have not so far become familiar to the public.^{23,24} However, the technology and methods used in health care facilities and laboratories can also help against potential bioterrorism agents that cause anthrax, smallpox, viral hemorrhagic fevers, pneumonic plague, glanders, tularemia, drug-resistant tuberculosis, influenza pandemics, and severe acute respiratory syndrome to mention a few.²⁴

Biological UV dosimeters

It is widely accepted that biological UV dosimetry is an indicative tool for assessing the UV radiation impact on health and ecosystems. The accumulated data indicates that standard UV treatments that are effective against *B. subtilis* spores are likely also to be sufficient to inactivate *B. anthracis* spores and that

the spores of standard *B. subtilis* strains could reliably be used as a biosimetry model for the UV inactivation of *B. anthracis* spores.⁴¹ There are several studies now utilizing the concept of “biological UV dosimeters” as indicators of UV exposure where bacteria such as *E. coli* and *B. subtilis* have been used as sensing elements.⁴²

UV radiation is estimated to be one of the most important risk factors for nonmelanoma and melanoma skin cancers. In a study Moehrle et al.⁴³ assessed the annual occupational UV exposure of mountain guides that were using spore film test chambers containing spores of *B. subtilis* as UV dosimeter-agents that have a spectral sensitivity profile similar to erythema-weighted data (calculated from spectroradiometric measurements). In the study nine mountain guide instructors carried dosimeters on the sides of their heads in a total of 1406 working days throughout a year. During the study period the dosimeters were changed monthly.⁴³ In another study by the same group⁴⁴ they tested the practical application of the “biological UV dosimeters” on 11 persons in a span of 43 d, under different UV exposure conditions that were spread over 5 different geographical regions. The mixed cohort included 4 professional lifeguards of a swimming pool who carried the dosimeters attached to their shoulders or to their head-caps for 11 d; 3 mountain guides that attached the dosimeters laterally to their heads on 27 different occasions of mountaineering activity in different mountain regions; and 4 ski instructors who carried lateral head dosimeters during 8 d of skiing in the Alps. The conclusion of the study was that *B. subtilis* spore film dosimeters can effectively be used as personal “solar UV exposure detectors”.

In a different study Vähävihi et al.⁴⁵ assessed the viability of personal UV dosimeters; where UVB dose exposure during a 13-d heliotherapy for atopic dermatitis using *B. subtilis* spore film dosimeters with UV meter, and diary records were used. In addition, correlation between personal UVB dose exposure and changes in serum 25-hydroxyvitamin D (25[OH]D) was studied over a set of 21 adult cohorts in the Canary Islands. The study concluded that the increase in serum 25(OH)D correlates with the UVB exposure length, and that spore films are feasible and reliable in vivo tools to be used as personal UV dosimeters in field conditions.⁴⁵

Bacterial resistance to UV irradiation: effective internal repair mechanism

Studies have been revealing that bacterial spores possess an enormous resistance to UV radiation⁴⁶⁻⁴⁹ which is a source of concern to some degree. Even more interestingly dormant spores of the various *Bacillus* species, including *B. subtilis*, are shown to be 5 to 50 times more resistant to UV radiation than are the corresponding growing cells.⁵⁰⁻⁵² This resistance arises largely due to the use of a unique DNA repair enzyme called spore photoproduct lyase (SP lyase) which apparently repairs specific UV-induced DNA lesions through a radical-based mechanism. The interesting thing about this repair mechanism is that, unlike DNA photolyases, SP lyase belongs to the emerging superfamily of radical S-adenosyl-L-methionine (SAM) enzymes and uses a (4Fe-4S)⁺ cluster and SAM to initiate the repair reaction (where the DNA lesion recognition and binding site involves a β -hairpin

structure).⁴⁶ It has been shown that SAM and the cysteine residue are perfectly positioned at the active and as such facilitate the hydrogen atom abstraction (from the dihydrothymine residue of the lesion) and subsequently donation to the α -thymine radical moiety. Based on structural and biochemical characterizations of mutant proteins, the researchers were able to substantiate the role of this cysteine residue in the enzymatic mechanism of action. The proposed structure reveals how SP lyase combines specific features of radical SAM and DNA repair enzymes, in enabling a complex radical-based repair reaction to occur.⁴⁶ In essence, the SP lyase repairs the UV-induced thymine dimer (a spore photoproduct [SP]) in germinating endospores and, as such, it is responsible for the strong UV resistance of the endospores. SP lyase is a radical S-adenosyl-L-methionine (SAM) enzyme that is using the (4Fe-4S)⁺ cluster in reducing SAM and generating the catalytic 5'-deoxyadenosyl radical (5'-dA[•]).⁵³ A very recent publication by Young et al. is revealing that two conserved tyrosines may be also critical for the enzymes catalytic activity. The one tyrosine in *B. subtilis* SPL, Y99(Bs), is downstream of the cysteine, suggesting that SP lyase uses a novel hydrogen atom transfer (HAT) pathway and with a pair of cysteine and tyrosine residues regenerates the SAM. The second tyrosine, Y97(Bs), has a structural role and serves to facilitate the SAM binding. In fact, the researchers think that it may also contribute to the SAM regeneration process by interacting with the putative Y99(Bs) and/or 5'-dA intermediates, and thus lowering the energy barrier for the second H abstraction step.⁵³

In essence, the observed remarkable resistance of the bacterial spores to chemical and physical stresses, including exposure to UV radiation, arises as a result of a unique photochemistry of spore DNA that generates and accumulates the spore photoproduct 5-thymine-5,6-dihydrothymine and coupled with the capabilities of efficient repair of the accumulated damage by the enzyme SP lyase this unique viability effect comes to life. As such the observed elevated spore UV resistance corner stones can be listed as:

- Photochemistry of the DNA within spores: UV generates few (if any) cyclobutane dimers, but rather the spore photoproduct 5-thymine-5,6-dihydrothymine. As such, it is an exclusive DNA photodamage product in bacterial endospores and a radical S-adenosylmethionine enzyme (SAM) and the SP lyase (at the bacterial early germination phase) repairs it.
- The DNA repair effect (in particular SP lyase repair), during spore germination process: the unique UV photochemistry of spore DNA is largely due to its saturation with a group of small, acid-soluble proteins (SASP) that are unique to spores and whose binding alters the DNA conformation and as such its photochemistry. This SP-specific repair is also unique to spores and is performed by a light-independent SP-lyase, an iron-sulfur protein that utilizes S-adenosylmethionine to catalyze SP monomerization without DNA backbone cleavage.^{47,50,52}

Resistance of vegetative bacteria to UV photoinactivation can also be developed. The bacterial growth rate strongly affects the sensitivity to UVC,⁵⁴ and bacteria isolated from a high-altitude extreme environment were more resistant to UV.⁵⁵ There are UV-inducible DNA repair systems such as those found in

E. coli mutants deficient in induction of mutations by UV light.⁵⁶ Nucleotide excision repair involving the products of the *uvrA*, *uvrB*, and *uvrC* genes, and the error-prone repair in association with the *umuDC* gene products is also known to occur.⁵⁷ The latter process, the SOS response is triggered by the activated RecA* protein, which facilitates the autocleavage of the UmuD protein to yield the active UmuD9 C-terminal fragment.

Clearly once the potential of UV light to kill microorganisms like bacteria, viruses, and fungi was understood, there has been an increasing interest to improve the light utilization. We highlight below some studies which used UV light to kill various microorganisms in water, air, food, or in experimental models and demonstrate that UV light can be a viable tool against a possible bioterrorist action using these microorganisms.

Germicidal UV for Infections

Although it has been known for the past 100 years that UVC irradiation is highly germicidal, the use of UVC irradiation for prevention and treatment of localized infections is still in the very early stages of development. Our laboratory has performed several studies designed to show that UVC irradiation can be used in vivo to treat mouse models of infections caused by virulent and pathogenic microorganisms.⁵⁸ UVC treatment (2.59 J/cm²) of partial thickness skin abrasions in mice infected with *Pseudomonas aeruginosa* increased the survival rate of mice by 58.3% ($P = 0.0023$).²⁵ When the same treatment was applied to mice with abrasions infected with *S. aureus*, the wound healing rate was increased by 31.2% ($P < 0.00001$). In mice with wounds and burns infected with a virulent strain of *Acinetobacter baumannii* isolated from US soldiers in Iraq, UVC was able to reduce the bacterial burden by >90%.⁵⁹ Although DNA lesions were observed by immunofluorescence in the surrounding mouse skin immediately after a UVC exposure of 3.24 J/cm², the lesions were extensively repaired within 72 h. UVC was also successfully employed to treat a cutaneous *Candida albicans* fungal infection in mouse burns.⁶⁰

Photocatalytic Inactivation of Biological Warfare Agents: Titania Photocatalysis


The ability of titanium dioxide (TiO₂) to act as a photocatalyst has been reported since 1929⁶¹ (and references therein). In 1972, Fujishima and Honda⁶² first reported the photoelectrolysis of water at a semiconductor electrode. This property was then utilized to catalyze the oxidation of pollutants.^{63,64} Photocatalytic surfaces can be manufactured into construction and building materials⁶⁵ and some of the commercial uses include self-cleaning windows and self-cleaning glass covers for road lights⁶¹

One of the most important aspects of TiO₂ photocatalysis is that the process, just like the photoelectric effect, depends entirely on the energy of the incident photons and not (to a first approximation) on their intensity.⁶⁶ This suggests that, if there are even just few photons of required energy, they can induce photocatalysis; a phenomenon that has enormous practical implications.

There are three main polymorphs of TiO₂: anatase, rutile, and brookite; in all the three forms, titanium (Ti⁴⁺) atoms are coordinated to 6 oxygen atoms (O²⁻) and are forming the TiO₆ octahedra. Typically TiO₂ is an n-type semiconductor because of its oxygen deficiency, a fact having a leading role in the photocatalytic processes and mechanisms. The bandgap energy (energy required to promote an electron) of TiO₂ is of 3.0 eV for the rutile, 3.2 eV for anatase, and ~3.2 eV for brookite polymorphs, which means that photocatalysis can be activated by photons with a wavelength shorter than 385 nm (i.e., UVA). The adsorption of a photon with sufficient energy promotes an electron from the valence band to the conduction band leaving a positively charged hole in the valence band. The hole may be filled by migration of an electron from an adjacent molecule, leaving that molecule with a hole, and so on. And when electrons reach the surface, they can react with O₂ to produce superoxide radical anion (O₂⁻), and the photogenerated holes can react with water to produce hydroxyl radicals (*OH). On the other hand, O₂⁻ can react further to form H₂O₂ and more *OH. As such, the photocatalytic process implies photon-assisted generation of catalytically active ROS rather than an action of the light as a catalyst in the reaction (Fig. 2).

The majority of studies have shown that anatase is the most effective photocatalyst while rutile is less active. Differences are probably due to differences in the extent of recombination of e⁻ and hole between the two forms.⁶⁷ However, studies have shown that mixtures of anatase and rutile were more effective photocatalysts than 100% anatase and were more efficient for inactivating viruses.⁶⁷

The mechanistic description of the TiO₂ photocatalysis process can be detailed as follows, where e⁻_{CB} is the electron generated at the conduction band, h⁺_{VB} is the hole generated (and left) at the valence band. A recent paper⁶⁸ suggests that the mechanism could be better characterized as “proton-coupled electron transfer”:



One can say that TiO₂ is a chemically stable and inert material, and can continuously exert antimicrobial effects when illuminated. The energy source could be even the solar light; therefore, TiO₂ photocatalysts are also useful in remote areas where electricity is insufficient. However, because of its large band gap for excitation, only biohazardous UV (UV) wavelengths can excite TiO₂, which limits its application in living environments. To circumvent this problem, impurity doping through metal coating and controlled calcination has been successfully used to modify the TiO₂ and to expand its absorption wavelengths to the visible light region (discussed below).

Matsunaga and colleagues^{69,70} were the first to use TiO₂ photocatalysis to kill microorganisms. This subject area has recently been comprehensively reviewed^{71,72} and the effect of key variables on the effectiveness has been studied.⁷³

Previous studies have investigated the antibacterial abilities of visible light-responsive photocatalysts using the model bacteria *Escherichia coli* and human pathogens. They have shown that modified TiO₂ photocatalysts significantly reduced the numbers of surviving bacterial cells in response to visible light illumination.

Bacterial inactivation studies have confirmed that even with significantly lower levels of TiO₂ generated radical scavengers, i.e., ROS, illumination with far-UV light can successfully promote microorganisms inactivation.⁷⁴ Spore-forming bacteria of *Bacillus* strains were investigated for demonstrating photocatalytic disinfection effects with relatively good results.⁷⁵ Armon et al. studied the photocatalytic inactivation of spores of *B. subtilis* and *B. cereus* (as a model for the main biological warfare element *B. anthracis*⁷⁶) where the spore-forming *B. cereus* is genetically very closely related to *B. anthracis* whereas *B. subtilis* is highly resistant to variety of stress factors.⁷⁷

It has been suggested that the photocatalytic killing mechanism initially damages the weak points at the bacterial cell surface before total breakage of the cell membranes. The internal bacterial components then leak from the cells through the damaged sites. And finally the photocatalytic reaction oxidizes all of the cell debris. In essence, the killing mechanism with TiO₂ involves degradation of the cell wall and cytoplasmic membrane due to the production of ROS such as hydroxyl radicals and hydrogen peroxide. This initially leads to leakage of cellular contents then cell lysis and may be followed by complete mineralization of the organism. Killing is most efficient when there is close contact between the organisms and the TiO₂ catalyst⁷¹ (Fig. 3).

Huang et al.⁷⁸ demonstrated with *E. coli* that TiO₂-treated cells continue to lose their viability even after the UV-irradiation stops, indicating that reactions in the media continue to propagate even after the UV-irradiation stops. Once the lethal oxidation reactions are initiated by the TiO₂ photocatalytic reaction, the damaging effects propagate in the dark via the Fenton reaction or free radical chain reactions of lipid peroxidation due to ROS.⁷⁹ The results suggest that initial oxidative damage happens on the cell wall (where the TiO₂ photocatalytic surface makes the first contact). The cells that sustained the initial oxidative damage insult on their cell walls are still viable, however, though localized, elimination of the cell-wall protection makes these cells susceptible to ensuing oxidative damages to the underlying cytoplasmic membrane. Overall, the photocatalytic action progressively increases the cell permeability ending in free efflux of the intracellular contents, thus, eventually leading to cell death. Also, it is plausible that TiO₂ can gain access into the membrane-damaged cells and generates a direct insult on the intracellular components, thus, accelerating the cell death.^{80,81}

In summary, visible light-responsive TiO₂ photocatalysts are more convenient than the traditional UV light-responsive TiO₂ photocatalysts because they do not require harmful UV light irradiation to function. These photocatalysts, thus, provide a




Figure 2. Photocatalytic effect of the TiO₂: a process where photon-assisted generation of catalytically active ROS is generated rather than an action of the light as a catalyst in the reaction.

promising and feasible approach for disinfection of pathogenic bacteria, facilitating the prevention of infectious diseases.⁸²

By contrast, recombination of the photogenerated charge carriers is a major limitation in the use of TiO₂ as a photocatalyst and an initiator in the photocatalytic process, and, as such, is an important agent in combating biowarfare. Since the excited e⁻ in the recombination process relaxes back to the valence band (either non-radiatively or radiatively, dissipating its energy as light or heat) without reacting with the possible biological sites (and thus not initiating the photocatalytic process—a bulk recombination process), there are several strategies developed to prevent this from happening and to improve the photocatalytic efficiency. To enhance the charge separation of the e⁻ and holes and to reduce the likelihood of bulk recombination, termed photoelectrocatalysis, it is possible to apply an electric field.^{83,84} Other approaches used to achieve improved efficiency include either chemical modifications (by incorporating additional components in the TiO₂ structure, termed as doping) or increasing the surface area and porosity of the photocatalyst.⁸⁵⁻⁸⁹

In some cases carbon has been used as a dopant and as such allowing not only visible light absorption but also “injecting” active trap sites within the TiO₂ bands, thus increasing the lifetime of the photogenerated charge carriers.⁸⁵

TiO₂ can be used in combination with some of the noble metals, such as Ag, Au, and Pt, which enhance the photocatalytic efficiency under visible light due to “injecting” traps for the electrons and promoting the interfacial charge transfer, and thus delaying the recombination process of the electron-hole pair.⁹⁰⁻⁹⁴

Data accumulated thus far shows that TiO₂ exhibits a strong visible-light induced anti-microbial activity when modified by doping or used in combination. Sulfur-doped TiO₂ is shown to have strong antibacterial effect.⁹⁵ Carbon-doped TiO₂ and TiO₂ modified with platinum (IV) chloride complexes used as suspension or immobilized at surfaces (infected with the microorganisms) show remarkable anti-bactericidal effects. The detrimental effect of the photocatalysts induced with visible light on various microorganism groups such as bacteria (i.e., *E. coli*, *S. aureus*, *Enterococcus faecalis*) or fungi (i.e., *Aspergillus niger*, *C. albicans*) and utilizing modified TiO₂ showed increased effect over these




Figure 3. Photocatalytic killing mechanism initially damages the weak points at the bacterial cells surfaces, and then total breakage of the cell membranes ensues, followed by of the internal bacterial components through the damaged sites. Finally, the photocatalytic reaction oxidizes all of the cell debris.

microorganisms in the order: *A. niger*, *C. albicans* > *E. faecalis*, *S. aureus* > *E. coli*.⁹⁶

TiO₂ photocatalysis with UV (UVA) light has proven to be a highly effective process for complete inactivation of airborne microbes. However, the overall efficiency of the technology needs to be improved to make it more attractive as a defense against bio-terrorism. Studies investigating the enhancement in the rate of destruction of bacterial spores on metal (aluminum) and fabric (polyester) substrates with metal (silver)-doped titanium dioxide (in comparison with conventional photocatalysis [TiO₂ P25/+UVA] and UVA photolysis), where *B. cereus* bacterial spores were used as an index to demonstrate the enhanced disinfection efficiency, showed complete inactivation of *B. cereus* spores with the enhanced photocatalyst effectiveness. The enhanced spore destruction rate may be attributed to the highly oxidizing radicals generated by the doped TiO₂.⁹⁷

According to Wong et al., anion-doped TiO₂ photocatalytic effect is with higher quantum efficiency under sunlight and as such showed inactivating effect on both spores and toxins of *B. anthracis* under irradiation by “ordinary” light source such as an incandescent lamp. Moreover, these carbon-doped and nitrogen-doped TiO₂ had a better performance in the presence of silver; the synergistic antibacterial effect resulted in approximately 5 logs reduction of *E. coli*, *S. pyogenes*, *S. aureus*, and *A. baumannii*. It appears, the presence of Ag enhances the bactericidal properties of various TiO₂ materials.⁹⁸ They also found that visible light illuminated nitrogen- or carbon-doped TiO₂ significantly reduces the viability of anthrax spores. Even

though the spore-killing efficiency is only approximately 25%, their data indicate that spores from photocatalyzed groups (not from untreated groups) have lower survival rate. In addition, their results indicated that the photocatalysis could directly inactivate a lethal toxin, the major virulence factor of *B. anthracis*. The study results show that the photocatalyzed spores have 10-fold less potency to induce mortality in mice in comparison with unexposed one. These results suggest that photocatalysis might be effective in injuring the spores through inactivating some spore components. In essence, photocatalysis may be a viable technique in inducing injuries to the spores than direct killing in order to reduce their pathogenicity in the host.^{99,100}

It has been shown that nano-sized titania particles exhibit better inactivation properties than the bulk-sized titania materials. Sunlight in the presence of nano-titania (mixture of anatase and rutile phases) displayed

better photocatalytic bactericidal activity of *B. anthracis* than sole treatment of sunlight.¹⁰¹

Studies on photocatalytic inactivation of spores of *B. anthracis* have been performed using nano-sized titania materials and UVA light or sunlight. Results demonstrated pseudo first order behavior of spore inactivation kinetics. The value of kinetic rate constant increased from 0.4 h⁻¹ to 1.4 h⁻¹ indicating photocatalysis facilitated by addition of nano-sized titania. Nano-sized titania exhibited superior inactivation kinetics on par with large sized titania. The value of kinetic rate constant increased from 0.02 h⁻¹ to 0.26 h⁻¹ on reduction of size from 1000 nm to 16 nm depicting the enhanced rate of inactivation of *B. anthracis* Sterne spores on the decrease of particle size.¹⁰²

These results signify that the excited TiO₂ nanoparticles potentiate the antimicrobial action of β lactams, cephalosporins, aminoglycosides, glycopeptides, macrolides, and lincosamides, making a possible synergistic combination of nano compound with antibiotics against MRSA.¹⁰³

Interestingly, Cheng et al. found that a mixture of anatase/rutile carbon doped TiO₂ nanoparticles show significantly enhanced bactericidal effect. Their experiments indicated that these nanoparticles (with higher bacterial interaction property), have significantly higher proportion of bacteria-killing effect over all tested pathogens (including *S. aureus*, *Shigella flexneri*, and *A. baumannii*). These findings suggest that developing materials with high bacterial interaction ability might be a useful strategy to improve the antimicrobial activity of visible-light-activated TiO₂.¹⁰⁴

In recent decades, incidences with antibiotic-resistant bacteria have shown sharp elevations, and as such, became one of the most significant problems in public health. TiO₂ has the potential to inactivate antibiotic-resistant bacteria. In the Tsai et al. study, UVA-activated TiO₂ was successfully used to inactivate the antibiotic-resistant bacteria MRSA, multidrug-resistant *A. baumannii* (MDRAB), and vancomycin-resistant *E. faecalis* (VRE) in suspension. Their results indicated that TiO₂ reaction time had the greatest influence on microbial survival, following the TiO₂ exposure in the presence of UVA. TiO₂ in the presence of UVA effectively reduced the number of antibiotic-resistant microbes in suspension by 1–3 logs.¹⁰⁵

Photo-activated TiO₂ is effective on microorganisms capable of killing a wide range of gram-negative and gram-positive bacteria, fungi (both unicellular and filamentous), protozoa, algae, mammalian viruses, and bacteriophages; the killing activity is enhanced by the presence of other antimicrobial agents, such as Cu and Ag.⁷¹

The level of UVA disinfection of *B. anthracis* and *B. brevis* vegetative cells increased with the presence of the TiO₂ and Ag photocatalysts, but had little effect on their spores. *Bacillus brevis* spores were slightly more sensitive to UVB and UVC than the spores of *Bacillus atrophaeus*. Photocatalytic sterilization against spores was strongest in UVC and UVB and weakest in UVA. The rate of inactivation of *Bacillus* spores was significantly increased by the presence of TiO₂ but was not markedly different from that induced by the presence of Ag. Therefore, TiO₂/Ag plus UVA can be used for the sterilization of vegetative cells, while TiO₂ and UVC are effective against spores.¹⁰⁶ However, in a study investigating the effects of toxin- and capsule-encoding plasmids on the kinetics of UV inactivation of various strains of *B. anthracis* it was found that the plasmids pXO1 and pXO2 had no effect on bacterial UV sensitivity or photoreactivation. Interestingly enough, vegetative cells were capable of photoreactivation whereas photo-induced repair of UV damage was absent in *B. anthracis* Sterne spores¹⁰⁷ which shows that *B. anthracis* makes highly stable and heat-resistant spores that can remain viable for decades.¹⁰⁸

Psoralens and UVA

Psoralens are a group of natural furanocoumarins, commercially derived from a plant found in Egypt, *Ammi majus*. They are also present in celery, carrots, parsley, parsnip, and other vegetables. It has been known since ancient times that consumption of these foodstuffs followed by sun exposure can lead to a phototoxic skin reaction similar to sunburn. The combination of psoralen with UVA light (known as PUVA) was first introduced as a medical treatment for psoriasis.¹⁰⁹ Patients orally ingested psoralen compounds or alternatively the psoralens were applied topically in a bath.¹¹⁰ The mechanism is that the psoralen molecule has the correct structure and shape to be able to intercalate between the two strands of DNA in the double helix, and upon illumination, induce the formation of covalent inter-strand cross-linking between opposite nucleic acid strands (Fig. 4).

Due to the DNA damaging action PUVA it has been used for the inactivation of bacteria, viruses, and protozoa in platelet and plasma blood component.¹¹¹ This photochemical inactivation using PUVA has the potential even to produce a new class of vaccines from whole microbes termed “Killed But Metabolically Active” (KBMA). KBMA vaccines are based on whole microbes that have been inactivated by defined genotoxic methods that leave the organism incapable of productive growth and of causing disease but preserve metabolic activity sufficient to induce immunity. These vaccines have two broad applications. First, recombinant KBMA vaccines encoding selected antigens relevant to infectious disease can be used to elicit a desired immune response. And when derived from attenuated forms of a targeted pathogen the entire antigenic repertoire is presented to the immune system, as here correlate of protection are unknown. In both applications the vaccine is inactivated by a distinct and limited disruption of the vaccine chromosome using photochemical treatment with a psoralen cross-linking agent, impacting an absolute block to DNA replication and possible vaccine outgrowth.¹¹² Initially this technology was developed for killing undetected microbes contaminating plasma and platelet blood products.^{113,114}

Brockstedt et al. performed a landmark study in KBMA vaccine approach demonstrating proof of concept for recombinant KBMA Lm vaccines in animal models of infectious disease and cancer.¹¹⁵ KBMA were developed by removing the genes required for nucleotide excision repair (uvrAB) and rendering microbial-based vaccines sensitive to photochemical inactivation with PUVA. Colony formation of these mutants was blocked by infrequent, randomly distributed psoralen crosslinks, though the bacterial population was able to express its genes, synthesize, and secrete proteins. Using the intracellular pathogen *Listeria monocytogenes* as a model platform, recombinant psoralen-inactivated Lm Δ uvrAB vaccines induced potent CD4⁺ and CD8⁺ T-cell responses and protected mice against virus challenge in an infectious disease model and provided therapeutic benefit in a mouse cancer model. Microbial KBMA vaccines used either as a recombinant vaccine platform or as a modified form of the pathogen itself may have broad application for the treatment of infectious disease and cancer. This was a new vaccine paradigm for eliciting effector T-cell responses and protective immunity.

In one study¹¹⁶ KBMA *B. anthracis* vaccines induced a broad and protective immunity against anthrax. In this approach a novel whole-bacterial-cell anthrax vaccine utilizing *B. anthracis* that was KBMA. Vaccine strains that are asporogenic and nucleotide excision repair deficient were engineered, rendering *B. anthracis* extremely sensitive to photochemical inactivation with amatosalen (S-59) psoralen (Fig. 5A) and UVA light. The workers also introduced point mutations, which allowed inactive but immunogenic toxins to be produced. These photochemically inactivated vaccine strains maintained a high degree of metabolic activity and secreted protective antigen, lethal factor, and edema factor. KBMA *B. anthracis* vaccines were found to be avirulent in mice and induced less injection site inflammation than recombinant protective antigen adsorbed to aluminum hydroxide gel. In animals KBMA *B. anthracis* vaccination produced antibodies against numerous anthrax antigens, including high levels of




Figure 4. Intercalation of the psoralen molecules between the strands of the double-stranded DNA helix or RNA where upon illumination with UVC light affects pyrimidines, purines, and flavins, thus promoting the formation of dimers in RNA (uracil and cytosine) and DNA (thymine and cytosine), a process which promotes inactivation of many microorganisms.

anti-protective antigen and toxin-neutralizing antibodies and fully protected mice against challenge with lethal doses of toxigenic unencapsulated Sterne 7702 spores and rabbits against challenge with lethal pneumonic doses of fully virulent Ames strain spores. Guinea pigs vaccinated with KBMA *B. anthracis* were partially protected against lethal Ames spore challenge, which was comparable to vaccination with the licensed vaccine anthrax vaccine adsorbed. Their data demonstrated that KBMA anthrax vaccines are well tolerated and elicit potent protective immune responses. The use of KBMA vaccines may be broadly applicable to bacterial pathogens, especially those for which the correlates of protective immunity are unknown.¹¹⁶ Toward the development of a KBMA *B. anthracis* vaccine candidate strain, in a different study a plasmid pMAD and a recombinase system Cre-loxP were used to knockout the *uvrAB* gene of *B. anthracis*

AP422, which lacks both of two plasmids pXO1 and pXO2. The results showed that the constructed *B. anthracis* AP422 Δ *uvrAB* was inactivated by photochemical treatment (including an exposure in a long-wavelength UVA light and a treatment of 8-Methoxypsoralen [8-MOP]). It was found that the killed *B. anthracis* AP422 Δ *uvrAB* maintained a highly metabolic activity for at least 4 h, showing a state of KBMA. Thus, the KBMA strain of *B. anthracis* AP422 Δ *uvrAB* provided the prospective vaccine candidate strain for anthrax.

Bruhn et al. demonstrated proof-of-concept for a KBMA vaccine based on a protozoan pathogen. This approach could be a new method for whole-cell vaccination against other complex intracellular pathogens. There are currently no effective vaccines for visceral leishmaniasis, the second most deadly parasitic infection in the world. This was a novel whole-cell vaccine approach using *Leishmania infantum chagasi* promastigotes treated with the psoralen compound amotosalen (S-59) and low doses of UVA radiation. This treatment generated permanent, covalent DNA cross-links within parasites and results in *Leishmania* KBMA. In this report, they characterized the in vitro growth characteristics of both KBMA *L. major* and KBMA *L. infantum chagasi*. Concentrations of S-59 that generated optimally attenuated parasites were identified. Like live *L. infantum chagasi*, KBMA *L. infantum chagasi* parasites were able to initially enter liver cells in vivo after intravenous infection. However, whereas live *L. infantum chagasi* infection leads to hepatosplenomegaly in mice after 6 mo, KBMA *L. infantum chagasi* parasites were undetectable in the organs of mice at this time point. In vitro, KBMA *L. infantum chagasi* retained the ability to enter macrophages and induce nitric oxide production. These characteristics of KBMA *L. infantum chagasi* correlated with the ability to prophylactically protect mice via subcutaneous vaccination at levels similar to vaccination with live, virulent organisms. Splenocytes from mice vaccinated with either live *L. infantum chagasi* or KBMA *L. infantum chagasi* displayed similar cytokine patterns in vitro. These results suggested that KBMA technology is a potentially safe and effective novel vaccine strategy against the intracellular protozoan *L. infantum chagasi*.¹¹⁷

Thus several groups have developed recombinant and pathogen-derived KBMA vaccine from whole microbes which have been shown to be harmless, immunogenic, and correlated with disease-specific prevention or reduction in preclinical animal models of infectious disease which gives a new hope in this direction.

Besides this PUVA has also been used for inactivation of diverse other viruses such as dengue virus, Chikungunya virus, etc. One of the groups used limes and synthetic psoralens to enhance solar disinfection of water. They performed a laboratory evaluation with norovirus, *E. coli*, and MS2. They concluded that psoralens and acidic lime extract both interact synergistically with UV radiation to accelerate inactivation of microbes.¹¹⁸ Most of the virus inactivation using psoralens has been done using platelets. In one of the studies transfusion of platelets was done during a Chikungunya virus epidemic in Ile de La Réunion that had been prepared with photochemical pathogen inactivation treatment. It was found that INTERCEPT-CPAs were well




Figure 5. List of some of the PS compounds discussed in the manuscript.

tolerated in a broad range of patients, including infants. The incidence of acute transfusion reactions (ATR) was low and when present ATRs were of mild severity.¹¹⁹

B19 is a clinically significant virus that can be transmitted through blood transfusion was also inactivated by photochemical treatment. It was stated that under defined conditions, photochemical treatment with amotosalen combined with UVA light could be used to inactivate B19.¹²⁰

Amotosalen (S-59) photochemical inactivation of severe acute respiratory syndrome coronavirus in human platelet concentrates was reported.¹²¹ Following photochemical treatment, SARS-CoV was consistently inactivated to the limit of detection in seven independent APC units. No infectious virus was detected after treatment when up to one-third of the APC unit was assayed, demonstrating a mean \log_{10} -reduction of >6.2 . Potent inactivation of SARS-CoV therefore extends the capability of the INTERCEPT Blood System in inactivating a broad spectrum of human pathogens including recently emerging respiratory viruses.

A transfusion trial was performed using platelets photochemically treated for pathogen inactivation using the synthetic psoralen amotosalen HCl.¹²² Patients with thrombocytopenia were randomly assigned to receive either photochemically treated or conventional (control) platelets for up to 28 d. Transfusion reactions were fewer following photochemically treated platelets (3.0% photochemically treated vs. 4.4% control, $P = 0.02$). The incidence of grade 2 bleeding was equivalent for photochemically treated and conventional platelets, although post-transfusion platelet count increments and days to next transfusion were decreased for photochemically treated compared with conventional platelets.

In one of the reports immunogenicity and protective efficacy of a psoralen was reported in which dengue-1 virus was inactivated which proved to be a vaccine candidate in *Aotus nancymae* monkeys. In this experiment the protective efficacy was tested of a psoralen-inactivated dengue vaccine candidate in non-human primates. Psoralen-inactivated DENV-1 was reported to be immunogenic in *Aotus nancymae* with a reduction in days of viremia following experimental challenge.¹²³ Evaluation has also been studied in a novel psoralen-inactivated dengue virus type 1 (DENV-1) vaccine candidate in *Mus musculus* mice which led to the conclusion that psoralen-inactivated DENV-1 is immunogenic in mice.¹²⁴ Poliovirus replication in HeLa cells was reported to be significantly inhibited in infected cells with 4,5,8-trimethylpsoralen plus long wavelength UV light. When infected cells were exposed to psoralen plus light during peak viral RNA synthesis, formation of virus-specific RNAs was inhibited. Viral RNA species that were either formed in vivo in the presence of or treated in vitro with psoralen plus light appeared to have become degraded. Treatment with psoralen plus light in vitro resulted in the loss of infectivity of single-stranded viral RNA.¹²⁵

It is known that excessive use of PUVA can cause skin cancer.¹²⁶ There has been concern expressed that psoralens themselves may be toxic and/or carcinogenic, but it should be emphasized that the use of PUVA to generate vaccines outside the body will not pose this risk of carcinogenicity. Indeed, the therapy known

as extracorporeal photophoresis (treating blood outside the body with psoralens and UVA) is widely used for graft-vs-host disease and other indications.¹²⁷

Blue Light Inactivation of Pathogens

The bacterial agents of bioweapons are often chosen from the bacteria that show antibiotic resistance or that form endospores and biofilms in order to be more resistant against available antibacterial treatment options. It is known that some bacteria can be converted into spore forms that may create deadly diseases in humans. Early symptoms of anthrax, for instance, can last 1 to 6 days and resemble the flu, but once the bugs multiply to large enough numbers, the body goes into shock and death can occur in 24 to 36 h. For these reasons successful phototherapy studies against virulent bacteria, fungi, and viruses are needed to defeat biological warfare.

UV light killing of bacteria is well understood, but this light-mediated antimicrobial effect may not be unique, since current studies indicate that blue light produces a somewhat similar effect. Even when compared with UV irradiation, blue light has been accepted to be much less detrimental to mammalian cells.^{128,129} Although effects of blue light seem to vary depending on wavelength, dose, and the nature of the bacteria, these wavelengths appear to exhibit a broad-spectrum antimicrobial effect against both gram-positive and gram-negative bacteria and have been suggested as an alternative treatment modality for treating some methicillin and penicillin resistance bacterial infections.¹³⁰

As an example, the 405- and 470-nm blue light showed dose-dependent bactericidal effects on *P. aeruginosa* and *S. aureus* in vitro. The results of this study indicated that the fluence of 5–15 J/cm² was the optimal dose of blue light for treatment of *P. aeruginosa* while for *S. aureus* a 470-nm light was used in a stronger dose (10–15 J/cm²).^{36,120,121} High-intensity 405-nm light may have application in the medical, military and agricultural fields to combat *B. anthracis* spore exposure which is known to have endospores of comparable robustness to *B. cereus* and *B. subtilis*.^{41,131,132}

The underlying proposed mechanism of action is that light may be absorbed by porphyrins produced by bacteria that result in increased free radicals, which may affect cytoplasmic membrane proteins and DNA,¹³³ or have a direct effect on photolabile pigments in bacteria.¹³⁴

Further studies support this opinion, indicating existence of a therapeutic window of blue light for bacterial infections where bacteria are selectively inactivated while host tissue cells are preserved.¹³⁵ Promising outcomes have been achieved when clinical trials have been conducted to investigate the use of blue light for *Helicobacter pylori*.^{136,137} Although the majority of the publications on the antimicrobial effect of blue light have been confined to in vitro studies,^{138–141} investigation by Dai et al. demonstrate potential effects of blue light shown effective in acute, potentially lethal *P. aeruginosa* burn infections in mice.¹³⁵

As mentioned above, blue light has recently attracted much attention in comparison to photodynamic therapy as an alternative antimicrobial approach¹⁴² due to its intrinsic antimicrobial

properties without the involvement of added exogenous photosensitizers.¹³⁰ As a result, the use of blue light inactivation is technically easier to carry out since the delivery of photosensitizers to the target microbes embedded deep within biofilms adherent to tissue has been somewhat challenging.

Bacterial spores are capable of extreme resistance to physical insults like heat, ionizing, UV and gamma radiation, osmotic pressure, and desiccation. The spores also protect the bacteria from chemical and biological disinfectants such as iodine, peroxides, and alkylating agents.¹⁴³ High-intensity, nonionizing blue light with wavelength of 405 nm and fluence of 1.73 kJ/cm² is capable of inactivate *B. cereus*, *Bacillus megaterium*, *B. subtilis*, and *Clostridium difficile* endospores of 4 log₁₀ colony-forming units.¹⁴⁴

The sporicidal effect of blue light seem to be an oxygen-dependent process since the efficacy of 405-nm blue light therapy explained by the presence of endogenous photoexcitation of intracellular chromophores such as coproporphyrin with Soret bands in the 400–420 nm regions of the visible spectrum and the subsequent generation of cytotoxic ROS such as singlet oxygen in *Bacillus* and *Clostridium* bacteria. Blue light can not only regulate bacterial motility, suppress biofilm formation, and potentiate light inactivation of bacteria, but it may also upregulate bacterial virulence factors.¹⁴⁵

In spite of the well understood inactivation of pathogenic microbial species used in bioweapons with UV light, visible light has a clear advantage due to well-recognized risk of UV in skin damage and cancer. To what extent UV light can be replaced with visible light in pilot studies and clinical application still remains questionable, but development of narrow-spectrum illumination of blue light could be lead to some application like air, contact surface,^{146,147} and medical instrument disinfection while in the presence of staff and patients which is much more important for disinfection of bacterial agents in bioweapons.

In comparison with UV, there is less concern about mutagenesis effects of the blue light over mammalian cells since the blue light absorption by DNA is weak. Although tissue penetration of the blue light is more efficient than UV, several studies have been conducted to further increase its penetration depth and make it compatible with the less common use of red light in antimicrobial PDT for eradication of Gram positive bacteria in vivo. Since the microbial cells shows some resistance to UV, one question that must be addressed is “Can microbial cells develop resistance to blue light inactivation?” To answer this question, the resistance of blue light in microbial cells must be considered.¹³⁰

Blue light inactivation with some known wavelengths (405, 415, or 470 nm) revealed antimicrobial effects activity as UV in photochemistry studies. For instance blue light with the wavelength of 405 nm showed strong bacterial killing against gram-positive and gram-negative bacteria in vitro.¹⁴⁸ As a result of this, investigation by Enwemeka and colleagues¹⁴⁹ has indicated that the consecutive delivery of a low light dose was more effective than a single high dose. This observation was suggested to be verified by in vivo studies.

In another study, inactivation of gram-positive bacteria like MRSA with blue light in 405 nm was found to be due to

photo-stimulation of porphyrin molecule in an oxygen-dependent process.¹⁴¹ Porphyrins are different in various bacteria; accordingly, slightly different wavelengths may be required to be absorbed by various porphyrins. These is no exogenous delivered photosensitizer involved in inactivation of bacterial using blue light which makes it easier to achieve. The wavelength of blue light use in infection treatment should be the wavelength that selectively absorbed by the chromophore located inside the pathogenic microbial cells. This idea further was supported since no activity revealed for inactivation of MRSA with blue light at 430 nm. Thus use of narrowband filters will provide more activity. Although some inactivation was observed at 420 nm, the best activity was found at 405 nm. In this wavelength, a blue light with absolute dose 23.5 J/cm² caused 2.4 log₁₀ reductions of methicillin-resistance *S. aureus*.¹⁵⁰

Enwemeka et al. in another study¹³⁸ worked with two different strains of *S. aureus*: MRSA US-300 (strain of CA-MRSA) and IS853 (strain of HA-MRSA) in vitro with different wavelength of blue light. The results showed that various wavelength produced a statistically significant dose-dependent reduction in both strains. However, maximum eradication of the CA-MRSA was achieved in 405 nm and HA-MRSA in 470 nm of blue light with 10 min irradiation. The eradication levels increased with increasing the light dose, albeit not linearly. The conclusion of the study was that phototherapy with low dose blue light may be an effective clinical tool for MRSA infections.

Blue light studies with the wavelength of 415 ± 10 nm in a mouse skin abrasion model infected with hospital-acquired MRSA was highly successful¹³⁵ and results in terms of log-reduction was more effective than that using bacterial suspensions in vitro. As found in this study, the required light fluence was 10- to 100-fold less than the light dose exposure needed for the equivalent bacterial inactivation in vitro.¹⁵¹⁻¹⁵³ One possible mechanism for this surprising finding would be that the metabolism of bacterial cells in vivo favored blue-light inactivation compared with broth cultured cells. Possibly in vivo growth promoted the biosynthesis of intracellular porphyrins, thus making the microbial cells in the tissue more sensitive to blue light than the identical cells growing in liquid growth medium.

The amount of light energy needed to kill biofilm¹⁵⁴ and endospores¹⁴⁴ is 10-fold higher than that needed to kill vegetative *B. cereus* and *C. difficile* cells; therefore, a blue light source with higher intensity is one important aspect of phototherapy. The efficacy of blue light is dependent on the wavelength, the irradiance, the duration of exposure, and the exposed body surface area. The phototherapy devices should not produce a lot of heat and should have a stable broad wavelength light output. Therefore, LED with greater efficacy and higher irradiance can be an ideal light source for the phototherapy.

A high intensity prototype blue gallium nitride LED phototherapy unit has been developed and its efficacy compared with commercially used phototherapy device by measuring both in vitro and in vivo bilirubin photodegradation.¹⁵⁵ In this study microhematocrit tubes (44 ± 7% vs. 35 ± 2%) were used for in vitro experiments and for in vivo experiments Gunn rats (30 ± 9% vs. 16 ± 8%) were applied. The LED device with two focused

arrays, each with 500 blue LEDs, showed a significantly higher efficacy of bilirubin photodegradation than the conventional phototherapy in both in vitro and in vivo experiment.

Photodynamic Inactivation (PDI) of Biological Warfare Agents

Photodynamic therapy (PDT) is a non-invasive procedure that uses a non-toxic photosensitizer (PS) and harmless visible or near-infrared (NIR) light to generate singlet oxygen and other reactive oxygen species (ROS) that react with biomolecules such as nucleic acids, proteins, and unsaturated lipids. In applications of PDT aimed at, for instance, curing cancer, the ROS cause damage to these crucial biomolecules within the tumor cells and initiate apoptosis leading to cell death. However, these previously referred to biomolecular targets of PDT (proteins, lipids, nucleic acids) are also major constituents of all the classes of biowarfare agents listed above. Hence PDT can destroy all known biowarfare agents.

This desirable property of destroying all classes of pathogen is not totally unique to PDT; certain other strong oxidizing agents such as boiling peracetic acid, chlorine dioxide, and cross-linking agents such as glutaraldehyde will also accomplish this feat. It is known that UV radiation and ionizing radiation will destroy bacteria, fungi, spores, and viruses, but not toxins. However, we believe that PDT has the potential to be the most versatile and certainly the most biocompatible strategy to combat biowarfare agents no matter if they are bacteria, viruses, fungi, spores, or even toxins.

PDT using the appropriate choice of photosensitizer and light could be used to destroy pathogens in water, on surfaces such as vehicles and equipment, in food, on skin, in wounds, and even when the agents have established localized infections in humans before systemic invasion has occurred. One important consideration in using PDT to decontaminate large surfaces (housing or vehicles) is that the PS can be efficiently activated by sunlight and after destroying all the microorganisms the residual PS will be harmlessly photobleached, and therefore would be considered environmentally friendly compared with alternative disinfectants. An additional advantage of PDT is its high level of selectivity, achieved through PS that selectively target specific cells or tissue types and the ability to control the illumination area.

Since mid-1990s, antimicrobial photodynamic-inactivation (PDI) and therapy has been developed as a prolific discovery and development platform, exploring many aspects of the microbial phenotype related to multidrug resistance such as efflux systems, biofilms, bacterial spores, and virulence determinants.¹⁵²

Bacteria

In the 1990s, it was observed that there was a fundamental difference in susceptibility to PDT between gram-positive and gram-negative bacteria. It was found that, in general, neutral or anionic PS molecules are efficiently bound to and photodynamically inactivate gram-positive bacteria, whereas they are bound, to a greater or lesser extent, only to the outer membrane of gram-negative bacterial cells but do not inactivate them after illumination.¹⁵⁶ In order to inactivate gram-negative bacteria it is

necessary to use PS with pronounced cationic charge or to take other measures to permeabilize the gram-negative cell wall.¹⁵⁷ The high susceptibility of gram-positive species is explained by their physiology, as their cytoplasmic membrane is surrounded by a relatively porous layer of peptidoglycan and lipoteichoic acid that allows PS to cross.^{156,158} The cell envelope of gram-negative bacteria consists of an inner cytoplasmic membrane and an outer membrane that are separated by the peptidoglycan-containing periplasm. The outer membrane forms a physical and functional barrier between the cell and its environment. In the outer membrane, several different proteins are present. Some of them function as pores to allow passage of nutrients, whereas others have an enzymatic function or are involved in maintaining the structural integrity of the outer membrane and the shape of the bacteria.¹⁵⁹

MRSA infections kill 19 000 hospitalized American patients annually; equivalent to the combined number of deaths due to AIDS, tuberculosis, and viral hepatitis.¹⁶⁰ In general, MRSA virulence factors are similar to those of *S. aureus*, with certain MRSA strains appearing to contain particular genetic backgrounds or factors that enhance their virulence and enabling particular clinical syndromes with net effect of creating havoc in the affected communities¹⁶¹⁻¹⁶⁴ (Fig. 6). There have been several previous reports on PDI of MRSA finding the drug-resistant strain to be as sensitive as the naïve strain¹⁶⁵ or be slightly less sensitive when compared with wild-type strains.¹⁶⁶ However, during the last few years, MRSA resistance has increased due to constant use of antimicrobials.¹⁶⁷

During PDI, PDT combined with photosensitizer (PS) toluidine blue O (TBO) (Fig. 5I), scientist were able to eliminate 100% of the MRSA in a sample obtained from a human wound with 3 laser exposures of 15 min duration.¹⁶⁸ Aluminum disulfonated phthalocyanine (AlPcS2) (Fig. 5B) was able to kill 3 logs of MRSA after gallium arsenide laser illumination (1.2 J, 11 mW) regardless the grow phase and the presence of horse or human serum as the medium.^{169,170} Tetrapyrrole-based photosensitizers, such as the porphyrin XF-73 (Fig. 5D) and the phthalocyanine RLP068/Cl (Fig. 5F), can kill multiple logs of MRSA respectively.^{171,172} In the same experimental conditions RLP068/Cl (but not TBO) was able to kill MRSA.¹⁷³ Sixteen epidemic strains of MRSA were subjected to PDI with AlPcS2 and all of them were susceptible to killing in a PS concentration-dependent manner.¹⁷⁴ PDI is effective in MRSA even when carried with non-coherent red light and polyethylenimine (PEI)-ce6 (2.7 logs of killing)¹⁷⁵ and is useful in wound models.¹⁷⁶ Treatment of local MRSA infections was improved when Hematoporphyrin (Hp) was encapsulated in liposomes or micelles.¹⁷⁷

S. pyogenes, also known as in group A streptococcus (GAS or group A strep), has been estimated to cause more than 500 000 deaths every year, making it one of the most harmful pathogens in the world.¹⁷⁸ Lethal photosensitization of *S. pyogenes* was performed with Indocyanine green (Fig. 5J), a negatively-charged polymethine dye, and a gallium-aluminum-arsenide (Ga-Al-As) NIR-Laser. Killing was 6.8 log, and even at the lowest concentration (25 µg/ml) killing was 4.7 log (99.99%).¹⁷⁹ PDI can be enhanced by PS entrapping.¹⁷⁷ A major difficulty in the inactivation of *S. pyogenes* is the formation of biofilms which are




Figure 6. Pathogenic factors of *S. aureus*, showing both the structural and the secreted products, playing roles as virulence factors. (A) Surface and secreted proteins; (B and C) are cross-sections of the cell envelope, from references 162 and 244 with modifications.

much more resistant to drug attack than isolated form of these bacteria. Hope and Wilson performed an interesting experiment which evaluated real-time PDI of *S. pyogenes* biofilms. They used Sn (IV) chlorin e6 (SnCe6) (Fig. 5L) as PS and illuminated with 488 nm argon and 543 nm HeNe lasers in a confocal microscope. Scanning the biofilm three times for 5 min each, they obtained significant reduction in biofilm fluorescence indicating the inactivation of the biofilm.¹⁸⁰

The gram-negative bacteria *Brucella abortus* and *F. tularensis* are responsible for extremely dangerous infections, brucellosis and tularaemia, respectively and are considered two of the most likely biowarfare agents. Both bacteria, in suspension with 0.1 mL of diluted methylene blue (MB) (Fig. 5E), with concentration 5 to 500 ppm were inactivated when illuminated with a 650 nm LED.¹⁸¹ *B. abortus* and *F. tularensis* were illuminated with 650 nm LED and saline and no killing effect was observed eliminating the possibility of photothermal damage.¹⁸¹

Recently, *Y. pestis*, a gram-negative bacterium, has gained attention as a possible biological warfare agent. A possible surrogate to study photoinactivation of *Y. pestis* is the gram-negative bacterium *Y. enterocolitica*. Using MB and several of its congeners against *Y. enterocolitica*, with illumination using a lamp emitting light in the waveband 615–645 nm, considerable bactericidal activity was noted using similar photosensitizer concentrations to those used elsewhere to inactivate blood-borne viruses. Two novel compounds in this area, the phenothiazinium new methylene blue N (Fig. 5M) and the phenoxazinium Brilliant Cresyl

Blue (Fig. 5N) exhibited bactericidal activity at lower concentrations than both of the established phenothiaziniums, MB and TBO and the recently published blood photovirucidal agent 1,9-dimethyl methylene blue (Fig. 5G). The photoactivity of these compounds was undiminished in the presence of red blood cells.¹⁸²

Macrophages are immune cells that play a pivotal role in the detection and elimination of pathogenic microorganisms by phagocytosis. Numerous pathogens, such as species of *Francisella*, *Legionella*, *Brucella*, and *Yersinia pestis*, parasitize macrophages, utilizing them as a host cell for their growth and replication, sometimes with disastrous effects. These infected macrophages therefore are a prime target for therapy and macrophage-targeted PDT may have a role to play especially when the infected macrophages are present in a localized granuloma.¹⁸³

Bacterial infections

Because PDI can have high selectivity for bacterial cells compared with host mammalian cells it is particularly suited as a treatment for localized infections.^{159,184} The PS is topically applied into the infected tissue which is then illuminated after a relatively short incubation time to ensure the PS is bound to the bacteria but has not had time to gain access to the host cells. The advantages of this approach compared with traditional antibiotics include its broad spectrum, rapid action, its equal effectiveness against multiply drug-resistant bacteria, and its ability to destroy bacteria in damaged tissue that has compromised blood perfusion. The effectiveness of PDI mediated by many of the PS described above

has been demonstrated in mouse models of wound infections (*E. coli*,¹⁸⁵ *P. aeruginosa*,¹⁸⁶ *Vibrio vulnificus*,¹⁸⁷ and MRSA¹⁵¹). PDI has also been studied in models of third degree burn infections by *S. aureus*¹⁸⁸ and *A. baumannii*.^{189,190} The effectiveness of PDI has also been demonstrated in deep established soft tissue abscesses caused by *S. aureus*.¹⁹¹

Bacterial endospores

B. anthracis is a gram-positive, endospore-forming bacterium that can grow under aerobic or anaerobic conditions. It is one of the major security and bioterrorism threats for this century since it cannot be easily inactivated by heat, radiation, antibiotics, or other antimicrobial agents.¹⁹² The experimental study of PDI of *B. anthracis* is difficult because of the biohazard risk involved.¹⁹³ Inhalation or ingestion will then cause a serious and frequently fatal disease, while entry of the spores into cuts and abrasions on the skin produces a less fatal but still serious disease, cutaneous anthrax. Anthrax is particularly deadly to humans due to the bacterium's ability to produce toxins with a sophisticated mechanism for killing mammalian cells.¹⁹⁴ Demidova and Hamblin¹⁹⁵ published a study demonstrating that a class of small cationic dyes known as phenothiazinium salts could photoinactivate 4 species of *Bacillus* spores that are surrogates to *B. anthracis*,¹⁹⁶ including *B. cereus* and *B. thuringiensis*, which are the same species as *B. anthracis*.¹⁹⁷ There were large differences in susceptibility to TBO-mediated PDI between spores of different *Bacillus* species. Spores of *B. cereus* and *B. thuringiensis* were the most susceptible. TBO (50 μ M) demonstrated a light-dose-dependent loss of viability of *B. cereus* and *B. thuringiensis* spores, with 40 J/cm² of 630 nm light leading to 99.999% killing. In contrast, *B. subtilis* and *B. atrophaeus* were much less sensitive and needed concentrations as high as 1.6 mM to achieve killing of >99.9% of cells and *B. megaterium*.¹⁹⁵ The relatively mild conditions needed for spore killing could have applications for treating wounds contaminated by anthrax spores, for which conventional sporicides would have unacceptable tissue toxicity.¹⁹⁵

Oliveira et al.¹⁹⁸ demonstrated that *B. cereus* endospores could be inactivated by porphyrin PS and light. There was a much smaller difference in sensitivity between spores and vegetative cells of *B. cereus* (the TBO concentration needed to kill spores was 3 to 4 times higher than that needed to kill vegetative cells) than between spores and vegetative cells of *B. subtilis* (>100 times the TBO concentration was needed to kill spores compared with vegetative cells).¹⁹⁵

B. atrophaeus has been used as a simulant for the biological warfare agent *B. anthracis* for decades. PDI of these spores was possible using an intense pulsed (period of 100 ms) visible light source in association with TMPyP (5, 10, 15, 20-Tetrakis [1-methylpyridinium-4-yl]-porphyrin tetra p-toluenesulfonate) (Fig. 5K). PDI induced oxidative damage which killed up to 6 log (>99.9999%) within a total treatment time of 10 s (fluencies from 20 J/cm² up to 80 J/cm²) using a TMPyP in a concentration range of 1–100 μ mol.¹⁹⁹ Similar experiment performed with only a single light flash (10 or 20 J/cm²) and 10 μ mol of TMPyP was able to kill more than 4 log of *B. atrophaeus*.²⁰⁰ These studies reinforce the application of PDI in military and national security for decontamination of anthrax spores.¹⁹⁸

Fungi

Fungi are eukaryotic cells that possess a cell wall outside the plasma membrane. *Coccidioides immitis* is the only fungal species present on the Select Agents Appendix A (biological warfare agents)²⁰¹: it is dimorphic, producing a mycelial form in nature that ages to produce spores (arthroconidia) that separate in a characteristic fashion via the disarticulation of the parent mycelium leaving the ruptured cell-wall fragments of adjacent cell remnants attached to opposing ends (Fig. 7). In vivo the spores enlarge to form spherules that are typically 20 microns or more in diameter when viewed in tissue sections of actively infected hosts. The spherules undergo internal divisions to yield endospores that are released upon maturation and go on to repeat the cycle of the infection.²⁰² Infection of normal hosts with spores of *C. immitis* can result in a spectrum of consequences ranging from minimal symptoms of disease or it can establish an active replicating cycle that can include profound pulmonary disease and dissemination from the pulmonary focus via the bloodstream to involve multiple systems of the body (typically meningitis, skin, bone, and internal organs). There are literature reports of PDI of a few species of fungus including both yeasts (*Saccharomyces*²⁰³ and *Candida* spp.^{204, 205}) and filamentous fungi (*Trichophyton*²⁰⁶ and *Aspergillus*²⁰⁷). As yet there have been no reports of PDT on actual *C. immitis* organisms but the successful eradication of related fungal species suggests that PDT should work well against this pathogen. Junqueira et al.²⁰⁸ reported on the use of a cationic nanoemulsion of zinc 2,9,16,23-tetrakis(phenylthio)-29H, 31H-phthalocyanine (Fig. 5C) to mediate PDI of biofilms formed by *Candida* spp. and the emerging pathogens *Trichosporon mucoides* and *Kodamaea ohmeri*.

Viruses

The short-lived ROS generated by PDI mechanisms are responsible for the damage induced to critical molecular targets in viruses.^{209,210} Different viral targets, such as the envelope lipids and proteins, capsid and core proteins, and the nucleic acid, can be attacked by singlet oxygen and/or other ROS (hydrogen peroxide, superoxide, and hydroxyl radicals) to achieve the loss of infectivity.²¹⁰ Viral DNA is one of the critical target structures for PDI by MB and light MB causing direct DNA damage and blockage of DNA replication which has been successfully used for HSV-1 treatment.²¹¹ It has been shown that enveloped viruses can be inactivated due to protein damage. However, while the same treatment is reported to be ineffective against some non-enveloped viruses,²¹² the results from Wong et al.²¹³ showed that even a non-enveloped virus can be efficiently inactivated due to the damage induced by PDI to its viral proteins. The efficiency of different types of PS in viral PDI has been proved for different types of mammalian viruses and bacteriophages, whether they are enveloped or non-enveloped, for either DNA or RNA viruses.²¹⁴ PDI of viruses has been of special interest for applications in blood banking sterilization.²¹⁵ Therefore, several types of virus have been tested for PDI.²¹⁵

Ebola

The filoviruses, Marburg and Ebola, are classified as Category A biowarfare agents by the Centers for Disease Control. Most known human infections with these viruses have been fatal

(fatality rates for humans of up to 90%) and no vaccines or effective therapies are currently available. They are enveloped, nonsegmented, negative-stranded RNA viruses.²¹⁶ Enveloped, RNA viruses from two different families, Semliki Forest virus (SFV, *Togaviridae*) and vesicular stomatitis virus (VSV, *Rhabdoviridae*), and can act as surrogate filovirus models for PDI. Using a suspension of 1 mg/ml fullerene C60 (buckyball) (Fig. 5O) as the PS and visible light (wavelengths higher than 495 nm) under constant stirring and flushing with oxygen, values of 7 log reduction for 5 h of illumination were obtained for both families of enveloped virus.²¹⁷ VSV plaque forming units were decreased by 5 log using methoxy-polyethylene glycol conjugated fullerene, illuminated by 120 J/cm² white light.²¹⁸

Smallpox: Variola major and Variola minor (Orthopoxviruses).

It is known that smallpox has been used as a biowarfare agent in the past. During World War II, scientists from the United Kingdom, the United States, and Japan were involved in research into producing a biological weapon from smallpox.²¹⁹ In 1992 Soviet defector Ken Alibek confirmed that the Soviet bioweapons program at Zagorsk had produced a large stockpile—as much as 20 tons—of weaponized smallpox (possibly engineered to resist vaccines), along with refrigerated warheads to deliver it. It is not known whether these stockpiles still exist in Russia. With the breakup of the Soviet Union and unemployment of many of the weapons program's scientists, there is concern that smallpox and the expertise to weaponize it may have become available to other governments or terrorist groups who might wish to use virus as means of biological warfare.²²⁰ The last occurrence of endemic smallpox was in Somalia in 1977, and the last human cases were laboratory-acquired infections in 1978. There are four types of *Variola major* smallpox: ordinary (the most frequent type, accounting for 90% or more of cases); modified (mild and occurring in previously vaccinated persons); flat; and hemorrhagic (both rare and very severe). Historically, *Variola major* has an overall fatality rate of about 30%; however, flat and hemorrhagic smallpox are usually fatal. Present laboratory examination of Variola virus requires high-containment (Biosafety Level 4).²²¹

Variola virus is the most notorious poxvirus, a member of a family of large, enveloped DNA viruses. It is generally accepted that enveloped viruses can be inactivated efficiently by singlet oxygen generating agents such as PDI. PDI of HIV-1 by MB/light treatment acts on HIV-1 at different target sites: the envelope and core proteins, and the inner core structures like RNA.²²²

Four PS (MB, rose bengal [RB] [Fig. 5H], uroporphyrin [UP], and aluminum phthalocyanine tetrasulphonate [AlPcS4]) could




Figure 7. *Coccidioides immitis* is the only fungal species present on the Select Agents Appendix A (biological warfare agents).²⁰¹ It is dimorphic, producing a mycelial form in nature that matures to produce spores (arthroconidia) that go on to repeat the cycle of the infection.²⁰²

inactivate adenovirus. Using MB (2.7 mM) and light (intensity of 106 mW/cm²) produced a complete inactivation of adenovirus after 1 min of exposure: 10 mM of RB was enough for just 0.5 log reduction after 20 min of illumination and complete inactivation was obtained after 30 min PDI with UP; however, AlPcS4 could not completely inactivate adenovirus even when used in 50 mM for 30 min.²²³

Nucleic acids may be important targets for photoinactivation of DNA viruses by MB and AlPcS4.²²⁴ Photoinactivation of DNA viruses are more efficiently induced by free than by DNA bound porphyrin. Photoreactions of TMPyP and TMPyMPP affect the structural integrity of DNA and also of viral proteins, despite their selective DNA binding.²²⁵ The binding of cationic porphyrins to DNA is presumably due to the electrostatic interaction between the positively-charged substituents in the porphyrin macrocycle and the negatively charged phosphate oxygen atoms of DNA.²²⁶

Lassa virus (Arenavirus) and RVF virus (Bunyaviridae)

Lassa virus (LASV) is an *Arenavirus* that causes Lassa hemorrhagic fever in human and non-human primates. Rift Valley fever (RVF) is caused by RVF virus belonging to *Bunyaviridae*, which is a family of negative-stranded, enveloped RNA viruses. Lassa virus and RVF virus are enveloped RNA viruses that are select agents requiring Biosafety Level 4-equivalent containment.²²¹

Dengue virus

Dengue and yellow fever viruses belong to the genus *Flavivirus* single-stranded RNA viruses. Dengue virus, an enveloped RNA virus, could be inactivated using MB in combination with a LED cluster (mid-peak bandwidth 29 nm, peak 664 nm). The amount of dengue virus remaining was evaluated by plaque forming assays. Dengue virus was completely inactivated within 5 min when the




Figure 8. Diagrammatic representation of the mode of action of several bacterial toxins. (A) Damage to cellular membranes by *Staphylococcus aureus* toxin. After binding and oligomerization, the stem of the mushroom-shaped toxin heptamer inserts into the target cell and disrupts membrane permeability as depicted by the influx and efflux of ions represented by red and green circles. (B) Inhibition of protein synthesis by Shiga toxins (Stx). Holotoxin, which consists of an enzymatically active (A) subunit and 5 binding (B) subunits, enters cells through the globotriaosylceramide (Gb3) receptor. The N-glycosidase activity of the (A) subunit then cleaves an adenosine residue from 28S rRNA, which halts protein synthesis. (C) Examples of bacterial toxins that activate secondary messenger pathways. Binding of the heat-stable enterotoxins (ST) to a guanylate cyclase receptor results in an increase in cyclic GMP (cGMP) that adversely effects electrolyte flux. By ADP-ribosylation or glucosylation respectively, the C3 exoenzyme (C3) of *Clostridium botulinum* and the *Clostridium difficile* toxins A and B (CdA and CdB) inactivate the small Rho GTP-binding proteins. Cytotoxic necrotizing factor (CNF) of *E. coli* and the dermonecrotic toxin (DNT) of *Bordetella* species activate Rho by deamidation.

MB concentration was higher than 1.0 μg/mL.²²⁷ Lin et al.²²⁸ compared light-dependent and light independent inactivation of dengue-2 and other enveloped viruses by the two regio-isomers of carboxyfullerene and found that asymmetric isomer had greater dark activity (even at much higher concentrations than needed for its PDT effect) due to its interaction with the lipid envelope of the virus.

Toxins

PDI is one of the few antimicrobial treatments that is also capable of inactivating toxins and secreted virulence factors produced by pathogens. The reactive oxygen species produced during photodynamic action (¹O₂ and HO^{*}) can attack molecular features susceptible to oxidation (sulfur atoms, aromatic rings, heterocyclic rings, unsaturated double bonds, amino groups, etc.) present on the toxin molecules themselves. These oxidative reactions can disturb the conformation or alter the functional groups of the toxins and abolish the biological function (Fig. 8). This approach has been well-demonstrated in the case of lipopolysaccharide (LPS, endotoxin from gram-negative bacteria). Komerik et al.²²⁹ first showed that TBO and red light could inactivate LPS from *E. coli* and they also were able to inactivate proteases from *P. aeruginosa*. Gianelli and colleagues²³⁰ used MB combined

with various light sources to inactivate *P. gingivalis* LPS adherent to titanium discs, cut from commercial dental implants. Tubby et al.²³¹ studied the ability of MB and red light to inactivate the following secreted virulence factors of *S. aureus*: V8 protease, α-hemolysin, and sphingomyelinase were shown to be inhibited in a dose-dependent manner by exposure to light in the presence of MB. Eubanks et al.²³² showed that an actual bio warfare agent, botulinum neurotoxin, could be photo-inactivated by exposure to riboflavin and white light. Our laboratory has obtained evidence that two additional microbial toxins, Shiga-like toxin from *E. coli* O157 and mycolactone from *Mycobacterium ulcerans* can be destroyed by exposure to benzoporphyrin derivative and red light (manuscript in preparation).

Anti-Microbial Effect of Femtosecond Lasers

It has been proposed that femtosecond lasers, or lasers that maintain a pulse duration of 10⁻¹⁵ s, break down transparent or semitransparent biological tissues due to nonlinear absorption of laser energy with minimal thermal and mechanical

effects.²³³ As a result of the adverse collateral damage possible with other laser systems, the femtosecond laser has been hypothesized to be a new approach for killing pathogens.

Recently, a series of studies reported the efficacy of a visible femtosecond laser or a near-infrared subpicosecond fiber laser on inactivation of a variety of viral species, including M13 bacteriophage, tobacco mosaic virus, human papillomavirus, and human immunodeficiency virus.²³⁴⁻²⁴⁰ M13 phages were inactivated by using a very low power (as low as 0.5 nJ/pulse) visible femtosecond laser with 425 nm wavelength, 100 fs pulse width, power density ≥ 50 MW/cm².²³⁷

One group reported²⁴¹ inactivation of an encephalomyocarditis virus, M13 bacteriophage, and *Salmonella* Typhimurium by a visible femtosecond diode-pumped continuous-wave (CW) mode-locked Ti-sapphire laser. The laser produced a continuous train of 60 fs pulses at a repetition rate of 80 MHz. The excitation laser was chosen to operate at a wavelength of λ = 425 nm and with an average power of about 50 mW. It has a pulse width of full-width at half maximum (FWHM) ≈ 100 fs. All the microorganisms were inactivated very efficiently, especially *S. Typhimurium*. There were different mechanisms of inactivation of different microorganism by femtosecond laser.

Inactivation of viruses involves the breaking of hydrogen/hydrophobic bonds or the separation of the weak protein links in the protein shell of a viral particle. On the contrary, inactivation of bacteria is related to the damage of their DNA due to irradiation of a visible femtosecond laser.

Another study²⁴² reported the inactivation of murine cytomegalovirus (MCMV), an enveloped, double-stranded DNA virus, by a visible (425 nm) femtosecond laser. The results showed that the laser irradiation caused a 5-log reduction in MCMV titer and caused selective aggregation of viral capsid and tegument proteins. However, the femtosecond laser did not cause significant changes to the global structure of MCMV virions including membrane and capsid, as assessed by electron microscopy; meanwhile, it could not produce the double-strand breaks or crosslinking in MCMV genomic DNA.

Manipulation of a near-infrared (NIR) femtosecond laser via impulsive stimulated raman scattering (ISRS) to produce damage (e.g., to the protein coat of a virus) is another method for selectively inactivating microorganisms.²³⁷

When NIR femtosecond laser induced the inactivation of virus and bacteria, its safety to the mammalian cells was considered. The relative research demonstrated that if the wavelength and pulse width of the femtosecond laser were appropriately selected, there was a window in power density that enabled them to achieve selective inactivation of target viruses and bacteria without causing cytotoxicity to mammalian cells. It was suggested that this strategy targeted the mechanical (vibrational) properties of microorganisms and thus its antimicrobial efficacy was likely to be unaffected by genetic differences in the microorganisms.²³⁸

In the view of the emerging threats from drug resistant pathogens and microorganisms, developing novel and more effective antimicrobial strategies is an absolute necessity. One such strategy is to develop the ultrashort pulsed (USP) laser technology as an effective and chemicals free inactivation technique that can be successfully used over broad spectrum of pathogens, both from bacterial and viral sources.²⁴³

In summary, the advantages of such novel laser technologies over the presently prevailing disinfection methods include: they are considered as noninvasive disinfection technologies, because no foreign materials are needed in the disinfection process; they are harmless environmental disinfection methods since no chemicals are used in the pathogen inactivation process; and they

are general methods for selective disinfection of pathogens with potentially minimal side effects.²⁴¹

Conclusion

Recent studies have highlighted the diversity of applications of light-mediated technology against pathogens of all known classes. Wavelengths from the short-UV to the near-infrared (either alone or combined with PS) can be used to kill or inactivate gram-positive and gram-negative bacteria, fungi, endospores, parasites, viruses, and even protein toxins. The mechanisms of action depend on the different microbial types and the wavelength and presence or not of a PS. The two broad target classes are nucleic acids for UVC and PUVA and oxidizable proteins for photocatalysis, PDT, and blue light. The broad occurrence of these biological targets in bioweapons agents means that the light-mediated technology is highly likely to be very broad-spectrum, thus avoiding the need to know the identity of the particular agent in any mass biological attack, and also suggests that the development of resistance to light-mediated inactivation is likely to non-existent. Furthermore, light is non-polluting and environmentally friendly, and even if PS need to be used, these compounds are likely to be photodegraded rapidly when the bio-threat has been neutralized thus leaving no lasting pollution. The use of light-based technology to prevent and treat actual infections suggests that they may be useful to decontaminate humans that have already received exposure to biological agents, without causing undue harm to host tissue. Lastly light-based inactivation may be particularly suitable to form vaccines as they kill pathogens while preserving their antigenicity.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Dedication

We dedicate this work to the great philanthropist Dr Orhan Ocalgiray for his profound contributions to the field of microbiology.

Acknowledgments

Research in the Hamblin laboratory is supported by US NIH grant R01AI050875.

References

1. Thavaselvam D, Vijayaraghavan R. Biological warfare agents. *J Pharm Bioallied Sci* 2010; 2:179-88; PMID:21829313; <http://dx.doi.org/10.4103/0975-7406.68499>
2. Oyston PC. *Francisella tularensis*: unravelling the secrets of an intracellular pathogen. *J Med Microbiol* 2008; 57:921-30; PMID:18628490; <http://dx.doi.org/10.1099/jmm.0.2008/000653-0>
3. McDevitt JJ, Milton DK, Rudnick SN, First MW. Inactivation of poxviruses by upper-room UVC light in a simulated hospital room environment. *PLoS One* 2008; 3:e3186; PMID:18781204; <http://dx.doi.org/10.1371/journal.pone.0003186>
4. Achtman M, Zurth K, Morelli G, Torrea G, Guiyoule A, Carniel E. *Yersinia pestis*, the cause of plague, is a recently emerged clone of *Yersinia pseudotuberculosis*. *Proc Natl Acad Sci U S A* 1999; 96:14043-8; PMID:10570195; <http://dx.doi.org/10.1073/pnas.96.24.14043>
5. Wuthiekanun V, Peacock SJ. Management of melioidosis. *Expert Rev Anti Infect Ther* 2006; 4:445-55; PMID:16771621; <http://dx.doi.org/10.1586/14787210.4.3.445>
6. Klevens RM, Morrison MA, Nadle J, Petit S, Gershman K, Ray S, Harrison LH, Lynfield R, Dumyati G, Townes JM, et al.; Active Bacterial Core surveillance (ABCs) MRSA Investigators. Invasive methicillin-resistant *Staphylococcus aureus* infections in the United States. *JAMA* 2007; 298:1763-71; PMID:17940231; <http://dx.doi.org/10.1001/jama.298.15.1763>
7. Musser JM, Hauser AR, Kim MH, Schlievert PM, Nelson K, Selander RK. *Streptococcus pyogenes* causing toxic-shock-like syndrome and other invasive diseases: clonal diversity and pyrogenic exotoxin expression. *Proc Natl Acad Sci U S A* 1991; 88:2668-72; PMID:1672766; <http://dx.doi.org/10.1073/pnas.88.7.2668>
8. Helgason E, Økstad OA, Caugant DA, Johansen HA, Fouet A, Mock M, Hegna I, Kolstø AB. *Bacillus anthracis*, *Bacillus cereus*, and *Bacillus thuringiensis*—one species on the basis of genetic evidence. *Appl Environ Microbiol* 2000; 66:2627-30; PMID:10831447; <http://dx.doi.org/10.1128/AEM.66.6.2627-2630.2000>
9. Sagripanti JL, Lytle CD. Sensitivity to ultraviolet radiation of Lassa, vaccinia, and Ebola viruses dried on surfaces. *Arch Virol* 2011; 156:489-94; PMID:21104283; <http://dx.doi.org/10.1007/s00705-010-0847-1>

10. Sugiyama H. Clostridium botulinum neurotoxin. *Microbiol Rev* 1980; 44:419-48; PMID:6252433
11. Meselson M, Guillemin J, Hugh-Jones M, Langmuir A, Popova I, Shelokov A, Yampolskaya O. The Sverdlovsk anthrax outbreak of 1979. *Science* 1994; 266:1202-8; PMID:7973702; <http://dx.doi.org/10.1126/science.7973702>
12. Manchee RJ, Broster MG, Melling J, Henstridge RM, Stagg AJ. Bacillus anthracis on Gruinard Island. *Nature* 1981; 294:254-5; PMID:6795509; <http://dx.doi.org/10.1038/294254a0>
13. Manchee RJ, Broster MG, Stagg AJ, Hibbs SE. Formaldehyde Solution Effectively Inactivates Spores of Bacillus anthracis on the Scottish Island of Gruinard. *Appl Environ Microbiol* 1994; 60:4167-71; PMID:16349444
14. Aldhous P. Biological warfare. Gruinard Island handed back. *Nature* 1990; 344:801; PMID:2109833; <http://dx.doi.org/10.1038/344801b0>
15. Grunow R, Finke EJ. A procedure for differentiating between the intentional release of biological warfare agents and natural outbreaks of disease: its use in analyzing the tularemia outbreak in Kosovo in 1999 and 2000. *Clin Microbiol Infect* 2002; 8:510-21; PMID:12197873; <http://dx.doi.org/10.1046/j.1469-0691.2002.00524.x>
16. Török TJ, Tauxe RV, Wise RP, Livengood JR, Sokolow R, Mauvais S, Birkness KA, Skeels MR, Horan JM, Foster LR. A large community outbreak of salmonellosis caused by intentional contamination of restaurant salad bars. *JAMA* 1997; 278:389-95; PMID:9244330; <http://dx.doi.org/10.1001/jama.1997.03550050051033>
17. Kolavic SA, Kimura A, Simons SL, Slutsker L, Barth S, Haley CE. An outbreak of Shigella dysenteriae type 2 among laboratory workers due to intentional food contamination. *JAMA* 1997; 278:396-8; PMID:9244331; <http://dx.doi.org/10.1001/jama.1997.03550050058034>
18. Hsu VP, Lukacs SL, Handzel T, Hayslett J, Harper S, Hales T, Semenova VA, Romero-Steiner S, Elic C, Quinn CP, et al. Opening a bacillus anthracis-containing envelope, Capitol Hill, Washington, D.C.: the public health response. *Emerg Infect Dis* 2002; 8:1039-43; PMID:12396912; <http://dx.doi.org/10.3201/eid0810.020332>
19. Centers for Disease Control and Prevention (CDC). Update: Investigation of bioterrorism-related anthrax, 2001. *MMWR Morb Mortal Wkly Rep* 2001; 50:1008-10; PMID:11724158
20. Centers for Disease Control and Prevention (CDC). Update: Cutaneous anthrax in a laboratory worker--Texas, 2002. *MMWR Morb Mortal Wkly Rep* 2002; 51:482; PMID:12064454
21. Cutler TD, Zimmerman JJ. Ultraviolet irradiation and the mechanisms underlying its inactivation of infectious agents. *Anim Health Res Rev* 2011; 12:15-23; PMID:21676338; <http://dx.doi.org/10.1017/S1466252311000016>
22. Jagger J. Introduction to research in ultraviolet photobiology. Prentice-Hall (Englewood Cliffs, N.J), 1967.
23. Reed NG. The history of ultraviolet germicidal irradiation for air disinfection. *Public Health Rep* 2010; 125:15-27; PMID:20402193
24. Brickner PW, Vincent RL, First M, Nardell E, Murray M, Kaufman W. The application of ultraviolet germicidal irradiation to control transmission of airborne disease: bioterrorism countermeasure. *Public Health Rep* 2003; 118:99-114; PMID:12690064; [http://dx.doi.org/10.1016/S0033-3549\(04\)50225-X](http://dx.doi.org/10.1016/S0033-3549(04)50225-X)
25. Dai T, Garcia B, Murray CK, Vrahas MS, Hamblin MR. UVC light prophylaxis for cutaneous wound infections in mice. *Antimicrob Agents Chemother* 2012; 56:3841-8; PMID:22564833; <http://dx.doi.org/10.1128/AAC.00161-12>
26. Ladd M. Introduction to Physical Chemistry. London, UK: Cambridge University Press, 1998.
27. Durban E, Grecz N. Resistance of spores of Clostridium botulinum 33A to combinations of ultraviolet and gamma rays. *Appl Microbiol* 1969; 18:44-50; PMID:4896102
28. Nourmoradi H, Nikaeen M, Stensvold CR, Mirhendi H. Ultraviolet irradiation: An effective inactivation method of Aspergillus spp. in water for the control of waterborne nosocomial aspergillosis. *Water Res* 2012; 46:5935-40; PMID:22985523; <http://dx.doi.org/10.1016/j.watres.2012.08.015>
29. Guerrero-Beltran J, Welti-Chanes J, Barbosa-Canovas G. UVC light processing of grape, cranberry and grapefruit juices to inactivate Saccharomyces Cerevisiae. *J Food Process Eng* 2009; 32:916-32; <http://dx.doi.org/10.1111/j.1745-4530.2008.00253.x>
30. Levy C, Aubert X, Lacour B, Carlin F. Relevant factors affecting microbial surface decontamination by pulsed light. *Int J Food Microbiol* 2012; 152:168-74; PMID:21924512; <http://dx.doi.org/10.1016/j.ijfoodmicro.2011.08.022>
31. Luna VA, Cannons AC, Amuso PT, Cattani J. The inactivation and removal of airborne Bacillus atrophaeus endospores from air circulation systems using UVC and HEPA filters. *J Appl Microbiol* 2008; 104:489-98; PMID:17927759
32. Dean SJ, Petty A, Swift S, McGhee JJ, Sharma A, Shah S, Craig JP. Efficacy and safety assessment of a novel ultraviolet C device for treating corneal bacterial infections. *Clin Experiment Ophthalmol* 2011; 39:156-63; PMID:21105972; <http://dx.doi.org/10.1111/j.1442-9071.2010.02471.x>
33. Umezawa K, Asai S, Inokuchi S, Miyachi H. A comparative study of the bactericidal activity and daily disinfection housekeeping surfaces by a new portable pulsed UV radiation device. *Curr Microbiol* 2012; 64:581-7; PMID:22447288; <http://dx.doi.org/10.1007/s00284-012-0110-y>
34. Al-Marir A. Ultraviolet C lethal effect on Brucella melitensis. *Microbiologica-Quarterly J of Microbiol Sci* 2008; 31:47-56
35. Rose LJ, O'Connell H. UV light inactivation of bacterial biothreat agents. *Appl Environ Microbiol* 2009; 75:2987-90; PMID:19270145; <http://dx.doi.org/10.1128/AEM.02180-08>
36. Miller SL, Linnes J, Luongo J. Ultraviolet germicidal irradiation: future directions for air disinfection and building applications. *Photochem Photobiol* 2013; 89:777-81; PMID:23581680; <http://dx.doi.org/10.1111/php.12080>
37. Lakretz A, Ron EZ, Mamane H. Biofouling control in water by various UVC wavelengths and doses. *Biofouling* 2010; 26:257-67; PMID:20024789; <http://dx.doi.org/10.1080/08927010903484154>
38. Kalisvaart BF. Re-use of wastewater: preventing the recovery of pathogens by using medium-pressure UV lamp technology. *Water Sci Technol* 2004; 50:337-44; PMID:15537023
39. Li J, Hirota K, Yumoto H, Matsuo T, Miyake Y, Ichikawa T. Enhanced germicidal effects of pulsed UV-LED irradiation on biofilms. *J Appl Microbiol* 2010; 109:2183-90; PMID:20854456; <http://dx.doi.org/10.1111/j.1365-2672.2010.04850.x>
40. Farrell NP, Garvey M, Cormican M, Laffey JG, Rowan NJ. Investigation of critical inter-related factors affecting the efficacy of pulsed light for inactivating clinically relevant bacterial pathogens. *J Appl Microbiol* 2010; 108:1494-508; PMID:19796119; <http://dx.doi.org/10.1111/j.1365-2672.2009.04545.x>
41. Nicholson WL, Galeano B. UV resistance of Bacillus anthracis spores revisited: validation of Bacillus subtilis spores as UV surrogates for spores of B. anthracis Sterne. *Appl Environ Microbiol* 2003; 69:1327-30; PMID:12571068; <http://dx.doi.org/10.1128/AEM.69.2.1327-1330.2003>
42. Rettberg P, Cockell CS. Biological UV dosimetry using the DLR-biofilm. *Photochem Photobiol Sci* 2004; 3:781-7; PMID:15295635; <http://dx.doi.org/10.1039/b315950g>
43. Moehrle M, Dennenmoser B, Garbe C. Continuous long-term monitoring of UV radiation in professional mountain guides reveals extremely high exposure. *Int J Cancer* 2003; 103:775-8; PMID:12516097; <http://dx.doi.org/10.1002/ijc.10884>
44. Moehrle M, Korn M, Garbe C. Bacillus subtilis spore film dosimeters in personal dosimetry for occupational solar ultraviolet exposure. *Int Arch Occup Environ Health* 2000; 73:575-80; PMID:11100953; <http://dx.doi.org/10.1007/s004200000183>
45. Vähävihi K, Ylianttila L, Kautiainen H, Tuohimaa P, Reunala T, Snellman E. Spore film dosimeters are feasible for UV dose monitoring during heliotherapy. *Photochem Photobiol* 2010; 86:1174-8; PMID:20573044; <http://dx.doi.org/10.1111/j.1751-1097.2010.00769.x>
46. Bendja A, Heil K, Barends TR, Carell T, Schlichting I. Structural insights into recognition and repair of UV-DNA damage by Spore Photoproduct Lyase, a radical SAM enzyme. *Nucleic Acids Res* 2012; 40:9308-18; PMID:22761404; <http://dx.doi.org/10.1093/nar/gks603>
47. Yang L, Lin G, Nelson RS, Jian Y, Telser J, Li L. Mechanistic studies of the spore photoproduct lyase via a single cysteine mutation. *Biochemistry* 2012; 51:7173-88; PMID:22906093; <http://dx.doi.org/10.1021/bi3010945>
48. Silver SC, Chandra T, Zilinskas E, Ghose S, Broderick WE, Broderick JB. Complete stereospecific repair of a synthetic dinucleotide spore photoproduct by spore photoproduct lyase. *J Biol Inorg Chem* 2010; 15:943-55; PMID:20405152; <http://dx.doi.org/10.1007/s00775-010-0656-8>
49. Friedel MG, Berteau O, Pieck JC, Atta M, Ollagnier-de-Choudens S, Fontecave M, Carell T. The spore photoproduct lyase repairs the 5S- and not the 5R-configured spore photoproduct DNA lesion. *Chem Commun (Camb)* 2006; 445-7; PMID:16493831; <http://dx.doi.org/10.1039/b514103f>
50. Setlow P. Resistance of spores of Bacillus species to ultraviolet light. *Environ Mol Mutagen* 2001; 38:97-104; PMID:11746741; <http://dx.doi.org/10.1002/em.1058>
51. Negro BL, Crafford D, Viljoen M. The effect of sympathomimetic medication on cardiovascular functioning of children with attention-deficit/hyperactivity disorder. *Cardiovasc J Afr* 2009; 20:296-9; PMID:19907802
52. Chandra T, Silver SC, Zilinskas E, Shepard EM, Broderick WE, Broderick JB. Spore photoproduct lyase catalyzes specific repair of the 5R but not the 5S spore photoproduct. *J Am Chem Soc* 2009; 131:2420-1; PMID:19178276; <http://dx.doi.org/10.1021/ja807375c>
53. Yang L, Nelson RS, Bendja A, Lin G, Telser J, Stoll S, Schlichting I, Li L. A radical transfer pathway in spore photoproduct lyase. *Biochemistry* 2013; 52:3041-50; PMID:23607538; <http://dx.doi.org/10.1021/bi3016247>
54. Bucheli-Witschel M, Bassin C, Egli T. UV-C inactivation in Escherichia coli is affected by growth conditions preceding irradiation, in particular by the specific growth rate. *J Appl Microbiol* 2010; 109:1733-44; PMID:20629801
55. Zenoff VF, Heredia J, Ferrero M, Siñeriz F, Farías ME. Diverse UV-B resistance of culturable bacterial community from high-altitude wetland water. *Curr Microbiol* 2006; 52:359-62; PMID:16604419; <http://dx.doi.org/10.1007/s00284-005-0241-5>
56. Steinborn G. Uvm mutants of Escherichia coli K12 deficient in UV mutagenesis. I. Isolation of uvm mutants and their phenotypical characterization in DNA repair and mutagenesis. *Mol Gen Genet* 1978; 165:87-93; PMID:362169; <http://dx.doi.org/10.1007/BF00270380>

57. Munoz-Najar U, Vijayakumar MN. An operon that confers UV resistance by evoking the SOS mutagenic response in streptococcal conjugative transposon Tn5252. *J Bacteriol* 1999; 181:2782-8; PMID:10217768
58. Dai T, Vrahas MS, Murray CK, Hamblin MR. Ultraviolet C irradiation: an alternative antimicrobial approach to localized infections? *Expert Rev Anti Infect Ther* 2012; 10:185-95; PMID:22339192; <http://dx.doi.org/10.1586/eri.11.166>
59. Dai T, Murray CK, Vrahas MS, Baer DG, Tegos GP, Hamblin MR. Ultraviolet C light for *Acinetobacter baumannii* wound infections in mice: potential use for battlefield wound decontamination? *J Trauma Acute Care Surg* 2012; 73:661-7; PMID:22929495; <http://dx.doi.org/10.1097/TA.0b013e31825c149c>
60. Dai T, Kharkwal GB, Zhao J, St Denis TG, Wu Q, Xia Y, Huang L, Sharma SK, d'Enfert C, Hamblin MR. Ultraviolet-C light for treatment of *Candida albicans* burn infection in mice. *Photochem Photobiol* 2011; 87:342-9; PMID:21208209; <http://dx.doi.org/10.1111/j.1751-1097.2011.00886.x>
61. Hashimoto K, Irie H, Fujishima A. TiO₂ Photocatalysis: A Historical Overview and Future Prospects. *Jpn J Appl Phys* 2005; 44:8269-85; <http://dx.doi.org/10.1143/JJAP.44.8269>
62. Fujishima A, Honda K. Electrochemical photolysis of water at a semiconductor electrode. *Nature* 1972; 238:37-8; PMID:12635268; <http://dx.doi.org/10.1038/238037a0>
63. Carey JH, Lawrence J, Tosine HM. Photodechlorination of PCB's in the presence of titanium dioxide in aqueous suspensions. *Bull Environ Contam Toxicol* 1976; 16:697-701; PMID:828069; <http://dx.doi.org/10.1007/BF01685575>
64. Frank SN, Bard AJ. Heterogeneous photocatalytic oxidation of cyanide ion in aqueous solutions at titanium dioxide powder. *J Am Chem Soc* 1977; 99:303-4; <http://dx.doi.org/10.1021/ja00443a081>
65. Chen J, Poon CS. Photocatalytic construction and building materials: from fundamentals to applications. *Build Environ* 2009; 44:1899-906; <http://dx.doi.org/10.1016/j.buildenv.2009.01.002>
66. Fujishima A, Rao TN, Tryk DA. Titanium dioxide photocatalysis. *J Photochem Photobiol Photochem Rev* 2000; 1:1-21; [http://dx.doi.org/10.1016/S1389-5567\(00\)00002-2](http://dx.doi.org/10.1016/S1389-5567(00)00002-2)
67. Lilja M, Forsgren J, Welch K, Astrand M, Engqvist H, Strømme M. Photocatalytic and antimicrobial properties of surgical implant coatings of titanium dioxide deposited through cathodic arc evaporation. *Biotechnol Lett* 2012; 34:2299-305; PMID:22941372; <http://dx.doi.org/10.1007/s10529-012-1040-2>
68. Schrauben JN, Hayoun R, Valdez CN, Braten M, Fridley L, Mayer JM. Titanium and zinc oxide nanoparticles are proton-coupled electron transfer agents. *Science* 2012; 336:1298-301; PMID:22679095; <http://dx.doi.org/10.1126/science.1220234>
69. Matsunaga T, Tomoda R, Nakajima T, Wake H. Photoelectrochemical sterilization of microbial cells by semiconductor powders. *FEMS Microbiol Lett* 1985; 29:211-4; <http://dx.doi.org/10.1111/j.1574-6968.1985.tb00864.x>
70. Matsunaga T, Tomoda R, Nakajima T, Nakamura N, Komine T. Continuous-sterilization system that uses photosemiconductor powder. *Appl Environ Microbiol* 1988; 54:1330-3; PMID:3046487
71. Foster HA, Ditta IB, Varghese S, Steele A. Photocatalytic disinfection using titanium dioxide: spectrum and mechanism of antimicrobial activity. *Appl Microbiol Biotechnol* 2011; 90:1847-68; PMID:21523480; <http://dx.doi.org/10.1007/s00253-011-3213-7>
72. McCullagh C, Robertson JM, Bahnemann DW, K. RP. The application of TiO₂ photocatalysis for disinfection of water contaminated with pathogenic micro-organisms: a review. *Res Chem Intermed* 2007; 33:359-75; <http://dx.doi.org/10.1163/15685670779238775>
73. Cushnie TP, Robertson PK, Officer S, Pollard PM, McCullagh C, Robertson JM. Variables to be considered when assessing the photocatalytic destruction of bacterial pathogens. *Chemosphere* 2009; 74:1374-8; PMID:19101016; <http://dx.doi.org/10.1016/j.chemosphere.2008.11.012>
74. Ireland JC, Klostermann P, Rice EW, Clark RM. Inactivation of *Escherichia coli* by titanium dioxide photocatalytic oxidation. *Appl Environ Microbiol* 1993; 59:1668-70; PMID:8390819
75. Pham HN, McDowell T, Wilkins E. Photocatalytically-mediated disinfection of water using TiO₂ as a catalyst and spore-forming *Bacillus pumilus* as a model. *J Environ Sci and Health* 1995; A30:627-36
76. Inglesby TV. Anthrax: A possible case history. *Emerg Infect Dis* 1999; 5:556-60; PMID:10458965; <http://dx.doi.org/10.3201/eid0504.990419>
77. Armon R, Welch-Cohen G, Bettane P. Disinfection of *Bacillus* spp. spores in drinking water by TiO₂ photocatalysis as a model for *Bacillus anthracis*. *Water Supply* 2004; 4:7-14
78. Maness PC, Smolinski S, Blake DM, Huang Z, Wolfrum EJ, Jacoby WA. Bactericidal activity of photocatalytic TiO₂(2) reaction: toward an understanding of its killing mechanism. *Appl Environ Microbiol* 1999; 65:4094-8; PMID:10473421
79. Maness PC, Smolinski S, Blake DM, Huang Z, Wolfrum EJ, Jacoby WA. Bactericidal activity of photocatalytic TiO₂(2) reaction: toward an understanding of its killing mechanism. *Appl Environ Microbiol* 1999; 65:4094-8; PMID:10473421
80. Pelaez M, Nolan NT, Pillai SC, Seery MK. A review on the visible light active titanium dioxide photocatalysis for environmental applications. *Appl Catal B* 2012; <http://dx.doi.org/10.1016/j.apcatb.2012.05.036>
81. Gogniat G, Thyssen M, Denis M, Pulgarin C, Dukan S. The bactericidal effect of TiO₂ photocatalysis involves adsorption onto catalyst and the loss of membrane integrity. *FEMS Microbiol Lett* 2006; 258:18-24; PMID:16630249; <http://dx.doi.org/10.1111/j.1574-6968.2006.00190.x>
82. Liou JW, Chang HH. Bactericidal effects and mechanisms of visible light-responsive titanium dioxide photocatalysts on pathogenic bacteria. *Arch Immunol Ther Exp (Warsz)* 2012; 60:267-75; PMID:22678625; <http://dx.doi.org/10.1007/s00005-012-0178-x>
83. Zhang H, Zhao H, Zhang S, Quan X. Photoelectrochemical manifestation of photoelectron transport properties of vertically aligned nanotubular TiO₂ photoanodes. *Chemphyschem* 2008; 9:117-23; PMID:18072232; <http://dx.doi.org/10.1002/cphc.200700406>
84. Harper JC, Christensen PA, Egerton TA. Effect of catalyst type on the kinetics of photoelectrical disinfection of water inoculated with *E. coli*. *J Appl Electrochem* 2000; 31:623-8; <http://dx.doi.org/10.1023/A:1017539328022>
85. Irie H, Watanabe Y, Hashimoto K. Carbon-doped Anatase TiO₂ Powders as a Visible-light Sensitive Photocatalyst. *Chem Lett* 2003; 32:772; <http://dx.doi.org/10.1246/cl.2003.772>
86. Rhee CH, Bae SW, Lee JS. Template-free Hydrothermal Synthesis of High Surface Area Nitrogen-doped Titania Photocatalyst Active under Visible Light. *Chem Lett* 2005; 34:660; <http://dx.doi.org/10.1246/cl.2005.660>
87. Lin L, Lin W, Zhu Y, Zhao B, Xie Y. Phosphor-doped Titania — a Novel Photocatalyst Active in Visible Light. *Chem Lett* 2005; 34:284; <http://dx.doi.org/10.1246/cl.2005.284>
88. Tokudome H, Miyauchi M. N-doped TiO₂ Nanotube with Visible Light Activity. *Chem Lett* 2004; 33:1108; <http://dx.doi.org/10.1246/cl.2004.1108>
89. Irie H, Miura S, Nakamura R, Hashimoto K. A Novel Visible-light-sensitive Efficient Photocatalyst, Cr(II)-grafted TiO₂. *Chem Lett* 2008; 37:252; <http://dx.doi.org/10.1246/cl.2008.252>
90. Li XZ, Li FB. Study of Au/Au(3+)-TiO₂ photocatalysts toward visible photooxidation for water and wastewater treatment. *Environ Sci Technol* 2001; 35:2381-7; PMID:11414049; <http://dx.doi.org/10.1021/es001752w>
91. Behar D, Rabani J. Kinetics of hydrogen production upon reduction of aqueous TiO₂ nanoparticles catalyzed by Pd(0), Pt(0), or Au(0) coatings and an unusual hydrogen abstraction; steady state and pulse radiolysis study. *J Phys Chem B* 2006; 110:8750-5; PMID:16640431; <http://dx.doi.org/10.1021/jp060971m>
92. Zeng Y, Wu W, Lee S, Gao J. Photocatalytic performance of plasma sprayed Pt-modified TiO₂ coatings under visible light irradiation. *Catal Commun* 2007; 8:906-12; <http://dx.doi.org/10.1016/j.catcom.2006.09.023>
93. Wang W, Zhang J, Chen F, He D, Anpo M. Preparation and photocatalytic properties of Fe³⁺-doped Ag@TiO₂ core-shell nanoparticles. *J Colloid Interface Sci* 2008; 323:182-6; PMID:18448112; <http://dx.doi.org/10.1016/j.jcis.2008.03.043>
94. You X, Chen F, Zhang J, Anpo M. A novel deposition precipitation method for preparation of Ag-loaded titanium dioxide. *Catal Lett* 2005; 102:247-50; <http://dx.doi.org/10.1007/s10562-005-5863-5>
95. Yu JC, Ho W, Yu J, Yip H, Wong PK, Zhao J. Efficient visible-light-induced photocatalytic disinfection on sulfur-doped nanocrystalline titania. *Environ Sci Technol* 2005; 39:1175-9; PMID:15773492; <http://dx.doi.org/10.1021/es035374h>
96. Mitoraj D, Jańczyk A, Strus M, Kisch H, Stochel G, Heczko PB, Macyk W. Visible light inactivation of bacteria and fungi by modified titanium dioxide. *Photochem Photobiol Sci* 2007; 6:642-8; PMID:17549266; <http://dx.doi.org/10.1039/b617043a>
97. Vohra A, Goswami DY, Deshpande DA, Block SS. Enhanced photocatalytic inactivation of bacterial spores on surfaces in air. *J Ind Microbiol Biotechnol* 2005; 32:364-70; PMID:16044291; <http://dx.doi.org/10.1007/s10295-005-0006-y>
98. Wong MS, Sun DS, Chang HH. Bactericidal performance of visible-light responsive titania photocatalyst with silver nanostructures. *PLoS One* 2010; 5:e10394; PMID:20454454; <http://dx.doi.org/10.1371/journal.pone.0010394>
99. Kau JH, Sun DS, Huang HH, Wong MS, Lin HC, Chang HH. Role of visible light-activated photocatalyst on the reduction of anthrax spore-induced mortality in mice. *PLoS One* 2009; 4:e4167; PMID:19132100; <http://dx.doi.org/10.1371/journal.pone.0004167>
100. Wong MS, Chu WC, Sun DS, Huang HS, Chen JH, Tsai PJ, Lin NT, Yu MS, Hsu SF, Wang SL, et al. Visible-light-induced bactericidal activity of a nitrogen-doped titanium photocatalyst against human pathogens. *Appl Environ Microbiol* 2006; 72:6111-6; PMID:16957236; <http://dx.doi.org/10.1128/AEM.02580-05>
101. Prasad GK, Agarwal GS, Singh B, Rai GP, Vijayaraghavan R. Photocatalytic inactivation of *Bacillus anthracis* by titania nanomaterials. *J Hazard Mater* 2009; 165:506-10; PMID:19056174; <http://dx.doi.org/10.1016/j.jhazmat.2008.10.009>
102. Prasad GK, Ramacharyulu PV, Merwyn S, Agarwal GS, Srivastava AR, Singh B, Rai GP, Vijayaraghavan R. Photocatalytic inactivation of spores of *Bacillus anthracis* using titania nanomaterials. *J Hazard Mater* 2011; 185:977-82; PMID:21035260; <http://dx.doi.org/10.1016/j.jhazmat.2010.10.001>

103. Roy AS, Parveen A, Koppalkar AR, Prasad M. Effect of Nano - Titanium Dioxide with Different Antibiotics against Methicillin-Resistant Staphylococcus Aureus. *J Biomater Nanobiotechnol* 2010; 1:37-41; <http://dx.doi.org/10.4236/jbnt.2010.11005>
104. Cheng CL, Sun DS, Chu WC, Tseng YH, Ho HC, Wang JB, Chung PH, Chen JH, Tsai PJ, Lin NT, et al. The effects of the bacterial interaction with visible-light responsive titania photocatalyst on the bactericidal performance. *J Biomed Sci* 2009; 16:7; PMID:19272171; <http://dx.doi.org/10.1186/1423-0127-16-7>
105. Tsai T-M, Chang H-H, Chang K-C, Liu Y-L, Tseng C-C. A comparative study of the bactericidal effect of photocatalytic oxidation by TiO₂ on antibiotic-resistant and antibiotic-sensitive bacteria. *J Chem Technol Biotechnol* 2010; 85:1642-53; <http://dx.doi.org/10.1002/jctb.2476>
106. Nhung TT, Nagata H, Takahashi A, Aihara M, Okamoto T, Shimohata T, Mawatari K, Akutagawa M, Kinouchi Y, Haraguchi M. Sterilization effect of UV light on Bacillus spores using TiO₂ films depends on wavelength. *J Med Invest* 2012; 59:53-8; PMID:22449993; <http://dx.doi.org/10.2152/jmi.59.53>
107. Knudson GB. Photoreactivation of ultraviolet-irradiated, plasmid-bearing, and plasmid-free strains of Bacillus anthracis. *Appl Environ Microbiol* 1986; 52:444-9; PMID:2429617
108. Baweja RB, Zaman MS, Mattoo AR, Sharma K, Tripathi V, Aggarwal A, Dubey GP, Kurupati RK, Ganguli M, Chaudhury NK, et al. Properties of Bacillus anthracis spores prepared under various environmental conditions. *Arch Microbiol* 2008; 189:71-9; PMID:17713759; <http://dx.doi.org/10.1007/s00203-007-0295-9>
109. Melski JW, Tanenbaum L, Parrish JA, Fitzpatrick TB, Bleich HL. Oral methoxsalen photochemotherapy for the treatment of psoriasis: a cooperative clinical trial. *J Invest Dermatol* 1977; 68:328-35; PMID:864273; <http://dx.doi.org/10.1111/1523-1747.ep12496022>
110. Salem SA, Barakat MA, Morcos CM. Bath psoralen+ultraviolet A photochemotherapy vs. narrow band-ultraviolet B in psoriasis: a comparison of clinical outcome and effect on circulating T-helper and T-suppressor/cytotoxic cells. *Photodermatol Photoimmunol Photomed* 2010; 26:235-42; PMID:20831697; <http://dx.doi.org/10.1111/j.1600-0781.2010.00525.x>
111. Cazenave JP, Waller C, Kientz D, Mendel I, Lin L, Jacquet M, Propst M, Liu W, Corash L, Sundin D, et al. An active hemovigilance program characterizing the safety profile of 7483 transfusions with plasma components prepared with amotosalen and UVA photochemical treatment. *Transfusion* 2010; 50:1210-9; PMID:20113450; <http://dx.doi.org/10.1111/j.1537-2995.2009.02579.x>
112. Dubensky TW Jr., Skoble J, Lauer P, Brockstedt DG. Killed but metabolically active vaccines. *Curr Opin Biotechnol* 2012; 23:917-23; PMID:22608846; <http://dx.doi.org/10.1016/j.copbio.2012.04.005>
113. Lai C, Cao H, Hearst JE, Corash L, Luo H, Wang Y. Quantitative analysis of DNA interstrand cross-links and monoadducts formed in human cells induced by psoralens and UVA irradiation. *Anal Chem* 2008; 80:8790-8; PMID:18947205; <http://dx.doi.org/10.1021/ac801520m>
114. Irsch J, Lin L. Pathogen Inactivation of Platelet and Plasma Blood Components for Transfusion Using the INTERCEPT Blood System™. *Transfus Med Hemother* 2011; 38:19-31; PMID:21779203; <http://dx.doi.org/10.1159/000323937>
115. Brockstedt DG, Bahjat KS, Giedlin MA, Liu W, Leong M, Luckett W, Gao Y, Schnupf P, Kapadia D, Castro G, et al. Killed but metabolically active microbes: a new vaccine paradigm for eliciting effector T-cell responses and protective immunity. *Nat Med* 2005; 11:853-60; PMID:16041382; <http://dx.doi.org/10.1038/nm1276>
116. Skoble J, Beaber JW, Gao Y, Lovchik JA, Sower LE, Liu W, Luckett W, Peterson JW, Calendar R, Portnoy DA, et al. Killed but metabolically active Bacillus anthracis vaccines induce broad and protective immunity against anthrax. *Infect Immun* 2009; 77:1649-63; PMID:19168734; <http://dx.doi.org/10.1128/IAI.00530-08>
117. Bruhn KW, Birnbaum R, Haskell J, Vanchinathan V, Greger S, Narayan R, Chang PL, Tran TA, Hickerson SM, Beverley SM, et al. Killed but metabolically active Leishmania infantum as a novel whole-cell vaccine for visceral leishmaniasis. *Clin Vaccine Immunol* 2012; 19:490-8; PMID:22323556; <http://dx.doi.org/10.1128/CVI.05660-11>
118. Harding AS, Schwab KJ. Using limes and synthetic psoralens to enhance solar disinfection of water (SODIS): a laboratory evaluation with norovirus, Escherichia coli, and MS2. *Am J Trop Med Hyg* 2012; 86:566-72; PMID:22492137; <http://dx.doi.org/10.4269/ajtmh.2012.11-0370>
119. Rasonglès P, Angelini-Tibert MF, Simon P, Currie C, Isola H, Kientz D, Slaedts M, Jacquet M, Sundin D, Lin L, et al. Transfusion of platelet components prepared with photochemical pathogen inactivation treatment during a Chikungunya virus epidemic in Ile de La Réunion. *Transfusion* 2009; 49:1083-91; PMID:19309473; <http://dx.doi.org/10.1111/j.1537-2995.2009.02111.x>
120. Sawyer L, Hanson D, Castro G, Luckett W, Dubensky TW Jr., Stassinopoulos A. Inactivation of parvovirus B19 in human platelet concentrates by treatment with amotosalen and ultraviolet A illumination. *Transfusion* 2007; 47:1062-70; PMID:17524098; <http://dx.doi.org/10.1111/j.1537-2995.2007.01237.x>
121. Pinna D, Sampson-Johannes A, Clementi M, Poli G, Rossini S, Lin L, Vicenzi E. Amotosalen photochemical inactivation of severe acute respiratory syndrome coronavirus in human platelet concentrates. *Transfus Med* 2005; 15:269-76; PMID:16101804; <http://dx.doi.org/10.1111/j.0958-7578.2005.00588.x>
122. McCullough J, Vesole DH, Benjamin RJ, Slichter SJ, Pineda A, Snyder E, Stadmauer EA, Lopez-Plaza I, Coutre S, Strauss RG, et al. Therapeutic efficacy and safety of platelets treated with a photochemical process for pathogen inactivation: the SPRINT Trial. *Blood* 2004; 104:1534-41; PMID:15138160; <http://dx.doi.org/10.1182/blood-2003-12-4443>
123. Maves RC, Oré RM, Porter KR, Kochel TJ. Immunogenicity and protective efficacy of a psoralen-inactivated dengue-1 virus vaccine candidate in Aotus nancymae monkeys. *Vaccine* 2011; 29:2691-6; PMID:21303709; <http://dx.doi.org/10.1016/j.vaccine.2011.01.077>
124. Maves RC, Castillo Oré RM, Porter KR, Kochel TJ. Immunogenicity of a psoralen-inactivated dengue virus type 1 vaccine candidate in mice. *Clin Vaccine Immunol* 2010; 17:304-6; PMID:20007362; <http://dx.doi.org/10.1128/CVI.00353-09>
125. Nair CN, Davis R. Photochemical inhibition of poliovirus replication by 4',5',8-trimethylpsoralen plus light. *Intervirology* 1978; 9:65-75; PMID:201583; <http://dx.doi.org/10.1159/000148924>
126. Stern RS; PUVA Follow-Up Study. The risk of squamous cell and basal cell cancer associated with psoralen and ultraviolet A therapy: a 30-year prospective study. *J Am Acad Dermatol* 2012; 66:553-62; PMID:22264671; <http://dx.doi.org/10.1016/j.jaad.2011.04.004>
127. Rangarajan HG, Ponzalzan RC, Camitta BM, Talano JA. The use of novel Therakos™ Cellex® for extracorporeal photopheresis in treatment of graft-versus-host disease in paediatric patients. *Br J Haematol* 2013; PMID:23961954; <http://dx.doi.org/10.1111/bjh.12535>
128. Kleinpenning MM, Smits T, Frunt MH, van Erp PE, van de Kerkhof PC, Gerritsen RM. Clinical and histological effects of blue light on normal skin. *Photodermatol Photoimmunol Photomed* 2010; 26:16-21; PMID:20070834; <http://dx.doi.org/10.1111/j.1600-0781.2009.00474.x>
129. Liebmann J, Born M, Kolb-Bachofen V. Blue-light irradiation regulates proliferation and differentiation in human skin cells. *J Invest Dermatol* 2010; 130:259-69; PMID:19675580; <http://dx.doi.org/10.1038/jid.2009.194>
130. Dai T, Gupta A, Murray CK, Vrahas MS, Tegos GP, Hamblin MR. Blue light for infectious diseases: Propionibacterium acnes, Helicobacter pylori, and beyond? *Drug Resist Updat* 2012; 15:223-36; PMID:22846406; <http://dx.doi.org/10.1016/j.drup.2012.07.001>
131. Helgason E, Okstad OA, Caugant DA, Johansen HA, Fouet A, Mock M, Hegna I, Kolstø AB. Bacillus anthracis, Bacillus cereus, and Bacillus thuringiensis—one species on the basis of genetic evidence. *Appl Environ Microbiol* 2000; 66:2627-30; PMID:10831447; <http://dx.doi.org/10.1128/AEM.66.6.2627-2630.2000>
132. Sagripanti JL, Carrera M, Insalaco J, Ziemski M, Rogers J, Zandomeni R. Virulent spores of Bacillus anthracis and other Bacillus species deposited on solid surfaces have similar sensitivity to chemical decontaminants. *J Appl Microbiol* 2007; 102:11-21; PMID:17184315; <http://dx.doi.org/10.1111/j.1365-2672.2006.03235.x>
133. Bertoloni G, Lauro FM, Cortella G, Merchat M. Photosensitizing activity of hematoporphyrin on Staphylococcus aureus cells. *Biochim Biophys Acta* 2000; 1475:169-74; PMID:10832032; [http://dx.doi.org/10.1016/S0304-4165\(00\)00071-4](http://dx.doi.org/10.1016/S0304-4165(00)00071-4)
134. Eraso AJ, Albessa I. Elevation of alanine amino transferase and aspartate amino transferase produced by pyoverdinin, a photolabile pigment of Pseudomonas fluorescens. *Nat Toxins* 1998; 6:61-5; PMID:9888631; [http://dx.doi.org/10.1002/\(SICI\)1522-7189\(199804\)6:2<61::AID-NT5>3.0.CO;2-H](http://dx.doi.org/10.1002/(SICI)1522-7189(199804)6:2<61::AID-NT5>3.0.CO;2-H)
135. Dai T, Gupta A, Huang YY, Sherwood ME, Murray CK, Vrahas MS, Kielian T, Hamblin MR. Blue Light Eliminates Community-Acquired Methicillin-resistant Staphylococcus aureus in Infected Mouse Skin Abrasions. *Photomed Laser Surg* 2013; PMID:23406384; <http://dx.doi.org/10.1089/pho.2012.3365>
136. Ganz RA, Viveiros J, Ahmad A, Ahmadi A, Khalil A, Tolkoft MJ, Nishioka NS, Hamblin MR. Helicobacter pylori in patients can be killed by visible light. *Lasers Surg Med* 2005; 36:260-5; PMID:15791671; <http://dx.doi.org/10.1002/lsm.20161>
137. Lembo AJ, Ganz RA, Sheth S, Cave D, Kelly C, Levin P, Kazlas PT, Baldwin PC 3rd, Lindmark WR, McGrath JR, et al. Treatment of Helicobacter pylori infection with intra-gastric violet light phototherapy: a pilot clinical trial. *Lasers Surg Med* 2009; 41:337-44; PMID:19533762; <http://dx.doi.org/10.1002/lsm.20770>
138. Enwemeka CS, Williams D, Enwemeka SK, Hollosi S, Yens D. Blue 470-nm light kills methicillin-resistant Staphylococcus aureus (MRSA) in vitro. *Photomed Laser Surg* 2009; 27:221-6; PMID:19196103; <http://dx.doi.org/10.1089/pho.2008.2413>
139. Enwemeka CS, Williams D, Hollosi S, Yens D, Enwemeka SK. Visible 405 nm SLD light photo-destroys methicillin-resistant Staphylococcus aureus (MRSA) in vitro. *Lasers Surg Med* 2008; 40:734-7; PMID:19065556; <http://dx.doi.org/10.1002/lsm.20724>
140. Guffey JS, Wilborn J. In vitro bactericidal effects of 405-nm and 470-nm blue light. *Photomed Laser Surg* 2006; 24:684-8; PMID:17199466; <http://dx.doi.org/10.1089/pho.2006.24.684>

141. Maclean M, MacGregor SJ, Anderson JG, Woolsey G. Inactivation of bacterial pathogens following exposure to light from a 405-nanometer light-emitting diode array. *Appl Environ Microbiol* 2009; 75:1932-7; PMID:19201962; <http://dx.doi.org/10.1128/AEM.01892-08>
142. Bush K, Courvalin P, Dantas G, Davies J, Eisenstein B, Huovinen P, Jacoby GA, Kishony R, Kreiswirth BN, Kutter E, et al. Tackling antibiotic resistance. *Nat Rev Microbiol* 2011; 9:894-6; PMID:22048738; <http://dx.doi.org/10.1038/nrmicro2693>
143. Nicholson WL, Munakata N, Horneck G, Melosh HJ, Setlow P. Resistance of *Bacillus* endospores to extreme terrestrial and extraterrestrial environments. *Microbiol Mol Biol Rev* 2000; 64:548-72; PMID:10974126; <http://dx.doi.org/10.1128/MMBR.64.3.548-572.2000>
144. Maclean M, Murdoch LE, MacGregor SJ, Anderson JG. Sporicidal effects of high-intensity 405 nm visible light on endospore-forming bacteria. *Photochem Photobiol* 2013; 89:120-6; PMID:22803813; <http://dx.doi.org/10.1111/j.1751-1097.2012.01202.x>
145. Gomelsky M, Hoff WD. Light helps bacteria make important lifestyle decisions. *Trends Microbiol* 2011; 19:441-8; PMID:21664820; <http://dx.doi.org/10.1016/j.tim.2011.05.002>
146. Bache SE, Maclean M, MacGregor SJ, Anderson JG, Gettinby G, Coia JE, et al. Clinical studies of the High-Intensity Narrow-Spectrum light Environmental Decontamination System (HINS-light EDS), for continuous disinfection in the burn unit inpatient and outpatient settings. *Burns: journal of the International Society for Burn Injuries* 2012; 38:69-76.
147. Maclean M, Macgregor SJ, Anderson JG, Woolsey GA, Coia JE, Hamilton K, Taggart I, Watson SB, Thakker B, Gettinby G. Environmental decontamination of a hospital isolation room using high-intensity narrow-spectrum light. *J Hosp Infect* 2010; 76:247-51; PMID:20864210; <http://dx.doi.org/10.1016/j.jhin.2010.07.010>
148. Enwemeka CS. Therapeutic blue light: a different ray of light on an age-old problem. *Photomed Laser Surg* 2006; 24:679; PMID:17199464; <http://dx.doi.org/10.1089/pho.2006.24.679>
149. Bumah VV, Masson-Meyers DS, Cashin SE, Enwemeka CS. Wavelength and Bacterial Density Influence the Bactericidal Effect of Blue Light on Methicillin-Resistant *Staphylococcus aureus* (MRSA). *Photomed Laser Surg* 2013; PMID:23621894; <http://dx.doi.org/10.1089/pho.2012.3461>
150. Maclean M, MacGregor SJ, Anderson JG, Woolsey G. High-intensity narrow-spectrum light inactivation and wavelength sensitivity of *Staphylococcus aureus*. *FEMS Microbiol Lett* 2008; 285:227-32; PMID:18557942; <http://dx.doi.org/10.1111/j.1574-6968.2008.01233.x>
151. Dai T, Tegos GP, Zhiyentayev T, Mylonakis E, Hamblin MR. Photodynamic therapy for methicillin-resistant *Staphylococcus aureus* infection in a mouse skin abrasion model. *Lasers Surg Med* 2010; 42:38-44; PMID:20077489; <http://dx.doi.org/10.1002/lsm.20887>
152. St Denis TG, Dai T, Izikson L, Astrakas C, Anderson RR, Hamblin MR, Tegos GP. All you need is light: antimicrobial photoinactivation as an evolving and emerging discovery strategy against infectious disease. *Virulence* 2011; 2:509-20; PMID:21971183; <http://dx.doi.org/10.4161/viru.2.6.17889>
153. Ragàs X, Dai T, Tegos GP, Agut M, Nonell S, Hamblin MR. Photodynamic inactivation of *Acinetobacter baumannii* using phenothiazinium dyes: in vitro and in vivo studies. *Lasers Surg Med* 2010; 42:384-90; PMID:20583252; <http://dx.doi.org/10.1002/lsm.20922>
154. Song HH, Lee JK, Um HS, Chang BS, Lee SY, Lee MK. Phototoxic effect of blue light on the planktonic and biofilm state of anaerobic periodontal pathogens. *J Periodontol Implant Sci* 2013; 43:72-8; PMID:23678390; <http://dx.doi.org/10.5051/jpis.2013.43.2.72>
155. Chang YS, Hwang JH, Kwon HN, Choi CW, Ko SY, Park WS, Shin SM, Lee M. In vitro and in vivo efficacy of new blue light emitting diode phototherapy compared to conventional halogen quartz phototherapy for neonatal jaundice. *J Korean Med Sci* 2005; 20:61-4; PMID:15716604; <http://dx.doi.org/10.3346/jkms.2005.20.1.61>
156. Malik Z, Ladan H, Nitzan Y. Photodynamic inactivation of Gram-negative bacteria: problems and possible solutions. *J Photochem Photobiol B* 1992; 14:262-6; PMID:1432395; [http://dx.doi.org/10.1016/1011-1344\(92\)85104-3](http://dx.doi.org/10.1016/1011-1344(92)85104-3)
157. Sharma SK, Dai T, Kharkwal GB, Huang YY, Huang L, De Arce VJ, Tegos GP, Hamblin MR. Drug discovery of antimicrobial photosensitizers using animal models. *Curr Pharm Des* 2011; 17:1303-19; PMID:21504410; <http://dx.doi.org/10.2174/138161211795703735>
158. Minnock A, Vernon DI, Schofield J, Griffiths J, Parish JH, Brown ST. Photoinactivation of bacteria. Use of a cationic water-soluble zinc phthalocyanine to photoinactivate both gram-negative and gram-positive bacteria. *J Photochem Photobiol B* 1996; 32:159-64; PMID:8622179; [http://dx.doi.org/10.1016/1011-1344\(95\)07148-2](http://dx.doi.org/10.1016/1011-1344(95)07148-2)
159. Hamblin MR, Hasan T. Photodynamic therapy: a new antimicrobial approach to infectious disease? *Photochem Photobiol Sci* 2004; 3:436-50; PMID:15122361; <http://dx.doi.org/10.1039/b311900a>
160. Klevens RM, Morrison MA, Nadle J, Petit S, Gershman K, Ray S, Harrison LH, Lynfield R, Dumyati G, Townes JM, et al. Active Bacterial Core surveillance (ABCs) MRSA Investigators. Invasive methicillin-resistant *Staphylococcus aureus* infections in the United States. *JAMA* 2007; 298:1763-71; PMID:17940231; <http://dx.doi.org/10.1001/jama.298.15.1763>
161. Baba T, Takeuchi F, Kuroda M, Yuzawa H, Aoki K, Oguchi A, Nagai Y, Iwama N, Asano K, Naimi T, et al. Genome and virulence determinants of high virulence community-acquired MRSA. *Lancet* 2002; 359:1819-27; PMID:12044378; [http://dx.doi.org/10.1016/S0140-6736\(02\)08713-5](http://dx.doi.org/10.1016/S0140-6736(02)08713-5)
162. Gordon RJ, Lowy FD. Pathogenesis of methicillin-resistant *Staphylococcus aureus* infection. *Clin Infect Dis* 2008; 46(Suppl 5):S350-9; PMID:18462090; <http://dx.doi.org/10.1086/533591>
163. Miko BA, Hafer CA, Lee CJ, Sullivan SB, Hackel MA, Johnson BM, Whittier S, Della-Latta P, Uhlemann AC, Lowy FD. Molecular characterization of methicillin-susceptible *Staphylococcus aureus* clinical isolates in the United States, 2004 to 2010. *J Clin Microbiol* 2013; 51:874-9; PMID:23284029; <http://dx.doi.org/10.1128/JCM.00923-12>
164. Lowy FD. How *Staphylococcus aureus* adapts to its host. *N Engl J Med* 2011; 364:1987-90; PMID:21612466; <http://dx.doi.org/10.1056/NEJMp1100251>
165. Maisch T, Bosl C, Szeimies RM, Lehn N, Abels C. Photodynamic effects of novel XF porphyrin derivatives on prokaryotic and eukaryotic cells. *Antimicrob Agents Chemother* 2005; 49:1542-52; PMID:15793136; <http://dx.doi.org/10.1128/AAC.49.4.1542-1552.2005>
166. Wainwright M, Phoenix DA, Laycock SL, Wareing DR, Wright PA. Photobactericidal activity of phenothiazinium dyes against methicillin-resistant strains of *Staphylococcus aureus*. *FEMS Microbiol Lett* 1998; 160:177-81; PMID:9532735; <http://dx.doi.org/10.1111/j.1574-6968.1998.tb12908.x>
167. Fu XJ, Fang Y, Yao M. Antimicrobial photodynamic therapy for methicillin-resistant *Staphylococcus aureus* infection. *Biomed Res Int* 2013; 2013:159157; PMID:23555074; <http://dx.doi.org/10.1155/2013/159157>
168. Hajim KI, Salih DS, Rassam YZ. Laser light combined with a photosensitizer may eliminate methicillin-resistant strains of *Staphylococcus aureus*. *Lasers Med Sci* 2010; 25:743-8; PMID:20552385; <http://dx.doi.org/10.1007/s10103-010-0803-z>
169. Tubby S, Wilson M, Nair SP. Inactivation of staphylococcal virulence factors using a light-activated antimicrobial agent. *BMC Microbiol* 2009; 9:211; PMID:19804627; <http://dx.doi.org/10.1186/1471-2180-9-211>
170. Wilson M, Pratten J. Lethal photosensitisation of *Staphylococcus aureus* in vitro: effect of growth phase, serum, and pre-irradiation time. *Lasers Surg Med* 1995; 16:272-6; PMID:7791501; <http://dx.doi.org/10.1002/lsm.1900160309>
171. Street C, Pedigo L, Gibbs A, Loebel N. Antimicrobial photodynamic therapy for the decolonization of methicillin-resistant *Staphylococcus aureus* from the anterior nares. 12th World Congress of the International Photodynamic Association: International Society for Optics and Photonics, 2009:73803B-B-16.
172. Maisch T, Bosl C, Szeimies R-M, Love B, Abels C. Determination of the antibacterial efficacy of a new porphyrin-based photosensitizer against MRSA ex vivo. *Photochem Photobiol Sci* 2007; 6:545-51; PMID:17487307; <http://dx.doi.org/10.1039/b614770d>
173. Vecchio D, Dai T, Huang L, Fantetti L, Roncucci G, Hamblin MR. Antimicrobial photodynamic therapy with RLP068 kills methicillin-resistant *Staphylococcus aureus* and improves wound healing in a mouse model of infected skin abrasion PDT with RLP068/Cl in infected mouse skin abrasion. *J Biophotonics* 2013; 6:733-42; PMID:22987338; <http://dx.doi.org/10.1002/jbio.201200121>
174. Griffiths MA, Wren BW, Wilson M. Killing of methicillin-resistant *Staphylococcus aureus* in vitro using aluminium disulphonated phthalocyanine, a light-activated antimicrobial agent. *J Antimicrob Chemother* 1997; 40:873-6; PMID:9462440; <http://dx.doi.org/10.1093/jac/40.6.873>
175. Dai T, Tegos GP, Zhiyentayev T, Mylonakis E, Hamblin MR. Photodynamic therapy for methicillin-resistant *Staphylococcus aureus* infection in a mouse skin abrasion model. *Lasers Surg Med* 2010; 42:38-44; PMID:20077489; <http://dx.doi.org/10.1002/lsm.20887>
176. Zolfaghari PS, Packer S, Singer M, Nair SP, Bennett J, Street C, Wilson M. In vivo killing of *Staphylococcus aureus* using a light-activated antimicrobial agent. *BMC Microbiol* 2009; 9:27; PMID:19193212; <http://dx.doi.org/10.1186/1471-2180-9-27>
177. Tsai T, Yang Y-T, Wang T-H, Chien H-F, Chen C-T. Improved photodynamic inactivation of gram-positive bacteria using hematoporphyrin encapsulated in liposomes and micelles. *Lasers Surg Med* 2009; 41:316-22; PMID:19347938; <http://dx.doi.org/10.1002/lsm.20754>
178. Cohen-Poradosu R, Kasper DL. Group A streptococcus epidemiology and vaccine implications. *Clin Infect Dis* 2007; 45:863-5; PMID:17806050; <http://dx.doi.org/10.1086/521263>
179. Omar GS, Wilson M, Nair SP. Lethal photosensitization of wound-associated microbes using indocyanine green and near-infrared light. *BMC Microbiol* 2008; 8:111; PMID:18593460; <http://dx.doi.org/10.1186/1471-2180-8-111>
180. Hope CK, Wilson M. Induction of lethal photosensitization in biofilms using a confocal scanning laser as the excitation source. *J Antimicrob Chemother* 2006; 57:1227-30; PMID:16549510; <http://dx.doi.org/10.1093/jac/dkl096>

181. Ulianova O, Ulianov S, Li P, Luo Q. Estimation of reactogenicity of preparations produced on the basis of photoactivated live vaccines against brucellosis and tularaemia on the organismic level. 1. Using the LASCA method. *Quantum Electron* 2011; 41:340-3; <http://dx.doi.org/10.1070/QE2011v041n04ABEH014598>
182. Wainwright M, Phoenix DA, Smillie TE, Wareing DR. Phenothiaziniums as putative photobactericidal agents for red blood cell concentrates. *J Chemother* 2001; 13:503-9; PMID:11760214
183. Demidova TN, Hamblin MR. Photodynamic therapy targeted to pathogens. *Int J Immunopathol Pharmacol* 2004; 17:245-54; PMID:15461858
184. Dai T, Huang YY, Hamblin MR. Photodynamic therapy for localized infections-State of the art. *Photodiagn Photodyn Ther* 2009; 6:170-88; <http://dx.doi.org/10.1016/j.pdpdt.2009.10.008>
185. Hamblin MR, O'Donnell DA, Murthy N, Contag CH, Hasan T. Rapid control of wound infections by targeted photodynamic therapy monitored by in vivo bioluminescence imaging. *Photochem Photobiol* 2002; 75:51-7; PMID:11837327; [http://dx.doi.org/10.1562/0031-8655\(2002\)075<0051:RCOWIB>2.0.CO;2](http://dx.doi.org/10.1562/0031-8655(2002)075<0051:RCOWIB>2.0.CO;2)
186. Hamblin MR, Zahra T, Contag CH, McManus AT, Hasan T. Optical monitoring and treatment of potentially lethal wound infections in vivo. *J Infect Dis* 2003; 187:1717-25; PMID:12751029; <http://dx.doi.org/10.1086/375244>
187. Wong TW, Wang YY, Sheu HM, Chuang YC. Bactericidal effects of toluidine blue-mediated photodynamic action on *Vibrio vulnificus*. *Antimicrob Agents Chemother* 2005; 49:895-902; PMID:15728881; <http://dx.doi.org/10.1128/AAC.49.3.895-902.2005>
188. Lambrechts SA, Demidova TN, Aalders MC, Hasan T, Hamblin MR. Photodynamic therapy for *Staphylococcus aureus* infected burn wounds in mice. *Photochem Photobiol Sci* 2005; 4:503-9; PMID:15986057; <http://dx.doi.org/10.1039/b502125a>
189. Ragàs X, Dai T, Tegos GP, Agut M, Nonell S, Hamblin MR. Photodynamic inactivation of *Acinetobacter baumannii* using phenothiazinium dyes: in vitro and in vivo studies. *Lasers Surg Med* 2010; 42:384-90; PMID:20583252; <http://dx.doi.org/10.1002/lsm.20922>
190. Dai T, Tegos GP, Lu Z, Huang L, Zhiyentayev T, Franklin MJ, Baer DG, Hamblin MR. Photodynamic therapy for *Acinetobacter baumannii* burn infections in mice. *Antimicrob Agents Chemother* 2009; 53:3929-34; PMID:19564369; <http://dx.doi.org/10.1128/AAC.00027-09>
191. Gad F, Zahra T, Francis KP, Hasan T, Hamblin MR. Targeted photodynamic therapy of established soft-tissue infections in mice. *Photochem Photobiol Sci* 2004; 3:451-8; PMID:15122362; <http://dx.doi.org/10.1039/b311901g>
192. Spotts Whitney EA, Beatty ME, Taylor TH Jr, Weyant R, Sobel J, Arduino MJ, Ashford DA. Inactivation of *Bacillus anthracis* spores. *Emerg Infect Dis* 2003; 9:623-7; PMID:12780999; <http://dx.doi.org/10.3201/eid0906.020377>
193. Cohen M, Whalen T. Implications of low level human exposure to respirable *B. anthracis*. *Applied Biosafety* 2007; 12:109
194. Lacy DB, Collier RJ. Structure and function of anthrax toxin. *Curr Top Microbiol Immunol* 2002; 271:61-85; PMID:12224524; http://dx.doi.org/10.1007/978-3-662-05767-4_4
195. Demidova TN, Hamblin MR. Photodynamic inactivation of *Bacillus* spores, mediated by phenothiazinium dyes. *Appl Environ Microbiol* 2005; 71:6918-25; PMID:16269726; <http://dx.doi.org/10.1128/AEM.71.11.6918-6925.2005>
196. Greenberg D, Busch J, Keim P, Wagner D. Identifying experimental surrogates for *Bacillus anthracis* spores: a review. *Invest Genet* 2010; 1.
197. Helgason E, Okstad OA, Caugant DA, Johansen HA, Fouet A, Mock M, Hegna I, Kolstø AB. *Bacillus anthracis*, *Bacillus cereus*, and *Bacillus thuringiensis*--one species on the basis of genetic evidence. *Appl Environ Microbiol* 2000; 66:2627-30; PMID:10831447; <http://dx.doi.org/10.1128/AEM.66.6.2627-2630.2000>
198. Oliveira A, Almeida A, Carvalho CM, Tomé JP, Faustino MA, Neves MG, Tomé AC, Cavaleiro JA, Cunha A. Porphyrin derivatives as photosensitizers for the inactivation of *Bacillus cereus* endospores. *J Appl Microbiol* 2009; 106:1986-95; PMID:19228253; <http://dx.doi.org/10.1111/j.1365-2672.2009.04168.x>
199. Maisch T, Spannberger F, Regensburger J, Felgenträger A, Bäumler W. Fast and effective: intense pulse light photodynamic inactivation of bacteria. *J Ind Microbiol Biotechnol* 2012; 39:1013-21; PMID:22354734; <http://dx.doi.org/10.1007/s10295-012-1103-3>
200. Morison WL. Variations of PUVa: practical and effective? *Arch Dermatol* 1998; 134:1286-8; PMID:9801689; <http://dx.doi.org/10.1001/archderm.134.10.1286>
201. Dixon DM. *Coccidioides immitis* as a Select Agent of bioterrorism. *J Appl Microbiol* 2001; 91:602-5; PMID:11576294; <http://dx.doi.org/10.1046/j.1365-2672.2001.01496.x>
202. Cox RA. Antigenic structure of *Coccidioides immitis*. *Immunol Ser* 1989; 47:133-70; PMID:2490076
203. Paardekooper M, De Bruijne AW, Van Steveninck J, Van den Broek PJ. Intracellular damage in yeast cells caused by photodynamic treatment with toluidine blue. *Photochem Photobiol* 1995; 61:84-9; PMID:7899497; <http://dx.doi.org/10.1111/j.1751-1097.1995.tb09247.x>
204. Chabrier-Roselló Y, Foster TH, Pérez-Nazario N, Mitra S, Haidaris CG. Sensitivity of *Candida albicans* germ tubes and biofilms to photofrin-mediated phototoxicity. *Antimicrob Agents Chemother* 2005; 49:4288-95; PMID:16189110; <http://dx.doi.org/10.1128/AAC.49.10.4288-4295.2005>
205. Lambrechts SA, Aalders MC, Van Marle J. Mechanistic study of the photodynamic inactivation of *Candida albicans* by a cationic porphyrin. *Antimicrob Agents Chemother* 2005; 49:2026-34; PMID:15855528; <http://dx.doi.org/10.1128/AAC.49.5.2026-2034.2005>
206. Smijs TG, van der Haas RN, Lugtenburg J, Liu Y, de Jong RL, Schuitmaker HJ. Photodynamic treatment of the dermatophyte *Trichophyton rubrum* and its microconidia with porphyrin photosensitizers. *Photochem Photobiol* 2004; 80:197-202; PMID:15244503; <http://dx.doi.org/10.1562/2004-04-22-RA-146.1>
207. Friedberg JS, Skema C, Baum ED, Burdick J, Vinogradov SA, Wilson DF, Horan AD, Nachamkin I. In vitro effects of photodynamic therapy on *Aspergillus fumigatus*. *J Antimicrob Chemother* 2001; 48:105-7; PMID:11418518; <http://dx.doi.org/10.1093/jac/48.1.105>
208. Junqueira JC, Jorge AO, Barbosa JO, Rossoni RD, Vilela SF, Costa AC, Primo FL, Gonçalves JM, Tedesco AC, Suleiman JM. Photodynamic inactivation of biofilms formed by *Candida* spp., *Trichosporon mucoides*, and *Kodamaea ohmeri* by cationic nano-emulsion of zinc 2,9,16,23-tetrakis(phenylthio)-29H, 31H-phthalocyanine (ZnPc). *Lasers Med Sci* 2012; 27:1205-12; PMID:22278349; <http://dx.doi.org/10.1007/s10103-012-1050-2>
209. Wainwright M. Local treatment of viral disease using photodynamic therapy. *Int J Antimicrob Agents* 2003; 21:510-20; PMID:12791463; [http://dx.doi.org/10.1016/S0924-8579\(03\)00035-9](http://dx.doi.org/10.1016/S0924-8579(03)00035-9)
210. Müller-Breitkreutz K, Mohr H, Briviba K, Sies H. Inactivation of viruses by chemically and photochemically generated singlet molecular oxygen. *J Photochem Photobiol B* 1995; 30:63-70; PMID:8558363; [http://dx.doi.org/10.1016/1011-1344\(95\)07150-Z](http://dx.doi.org/10.1016/1011-1344(95)07150-Z)
211. Müller-Breitkreutz K, Mohr H. Infection cycle of herpes viruses after photodynamic treatment with methylene blue and light. *Beitr Infusionsther Transfusionsmed* 1997; 34:37-42; PMID:9356656
212. Smetana Z, Ben-Hur E, Mendelson E, Salzberg S, Wagner P, Malik Z. Herpes simplex virus proteins are damaged following photodynamic inactivation with phthalocyanines. *J Photochem Photobiol B* 1998; 44:77-83; PMID:9745730; [http://dx.doi.org/10.1016/S1011-1344\(98\)00124-9](http://dx.doi.org/10.1016/S1011-1344(98)00124-9)
213. Wong T-W, Huang H-J, Wang Y-F, Lee Y-P, Huang C-C, Yu C-K. Methylene blue-mediated photodynamic inactivation as a novel disinfectant of enterovirus 71. *J Antimicrob Chemother* 2010; 65:2176-82; PMID:20719762; <http://dx.doi.org/10.1093/jac/dkq301>
214. Costa L, Faustino MA, Neves MG, Cunha A, Almeida A. Photodynamic inactivation of mammalian viruses and bacteriophages. *Viruses* 2012; 4:1034-74; PMID:22852040; <http://dx.doi.org/10.3390/v4071034>
215. Matthews JL, Newman JT, Sogandares-Bernal F, Judy MM, Skiles H, Leveson JE, Marengo-Rowe AJ, Chanh TC. Photodynamic therapy of viral contaminants with potential for blood banking applications. *Transfusion* 1988; 28:81-3; PMID:2829396; <http://dx.doi.org/10.1046/j.1537-2995.1988.2818127963.x>
216. Bray M. Defense against filoviruses used as biological weapons. *Antiviral Res* 2003; 57:53-60; PMID:12615303; [http://dx.doi.org/10.1016/S0166-3542\(02\)00200-0](http://dx.doi.org/10.1016/S0166-3542(02)00200-0)
217. Käsermann F, Kempf C. Photodynamic inactivation of enveloped viruses by buckminsterfullerene. *Antiviral Res* 1997; 34:65-70; PMID:9107386; [http://dx.doi.org/10.1016/S0166-3542\(96\)01207-7](http://dx.doi.org/10.1016/S0166-3542(96)01207-7)
218. Hirayama J, Abe H, Kamo N, Shinbo T, Ohnishi-Yamada Y, Kurosawa S, Ikebuchi K, Sekiguchi S. Photoinactivation of vesicular stomatitis virus with fullerene conjugated with methoxy polyethylene glycol amine. *Biol Pharm Bull* 1999; 22:1106-9; PMID:10549864; <http://dx.doi.org/10.1248/bpb.22.1106>
219. Frischknecht F. The history of biological warfare. Human experimentation, modern nightmares and lone madmen in the twentieth century. *EMBO Rep* 2003; 4(Spec No):S47-52; PMID:12789407; <http://dx.doi.org/10.1038/sj.embor.embor849>
220. Whitby M, Street AC, Ruff TA, Fenner F. Biological agents as weapons 1: smallpox and botulism. *Med J Aust* 2002; 176:431-3; PMID:12056996
221. Nulens E, Voss A. Laboratory diagnosis and biosafety issues of biological warfare agents. *Clin Microbiol Infect* 2002; 8:455-66; PMID:12789868; <http://dx.doi.org/10.1046/j.1469-0691.2002.00528.x>
222. Bachmann B, Knüver-Hopf J, Lambrecht B, Mohr H. Target structures for HIV-1 inactivation by methylene blue and light. *J Med Virol* 1995; 47:172-8; PMID:8830122; <http://dx.doi.org/10.1002/jmv.1890470211>
223. Schagen FH, Moor AC, Cheong SC, Cramer SJ, van Ormondt H, van der Eb AJ, Dubbelman TM, Hoebe RC. Photodynamic treatment of adenoviral vectors with visible light: an easy and convenient method for viral inactivation. *Gene Ther* 1999; 6:873-81; PMID:10505113; <http://dx.doi.org/10.1038/sj.gt.3300897>
224. Abe H, Wagner SJ. Analysis of viral DNA, protein and envelope damage after methylene blue, phthalocyanine derivative or merocyanine 540 photosensitization. *Photochem Photobiol* 1995; 61:402-9; PMID:7740085; <http://dx.doi.org/10.1111/j.1751-1097.1995.tb08630.x>
225. Zupán K, Egyeki M, Tóth K, Fekete A, Herényi L, Módos K, Csik G. Comparison of the efficiency and the specificity of DNA-bound and free cationic porphyrin in photodynamic virus inactivation. *J Photochem Photobiol B* 2008; 90:105-12; PMID:18222092; <http://dx.doi.org/10.1016/j.jphotobiol.2007.11.007>

226. Caminos D, Durantini E. Interaction and photodynamic activity of cationic porphyrin derivatives bearing different patterns of charge distribution with GMP and DNA. *J Photochem Photobiol Chem* 2008; 198:274-81; <http://dx.doi.org/10.1016/j.jphotochem.2008.04.005>
227. Huang Q, Fu WL, Chen B, Huang JF, Zhang X, Xue Q. Inactivation of dengue virus by methylene blue/narrow bandwidth light system. *J Photochem Photobiol B* 2004; 77:39-43; PMID:15542360
228. Lin Y-L, Lei H-Y, Wen Y-Y, Luh T-Y, Chou C-K, Liu H-S. Light-independent inactivation of dengue-2 virus by carboxyfullerene C3 isomer. *Virology* 2000; 275:258-62; PMID:10998325; <http://dx.doi.org/10.1006/viro.2000.0490>
229. Kömerik N, Wilson M, Poole S. The effect of photodynamic action on two virulence factors of gram-negative bacteria. *Photochem Photobiol* 2000; 72:676-80; PMID:11107854; [http://dx.doi.org/10.1562/0031-8655\(2000\)072<0676:TEOPAO>2.0.CO;2](http://dx.doi.org/10.1562/0031-8655(2000)072<0676:TEOPAO>2.0.CO;2)
230. Giannelli M, Pini A, Formigli L, Bani D. Comparative in vitro study among the effects of different laser and LED irradiation protocols and conventional chlorhexidine treatment for deactivation of bacterial lipopolysaccharide adherent to titanium surface. *Photomed Laser Surg* 2011; 29:573-80; PMID:21438842; <http://dx.doi.org/10.1089/pho.2010.2958>
231. Tubby S, Wilson M, Nair SP. Inactivation of staphylococcal virulence factors using a light-activated antimicrobial agent. *BMC Microbiol* 2009; 9:211; PMID:19804627; <http://dx.doi.org/10.1186/1471-2180-9-211>
232. Eubanks LM, Dickerson TJ, Janda KD. Vitamin B2-mediated cellular photoinhibition of botulinum neurotoxin A. *FEBS Lett* 2005; 579:5361-4; PMID:16198354; <http://dx.doi.org/10.1016/j.febslet.2005.08.072>
233. Shen N. *Photodisruption in biological tissues using femtosecond laser pulses* Harvard University Press, Cambridge, 2003.
234. Tsen KT, Tsen SW, Chang CL, Hung CF, Wu TC, Kiang JG. Inactivation of viruses by coherent excitations with a low power visible femtosecond laser. *Virology* 2007; 4:50; PMID:17550590; <http://dx.doi.org/10.1186/1743-422X-4-50>
235. Tsen KT, Tsen SW, Chang CL, Hung CF, Wu TC, Kiang JG. Inactivation of viruses by laser-driven coherent excitations via impulsive stimulated Raman scattering process. *J Biomed Opt* 2007; 12:064030; PMID:18163846; <http://dx.doi.org/10.1117/1.2821713>
236. Tsen KT, Tsen SW, Fu Q, Lindsay SM, Kibler K, Jacobs B, Wu TC, Karanam B, Jagu S, Roden RB, et al. Photonic approach to the selective inactivation of viruses with a near-infrared subpicosecond fiber laser. *J Biomed Opt* 2009; 14:064042; PMID:20059280; <http://dx.doi.org/10.1117/1.3275477>
237. Tsen K, Tsen S, Chang C, Hung C, Wu T, Kiang J. Inactivation of viruses with a very low power visible femtosecond laser. *J Phys Condens Matter* 2007; 19:322102; <http://dx.doi.org/10.1088/0953-8984/19/32/322102>
238. Tsen K, Tsen SD, Sankey O, Kiang J. Selective inactivation of microorganisms with near-infrared femtosecond laser pulses. *J Phys Condens Matter* 2007; 19:472201; <http://dx.doi.org/10.1088/0953-8984/19/47/472201>
239. Tsen K, Tsen S, Chang C, Hung C, Wu T, Ramakrishna B, et al. Inactivation of viruses with a femtosecond laser via impulsive stimulated Raman scattering. *Proc SPIE* 2008; 6854:68540N; <http://dx.doi.org/10.1117/12.762324>
240. Tsen S, Tsen Y, Tsen K, Wu T. Selective inactivation of viruses with femtosecond laser pulses and its potential use for in vitro therapy. *J Healthcare Engineering* 2010; 1:185-96; <http://dx.doi.org/10.1260/2040-2295.1.2.185>
241. Tsen KT, Tsen SW, Fu Q, Lindsay SM, Li Z, Cope S, Vaiana S, Kiang JG. Studies of inactivation of encephalomyocarditis virus, M13 bacteriophage, and Salmonella typhimurium by using a visible femtosecond laser: insight into the possible inactivation mechanisms. *J Biomed Opt* 2011; 16:078003; PMID:21806295; <http://dx.doi.org/10.1117/1.3600771>
242. Tsen SW, Chapa T, Beatty W, Tsen KT, Yu D, Achilefu S. Inactivation of enveloped virus by laser-driven protein aggregation. *J Biomed Opt* 2012; 17:128002; PMID:23224114; <http://dx.doi.org/10.1117/1.JBO.17.12.128002>
243. Tsen SW, Wu TC, Kiang JG, Tsen KT. Prospects for a novel ultrashort pulsed laser technology for pathogen inactivation. *J Biomed Sci* 2012; 19:62; PMID:22768792; <http://dx.doi.org/10.1186/1423-0127-19-62>
244. Lowy FD. Staphylococcus aureus infections. *N Engl J Med* 1998; 339:520-32; PMID:9709046; <http://dx.doi.org/10.1056/NEJM199808203390806>
245. Dai T, Tegos GP, Rolz-Cruz G, Cumbie WE, Hamblin MR. Ultraviolet C inactivation of dermatophytes: implications for treatment of onychomycosis. *Br J Dermatol* 2008; 158:1239-46; PMID:18410410; <http://dx.doi.org/10.1111/j.1365-2133.2008.08549.x>
246. Field Manual: Treatment of Biological Warfare Agents Casualties. Field Headquarters, Departments of the Army, the Navy, and the Air Force, and Commandant, Marine Corps; Army FM 8-284, Navy NAVMED P-5042, Air Force AFMAN (I) 44-156, Marine Corps MCRP 4-11.1C.