



ELSEVIER

Contents lists available at ScienceDirect

Data in brief

journal homepage: www.elsevier.com/locate/dib



Data Article

Data on the yield and quality of organically hybrids of tropical tomato fruits at two stages of fruit maturation



Syariful Mubarak^{a,*}, Fathimah Fathinatunnabilah Farhah^a,
Anas^a, Nursuhud Suwali^a, Dikdik Kurnia^b, Kusumiyati^a,
Erni Suminar^a, Hiroshi Ezura^c

^a Department of Agronomy, Faculty of Agriculture, Universitas Padjadjaran, Bandung, 45363, Indonesia

^b Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Padjadjaran, Bandung, 45363, Indonesia

^c Faculty of Life and Environmental Sciences, University of Tsukuba, Tsukuba, 305-8572, Japan

ARTICLE INFO

Article history:

Received 1 April 2019

Received in revised form 2 May 2019

Accepted 14 May 2019

Available online 30 May 2019

ABSTRACT

Organic and traditional cultivation techniques significantly affect the yield and quality of tomato fruit. To achieve the highest possible production of hybrid lines, the appropriate cultivation system is needed. The application of different cultivation systems was expected to improve the yield and fruit quality of three new tropical hybrid tomatoes varieties that prolong fruit shelf life. This experiment was conducted to identify the effect of the different cultivation systems on the yield and fruit quality of three hybrid tomatoes from different tropical parental backgrounds ('Mutiar', 'Intan' and 'Ratna'). Those hybrid lines were cultivated with two farming systems (organic and conventional cultivation system), and the fruit quality was analysed at two stages of fruit maturation (Breaker and Red).

© 2019 The Author(s). Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

* Corresponding author.

E-mail address: syariful.mubarak@unpad.ac.id (S. Mubarak).

Specifications Table

Subject area	Agriculture
More specific subject area	Postharvest, Horticulture
Type of data	Figure
How data was acquired	Refractometer model 3810 Pal-1, Advanced Bench pH Meter 3510, AquaMate 8000 UV–Vis Spectrophotometer
Data format	Analysed
Experimental factors	This experiment consisted of three factors (hybrid/F1 varieties, cultivation methods, and stage of fruit maturation) and was repeated four times. Three hybrid lines of tropical tomatoes varieties (Mutiar F1, Intan F1, and Ratna F1) were cultivated using two methods of cultivation (organic and conventional cultivation systems), and the fruits were harvested at two stages of fruit maturation (Breaker and Red) in triplicate.
Experimental features	Determination of fruit yield, fruit quality i.e. fruit water content, titratable acidity, total soluble solid, pH, β -carotene and lycopene, and also data of micro-climate
Data source location	Sumedang, Indonesia.
Data accessibility	The data are obtainable within this article and publicly accessible.
Related research article	Mubarak, S., Okabe, Y., Fukuda, N., Ariizumi, T., and Ezura, H. 2015. Potential use of a weak ethylene receptor mutant <i>Sletr1-2</i> , as breeding material to extend fruit shelf life of tomato. <i>Journal of Agricultural and Food Chemistry</i> . 63: 7995–8007 [1]. Vinha, A.F., Barreira, S.V., Costa, A.S., Alves, R.C., and Oliveira, M.B. 2014. Organic versus conventional tomatoes: influence on physicochemical parameters, bioactive compounds and sensorial attributes. <i>Food and Chemical Toxicology</i> . 67: 139–144 [2]. Makinde, A.I., Jekanola O.O., Adedeji J.A., Awogbade A.L., and Adekunle A.F. 2016. Impact of organic and inorganic fertilizers on the yield, lycopene, and some minerals in tomato (<i>Lycopersicum esculentum</i> Mill.) fruit. <i>European Journal of Agriculture and Forestry Research</i> . 4: 18–26 [3].

Value of the data
<ul style="list-style-type: none">• The data obtained here will contribute to our understanding of the relationships between cultivation systems and changes in tomato quality.• The data could be used for practitioner and as basic data of further research.

1. Data

The data report the fruit yield and quality of three hybrid tomato lines grown under two cultivation systems, organic and conventional. We analysed several parameters related to micro-climate and the fruit yield and quality (fruit weight per plant, fruit number per plant, fruit diameter, fruit weight, titratable acidity, fruit water content, total soluble solid, and fruit pH). Fig. 1 shows micro-climate data, and Fig. 2 shows the effects of two cultivation systems on fruit weight per plant, fruit diameter, fruit number per plant, and fruit length data. Fig. 3 shows data on fruit water content and total soluble solid (TSS). Fig. 4 reports data on fruit titratable acidity (TA) and pH. Fig. 5 shows data on β -carotene and lycopene.

2. Experimental design, materials, and methods

2.1. Plant preparation and cultivation

Three hybrid tomato cultivars (Mutiar F1, Intan F1, and Ratna F1) were cultivated on two cultivation systems (organic and conventional system) within a 100 m² of tunnel type screen house at the Field of Research Center of Faculty of Agriculture, Universitas Padjadjaran, Indonesia from June 2018 to August 2018. The micro-climate condition of the greenhouse is crucial to the impact of the yield and fruit quality [4], therefore, during experimental period, air temperature and humidity were measured using an HTC-2 digital thermo-hygrometer (HTC Instruments, India). Seeds were sown in a seed tray at the beginning of the plants' cultivation with soil and cocopeat (1:1/v:v) as a growing medium. After 3–4 weeks, the tomato plants were transplanted into a 30-cm polybag with a growing medium of compost and husk charcoal (1:1/v:v). The differences between organic and conventional cultivation system were in the used of organic or inorganic fertilizer and also pest/disease control. The fertilization




Fig. 1. Micro-climate data (temperature and humidity) during three months of experimental period.




Fig. 2. Fruit yield responses of tomato fruits from organic and conventional cultivation system. Mean \pm Standard Error (SE, $n = 4$) followed by one and two asterisk are significantly different compare to conventional cultivation system in each hybrid line according to the Student's T-Test at $p < 0,05$ and $p < 0,01$, respectively.

was applied every week with 1 ml/L of liquid organic fertilizer for organic cultivation system and 1 g/L of NPK fertilizer for conventional cultivation system. No chemical fertilizer and pesticide were applied under the organic cultivation system. To analyse the fruit quality, the fruits were harvested at two stages of fruit maturation, namely Breaker (Br) and Red (Br+7).

2.2. Fruit yield analysis

The same fruit maturation stage on Br and Br+7 were harvested to be used for fruit yield and quality analyses. Several parameters related to fruit yield were analysed (i.e., fruit weight per plant, fruit diameter, fruit number per plant, and fruit length).

2.3. Analysis of the fruit water content

The fruit water content was analysed according to the method described by Aveni [5]. Briefly, fruit samples were dried using an oven with a temperature of 60 °C until they reached stable, dried fruit weight. Fruit water content was measured using the following equation:




Fig. 3. Fruit water content and total soluble solid (TSS) of two stages of fruit maturation from organic and conventional cultivation system. Mean \pm Standard Error (SE, $n = 4$) followed by two asterisk are significantly different compare to conventional cultivation system in each hybrid line according to the Student's T-Test at $p < 0.01$.

$$\% \text{ Fruit water content} = \frac{a - b}{b} \times 100\%$$

a: fresh fruit weight (g)

b: dried fruit weight (g)

2.4. Analysis of the titratable acidity (TA), pH, and total soluble solid (TSS)

TA was measured using the modified titration method described by Mubarak et al. [6,7]. Briefly, 5 g of fresh fruit was homogenised with 50 mL distilled water and then titrated up to pH 8.1 with 0.1 N sodium hydroxide. TA and pH were determined using an Advanced Bench pH Meter 3510 (Jenway, United Kingdom). TSS was used to estimate the fruit sugar content and measured with a refractometer model 3810 Pal-1 (ATAGO CO., LTD., Japan).

2.5. Analysis of β -carotene and lycopene

The β -carotene and lycopene contents were analysed at the breaker and red stages of fruit maturation, as done by Mubarak et al. [1] with some modification. Briefly, 7 mL of hexane/acetone (6:4 v/v)




Fig. 4. Fruit titratable acidity (TA) and pH of two stages of fruit maturation from organic and conventional cultivation system. Mean \pm Standard Error (SE, $n = 4$) followed by two asterisk are significantly different compare to conventional cultivation system in each hybrid line according to the Student's T-Test at $p < 0.01$.

was used to extract the 700 mg tomato fruit pericarp. Then, the absorbance of the clear supernatant was measured using an AquaMate 8000 UV–Vis Spectrophotometer (Thermo Scientific, United States of America) at four absorbances namely 663 (A_{663}), 645 (A_{645}), 505 (A_{505}), and 453 (A_{453}). The following equations were used to measure lycopene (C_{LYC}) and β -carotene (C_{CAR}):

$$C_{LYC} = -0.0458 (A_{663}) + 0.204(A_{645}) + 0.372(A_{505}) - 0.0806(A_{453})$$

$$C_{CAR} = 0.216(A_{663}) - 1.22(A_{645}) - 0.304(A_{505}) + 0.452(A_{453})$$

The content of lycopene and β -carotene were represented in micro-grams per grams of fruit fresh weight ($\mu\text{g/g}$ FW).

2.6. Data analysis

This experiment was carried out with four replicates in a randomised block design. The data were represented as the mean values \pm SE, and a student's t-test was performed to compare the effects of the organic and conventional cultivation systems.




Fig. 5. β -carotene and lycopene of two stages of fruit maturation from organic and conventional cultivation system. Mean \pm Standard Error (SE, n = 4) followed by two asterisk are significantly different compare to conventional cultivation system in each hybrid line according to the Student's T-Test at $p < 0.01$.

Acknowledgments

We thank the Universitas Padjadjaran for providing the funding of RFU research project (No. 855/UN6.3.1/PL/2017). We also thank all of the laboratory members for helpful discussions throughout the work.

Conflict of interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

- [1] S. Mubarak, Y. Okabe, N. Fukuda, T. Ariizumi, H. Ezura, Potential use of a weak ethylene receptor mutant *Sletr1-2*, as breeding material to extend fruit shelf life of tomato, *J. Agric. Food Chem.* 63 (2015) 7995–8007. <https://doi.org/10.1021/acs.jafc.5b02742>.
- [2] A.F. Vinha, S.V. Barreira, A.S. Costa, R.C. Alves, M.B. Oliveira, Organic versus conventional tomatoes: influence on physicochemical parameters, bioactive compounds and sensorial attributes, *Food Chem. Toxicol.* 67 (2014) 139–144.
- [3] A.I. Makinde, O.O. Jekanola, J.A. Adedeji, A.L. Awogbade, A.F. Adekunle, Impact of organic and inorganic fertilizers on the yield, lycopene, and some minerals in tomato (*Lycopersicon esculentum* Mill.) fruit, *European Journal of Agriculture and Forestry Research* 4 (2016) 18–26.

- [4] R.R. Shamshiri, J.W. Jones, K.R. Thorp, D. Ahmad, H.C. Man, S. Taheri, Review of optimum temperature, humidity, and vapour pressure deficit for micro-climate evaluation and control in greenhouse cultivation of tomato: a review, *Int. Agrophys.* 32 (2018) 287–302.
- [5] A. Aventi, Penelitian Pengukuran Kadar Air Buah, *Proceeding Seminar Nasional Cendekiawan*, 2015, pp. 2–27 (In Indonesian).
- [6] S. Mubarak, K. Hoshikawa, Y. Okabe, R. Yano, M.D. Tri, T. Ariizumi, H. Ezura, Evidence of the functional role of the ethylene receptor genes *SlETR4* and *SlETR5* in ethylene signal transduction in tomato, *Mol. Genet. Genom.* 294 (2019) 301–313.
- [7] S. Mubarak, S. Dahlania, N. Suwali, Dataset on the change of postharvest quality of *Physalis peruviana* L. as an effect of ethylene inhibitor, *Data in Brief* 24 (2019) 103849.