



Recurrence of spinal schwannomas in a patient with neurofibromatosis type 1: a case report

Hardik Bhandari, MBBS*, Alok Dahal, MS, Grishma Khadka, MBBS

Introduction and importance: Spinal schwannomas are benign tumors usually attached to peripheral nerves, consisting of a clonal population of Schwann cells. Neurofibromatosis type 1 is an autosomal dominant neurocutaneous disorder that predominantly affects the skin, bone and nervous system. Neurofibromatosis type 1 is a clinically and genetically distinct from neurofibromatosis type 2. This case report highlights the rare association between spinal schwannoma and neurofibromatosis type 1.

Case presentation: The patient with a past medical history of spinal schwannoma, operated 1 year back, presented with back pain, weakness of lower limbs and urge incontinence. On examination, she had cutaneous features suggestive of neurofibromatosis type 1 and there was impairment of all sensory modalities below hip region. MRI revealed spinal schwannoma at D9 level for which laminectomy with removal of schwannoma was performed.

Clinical discussion: The occurrence and recurrence of spinal schwannoma in neurofibromatosis type 2 is a common finding. But such an association has not been established between spinal schwannoma and neurofibromatosis type 1. In this case, the recurrence of spinal schwannoma has been linked to neurofibromatosis type 1 in the absence of other well-defined etiologies.

Conclusion: The occurrence of spinal schwannoma can be genetic or sporadic. The recurrence is usually associated with familial tumor syndrome. The available literature has not established association between neurofibromatosis type 1 and spinal schwannoma, thus, emphasizing the need of more focused studies.

Keywords: neurofibromatosis type 1, schwannoma, spinal

Introduction

Schwannomas are benign, well-circumscribed tumors usually attached to peripheral nerves, consisting of a clonal population of Schwann cells^[1]. Spinal schwannomas account for about 25% of primary intradural spinal cord tumors in adults^[2]. In the vast majority of cases, schwannomas are solitary and sporadic, while multiple schwannomas are associated with neurofibromatosis type 2 (NF2) and schwannomatosis. Neurofibromatosis type 1 (NF1) is an autosomal dominant neurocutaneous disorder that predominantly affects the skin, bone and nervous system. NF1 is distinct on clinical and genetic grounds from neurofibromatosis type 2, a rare disorder characterized by bilateral vestibular schwannomas and other benign nervous system tumors^[3]. The

HIGHLIGHTS

- Recurrence of spinal schwannomas is common in neurofibromatosis type 2.
- The occurrence of spinal schwannoma in neurofibromatosis type 1 is a rare phenomenon.
- Further studies is necessary to establish the association between spinal schwannoma and neurofibromatosis type 1.

gold standard treatment for symptomatic spinal schwannomas is complete surgical resection, which stops symptoms progression, helps recovery in most patients, and decreases the rate of recurrence. Radiotherapy can be considered as second-choice treatment in patients who are not good candidates for surgery, or for recurrent tumors^[4]. The recurrence rate for spinal schwannomas ranges from 4.08 to 9.4% and depends on size, extent of tumor and genetic associations rather than the location of the tumor^[5,6]. This case has been reported in line with the SCARE 2023 criteria^[7].

Case presentation

A 50-year-old female presented to OPD with complains of back pain and weakness of lower limbs for 3 months. The pain was insidious in onset, gradually progressing and later radiating to bilateral lower limbs. It was associated with weakness of lower limbs in the form of difficulty in walking and inability to stand for prolonged periods of time that had increased to the extent that she had been unable to walk without support. She had urge incontinence as she could not control her bladder and bowel once

Department of Neurosurgery, BP Koirala Institute of Health Sciences, Dharan, Nepal
Sponsorships or competing interests that may be relevant to content are disclosed at the end of this article.

*Corresponding author. Address: Department of Neurosurgery, BP Koirala Institute of Health Sciences, Dharan, Nepal. Tel.: +977 984 745 8290.
E-mail: hbbhandari412@gmail.com (H. Bhandari).

Copyright © 2024 The Author(s). Published by Wolters Kluwer Health, Inc. This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

Annals of Medicine & Surgery (2024) 86:6169–6172

Received 14 May 2024; Accepted 31 July 2024

Published online 14 August 2024

<http://dx.doi.org/10.1097/MS9.0000000000002457>

she had sensation. There is also a past history of being operated 1 year back after she was diagnosed with spinal schwannoma; she had undergone laminectomy with total resection for the spinal schwannoma extending from L5–S1.

On examination, there was skin patch over her left hand measuring 2 cm, axillary freckling along with multiple fibromas over her back; suggestive of NF type 1. Genetic studies were not performed. The diagnostic criteria utilized to make a diagnosis is NIH criteria for neurofibromatosis 1^[8]. Her higher mental function was intact, cranial nerves were intact. There was no motor or sensory deficit over both upper limbs. The power of the right lower and left lower limb were 3/5 and 4/5, respectively. There was sensory impairment of all modalities below the hip region (right > left).

Her recent MRI revealed a single right-sided intradural-extramedullary, homogeneously enhancing, well-marginated mass at D9 level (Fig. 1). Other laboratory parameters were within normal range. She underwent bilateral laminectomies of D9, following the durotomy. A well-defined solid yellowish lesion was visualized adjacent to the D9 vertebrae; it was then completely resected (Fig. 2). Neuromonitoring potentials remained unchanged. The tumor was sent for histopathology and was later confirmed as benign spinal schwannoma. The patient recovered postoperatively with a gradual increase in the strength of lower limbs and was discharged on 3rd postoperative day.

Discussion

Recurrence is defined as reappearance after definite surgical excision or symptomatic increase in size of a residual tumor on follow-up imaging studies. Local recurrence means reappearance in the same place as the original tumor or very close to it. Regional recurrence means reappearance into lymph nodes or tissues near the original tumor. Distant recurrence means reappearance in organs or tissues far from the original tumor^[9].

Spinal schwannoma is a solid, encapsulated, mostly benign and intradural-extramedullary tumor. They may occur spontaneously, or in the context of a familial tumor syndrome such as NF2, schwannomatosis and Carney's complex^[1]. However, the occurrence of spinal schwannomas in neurofibromatosis type 1 is not well established and such phenomenon is rare finding in itself. But as in other causes of spinal schwannomas, gross total resection (GTR) with preservation of neurological functions is the best treatment to relieve patients' complaints and to reduce the recurrence rate^[6].

Analyzing our case, the patient developed spinal schwannomas at two different locations at different periods of time and at both instances, she underwent laminectomy with removal of schwannoma with good postoperative recovery. Recurrence is common in schwannoma if the residual tumor is left during excision but at the same site of the initial lesion^[10]. There should be high degree of suspicion to consider the recurrence of spinal schwannoma,




Figure 1. MRI revealed intradural-extramedullary lesion extending from D9 to D10 level and compressing the cord (marked by arrow).



Figure 2. Postoperative tumor (schwannoma) after excision.

especially if the patient has been previously operated for the same condition. Chordomas, Giant cell tumors, Meningioma and Chondroblastoma are part of the differential diagnoses for spinal schwannomas, and each should be carefully excluded to reach a diagnosis of spinal schwannoma. Laboratory tests combined with MRI have a good sensitivity and specificity for spinal schwannoma. Laminectomy has normally been used as a standard approach for intradural spinal tumors but this procedure is associated with spinal instability and deformity. Laminoplasty was developed to overcome these complications^[11]. However, studies have not demonstrated laminoplasty to have any better functional outcome compared to laminectomy in the resection of intradural lesions^[12,13].

Since schwannoma is a slow-growing, benign neoplasm, the expected interval between two episodes can be years but the patient developed two different neoplasms within a period of 1 year; the cause of which cannot be linked to other than neurofibromatosis type 1. The neurofibromatosis type 1 is typically associated with the occurrence of plexiform neurofibromas and less commonly central nervous system gliomas^[14]. There is no literature describing the recurrence of spinal schwannomas in neurofibromatosis type 1. Thereby, the spinal schwannoma is a rare occurrence in neurofibromatosis type 1 and needs further study to establish a link between these two entities.

Thus, although there is well-established association between spinal schwannomas and neurofibromatosis type 2; the diagnosis should be considered in a patient with neurofibromatosis type 1 whenever the patient develops symptoms or the symptoms worsens followed by proper investigation to identify/ rule out the diagnosis.

Conclusion

The occurrence of spinal schwannomas in a patient with neurofibromatosis type 1 is a rare phenomenon and in the absence of

well-established association between spinal schwannoma and neurofibromatosis type 1, it becomes difficult for clinicians to consider the diagnosis especially when there is recurrence following the initial surgery. Thus, further study is needed between neurofibromatosis type 1 and spinal schwannomas to establish an association between them so that it would help clinicians to consider the diagnosis when the patient presents with such symptoms.

Ethical approval

Ethical approval is not required for case report from ethics committee at our institution.

Consent

Written informed consent was obtained from the patient for publication and any accompanying images. A copy of the written consent is available for review by the Editor-in-Chief of this journal on request.

Source of funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Author contribution

H.B.: conceptualization, writing—original draft preparation, literature review, and final manuscript approval. A.D.: conceptualization, literature review, supervision and final manuscript approval. G.K. supervision and final manuscript approval.

Conflicts of interest disclosure

The authors declare no conflicts of interest.

Research registration unique identifying number (UIN)

1. Name of the registry: not applicable.
2. Unique identifying number of the study: not applicable.
3. Hyperlink to your specific registration (must be publicly accessible and will be checked): not applicable.

Guarantor

Hardik Bhandari is the guarantor.

Data availability statement

Not applicable.

Provenance and peer review

Not commissioned, externally peer-reviewed.

References

- [1] Hilton DA, Hanemann CO. Schwannomas and their pathogenesis. *Brain Pathol* 2014;24:205–20.
- [2] Senapati S, Mishra S, Dhir M, *et al*. Recurrence of spinal schwannoma: is it preventable? *Asian J Neurosurg* 2016;11:451–2.
- [3] Ahlawat S, Blakeley J, Montgomery E, *et al*. Schwannoma in neurofibromatosis type 1: a pitfall for detecting malignancy by metabolic imaging. *Skeletal Radiol* 2017;42:1317–22.
- [4] Lenzi J, Anichini G, Landi A, *et al*. Spinal Nerves Schwannomas: Experience on 367 Cases - Historic Overview on How Clinical, Radiological, and Surgical Practices Have Changed over a Course of 60 Years. *Neurol Res Int* 2017;2017:3568359.
- [5] Sowash M, Barzilai O, Kahn S, *et al*. Clinical outcomes following resection of giant spinal schwannomas: a case series of 32 patients. *J Neurosurg Spine* 2017;26:494–500.
- [6] Emel E, Abdallah A, Sofuoglu OE, *et al*. Long-term surgical outcomes of spinal schwannomas: Retrospective analysis of 49 consecutive cases. *Turk Neurosurg* 2017;27:217–25.
- [7] Sohrabi C, Mathew G, Maria N, *et al*. The SCARE 2023 guideline: updating consensus Surgical CAse REport (SCARE) guidelines. *Int J Surg* 2023;109:1136–40.
- [8] Legius E, Messiaen L, Wolkenstein P, *et al*. Revised diagnostic criteria for neurofibromatosis type 1 and Legius syndrome: an international consensus recommendation. *Genet Med* 2021;23:1506–13.
- [9] Recurrent Cancer - NCI [Internet]. Accessed 18 July 2024. <https://www.cancer.gov/types/recurrent-cancer>
- [10] Seppala MT, Haltia MJJ, Sankila RJ, *et al*. Long-term outcome after removal of spinal schwannoma: a clinicopathological study of 187 cases. *J Neurosurg* 1995;83:621–6.
- [11] Montano N, Trevisi G, Cioni B, *et al*. The role of laminoplasty in preventing spinal deformity in adult patients submitted to resection of an intradural spinal tumor. *Case Ser Literature Rev* 2014;125:69–74.
- [12] Onyia C, Menon SK. Laminectomy versus laminoplasty in the surgical management of long-segment intradural spinal tumors: Any difference in neurological outcomes?. *Asian J Neurosurg* 2018;13:1128–33.
- [13] Lao L, Zhong G, Li X, *et al*. Laminoplasty versus laminectomy for multi-level cervical spondylotic myelopathy: a systematic review of the literature. *Asian J Neurosurg and research*. 2013;1–9.
- [14] Rasmussen SA, Friedman JM. *NF1* Gene and Neurofibromatosis 1. *Am j Epidemiol* 2000;151:33–40.