

Hypothesis & Experience  
Case Report



# Multiple food allergy – unexpected culprits

**Cristina Isabel de Figueiredo Ornelas** <sup>1\*</sup>, **Fátima Cabral Duarte**<sup>1</sup>,  
**Maria Conceição Galvão Pereira dos Santos**<sup>2,3</sup>,  
and **Manuel Augusto de Castro Pereira Barbosa**<sup>1,3</sup>

<sup>1</sup>Serviço de Imunoalergologia, Hospital de Santa Maria – Centro Hospitalar Lisboa Norte, Lisbon, Portugal

<sup>2</sup>Unidade de Imunologia Clínica – Instituto de Medicina Molecular, Faculdade de Medicina – Universidade de Lisboa, Lisbon, Portugal

<sup>3</sup>Clínica Universitária de Imunoalergologia, Faculdade de Medicina – Universidade de Lisboa, Lisbon, Portugal



Received: Mar 28, 2018  
Accepted: Jul 20, 2018

### \*Correspondence to

**Cristina Isabel de Figueiredo Ornelas**

Serviço de Imunoalergologia, Hospital de Santa Maria – Centro Hospitalar Lisboa Norte Avenida Professor Egas Moniz, 1649-035 Lisboa, Portugal.

Tel: +351914349064

Fax: +351217999527

E-mail: cristina.ornelas@gmail.com

Copyright © 2018. Asia Pacific Association of Allergy, Asthma and Clinical Immunology. This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

### ORCID iDs

Cristina Isabel de Figueiredo Ornelas   
<https://orcid.org/0000-0002-2296-9643>

### Author Contributions

Conceptualization: Cristina Isabel de Figueiredo Ornelas, Maria de Fátima Cabral Duarte de Moura Portugal, Maria Conceição Galvão Pereira dos Santos. Data curation: Cristina Isabel de Figueiredo Ornelas, Maria de Fátima Cabral Duarte de Moura Portugal. Investigation: Cristina Isabel de Figueiredo Ornelas, Maria de Fátima Cabral Duarte de Moura Portugal, Maria Conceição Galvão Pereira dos Santos. Supervision: Maria Conceição Galvão Pereira dos Santos, Manuel

## ABSTRACT

Food allergy has an estimated prevalence of 6%–8% in children. Meat allergy and multiple food allergy due to sensitization to cross-reactive components in infancy is, however, less frequent. A 5-year-old girl was referred to our department with a multiple food allergy history. She had severe immediate worsening of her atopic dermatitis with hen's egg (6 months) and cow's milk introduction (7 months). At the age of 9 months, she presented with recurrent and reproducible atopic dermatitis' worsening and lip edema with the introduction of different meats (chicken, turkey, cow, pork, and rabbit), having the same complaints with fish at 12 months (salmon and hake). At her first appointment she was avoiding hen's egg, cow's milk, meat, and fish (except fresh tuna, codfish, and pollock). We performed skin prick tests (commercial extract and prick-to-prick with whole food) and specific IgE, which revealed sensitization to hen's egg, raw meat (cow, pork, chicken, turkey, duck, lamb, goat, and rabbit; negative for cooked meat), codfish and cow's milk (mild). ISAC was performed, revealing sensitization to 3 cross-reactive components (serum albumins Bosd6, Canf3, and Feld2) and specific food components of chicken's egg/meat (Gald1, 2, 3, and 5), cod (Gadcl1), hazelnut (Cora9), and kiwi (Actd1). We present a rare case of multiple food allergy in infancy, where sensitization to cross-reactive components was responsible for most of the children complaints. The detection of serum albumins' involvement was especially important, because it can possibly mean tolerance to these foods in well-cooked forms, substantially improving patient and family's quality of life.

**Keywords:** Food allergy; Serum albumin; Cross-reactions

## INTRODUCTION

Food allergy has an overall estimated prevalence of 6%–8% in children, being cow's milk, hen's egg, fish, wheat, soy, peanuts, and tree nuts the most frequent allergens involved [1]. Meat allergy and multiple food allergy due to sensitization to cross-reactive components in infancy is, however, less frequent.

Serum albumins (SAs) are  $\alpha$ -helical globular proteins synthesized in the liver, being abundant in plasma and contributing to regulation of oncotic pressure. They're present in dander, saliva, milk, and meat of animals, being denatured by heating due to their thermolabile properties

Augusto de Castro Pereira Barbosa. Validation: Cristina Isabel de Figueiredo Ornelas, Maria de Fátima Cabral Duarte de Moura Portugal, Maria Conceição Galvão Pereira dos Santos, Manuel Augusto de Castro Pereira Barbosa. Writing - original draft: Cristina Isabel de Figueiredo Ornelas, Maria de Fátima Cabral Duarte de Moura Portugal, Maria Conceição Galvão Pereira dos Santos. Writing - review & editing: Cristina Isabel de Figueiredo Ornelas, Maria de Fátima Cabral Duarte de Moura Portugal, Maria Conceição Galvão Pereira dos Santos, Manuel Augusto de Castro Pereira Barbosa.

[2, 3]. SAs are usually considered minor allergens, with seven being officially registered as allergens by World Health Organization and International Union of Immunological Societies Allergen Nomenclature Subcommittee: Bos d 6 (bovine), Can f 3 (dog), Cav p 4 (guinea pig), Equ c 3 (horse), Fel d 2 (cat), Gal d 5 (chicken), and Sus s 1 (pig). Mammalian SAs have high sequence identities and similarities (72%–88%) relative to human SA (HSA) and between them, while aviarian SA presents lower identities (42%–49%) to HSA and other mammalian SA [2]. Due to their high sequence identity (>70%) mammalian SAs are highly cross-reactive [2], while cross-reactivity between mammalian and aviarian SA is rare but already described [4].

## CASE REPORT

We report the case of a 5-year-old girl that was referred by her pediatrician to our Imunoallergology Department for investigation of a multiple food allergy. She had previous clinical history of moderate atopic dermatitis (AD) since 3 months, moderate to severe persistent allergic rhinitis since 3 years and recurrent wheezing since the age of 4, with sensitization to house dust mites, mold, grass pollen, cat epithelium and feathers. She was medicated with daily emollient, nasal corticosteroid, oral antihistamine, leukotriene antagonist, and inhaled long-acting  $\beta$ -2 agonist/corticosteroid association. Her father had AD and her sister of 12-years-old has AD, allergic rhinitis and asthma. She lives in an urban environment without domestic pets but with frequent contact with pigeons outside and neighbor's cat. She had severe immediate worsening of her AD with hen's egg (white and yolk simultaneously) and cow's milk introduction (without previous complaints with formula milk) at 6 and 7 months, respectively. At the age of 9 months she presented with recurrent and reproducible AD's worsening and lip edema with the introduction of different meats (chicken, turkey, cow, pork, and rabbit). The same happened at 12 months with fish (salmon and hake). Her mother denies AD's worsening or other complaints outside these episodes related with food intake. She was referred several times to our department before but she missed all the appointments. At her first appointment she was 5-years-old and she was avoiding hen's egg, cow's milk (tolerated milk traces), meat, and fish (except fresh tuna, codfish, and pollock).

We performed skin prick tests (SPT) (commercial extract and prick-to-prick with whole food) with all the suspected foods (Fig. 1). They were considered positive when showing a wheal with the largest diameter at least 3 mm larger than that of the negative control. SPT were positive (Fig. 2) for hen's egg (white and yolk – commercial extract and food in nature, raw and cooked forms; ovomucoid; ovalbumin) and raw meat (cow, pork, chicken, turkey, duck, lamb, goat, and rabbit; negative for cooked meat). SPT were negative for all of the other foods suspected.

She had total IgE of 1,078 kU/L. Specific IgE (ImmunoCAP system, Thermo Fisher, Uppsala, Sweden) was strongly positive for hen's egg (all components) and chicken meat, slightly positive for pork meat, codfish and cow's milk and fractions and negative for alpha-gal (Fig. 3A). ISAC (Thermo Fisher) was performed (Fig. 3B), revealing sensitization to 3 cross-reactive components (SAs Bos d 6, Can f 3, and Fel d 2) and specific food components of chicken's egg/meat (Gal d 1-3 and 5), codfish (Gad c 1), hazelnut (Cor a 9), and kiwi (Act d 1), the last two without associated clinical manifestations.

Until present time she still maintains sensitization and is still avoiding hen's egg but she can now tolerate lamb, goat, and duck meat and cow's milk in very well cooked form. Fish allergy is now overcome.



**Fig. 1.** Picture of skin prick tests (commercial extract and prick-to-prick with whole food) with cow's milk, hen's egg, fish, and meat (raw and cooked).



**Fig. 2.** Picture of skin prick tests results - positive results for hen's egg (white and yolk) and raw meat (cow, pork, turkey, and rabbit); negative results for cooked meat, cow's milk, and codfish.

## DISCUSSION

In this case sensitization to bovine SA (BSA) Bos d 6 seems to be responsible for cow's milk and beef allergy. In fact BSA has been identified as one of major beef allergens, particularly in children, being responsible for clinical cross-reactivity between raw beef and cow's milk (sensitization independent of other cow's milk proteins, as seen in this case) [5-7]. Sensitization to other SAs may be explained by the high sequence homology between mammalian SAs, being possibly responsible for the allergic symptoms reported with pork and rabbit meat (SAs not available). It is important, however, to note that patient presented sensitization to cat epithelium and this may also have contributed to pork meat sensitization, by cross-reactivity between cat and pork SAs (cat-pork syndrome), as already been reported in literature [8, 9]. Regarding complaints with chicken and turkey meat, since cross-reactivity

<b>A</b>			
Food allergen	Specific IgE (kU/L)		
Cow's milk	1.50		
$\alpha$ -lactalbumin	0.63		
B-lactoglobulin	0.69		
Casein	0.62		
White egg	50.30		
Yolk egg	70.40		
Ovalbumin	44.10		
Ovomucoid	10.30		
Codfish	5.16		
Pork meat	5.05		
Chicken meat	51.40		
$\alpha$ -gal	0.06		

  

<b>B</b>			
Summary of positive IgE results			
Mainly species-specific food components			
Egg white	nGal d 1	Ovomucoid	1, 9 ISU-E
	nGal d 2	Ovalbumin	1, 7 ISU-E
	nGal d 3	Conalbumin/Ovotransferrin	4, 6 ISU-E
Egg yolk/chicken meat	nGal d 5	Livetin/Serum albumin	9, 4 ISU-E
Cod	rGad c 1	Parvalbumin	0, 6 ISU-E
Hazelnut	nCor a 9	Storage protein, 11S globulin	0, 4 ISU-E
Kiwi	nAct d 1	Cysteine protease	0, 5 ISU-E
Mainly species-specific aeroallergen components			
<b>Grass pollen</b>			
Timothy grass	nPhl 4	Berberine bridge enzyme	0, 4 ISU-E
<b>Mold</b>			
Aspergillus	rAsp f 6	Mn superoxide dismutase	0, 4 ISU-E
<b>Mite</b>			
<i>B. tropicalis</i> (HDM)	rBlo t 5	Mite group 5	0, 7 ISU-E
<i>D. farinae</i> (HDM)	nDer f 1	Cysteine protease	12 ISU-E
	rDer f 2	NPC2 family	20 ISU-E
	nDer p 1	Cysteine protease	17 ISU-E
<i>D. pteronyssinus</i> (HDM)	rDer p 2	NPC2 family	22 ISU-E
	rLep d 2	NPC2 family	0, 4 ISU-E
<i>L. destructor</i> (storage mite)			
Cross-reactive components			
<b>Serum albumin</b>			
Cow's milk/meat	nBos d 6	Serum albumin	2, 3 ISU-E
Dog	nCan f 3	Serum albumin	0, 8 ISU-E
Cat	nFel d 2	Serum albumin	1, 2 ISU-E

**Fig. 3.** *In vitro* workup. (A) Specific IgE (ImmunoCAP system, Thermo Fisher, Uppsala, Sweden) results - only positive results were presented. (B) ISAC (Thermo Fisher) results. *B. tropicalis*, *Blomia tropicalis*; *D. farinae*, *Dermatophagoides farinae*; *D. pteronyssinus*, *Dermatophagoides pteronyssinus*; *L. destructor*, *Lepidoglyphus destructor*.

between aviarian and mammalian SA is rare [2], these are probably due to sensitization to Gal d 5. Primary sensitization to Gal d 5, a food and inhalant allergen, may have occurred in this case through 2 different routes: respiratory, with evidence of sensitization to feathers (bird-egg syndrome) or alimentary due to hen's egg allergy, with evidence of sensitization to Gal d 1-3 (egg-bird syndrome) [10].

This case report has some limitations, such as some degree of subjectivity in clinical symptoms (mainly AD' worsening) and the inability to perform the oral food challenge until present time due to different reasons.

In conclusion we present a rare case of multiple food allergy in infancy, where sensitization to cross-reactive components was responsible for most of the children complaints. The discovery of SAs' involvement is especially important in similar cases as the one we presented because it may enable a possible tolerance to well-cooked meat and boiled milk, as described in this case, which will lead to a substantial improvement of the patient and family's quality of life.

## REFERENCES

1. Sicherer SH, Sampson HA. Food allergy: epidemiology, pathogenesis, diagnosis, and treatment. *J Allergy Clin Immunol* 2014;133:291-307.  
[PUBMED](#) | [CROSSREF](#)
2. Chruszcz M, Mikolajczak K, Mank N, Majorek KA, Porebski PJ, Minor W. Serum albumins-unusual allergens. *Biochim Biophys Acta* 2013;1830:5375-81.  
[PUBMED](#) | [CROSSREF](#)
3. Matricardi PM, Kleine-Tebbe J, Hoffmann HJ, Valenta R, Hilger C, Hofmaier S, Aalberse RC, Agache I, Asero R, Ballmer-Weber B, Barber D, Beyer K, Biedermann T, Bilò MB, Blank S, Bohle B, Bosshard PP, Breiteneder H, Brough HA, Caraballo L, Caubet JC, Cramer R, Davies JM, Douladiris N, Ebisawa M, Eigenmann PA, Fernandez-Rivas M, Ferreira F, Gadermaier G, Glatz M, Hamilton RG, Hawranek T, Hellings P, Hoffmann-Sommergruber K, Jakob T, Jappe U, Jutel M, Kamath SD, Knol EF, Korosec P, Kuehn A, Lack G, Lopata AL, Mäkelä M, Morisset M, Niederberger V, Nowak-Węgrzyn AH, Papadopoulos NG, Pastorello EA, Pauli G, Platts-Mills T, Posa D, Poulsen LK, Raulf M, Sastre J, Scala E, Schmid JM, Schmid-Grendelmeier P, van Hage M, van Ree R, Vieths S, Weber R, Wickman M, Muraro A, Ollert M. EAACI molecular allergology user's guide. *Pediatr Allergy Immunol* 2016;27 Suppl 23:1-250.  
[PUBMED](#) | [CROSSREF](#)
4. Hilger C, Swiontek K, Hentges F, Donnay C, de Blay F, Pauli G. Occupational inhalant allergy to pork followed by food allergy to pork and chicken: sensitization to hemoglobin and serum albumin. *Int Arch Allergy Immunol* 2010;151:173-8.  
[PUBMED](#) | [CROSSREF](#)
5. Fiocchi A, Restani P, Riva E. Beef allergy in children. *Nutrition* 2000;16:454-7.  
[PUBMED](#) | [CROSSREF](#)
6. Martelli A, De Chiara A, Corvo M, Restani P, Fiocchi A. Beef allergy in children with cow's milk allergy; cow's milk allergy in children with beef allergy. *Ann Allergy Asthma Immunol* 2002;89 6 Suppl 1:38-43.  
[PUBMED](#) | [CROSSREF](#)
7. Restani P, Ballabio C, Tripodi S, Fiocchi A. Meat allergy. *Curr Opin Allergy Clin Immunol* 2009;9:265-9.  
[PUBMED](#) | [CROSSREF](#)
8. Posthumus J, James HR, Lane CJ, Matos LA, Platts-Mills TA, Commins SP. Initial description of pork-cat syndrome in the United States. *J Allergy Clin Immunol* 2013;131:923-5.  
[PUBMED](#) | [CROSSREF](#)
9. Hilger C, Kohnen M, Grigioni F, Lehnert C, Hentges F. Allergic cross-reactions between cat and pig serum albumin. Study at the protein and DNA levels. *Allergy* 1997;52:179-87.  
[PUBMED](#) | [CROSSREF](#)
10. Hemmer W, Klug C, Swoboda I. Update on the bird-egg syndrome and genuine poultry meat allergy. *Allergo J Int* 2016;25:68-75.  
[PUBMED](#) | [CROSSREF](#)