

RESEARCH ARTICLE

# Interleukin-33 from Monocytes Recruited to the Lung Contributes to House Dust Mite-Induced Airway Inflammation in a Mouse Model

Hiroki Tashiro<sup>1</sup>, Koichiro Takahashi<sup>1\*</sup>, Shinichiro Hayashi<sup>1</sup>, Go Kato<sup>1</sup>, Keigo Kurata<sup>2</sup>, Shinya Kimura<sup>1</sup>, Naoko Sueoka-Aragane<sup>1</sup>

**1** Division of Hematology, Respiratory Medicine and Oncology, Department of Internal Medicine, Faculty of Medicine, Saga University, Saga, Japan, **2** Institute of Tokyo Environmental Allergy, Tokyo, Japan

\* [takahak@cc.saga-u.ac.jp](mailto:takahak@cc.saga-u.ac.jp)



## Abstract

### Background

Interleukin-33 (IL-33) activates group 2 innate lymphoid cells (ILC2), resulting in T-helper-2 inflammation in bronchial asthma. Airway epithelial cells were reported as sources of IL-33 during apoptosis and necrosis. However, IL-33 is known to be from sources other than airway epithelial cells such as leukocytes, and the mechanisms of IL-33 production and release are not fully understood. The aim of this study was to clarify the role of IL-33 production by monocytes in airway inflammation.

### Methods

BALB/c mice were sensitized and challenged with a house dust mite (HDM) preparation. Airway inflammation was assessed by quantifying inflammatory cells in bronchoalveolar lavage (BAL) fluid, and IL-25, IL-33, and thymic stromal lymphopoietin (TSLP) levels in lung. Immunohistochemistry for IL-33 in lung sections was also performed. Ly6c, CD11b, and CD11c expression was examined by flow cytometry. Clodronate liposomes were used in the HDM-airway inflammation model to deplete circulating monocytes.

### Results

The IL-33, but not IL-25 or TSLP, level in lung homogenates was markedly increased in HDM mice compared to control mice. IL-33-positive cells in the lungs were identified using immunohistochemistry and were increased in areas surrounding bronchi and vasculature. Furthermore, IL-33 levels were increased in mononuclear cells derived from lungs of HDM mice compared to controls. The expression of Ly6c in mononuclear cells was significantly higher in HDM mice than in controls. Treatment with clodronate liposomes led to inhibition of not only inflammatory cells in BAL fluid, airway hyper reactivity and Th2 cytokines in lung, but also IL-33 in lung.

## OPEN ACCESS

**Citation:** Tashiro H, Takahashi K, Hayashi S, Kato G, Kurata K, Kimura S, et al. (2016) Interleukin-33 from Monocytes Recruited to the Lung Contributes to House Dust Mite-Induced Airway Inflammation in a Mouse Model. PLoS ONE 11(6): e0157571. doi:10.1371/journal.pone.0157571

**Editor:** Bernhard Ryffel, French National Centre for Scientific Research, FRANCE

**Received:** February 27, 2016

**Accepted:** June 1, 2016

**Published:** June 16, 2016

**Copyright:** © 2016 Tashiro et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** All relevant data are within the paper and its Supporting Information files.

**Funding:** The authors have no support or funding to report.

**Competing Interests:** The authors have declared that no competing interests exist.

## Conclusion

IL-33 from monocytes recruited to the lung may contribute to the pathogenesis of HDM-induced airway inflammation.

## Background

Bronchial asthma is an airway inflammatory disease characterized by bronchoconstriction, airway hyper-responsiveness, and airway remodeling [1]. Airway eosinophilia, mediated mainly by T helper 2 (Th2)-type lymphocytes, has been reported to play an essential role in bronchial asthma [2,3].

Recently the genes encoding IL-33 and ST2 (also known as interleukin-1 receptor-like 1, IL-1RL1), have been identified as important factors for human asthma in several genome-wide association studies that included thousands of patients from diverse ethnic groups having different forms of asthma [4–6]. Other studies have suggested that early severe exacerbation of childhood asthma is closely correlated to the IL-33 gene [7]. Furthermore, high serum IL-33 was found to be related to severity of asthma [8], and IL-33 also caused airway remodeling in severe steroid-resistant asthma cases [9,10].

IL-33 is a member of the IL-1 family of cytokines and a specific ligand of the ST2/IL-1 receptor accessory protein (IL-1RAP) receptor complex [11]. IL-33 activates group 2 innate lymphoid cells (ILC2) and induces a large amount of Th2 cytokines such as IL-5 and IL-13 [12]. Th2 cytokines play a crucial role in bronchial asthma, which is characterized by eosinophilic airway inflammation and goblet cell hyperplasia [1,13,14]. The presence of IL-33 has been reported during necrosis or apoptosis of various cells including bronchial epithelial cells [15], alveolar type II cells [16], mast cells [17], dendritic cells [18], and vascular smooth muscle cells [19]. Previous studies have referred to bronchial epithelial cells as a major source of IL-33 in asthmatic airways. However, these findings remain controversial.

When pathogens or allergens invade their hosts, circulating monocytes mature into macrophages in specific organs [20]. Macrophages are classified into residential macrophages in tissues and recruited monocytes from the circulation [21,22]. Circulating monocytes, rather than residential alveolar macrophages, play a critical role in allergic airway inflammation [23].

Using a house dust mite (HDM)-induced airway inflammation mouse model, we demonstrated, in vivo and in vitro, the possibility that IL-33 from monocytes recruited to the lung played an important role.

## Materials and Methods

### Allergen and chemicals

Two batches of house dust mite (HDM) extract from *Dermatophagoides farina* (Der f) were provided by ITEA Inc. (Tokyo, Japan) as a lyophilized preparation of milled mites. Clophosome-A, liposomal clodronate and plain control liposomes were purchased from FormuMax Scientific Inc. (Palo Alto, CA, USA).

### Animals

Female BALB/c mice (Japan SLC Inc., Hamamatsu, Japan) aged 6–8 weeks were maintained at the Saga University animal facility under specific pathogen-free conditions. Animal experiments were undertaken following the guidelines for care and use of experimental animals of

the Japanese Association for Laboratory Animals Science (1987) and were approved by the Saga University Animal Care and Use Committee.

### Protocol for HDM-induced airway inflammation and treatment of mice with clophosome-A and liposomal clodronate

Mice were sensitized intranasally with 25  $\mu$ g HDM or vehicle on days 1, 8, and 15. Mice were challenged intranasally with 5  $\mu$ g HDM on days 22, 23, and 24 [24]. Four hours after the final challenge, mice were euthanized by intraperitoneal injection with sodium pentobarbital. Serum, bronchoalveolar lavage fluid (BALF) and lung tissue were harvested for further analysis. Clodronate and control liposomes (150  $\mu$ l / animal) were administered by intravenous injection on days 0, 7, 14, 21, 22, and 23. On the days when HDM was given, clodronate and control liposomes were given 30 minutes before the HDM inoculation.

### Collection of BALF

BALF samples were collected as described previously [24,25]. Briefly, a 20-gauge tube was inserted into the trachea and the lungs were lavaged with 1 ml of saline twice. The cell suspension was centrifuged at  $100 \times g$  for 5 min at 4°C. The total number of cells was counted using a hemocytometer. Cytospin samples were prepared from the cell suspension. Cell differentiation was determined by counting at least 300 leukocytes in samples stained with Diff-Quik (Siemens, Germany).

### H&E and PAS Histology

Histological examination was performed as previously reported. Lungs were fixed with 10% neutral buffered formalin (Wako, Japan) and embedded in paraffin. Lung sections were stained with Hematoxylin and Eosin (H&E) and periodic acid-Schiff (PAS) stains.

### Immunohistochemistry and Immunofluorescence

Paraffin sections of lung were deparaffinized in xylene and hydrated through graded alcohols. Sections were incubated overnight at 4°C with anti-IL-33 antibody or anti-normal goat IgG as the control (R&D Systems Inc., Minneapolis, MN, USA). Streptavidin-biotin amplification (Dako, Denmark) was carried out for 30 minutes, and hematoxylin was used to counterstain. For immunofluorescence, sections were incubated overnight at 4°C with anti-IL-33 antibody, and Northern Light 557 conjugated anti-goat IgG (R&D Systems Inc.) was used as the secondary antibody. FITC-conjugated anti-F4/80 was used to identify tissue macrophages (eBioscience, San Diego, CA). Fluorescent images were visualized using a bio-imaging navigator microscope system (Olympus, Tokyo, Japan).

### Preparation of lung homogenates

After bronchoalveolar lavage, the left lung was isolated and homogenized in 50 mM Tris buffered saline (pH 7.4) containing 1 mM EDTA, 1 mM PMSF, 1  $\mu$ g/ml aprotinin, 1  $\mu$ g/ml leupeptin, 1 mM  $\text{Na}_3\text{VO}_4$ , and 1 mM NaF. The lung homogenates were centrifuged at  $10,000 \times g$  for 15 min, and then supernatants were collected and stored at -80°C until required.

### Isolation of single cells from lung tissue

The peripheral lung tissue was cut into small pieces with scissors, transferred through 70  $\mu$ m mesh and processed in digestion buffer which included deoxyribonuclease I (Invitrogen, Waltham, MA, USA) and collagenase type 2 (Worthington Inc., Lakewood, NJ). The remaining

red cells were lysed using BD Pharm Lysis (BD Biosciences, San Jose CA), and single cells suspensions were obtained.

### Separation and analyzation of lung mononuclear cells from lung single cells

To separate lung mononuclear cells, the single cell suspensions of lung tissue were cultured at 37°C in a humidified atmosphere containing 5% CO<sub>2</sub> for 24 hours. After 24 hours, culture medium was removed, plates were washed with PBS, and then adherent cells were harvested. More than 95% of the cells were recognized as monocytes by staining with Diff-Quik (Siemens, Germany). The cells were lysed by freezing and thawing to extract proteins and then frozen for further analysis.

### Quantification of cytokines using ELISA

IL-4, IL-5, IL-13, TNF- $\alpha$ , IL-1 $\beta$ , IL-6, IL-33, and IL-25 were measured using ELISA Kits (R&D Systems Inc.) according to the manufacturer's instructions. All samples were tested in duplicate.

### Flow cytometry

Cells were stained with F4/80, CD11b, CD11c, CD45, CD115, and Ly6c (eBioscience, San Diego, CA, USA). Cells were collected on a FACS caliber flow cytometer (BD Bioscience, Franklin Lakes, NJ, USA) and analyzed with FlowJo 8.3.3 software.

### Airway hyper reactivity to methacholine

Mice were anesthetized by an intraperitoneal injection of pentobarbital and 18-gauge metal needle was inserted into exposed trachea. Airway hyper reactivity responses were measured by a forced oscillation technique (FlexiVent, SCIREQ, Montreal, Canada). Airway resistance (R) was investigated after exposures of increasing methacholine concentration.

### RNA extraction and real-time RT PCR

RNA was extracted from lungs, blood, and monocytes using the RNeasy<sup>®</sup> Protect Mini Kit (QIAGEN, The Netherlands). RNA quantity and quality were determined using a NanoDrop 1000A spectrophotometer (NanoDrop Products, Wilmington, DE, USA). RNA was reverse transcribed to cDNA, and Taqman gene expression assays were used to detect TSLP (Mm00498739-m1 Tslp), arginase-1 (Mm00475988-m1 arg1), Ym-1 (Mm0065789-mH Chil3), TNF (Mm00443260-g1 Tnf), IL-1 $\beta$  (Mm01336189-m1 Il1b), and 18S RNA (Mm03928990-g1 Rn18s). Messenger RNA expression levels were normalized using 18S RNA expression.

### Statistical analysis

The data are shown as mean  $\pm$  standard deviation (SD). Analysis of variance (ANOVA) was used for multiple comparison of continuous variables. When there was a significant difference, the difference between each group was tested using Scheffe's test. All tests were 2-sided, and significance was set at 0.05.

## Results

### IL-33 was significantly increased in lungs during HDM-induced airway inflammation

We investigated the involvement of cytokines in HDM-treated mice. The cytokines in BALF were undetectable by ELISA (data not shown). The concentration of IL-33 in lung tissue was

significantly increased in HDM mice compared with control mice (Fig 1A). In contrast, IL-25 and TSLP in lung tissue were not increased in HDM mice (Fig 1B and 1C). Immunohistochemistry showed that IL-33-positive cells were located in the peribronchial regions of the lung. Bronchial epithelial cells, which were reported as a source of IL-33, were weakly stained (Fig 1D). The total positive cell counts in the IL-33-rich peribronchial regions were significantly increased in HDM mice compared with control mice (Fig 1E). The distribution and morphology of IL-33-positive cells by immunohistochemistry suggested that the cells were monocytes recruited from the bloodstream to the peribronchial regions. To identify monocytes in the lung, we examined the lung single cell suspensions by flow cytometry. After the exclusion of doublets and debris, leucocytes were separated using the pan-hematopoietic marker CD45. According to previous reports, Ly6c-positive monocytes were characterized by the expression of CD11b, Ly6c, and the absence of CD11c (Fig 2A) and alveolar macrophages by the expression of CD11c and the absence of CD11b (S1A Fig) [26]. Ly6c<sup>+</sup> monocytes were increased in lungs of HDM mice compared with control mice (Fig 2B and 2C). In contrast, alveolar macrophages were not significantly increased (S1B Fig).


### Ly6c-positive lung mononuclear cells could be a source of IL-33 in vivo and ex vivo

To demonstrate that IL-33 was present within monocytes, we used mononuclear cells isolated from lung. These cells were confirmed to be more than 95% monocytes using Diff-Quik (data not shown). Using immunofluorescence, IL-33-positive cells were merged with F4/80-positive cells to identify monocytes (Fig 3A). IL-33 levels in these cells were increased in HDM mice compared with control mice (Fig 3B). To clarify whether these cells were recruited from blood, the number of Ly6c-positive cells was determined by flow cytometry and was found to be significantly increased in HDM mice (Fig 3C and 3D). These results suggest that monocytes recruited to the lung could be one of the sources of IL-33.

### Clodronate-liposomes attenuated HDM-induced airway inflammation and airway hyper reactivity

To clarify whether the IL-33-producing cells were monocytes recruited to the lung, we used clodronate-liposomes *in vivo*. Clodronate is a bisphosphonate compound which causes apoptosis by intracellular delivery to macrophages via liposomes. The effect is specific to phagocytic cells such as macrophages [27]. In mice with airway inflammation the administration route is critical for the type of macrophage depleted in the lung. If clodronate was administered intratracheally, alveolar macrophages were depleted but not monocytes recruited from the bloodstream to the tissue. In contrast, if clodronate was administered intravenously, monocytes from the blood were depleted but not alveolar macrophages [23]. We investigated the effect of clodronate administered intratracheally to HDM-induced airway inflammation mice (Fig 4). More than 90% of monocytes were depleted after 24 hours and numbers were unchanged after 48 hours in the spleen. In bone marrow, 70% of monocytes were depleted after 24 hours of administration, but numbers were recovered after 48 hours (S2 Fig). Lymphocytes in the spleen were unchanged after administration of clodronate (data not shown).

Total and inflammatory cells including neutrophils and eosinophils, but not macrophages, in BALF were significantly decreased after clodronate treatment (Fig 5A). Airway hyper reactivity of HDM mice was significantly increased compared with control mice and this elevation was significantly attenuated by administration of clodronate (Fig 5B). Inflammatory cells and goblet cell hyperplasia were also inhibited by clodronate treatment compared with untreated HDM mice (Fig 5C). Th2 cytokines such as IL-4, IL-5, and IL-13 in lung tissues were also



**Fig 1. IL-33 was significantly increased in lungs of HDM mouse model.** Concentrations of (a) IL-33, (b) IL-25 in the lung tissue were measured by ELISA (n = 6 in each group). (c) Thymic stromal lymphopoietin (TSLP) level in the lung tissue was measured by real-time RT-PCR assay (n = 6 in each group), because TSLP protein concentration was below the detection level. (d) Immunohistochemistry (IHC)


examination of anti-normal goat IgG control and anti-IL-33 antibody in the lung section. Original magnifications were  $\times 200$  and  $\times 500$ . (e) The number of IL-33 positive cells by IHC (6 sections were counted by two pulmonologists).

doi:10.1371/journal.pone.0157571.g001

inhibited in clodronate-treated mice (Fig 5D–5F). However, TNF- $\alpha$ , IL-1 $\beta$  and IL-6 were not attenuated by administration of clodronate (S3A–S3C Fig). We next examined TNF and IL-1 $\beta$  expression as markers of M1 macrophages and arginase-1 and Ym1 as markers of M2 macrophages. Interestingly, M2 markers were decreased by clodronate treatment compared with HDM mice, while M1 markers were not affected by clodronate treatment (Fig 6A–6D). These results suggest that HDM-induced airway inflammation depends on recruited monocytes, which polarize the response toward the M2 phenotype.


### Monocytes recruited to lung are a source of IL-33

IL-33 in lung tissue was significantly decreased in clodronate-treated mice compared with HDM mice (Fig 7A). IL-33-positive cells in immunohistochemistry were clearly decreased in clodronate-treated mice (Fig 7B). The number of IL-33-positive cells was significantly decreased in the peribronchial regions of clodronate-treated mice compared with HDM mice



**Fig 2. Flow cytometry analysis of Ly6c-positive monocytes in lungs of HDM mice and control mice.** (a) Cells were identified from digested lungs, and after exclusion of doublets and debris, leukocytes were separated by CD45 staining. CD11c-negative cells were identified to investigate Ly6c-positive monocytes. (b) Flow cytometry analysis of Ly6c-positive monocytes (CD45<sup>+</sup>, CD11c<sup>-</sup>, CD11b<sup>+</sup>, Ly6c<sup>+</sup>) in lungs of HDM mice and control mice. (c) The percentage of Ly6c-positive monocytes in lung were compared in HDM mice and control mice (n = 6 in each group). \*\*P<0.01

doi:10.1371/journal.pone.0157571.g002



**Fig 3. Lung mononuclear cells which positively expressed Ly6c are a source of IL-33.** Lung mononuclear cells were collected from lung tissue as described in the text. Briefly, isolated single cells from lung tissue were resuspended at  $2.0 \times 10^6$  cells/ml in RPMI1640 medium supplemented with 10% FCS for 24 hours. Adherent cells were harvested and used as mononuclear cells in the experiments. (a) Immunofluorescence examination of lung mononuclear cells in HDM mice. Cells were stained with anti-F4/80 antibody (right upper panels) and anti-IL-33 antibody (left lower panels) and images were merged (right lower panels). Original magnification was  $\times 800$ . (b) Lung mononuclear cells were lysed by freezing and thawing to extract protein. Concentrations of IL-33 in the lung mononuclear cells were measured by ELISA ( $n = 6$  in each group). (c) Lung mononuclear cells were stained for Ly6c and analyzed by flow cytometry. Representative histograms of Ly6c<sup>+</sup> cells in control mice and HDM mice are shown. (d) The percentage of Ly6c<sup>+</sup> cells were counted by flow cytometry. \*\* $P < 0.01$ .


doi:10.1371/journal.pone.0157571.g003

(Fig 7C). These results suggest that clodronate attenuated IL-33 in HDM mice and that recruited monocytes, but not residential alveolar macrophages, were the sources of IL-33.

## Discussion

The results of this study demonstrated that monocytes recruited to lung are a potent source of IL-33 and enhance Th2 airway inflammation induced by HDM. Immunohistology showed the presence of intranuclear IL-33 in pulmonary monocytes. IL-33 was released from pulmonary






**Fig 4. The protocols for HDM-induced airway inflammation and clodronate treatment of mice.** Mice were sensitized intranasally with 25 µg HDM or vehicle on days 1, 8, and 15. Mice were challenged intranasally with 5 µg HDM on days 22, 23, and 24. Four hours later after final challenge, bronchoalveolar lavage (BAL) fluid and lung tissue were harvested for further analysis. Clophosome-A, liposomal clodronate, and plain control liposomes for clophosome (150µl / animal) were administered by intravenous injection on days 0, 7, 14, 21, 22, and 23. On the day when HDM was given, clodronate and control liposomes were administered 30 minutes before the HDM inoculation.

doi:10.1371/journal.pone.0157571.g004

mononuclear cells, which were shown to be mostly monocytes. Depletion of circulating monocytes by clodronate liposomes led to inhibition of HDM-induced airway inflammation, meaning that recruited monocytes were essential for airway inflammation. To our knowledge, this is the first report to identify that IL-33 from inflammatory monocytes recruited to the lung contributes to airway inflammation.

IL-33 has emerged as a danger signal that is rapidly released from various cells by cellular damage or stress [28]. Biologically active full length IL-33 was released into the extracellular space after cell necrosis or mechanical injury [29]. IL-33 induces the production of Th2 cytokines such as IL-5 and IL-13 through activation of ILC2s, leading to allergic inflammation [11]. Bronchial epithelial cells were reported as a major source of IL-33 in previous studies [30,31]. In airway diseases, IL-33 was shown to be released into the airways after stimulation with influenza viruses [32], alternaria [15], and papain [33]. These responses occurred in the early phase after stimulation because IL-33 concentration reached a maximum at 6 to 24 hours. Thus IL-33 is released via the innate immune system as an initial mediator in allergic airway inflammation following attack from exogenous allergens and pathogens [14]. In this study, IL-33 in BALF was not detected in our mouse model, while markedly increased IL-33 was noted in lung homogenates from mice challenged with HDM. We found that IL-33 stained cells that shared morphological and immunological characteristics with macrophages were present around the bronchovascular areas (Fig 1D). We were able to detect only a few bronchial epithelial cells stained with IL-33 in the immunohistological findings. We considered the possibility that IL-33 within bronchial epithelial cells might have been released in an earlier phase of HDM sensitization.

Macrophages are the most abundant pulmonary immune cells and are responsible for antigen and allergen recognition and presentation [34,35]. Alveolar macrophages in patients with severe asthma were decreased in LPS-responsiveness that is manifested by defective apoptotic cell uptake and reduced secretion of inflammatory mediators [36]. Pulmonary macrophages have also been reported to be associated with severe asthma characterized by glucocorticoid-resistance and airway remodeling [10]. Li and colleagues reported that macrophages are




**Fig 5. Clodronate-liposome attenuated the HDM induced airway inflammation, airway hyper reactivity and production of Th2 cytokines.** (a) Total and differential cell counts in bronchoalveolar lavage fluids (n = 6 in each group). (b) Airway hyper reactivity was measured by assaying airway resistance during graded concentration of methacholine (n = 6 in each group). \*p<0.05 HDM+emp. lipo group compared with PBS+emp. lipo group. ##p<0.05 HDM+CL. lipo group compared with HDM+emp. lipo group. (c) Histological examination of airway inflammation. Sections were stained with H&E (upper panels) or PAS (lower panels). Original magnification was × 200. Empty liposome control (emp. lipo) and clodronate liposome (CL) results are shown. Concentrations of (d) IL-4, (e) IL-5, and (f) IL-13 in the lung tissue were measured by ELISA (n = 6 in each group). \*\*P<0.01, \*P<0.05

doi:10.1371/journal.pone.0157571.g005


involved in glucocorticoid-resistant airway hyperresponsiveness (AHR) through a MyD88-dependent mechanism [37].

We demonstrated that clodronate treatment inhibited airway inflammation and IL-33 release, which indicated that inflammatory monocytes recruited to the lung had a crucial role in HDM-induced airway inflammation. Ly6c has been reported as a marker of inflammatory monocytes in blood that cause tissue inflammation [20,38,39]. We showed that Ly6c-positive



**Fig 6. Lung mononuclear cells expressed M2 markers but not M1 markers.** The expression of TNF (a), IL-1β (b), arginase-1 (c), and YM-1 (d) levels in the lung mononuclear cells was measured by real-time RT-PCR assay.

doi:10.1371/journal.pone.0157571.g006



**Fig 7. Clodronate attenuated the IL-33 in lungs of HDM mice.** (a) Concentrations of IL-33 in the lung tissue were measured by ELISA in 3 groups (intranasal PBS and intravenous empty liposomes, HDM and empty liposomes, HDM and clodronate liposomes:  $n = 6$  in each group). (b) Immunohistochemistry (IHC) examination of IL-33 in the lung sections. Original magnification was  $\times 200$ . (c) IL-33-positive cell counts by IHC (6 sections were counted by two pulmonologists). **\*\*** $P < 0.01$ .

doi:10.1371/journal.pone.0157571.g007

monocytes, but not alveolar macrophages, were increased in lungs of HDM mice. These data suggest that recruited monocytes were more important in HDM-induced airway inflammation than alveolar macrophages.

We also showed that the expression of Ly6c in lung mononuclear cells was increased in HDM mice compared with control mice (Fig 2C and 2D). These results suggest that monocytes recruited to lung were one of the sources of IL-33.

Macrophages have been classified into M1 and M2 subtypes based on their expression of cytokines, especially chemokines, and other specific markers [40]. M1 macrophages are characterized by a pro-inflammatory phenotype that promotes the Th1 immune response and anti-tumor activity. M2 macrophages play a role in regulatory functions in tissue repair, remodeling, and promotion of the Th2 immune response [41–43]. We showed that M2 macrophage markers decreased following clodronate treatment compared to HDM mice, suggesting that recruited monocytes were involved in HDM-airway inflammation. These data are consistent with previously reported functions of M2 macrophages in allergic disease [40].

We used a first generation bisphosphonate, clodronate, in liposomes, to deplete macrophages in this study [27]. A large dose of a second generation bisphosphonate, alendronate, attenuated eosinophilic airway inflammation induced by ovalbumin in mice [44]. In this study, we performed airway sensitization and challenge with HDM in mice. Our data indicated that circulating cells such as monocytes had a crucial role in airway inflammation, suggesting that bronchial asthma involves not only the airways but also the systemic circulation. Targeted therapy against systemic monocytes might be a potential treatment for severe, treatment-resistant asthma. It is expected that new agents with higher specificity for monocytes will be developed.

In conclusion, IL-33 derived from monocytes recruited to the lung may contribute to HDM-induced airway inflammation in vivo and in vitro. This study describes, to our knowledge for the first time, mechanisms of IL-33-related airway inflammation through recruited monocytes.

## Supporting Information

**S1 Fig. Flow cytometry analysis of alveolar macrophages were not increased in lungs of HDM mouse model.** Lung cells from control and HDM mice were separated by flow cytometry and examined. (a) Lung single cells were examined after the exclusion of doublets and debris and CD45<sup>+</sup> cells were isolated. (b) Alveolar macrophages (CD45<sup>+</sup>, CD11b<sup>-</sup>, CD11c<sup>+</sup>) in lungs of HDM and control mice were compared.  
(TIF)

**S2 Fig. Depletion of bone marrow monocytes and splenic monocytes after administration of clodronate liposome in a mouse model.** (a) Depletion of CD11b<sup>+</sup> F4/80<sup>+</sup> bone marrow monocytes and CD115<sup>+</sup> F4/80<sup>+</sup> splenic monocytes at 24 hours after intravenous administration of clodronate liposomes compared to mice treated with control liposomes, analyzed by flow cytometry. Bone marrow cells were isolated from femur. Spleen was digested and separated into single cells. (b) The kinetics and compartment-specific depletion of monocytes after clodronate liposome administration, determined by flow cytometry. Solid line depicts percentage of CD11b<sup>+</sup> F4/80<sup>+</sup> bone marrow monocytes. Dashed line depicts percentage of CD115<sup>+</sup> F4/80<sup>+</sup> splenic monocytes. Data points are means of 3 mice at each time point.  
(TIF)

**S3 Fig. Clodronate did not attenuate the TNF- $\alpha$ , IL-1 $\beta$  and IL-6 in lung of HDM mice.** Concentrations of (a) TNF- $\alpha$ , (b) IL-1 $\beta$  and (c) IL-6 in lung tissue were measured by ELISA (n = 6 in each group).  
(TIF)

## Author Contributions

Conceived and designed the experiments: KT SH. Performed the experiments: HT GK. Analyzed the data: HT KT. Contributed reagents/materials/analysis tools: KK. Wrote the paper: HT KT SK NSA.

## References

1. Tattersfield AE, Knox AJ, Britton JR, Hall IP (2002) Asthma. *Lancet* 360: 1313–1322. PMID: [12414223](#)
2. Lambrecht BN, Hammad H (2015) The immunology of asthma. *Nat Immunol* 16: 45–56. doi: [10.1038/ni.3049](#) PMID: [25521684](#)
3. Fahy JV (2015) Type 2 inflammation in asthma—present in most, absent in many. *Nat Rev Immunol* 15: 57–65. doi: [10.1038/nri3786](#) PMID: [25534623](#)
4. Gudbjartsson DF, Bjornsdottir US, Halapi E, Helgadóttir A, Sulem P, Jonsdóttir GM, et al. (2009) Sequence variants affecting eosinophil numbers associate with asthma and myocardial infarction. *Nat Genet* 41: 342–347. doi: [10.1038/ng.323](#) PMID: [19198610](#)
5. Torgerson DG, Ampleford EJ, Chiu GY, Gauderman WJ, Gignoux CR, Graves PE, et al. (2011) Meta-analysis of genome-wide association studies of asthma in ethnically diverse North American populations. *Nat Genet* 43: 887–892. doi: [10.1038/ng.888](#) PMID: [21804549](#)
6. Moffatt MF, Gut IG, Demenais F, Strachan DP, Bouzigon E, Heath S, et al. (2010) A large-scale, consortium-based genome-wide association study of asthma. *N Engl J Med* 363: 1211–1221. doi: [10.1056/NEJMoa0906312](#) PMID: [20860503](#)
7. Bonnelykke K, Sleiman P, Nielsen K, Kreiner-Moller E, Mercader JM, Belgrave D, et al. (2014) A genome-wide association study identifies CDHR3 as a susceptibility locus for early childhood asthma with severe exacerbations. *Nat Genet* 46: 51–55. doi: [10.1038/ng.2830](#) PMID: [24241537](#)
8. Guo Z, Wu J, Zhao J, Liu F, Chen Y, Bi L, et al. (2014) IL-33 promotes airway remodeling and is a marker of asthma disease severity. *J Asthma* 51: 863–869. doi: [10.3109/02770903.2014.921196](#) PMID: [24796648](#)
9. Saglani S, Lui S, Ullmann N, Campbell GA, Sherburn RT, Mathie SA, et al. (2013) IL-33 promotes airway remodeling in pediatric patients with severe steroid-resistant asthma. *J Allergy Clin Immunol* 132: 676–685 e613. doi: [10.1016/j.jaci.2013.04.012](#) PMID: [23759184](#)
10. Yang M, Kumar RK, Hansbro PM, Foster PS (2012) Emerging roles of pulmonary macrophages in driving the development of severe asthma. *J Leukoc Biol* 91: 557–569. doi: [10.1189/jlb.0711357](#) PMID: [22293472](#)
11. Schmitz J, Owyang A, Oldham E, Song Y, Murphy E, McClanahan TK, et al. (2005) IL-33, an interleukin-1-like cytokine that signals via the IL-1 receptor-related protein ST2 and induces T helper type 2-associated cytokines. *Immunity* 23: 479–490. PMID: [16286016](#)
12. Moro K, Yamada T, Tanabe M, Takeuchi T, Ikawa T, Kawamoto H, et al. (2010) Innate production of T(H)2 cytokines by adipose tissue-associated c-Kit(+)Sca-1(+) lymphoid cells. *Nature* 463: 540–544. doi: [10.1038/nature08636](#) PMID: [20023630](#)
13. Galli SJ, Tsai M, Piliponsky AM (2008) The development of allergic inflammation. *Nature* 454: 445–454. doi: [10.1038/nature07204](#) PMID: [18650915](#)
14. Oboki K, Ohno T, Kajiwara N, Arae K, Morita H, Ishii A, et al. (2010) IL-33 is a crucial amplifier of innate rather than acquired immunity. *Proc Natl Acad Sci U S A* 107: 18581–18586. doi: [10.1073/pnas.1003059107](#) PMID: [20937871](#)
15. Kouzaki H, Iijima K, Kobayashi T, O'Grady SM, Kita H (2011) The danger signal, extracellular ATP, is a sensor for an airborne allergen and triggers IL-33 release and innate Th2-type responses. *J Immunol* 186: 4375–4387. doi: [10.4049/jimmunol.1003020](#) PMID: [21357533](#)
16. Llop-Guevara A, Chu DK, Walker TD, Goncharova S, Fattouh R, Silver JS, et al. (2014) A GM-CSF/IL-33 pathway facilitates allergic airway responses to sub-threshold house dust mite exposure. *PLoS One* 9: e88714. doi: [10.1371/journal.pone.0088714](#) PMID: [24551140](#)
17. Hsu CL, Bryce PJ (2012) Inducible IL-33 expression by mast cells is regulated by a calcium-dependent pathway. *J Immunol* 189: 3421–3429. PMID: [22922818](#)
18. Yanagawa Y, Matsumoto M, Togashi H (2011) Adrenoceptor-mediated enhancement of interleukin-33 production by dendritic cells. *Brain Behav Immun* 25: 1427–1433. doi: [10.1016/j.bbi.2011.04.012](#) PMID: [21536121](#)
19. Martinez-Martinez E, Miana M, Jurado-Lopez R, Rousseau E, Rossignol P, Zannad F, et al. (2013) A role for soluble ST2 in vascular remodeling associated with obesity in rats. *PLoS One* 8: e79176. doi: [10.1371/journal.pone.0079176](#) PMID: [24265755](#)

20. Epelman S, Lavine KJ, Randolph GJ (2014) Origin and functions of tissue macrophages. *Immunity* 41: 21–35. doi: [10.1016/j.immuni.2014.06.013](https://doi.org/10.1016/j.immuni.2014.06.013) PMID: [25035951](https://pubmed.ncbi.nlm.nih.gov/25035951/)
21. Serbina NV, Jia T, Hohl TM, Pamer EG (2008) Monocyte-mediated defense against microbial pathogens. *Annu Rev Immunol* 26: 421–452. doi: [10.1146/annurev.immunol.26.021607.090326](https://doi.org/10.1146/annurev.immunol.26.021607.090326) PMID: [18303997](https://pubmed.ncbi.nlm.nih.gov/18303997/)
22. Landsman L, Jung S (2007) Lung macrophages serve as obligatory intermediate between blood monocytes and alveolar macrophages. *J Immunol* 179: 3488–3494. PMID: [17785782](https://pubmed.ncbi.nlm.nih.gov/17785782/)
23. Zaslona Z, Przybranowski S, Wilke C, van Rooijen N, Teitz-Tennenbaum S, Osterholzer JJ, et al. (2014) Resident alveolar macrophages suppress, whereas recruited monocytes promote, allergic lung inflammation in murine models of asthma. *J Immunol* 193: 4245–4253. doi: [10.4049/jimmunol.1400580](https://doi.org/10.4049/jimmunol.1400580) PMID: [25225663](https://pubmed.ncbi.nlm.nih.gov/25225663/)
24. Kato G, Takahashi K, Tashiro H, Kurata K, Shirai H, Kimura S, et al. (2014) beta2 adrenergic agonist attenuates house dust mite-induced allergic airway inflammation through dendritic cells. *BMC Immunol* 15: 39. doi: [10.1186/s12865-014-0039-y](https://doi.org/10.1186/s12865-014-0039-y) PMID: [25359462](https://pubmed.ncbi.nlm.nih.gov/25359462/)
25. Takahashi K, Koga K, Linge HM, Zhang Y, Lin X, Metz CN, et al. (2009) Macrophage CD74 contributes to MIF-induced pulmonary inflammation. *Respir Res* 10: 33. doi: [10.1186/1465-9921-10-33](https://doi.org/10.1186/1465-9921-10-33) PMID: [19413900](https://pubmed.ncbi.nlm.nih.gov/19413900/)
26. Misharin AV, Morales-Nebreda L, Mutlu GM, Budinger GR, Perlman H (2013) Flow cytometric analysis of macrophages and dendritic cell subsets in the mouse lung. *Am J Respir Cell Mol Biol* 49: 503–510. doi: [10.1165/rcmb.2013-0086MA](https://doi.org/10.1165/rcmb.2013-0086MA) PMID: [23672262](https://pubmed.ncbi.nlm.nih.gov/23672262/)
27. van Rooijen N, Sanders A, van den Berg TK (1996) Apoptosis of macrophages induced by liposome-mediated intracellular delivery of clodronate and propamidine. *J Immunol Methods* 193: 93–99. PMID: [8690935](https://pubmed.ncbi.nlm.nih.gov/8690935/)
28. Cayrol C, Girard JP (2014) IL-33: an alarmin cytokine with crucial roles in innate immunity, inflammation and allergy. *Curr Opin Immunol* 31: 31–37. doi: [10.1016/j.coi.2014.09.004](https://doi.org/10.1016/j.coi.2014.09.004) PMID: [25278425](https://pubmed.ncbi.nlm.nih.gov/25278425/)
29. Luthi AU, Cullen SP, McNeela EA, Duriez PJ, Afonina IS, Sheridan C, et al. (2009) Suppression of interleukin-33 bioactivity through proteolysis by apoptotic caspases. *Immunity* 31: 84–98. doi: [10.1016/j.immuni.2009.05.007](https://doi.org/10.1016/j.immuni.2009.05.007) PMID: [19559631](https://pubmed.ncbi.nlm.nih.gov/19559631/)
30. Prefontaine D, Nadigel J, Chouiali F, Audusseau S, Semlali A, Chakir J, et al. (2010) Increased IL-33 expression by epithelial cells in bronchial asthma. *J Allergy Clin Immunol* 125: 752–754. doi: [10.1016/j.jaci.2009.12.935](https://doi.org/10.1016/j.jaci.2009.12.935) PMID: [20153038](https://pubmed.ncbi.nlm.nih.gov/20153038/)
31. Lambrecht BN, Hammad H (2014) Allergens and the airway epithelium response: gateway to allergic sensitization. *J Allergy Clin Immunol* 134: 499–507. doi: [10.1016/j.jaci.2014.06.036](https://doi.org/10.1016/j.jaci.2014.06.036) PMID: [25171864](https://pubmed.ncbi.nlm.nih.gov/25171864/)
32. Chang YJ, Kim HY, Albacker LA, Baumgarth N, McKenzie AN, Smith DE, et al. (2011) Innate lymphoid cells mediate influenza-induced airway hyper-reactivity independently of adaptive immunity. *Nat Immunol* 12: 631–638. doi: [10.1038/ni.2045](https://doi.org/10.1038/ni.2045) PMID: [21623379](https://pubmed.ncbi.nlm.nih.gov/21623379/)
33. Kamijo S, Takeda H, Tokura T, Suzuki M, Inui K, Hara M, et al. (2013) IL-33-mediated innate response and adaptive immune cells contribute to maximum responses of protease allergen-induced allergic airway inflammation. *J Immunol* 190: 4489–4499. doi: [10.4049/jimmunol.1201212](https://doi.org/10.4049/jimmunol.1201212) PMID: [23547117](https://pubmed.ncbi.nlm.nih.gov/23547117/)
34. Pappas K, Papaioannou AI, Kostikas K, Tzanakis N (2013) The role of macrophages in obstructive airways disease: chronic obstructive pulmonary disease and asthma. *Cytokine* 64: 613–625. doi: [10.1016/j.cyto.2013.09.010](https://doi.org/10.1016/j.cyto.2013.09.010) PMID: [24084332](https://pubmed.ncbi.nlm.nih.gov/24084332/)
35. Donnelly LE, Barnes PJ (2012) Defective phagocytosis in airways disease. *Chest* 141: 1055–1062. doi: [10.1378/chest.11-2348](https://doi.org/10.1378/chest.11-2348) PMID: [22474147](https://pubmed.ncbi.nlm.nih.gov/22474147/)
36. Huynh ML, Malcolm KC, Kotaru C, Tilstra JA, Westcott JY, Fadok VA, et al. (2005) Defective apoptotic cell phagocytosis attenuates prostaglandin E2 and 15-hydroxyeicosatetraenoic acid in severe asthma alveolar macrophages. *Am J Respir Crit Care Med* 172: 972–979. PMID: [16040786](https://pubmed.ncbi.nlm.nih.gov/16040786/)
37. Li JJ, Wang W, Baines KJ, Bowden NA, Hansbro PM, Gibson PG, et al. (2010) IL-27/IFN-gamma induce MyD88-dependent steroid-resistant airway hyperresponsiveness by inhibiting glucocorticoid signaling in macrophages. *J Immunol* 185: 4401–4409. doi: [10.4049/jimmunol.1001039](https://doi.org/10.4049/jimmunol.1001039) PMID: [20817868](https://pubmed.ncbi.nlm.nih.gov/20817868/)
38. Chen L, Zhang Z, Barletta KE, Burdick MD, Mehrad B (2013) Heterogeneity of lung mononuclear phagocytes during pneumonia: contribution of chemokine receptors. *Am J Physiol Lung Cell Mol Physiol* 305: L702–711. doi: [10.1152/ajplung.00194.2013](https://doi.org/10.1152/ajplung.00194.2013) PMID: [24056971](https://pubmed.ncbi.nlm.nih.gov/24056971/)
39. Gordon S, Taylor PR (2005) Monocyte and macrophage heterogeneity. *Nat Rev Immunol* 5: 953–964. PMID: [16322748](https://pubmed.ncbi.nlm.nih.gov/16322748/)
40. Murray PJ, Allen JE, Biswas SK, Fisher EA, Gilroy DW, Goerdt S, et al. (2014) Macrophage activation and polarization: nomenclature and experimental guidelines. *Immunity* 41: 14–20. doi: [10.1016/j.immuni.2014.06.008](https://doi.org/10.1016/j.immuni.2014.06.008) PMID: [25035950](https://pubmed.ncbi.nlm.nih.gov/25035950/)

41. Wynn TA, Barron L (2010) Macrophages: master regulators of inflammation and fibrosis. *Semin Liver Dis* 30: 245–257. doi: [10.1055/s-0030-1255354](https://doi.org/10.1055/s-0030-1255354) PMID: [20665377](https://pubmed.ncbi.nlm.nih.gov/20665377/)
42. Barros MH, Hauck F, Dreyer JH, Kempkes B, Niedobitek G (2013) Macrophage polarisation: an immunohistochemical approach for identifying M1 and M2 macrophages. *PLoS One* 8: e80908. doi: [10.1371/journal.pone.0080908](https://doi.org/10.1371/journal.pone.0080908) PMID: [24260507](https://pubmed.ncbi.nlm.nih.gov/24260507/)
43. Gordon S, Martinez FO (2010) Alternative activation of macrophages: mechanism and functions. *Immunity* 32: 593–604. doi: [10.1016/j.immuni.2010.05.007](https://doi.org/10.1016/j.immuni.2010.05.007) PMID: [20510870](https://pubmed.ncbi.nlm.nih.gov/20510870/)
44. Sasaki O, Imamura M, Yamazumi Y, Harada H, Matsumoto T, Okunishi K, et al. (2013) Alendronate attenuates eosinophilic airway inflammation associated with suppression of Th2 cytokines, Th17 cytokines, and eotaxin-2. *J Immunol* 191: 2879–2889. doi: [10.4049/jimmunol.1300460](https://doi.org/10.4049/jimmunol.1300460) PMID: [23935198](https://pubmed.ncbi.nlm.nih.gov/23935198/)