



Osteochondral Allograft Reconstruction of the Tibia Plateau for Posttraumatic Defects—A Novel Computer-Assisted Method Using 3D Preoperative Planning and Patient-Specific Instrumentation

Martin Zaleski, MD¹ Sandro Hodel, MD¹ Philipp Fürnstahl² Lazaros Vlachopoulos, PD, MD¹
Sandro F. Fucentese, PD, MD¹

¹Department of Orthopaedics, Balgrist University Hospital, University of Zurich, Zurich, Switzerland

²Research in Orthopedic Computer Science, Balgrist University Hospital, University of Zurich, Zurich, Switzerland

Address for correspondence Martin Zaleski, MD, Department of Orthopaedics, Balgrist University Hospital, University of Zurich, Forchstrasse 340, 8008 Zurich, Switzerland (e-mail: Martin.Zaleski@balgrist.ch).

Surg J (NY) 2021;7:e289–e296.

Abstract

Background Surgical treatment of posttraumatic defects of the knee joint is challenging. Osteochondral allograft reconstruction (OCAR) is an accepted procedure to restore the joint congruity and for pain relief, particularly in the younger population. Preoperative three-dimensional (3D) planning and patient-specific instrumentation (PSI) are well accepted for the treatment of posttraumatic deformities for several pathologies. The aim of this case report was to provide a guideline and detailed description of the preoperative 3D planning and the intraoperative navigation using PSI in OCAR for posttraumatic defects of the tibia plateau. We present the clinical radiographic results of a patient who was operated with this new technique with a 3.5-year follow-up.

Materials and Methods 3D-triangular surface models are created based on preoperative computer tomography (CT) of the injured side and the contralateral side. We describe the preoperative 3D-analysis and planning for the reconstruction with an osteochondral allograft (OCA) of the tibia plateau. We describe the PSI as well as cutting and reduction techniques to show the intraoperative possibilities in posttraumatic knee reconstructions with OCA.

Results Our clinical results indicate that 3D-assisted osteotomy and OCAR for posttraumatic defects of the knee may be beneficial and feasible. We illustrate the planning and execution of the osteotomy for the tibia and the allograft using PSI, allowing an accurate anatomical restoration of the joint congruency.

Discussion With 3D-planning and PSI the OCAR might be more precise compared with conventional methods. It could improve the reproducibility and might allow less experienced surgeons to perform the precise and technically challenging osteotomy

Keywords

- ▶ knee
- ▶ tibia plateau
- ▶ trauma
- ▶ patient-specific instrumentation
- ▶ allograft reconstruction

received
May 17, 2020
accepted after revision
July 23, 2021

DOI <https://doi.org/10.1055/s-0041-1735602>.
ISSN 2378-5128.

© 2021. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited. (<https://creativecommons.org/licenses/by/4.0/>)
Thieme Medical Publishers, Inc., 333 Seventh Avenue, 18th Floor, New York, NY 10001, USA

cuts of the tibia and the allograft. Further, this technique might shorten operating time because time consuming intraoperative steps such as defining the osteotomy cuts of the tibia and the allograft during surgery are not necessary.

Conclusion OCAR of the tibia plateau for posttraumatic defects with 3D preoperative planning and PSI might allow for the accurate restoration of anatomical joint congruency, improve the reproducibility of surgical technique, and shorten the surgery time.

Fractures of the proximal tibia occur in 27 per 100.000 per year and are associated with high-energy trauma, especially in younger patients.¹ The disease burden of posttraumatic osteoarthritis (PTOA) is estimated to be 12% of all symptomatic osteoarthritis (OA) of the hip, knee, and ankle.² PTOA of the knee occurs at high rates after intra-articular and extra-articular fracture of either the distal femur or the proximal tibia, with the incidence in the literature ranging from 21 to 44%, and can also be seen after ligamentous, meniscal, and high-impact injuries.³⁻⁵ It has been shown that total knee arthroplasty (TKA) for PTOA is associated with a lower function, a lower quality of life, and a lower survival rate than for primary OA.⁶ Joint-preserving procedures remain the treatment of choice for young patients, because of critical results for UKA (unicondylar knee arthroplasty) and TKA in the long-term with high revision rates.⁷ Several options exist to address posttraumatic defects of the knee directly after trauma. Fractures can be treated conservatively for minimally displaced fragments, otherwise surgical management is recommended.^{8,9} Surgical possibilities are open reduction and internal fixation (ORIF), external fixation, or arthroscopically assisted osteosynthesis.¹⁰⁻¹⁶ However, operative reconstruction management of posttraumatic larger damages in the knee joint is challenging and may be associated with malunion if reduction is imprecise, which can lead to following operations and progressive arthrosis.^{8,17} Joint preserving options like osteochondral autograft transplantation surgery, autologous chondrocyte implantation, autologous matrix-induced chondrogenesis, and bulk allograft, depending on the size and location of the defect are supposed to be an alternative.¹⁸ Although these procedures have been implemented with varying degrees of success, no consensus exists on the gold-standard treatment. Additionally, the reconstruction after traumatic damages around the knee using osteochondral allograft is an established technique procedure. Previous studies demonstrated the benefit of fresh OCA in the reconstruction of posttraumatic defects of the knee with satisfying long-term results for the tibial plateau.^{19,20} The superiority of survival in vivo with higher chondrocyte viability of fresh OCA compared with frozen OCA was shown.^{19,20} However, big posttraumatic defects of the knee are a complex three-dimensional (3D) problem and the exact restoration of the joint line, anatomical axis, and tibial slope are challenging for the treating surgeon. The benefit of computer-assisted corrective osteotomies or allograft reconstruction in tumor surgery around the knee has

already been emphasized.^{21,22} The main advantage of computer-assisted surgery is the precise 3D analysis of the deformity. Therefore, a facilitated surgical planning of the osteochondral allograft reconstruction (OCAR) with accurate reconstruction results can be expected, using 3D computer-assisted planning. To our knowledge the use of patient-specific instrumentation (PSI) to treat posttraumatic defects of the tibia plateau with an OCA has not been reported so far. The aim of this case report is to present a step-by-step guideline for posttraumatic OCAR of the tibia plateau using a novel computer-assisted method with 3D preoperative planning and PSI.

Patients, Materials, and Methods

Patients

Informed consent for the publication of this case report and the use of the photographs was obtained from the patient. The local ethical committee approved this study (Zurich Cantonal Ethics Commission, KEK-ZH 2015-0186). The case (► **Fig. 1**) is a 31-year-old female office employee who suffered at the age of 28 a lateral tibial plateau fracture (Schatzker classification Type II) of the left knee after high-energy trauma. Initially she received an ORIF with lateral and posterior plate via lateral access in an external hospital. Nine months later she was assigned to us for a second opinion by persistent knee pain. We diagnosed a malunion through a computed tomography (CT) and performed 3 months later the removal of the osteosynthesis material. An infection was ruled out after taking intraoperative samples. Four months later 3D computer-assisted planning was performed and 7 months later OCAR with the technique described below was accomplished.

Materials and Methods

Preoperative 3D Deformity Analysis and Planning

The analysis of the deformity and the planned reconstruction were performed based on a reconstruction template. This approach has two main advantages. Namely, additional information is available about the ideal size of the allograft need. Furthermore, the time to complete the preoperative planning is reduced when the OCA is delivered (Neutromedics AG Ortho-Biologics & Implants, Cham, Switzerland), as the allograft adjustment has only to be performed. Ideally the delivery of the allograft and the production of the guides



Fig. 1 (a) Initial posttraumatic X-rays and CT (b) 9 months after ORIF (c) 4 months after hardware removal. CT, computer tomography; ORIF, open reduction and internal fixation.

should be performed quickly. A 3D triangular surface model of the pathological and the contralateral side is generated based on CT scans (slice thickness, 1 mm, 120 kv: Philips Brilliance 40 CT, Philips Healthcare, Eindhoven, the Netherlands) using thresholding, region growing, and the

marching cubes algorithm to identify the cortical bone layer and for separating the tibia from the surrounding bone anatomy as previously described.^{21,23–25} The contralateral tibia is supposed to be an accurate three-dimensional reconstruction template,²⁶ in patients without a history of trauma



Fig. 2 Reference guide with drill sleeves and inserted K-wires. Note the additional wing to improve the contact with the bone.

or pathological condition. Therefore, it is currently our preferred template. The 3D models are imported into the planning software Computer Assisted Surgery Planning Application (CASPA) (Balgrist CARD AG, Zürich, Switzerland). The model of the contralateral tibia is mirrored and subsequently aligned to the pathologic model using a surface registration algorithm. As in similar approaches, the iterative closest point surface registration algorithm is used for bone alignment.^{27,28}

This method superimposes the undeformed regions of the bone surfaces in an automatic fashion by minimizing the sum of quadratic distances between surface points.^{29,30} Defining the resection margins is possible by visualizing the exact 3D relation of the bone defect. After defining the resection planes and the fixation device, patient-specific guides can be designed to transfer the preoperative plan to the surgery.

Reference and Osteotomy Guides

The reference guides are used to allow the later positioning of the osteotomy guides as accurate as possible (►Fig. 2). They are serving as a registration tool between the 3D planning and the intraoperative situation. The reference guides correspond to the negative contour of the bony anatomy of the tibia. To control the fitting accuracy, a 3D printout of the native bone is regularly used. Meticulous intraoperative




Fig. 3 Osteotomy guide (left). The planned osteotomy of the lateral epicondyle is marked in purple (right).

positioning of these guides is critical to define the correct osteotomy position that has been preplanned. To avoid a soft tissue interruption the whole periosteum has to be removed. This is particularly possible in the area of the bone that has to be resected without influencing the vascularization of the remaining proximal tibia.

The reference guide has an additional wing to allow a maximum contact with prominent bony anatomy and to give additional rotational and translational stability. In earlier approaches it was shown that these wings lead to more precise osteotomies.³¹ Two K-wires (2.5 or 3.0 mm) are drilled through the predefined drill sleeves attached to the reference guide.

A guide design which constrains the saw blade is normally used as described in earlier approaches.²⁴ It is important for the osteotomy to calculate the corresponding cuts with an offset to consider the offset. The reference guide is removed while the 2 K-wires are left in place and the osteotomy guide is positioned over the 2 K-wires. After confirmation of the adequate fit of the osteotomy guide using the 3D-printed models of the preoperative planning, definitive osteotomy is performed (►Figs. 3, 4).

Preoperative Planning of the Osteochondral Allograft

The size of the OCA needs to be considered for the substitute of the bone defect. In an ideal situation, an equivalent bone (i.e., same bone in the same dimensions) should be used. The advantage of using an equivalent allograft bone is that the allograft needed for insertion can be used in one single piece and complex constructs, which reduce stability, can be avoided. A 3D surface model is created from CT scans of the fresh OCA, similar to what was previously described (►Fig. 5).

Allograft adjustment guides have to be prepared for the preparation of the allograft to customize it to the required shape. Therefore, the guides need to be applied on the allograft, as shown (►Figs. 6, 7). They have to be fixed by K-wires through the predefined drill sleeves attached. With the cutting slits and the drill sleeves, the saw blade is




Fig. 4 Intraoperative view from lateral of the resected lateral epicondyle after osteotomy with osteotomy guide.




Fig. 7 Allograft adjustment guide for the horizontal cut with view from lateral.




Fig. 5 3D surface model of the OCA before preparation. The planned allograft for implantation is marked in purple. 3D, three-dimensional; OCA, osteochondral allograft.

constrained to the planned osteotomy planes, and proper customizing of the allograft will be achieved. The consideration of the offcut in the preparation of the allograft is important as well.

Insertion and Surgical Technique

The intraoperative insertion can be performed either by manipulating the fragments directly, or indirectly using the final implant as a reduction guide. A K-wire-based-reduction guide or a fragment reduction guide as previously described might be used for the direct reduction of the prepared allograft.²⁴ For this purpose, one part of the undersurface of the guide body matches to the fragment surface in its reduced position and the other part to the reference fragment. Alternatively, the planned screw holes of the definitive plate fixation can be predrilled in the allograft. This allows an intraoperative reduction through the plate holes. In addition, a combination of these reduction guides might be applicable depending on the individual case. Using an anterior midline approach allows adequate visualization of the joint and the proximal tibia. The meniscus was preserved. An osteotomy of the medial and lateral epicondyle should be considered to allow better visualization of the joint for allograft insertion. After completion of the cuts in the tibia and the allograft, the allograft is fitted in the defect. Adequate plate fit and screw orientation for fixation of the allograft should be considered, when choosing the appropriate fixation plate. Fixation of the allograft is then performed using the preoperative planned plate (► **Fig. 8**). The soft tissue was fixed.




Fig. 6 Allograft adjustment guides for the vertical cut (a), for horizontal cut with view from anterior (b) and from lateral (c).




Fig. 8 Intraoperative allograft insertion (*left*). Plate fixation (*right*).

Results

The postoperative aftercare for the presented patient after OCAR was mobilization on walking sticks with partial load of 5 kg for 8 weeks. The patient initially wore a blocked brace for 2 weeks but with free flexion and extension out of the brace. After 8 weeks the load was subsequently increased over

6 weeks to full load. The patient had a low stress pain and a stable joint in the clinical examination 15 months after operation. She was able to work full time as office employee. Flexion/extension was 135/0/0 degrees. The CT 15 months after operation showed progressive consolidation of the tibial osteotomy gap and in the conventional radiographs 3.5 years postoperative a regular position of the osteosynthesis material was visible (→ **Fig. 9**). In the clinical examination 2 and 3.5 years after operation the patient was still satisfied, had no pain at and was not limited in daily life.

Discussion

We present in this study, as far as we know, the first performance of an OCAR of the tibia plateau after a posttraumatic defect using preoperative 3D planning and PSI.

In the younger population a joint-preserving technique is necessary to restore the function and to avoid an early onset of




Fig. 9 (a) Postoperative X-rays after 2 months, (b) after 15 months, (c) CT-scans after 15 months, (d) X-rays after 3.5 years after posttraumatic OCAR using preoperative 3D planning and PSI. 3D, three-dimensional; CT, computer tomography; OCAR, osteochondral allograft reconstruction; PSI, patient-specific instrumentation.

PTOA which is associated with severe functional impairment.³⁻⁶ A simple 2D analysis based on conventional X-ray might not be sufficient to restore the anatomy of complex posttraumatic deformities adequately. OCAR of extensive posttraumatic defects of the knee joint is a complex 3D technical challenge which requires considerable experience of the attending surgeon to reconstruct a precise anatomical joint congruency. Excellent accuracy of 3D planning has been proven for several anatomical locations.^{24,29,32,33} In knee surgery, its indication has been shown useful for tumor surgery or secondary interventions of malunions after osteotomies.^{21,22} This novel computer-assisted method using 3D preoperative planning and PSI in OCAR of the tibia plateau for posttraumatic defects might have several advantages compared with the conventional surgical methods. So far one of the main factors related to failure after OCAR is malalignment, which can lead to higher weight stress for the graft, a chondral destruction, a collapse of the allograft, residual pain, and dysfunction of the knee joint.³⁴ With 3D preoperative planning and PSI the reconstruction of the anatomical alignment including the joint line, leg axis, and tibial slope might be more precise. In addition, this novel technology could improve the reproducibility, because with preoperative 3D planning and PSI also less experienced surgeons might perform precisely the technical challenging osteotomy cuts of the tibia and the allograft. Another advantage might be a shortened operating time, because of previously performed preoperative 3D planning and PSI time consuming intraoperative steps such as defining the osteotomy cuts of the tibia and the allograft are not necessary. The availability of the allograft is a limiting factor due to the fact that it has to be ordered in advance before a surgery can be accomplished. With this new technique the establishment of an international 3D-database with an allograft storage would be a possibility for the future improving the graft availability. After CT scans of a patient with a posttraumatic defect had been made in an external hospital the 3D planning might be performed in the 3D database where the exactly calculated size of the required allograft could be prepared and delivered. Optional could be the preparation for PSI for tibia osteotomy as well. This might enable the younger patients with posttraumatic defects of the tibia to receive a challenging reconstruction outside of specialized centers, which have a stock of allografts. However, this new operative assessment should be in focus of further studies and analysis with larger number of patients and the clinical results of this new technology have to be compared with the conventional surgical methods.

Conclusion

OCAR of the tibia plateau for posttraumatic defects with 3D preoperative planning and PSI might restore the anatomical joint congruency accurate, improve the reproducibility, and shorten the surgery time.

Funding
None.

Conflict of Interest
None declared.

References

- Wennergren D, Bergdahl C, Ekelund J, Juto H, Sundfeldt M, Möller M. Epidemiology and incidence of tibia fractures in the Swedish Fracture Register. *Injury* 2018;49(11):2068–2074
- Brown TD, Johnston RC, Saltzman CL, Marsh JL, Buckwalter JA. Posttraumatic osteoarthritis: a first estimate of incidence, prevalence, and burden of disease. *J Orthop Trauma* 2006;20(10):739–744
- Papadopoulos EC, Parvizi J, Lai CH, Lewallen DG. Total knee arthroplasty following prior distal femoral fracture. *Knee* 2002;9(04):267–274
- Weiss NG, Parvizi J, Trousdale RT, Bryce RD, Lewallen DG. Total knee arthroplasty in patients with a prior fracture of the tibial plateau. *J Bone Joint Surg Am* 2003;85(02):218–221
- Weiss NG, Parvizi J, Hanssen AD, Trousdale RT, Lewallen DG. Total knee arthroplasty in post-traumatic arthrosis of the knee. *J Arthroplasty* 2003;18(3, suppl 1):23–26
- Lunenburg A, Parratte S, Gay A, Ollivier M, Garcia-Parra K, Argenson JN. Lower function, quality of life, and survival rate after total knee arthroplasty for posttraumatic arthritis than for primary arthritis. *Acta Orthop* 2015;86(02):189–194
- Parratte S, Boyer P, Piriou P, Argenson JN, Deschamps G, Massin PSFHG (French Hip and Knee Society) Total knee replacement following intra-articular malunion. *Orthop Traumatol Surg Res* 2011;97(Suppl 6):S118–S123
- Schatzker J, McBroom R, Bruce D. The tibial plateau fracture. The Toronto experience 1968–1975. *Clin Orthop Relat Res* 1979;(138):94–104
- Papagelopoulos PJ, Partsinevelos AA, Themistocleous GS, Mavrogenis AF, Korres DS, Soucacos PN. Complications after tibia plateau fracture surgery. *Injury* 2006;37(06):475–484
- Holzach P, Matter P, Minter J. Arthroscopically assisted treatment of lateral tibial plateau fractures in skiers: use of a cannulated reduction system. *J Orthop Trauma* 1994;8(04):273–281
- Chiu CH, Cheng CY, Tsai MC, et al. Arthroscopy-assisted reduction of posteromedial tibial plateau fractures with buttress plate and cannulated screw construct. *Arthroscopy* 2013;29(08):1346–1354
- Hung SS, Chao EK, Chan YS, et al. Arthroscopically assisted osteosynthesis for tibial plateau fractures. *J Trauma* 2003;54(02):356–363
- Dall'oca C, Maluta T, Lavini F, Bondi M, Micheloni GM, Bartolozzi P. Tibial plateau fractures: compared outcomes between ARIF and ORIF. *Stratag Trauma Limb Reconstr* 2012;7(03):163–175
- Musahl V, Tarkin I, Kobbe P, Tzioupis C, Siska PA, Pape HC. New trends and techniques in open reduction and internal fixation of fractures of the tibial plateau. *J Bone Joint Surg Br* 2009;91(04):426–433
- Manidakis N, Dosani A, Dimitriou R, Stengel D, Matthews S, Giannoudis P. Tibial plateau fractures: functional outcome and incidence of osteoarthritis in 125 cases. *Int Orthop* 2010;34(04):565–570
- Rademakers MV, Kerkhoffs GM, Siersevelt IN, Raaymakers EL, Marti RK. Operative treatment of 109 tibial plateau fractures: five- to 27-year follow-up results. *J Orthop Trauma* 2007;21(01):5–10
- Kettelkamp DB, Hillberry BM, Murrish DE, Heck DA. Degenerative arthritis of the knee secondary to fracture malunion. *Clin Orthop Relat Res* 1988;(234):159–169
- Richter DL, Schenck RC Jr, Wascher DC, Treme G. Knee articular cartilage repair and restoration techniques: a review of the literature. *Sports Health* 2016;8(02):153–160

- 19 Gross AE, Kim W, Las Heras F, Backstein D, Safir O, Pritzker KP. Fresh osteochondral allografts for posttraumatic knee defects: long-term follow-up. *Clin Orthop Relat Res* 2008;466(08):1863–1870
- 20 Shasha N, Krywulak S, Backstein D, Pressman A, Gross AE. Long-term follow-up of fresh tibial osteochondral allografts for failed tibial plateau fractures. *J Bone Joint Surg Am* 2003;85-A(2, suppl 2):33–39
- 21 Fűrnstahl P, Vlachopoulos L, Schweizer A, Fucentese SF, Koch PP. Complex osteotomies of tibial plateau malunions using computer-assisted planning and patient-specific surgical guides. *J Orthop Trauma* 2015;29(08):e270–e276
- 22 Jud L, Müller DA, Fűrnstahl P, Fucentese SF, Vlachopoulos L. Joint-preserving tumour resection around the knee with allograft reconstruction using three-dimensional preoperative planning and patient-specific instruments. *Knee* 2019;26(03):787–793
- 23 Jentzsch T, Vlachopoulos L, Fűrnstahl P, Müller DA, Fuchs B. Tumor resection at the pelvis using three-dimensional planning and patient-specific instruments: a case series. *World J Surg Oncol* 2016;14(01):249
- 24 Vlachopoulos L, Schweizer A, Meyer DC, Gerber C, Fűrnstahl P. Three-dimensional corrective osteotomies of complex malunited humeral fractures using patient-specific guides. *J Shoulder Elbow Surg* 2016;25(12):2040–2047
- 25 Lorensen WE, Cline HE. Marching cubes: a high resolution 3D surface construction algorithm. *Comput Graph* 1987;21:163–169
- 26 Schenk P, Vlachopoulos L, Hingsammer A, Fucentese SF, Fűrnstahl P. Is the contralateral tibia a reliable template for reconstruction: a three-dimensional anatomy cadaveric study. *Knee Surg Sports Traumatol Arthrosc* 2018;26(08):2324–2331
- 27 Chen Y. Object modeling by registration of multiple range images. Paper presented at: IEEE International Conference on Robotics and Automation. Piscataway, NJ: Institute of Electrical and Electronics Engineers; 1991:2724–2729
- 28 Audette MA, Ferrie FP, Peters TM. An algorithmic overview of surface registration techniques for medical imaging. *Med Image Anal* 2000;4(03):201–217
- 29 Schweizer A, Fűrnstahl P, Harders M, Székely G, Nagy L. Complex radius shaft malunion: osteotomy with computer-assisted planning. *Hand (N Y)* 2010;5(02):171–178
- 30 Besl PJ, McKay ND. A method for registration of 3-D shapes. *IEEE Trans Pattern Anal Mach Intell* 1992;14:239–256
- 31 Jud L, Fűrnstahl P, Vlachopoulos L, Gotschi T, Leoty LC, Fucentese SF. Malpositioning of patient-specific instruments within the possible degrees of freedom in high-tibial osteotomy has no considerable influence on mechanical leg axis correction. *Knee Surg Sports Traumatol Arthrosc* 2020;28(05):1356–1364
- 32 Vlachopoulos L, Schweizer A, Graf M, Nagy L, Fűrnstahl P. Three-dimensional postoperative accuracy of extra-articular forearm osteotomies using CT-scan based patient-specific surgical guides. *BMC Musculoskelet Disord* 2015;16:336
- 33 Vlachopoulos L, Schweizer A, Meyer DC, Gerber C, Fűrnstahl P. Computer-assisted planning and patient-specific guides for the treatment of midshaft clavicle malunions. *J Shoulder Elbow Surg* 2017;26(08):1367–1373
- 34 Ghazavi MT, Pritzker KP, Davis AM, Gross AE. Fresh osteochondral allografts for post-traumatic osteochondral defects of the knee. *J Bone Joint Surg Br* 1997;79(06):1008–1013