

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Chlorido[1-(2-oxidophenyl)ethylidene]-[tris(3,5-dimethylpyrazol-1-yl)hydroborato]iridium(III) chloroform monosolvate

Laura L. Santos,^a Margarita Paneque^a and Kurt Mereiter^{b*}

^aInstituto de Investigaciones Químicas (IIQ) and Departamento de Química Inorgánica, Consejo Superior de Investigaciones Científicas (CSIC) and Universidad de Sevilla, Avenida Américo Vespucio 49, 41092 Sevilla, Spain, and ^bInstitute of Chemical Technologies and Analytics, Vienna University of Technology, Getreidemarkt 9/164SC, A-1060 Vienna, Austria
Correspondence e-mail: kurt.mereiter@tuwien.ac.at

Received 6 March 2013; accepted 18 March 2013


Key indicators: single-crystal X-ray study; $T = 173$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; R factor = 0.028; wR factor = 0.071; data-to-parameter ratio = 23.6.

In the title compound, $[\text{Ir}(\text{C}_{15}\text{H}_{22}\text{BN}_6)(\text{C}_8\text{H}_7\text{O})\text{Cl}]\cdot\text{CHCl}_3$, the Ir atom is formally trivalent and is coordinated in a slightly distorted octahedral geometry by three facial N atoms, one C atom, one O atom and one Cl atom. The $\text{Ir}=\text{C}_{\text{carbene}}$ bond is strong and short and exerts a notable effect on the *trans*-Ir—N bond, which is about 0.10 Å longer than the two other Ir—N bonds. The chloroform solvent molecule is anchored *via* a weak $\text{C}-\text{H}\cdots\text{Cl}$ hydrogen bond to the Cl atom of the Ir complex molecule. In the crystal, the constituents adopt a layer-like arrangement parallel to (010) and are held together by weak intermolecular $\text{C}-\text{H}\cdots\text{Cl}$ hydrogen bonds, as well as weak $\text{Cl}\cdots\text{Cl}$ [3.498 (2) Å] and $\text{Cl}\cdots\pi$ [3.360 (4) Å] interactions. A weak intramolecular $\text{C}-\text{H}\cdots\text{O}$ hydrogen bond is also observed.

Related literature

The title compound represents a well crystallizing air-stable chloroform solvate of a mononuclear iridium complex based on the (hydrogen tris(3,5-dimethylpyrazolyl)borate- N,N',N'')-iridium moiety $\text{Ir}[\text{Tp}^{\text{Me}_2}]$. Its formation from $[(\text{Tp}^{\text{Me}_2})\text{Ir}(\text{C}_6\text{H}_5)_2(k^1\text{-N}_2)]$ ($\text{C}_6\text{H}_5 = \text{phenyl}$, $\text{N}_2 = \text{dinitrogen}$) and ethoxybenzene involved multiple $\text{C}-\text{C}, \text{H}, \text{O}, \text{Cl}$ bond transformations by the outstanding activity of the $\text{Ir}[\text{Tp}^{\text{Me}_2}]$ moiety. For general information on $\text{C}-\text{H}$ and $\text{C}-\text{C}$ activation, see: Lin & Yamamoto (1999); Dyker (1999); Labinger & Bercaw (2002). For $\text{C}-\text{H}$ bond activation reactions of ethers by $\text{Ir}[\text{Tp}^{\text{Me}_2}]$ complexes, see: Lara *et al.* (2009); Conejero *et al.* (2010); Santos *et al.* (2013). For the synthesis of the complex and related crystal structures, see: Gutiérrez-Puebla *et al.* (1998); Lara *et al.* (2009). For a description of the Cambridge

Structural Database, see: Allen (2002). For bond-length data, see: Allen *et al.* (1987).



Experimental

Crystal data

$[\text{Ir}(\text{C}_{15}\text{H}_{22}\text{BN}_6)(\text{C}_8\text{H}_7\text{O})\text{Cl}]\cdot\text{CHCl}_3$ $V = 2777.7$ (2) Å³
 $M_r = 763.35$ $Z = 4$
 Monoclinic, $P2_1/c$ Mo $K\alpha$ radiation
 $a = 10.1271$ (4) Å $\mu = 5.22$ mm⁻¹
 $b = 19.1711$ (8) Å $T = 173$ K
 $c = 14.3154$ (6) Å $0.32 \times 0.15 \times 0.10$ mm
 $\beta = 91.956$ (2)°

Data collection

Bruker SMART APEX CCD diffractometer 52411 measured reflections
 8053 independent reflections
 Absorption correction: multi-scan (SADABS; Bruker, 2003) 6999 reflections with $I > 2\sigma(I)$
 $T_{\text{min}} = 0.343$, $T_{\text{max}} = 0.593$ $R_{\text{int}} = 0.037$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.028$ 341 parameters
 $wR(F^2) = 0.071$ H-atom parameters constrained
 $S = 1.02$ $\Delta\rho_{\text{max}} = 1.29$ e Å⁻³
 8053 reflections $\Delta\rho_{\text{min}} = -1.43$ e Å⁻³

Table 1

Selected bond lengths (Å).

Ir1—C22	1.937 (3)	Ir1—O1	2.063 (2)
Ir1—N3	2.056 (3)	Ir1—N5	2.155 (3)
Ir1—N1	2.059 (3)	Ir1—Cl1	2.3500 (8)

Table 2

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C24—H24 \cdots Cl1	1.00	2.55	3.488 (5)	156
C11—H11A \cdots O1	0.98	2.37	3.230 (4)	146
C11—H11C \cdots Cl3 ⁱ	0.98	2.65	3.609 (4)	166

Symmetry code: (i) $x, -y + \frac{1}{2}, z - \frac{1}{2}$.

Data collection: SMART (Bruker, 2003); cell refinement: SAINT (Bruker, 2003); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine

structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *Mercury* (Macrae *et al.*, 2006); software used to prepare material for publication: *PLATON* (Spek, 2009) and *pubCIF* (Westrip, 2010).

Financial support (FEDER) from the Spanish Ministry of Science (projects CTQ2010–17476 and Consolider-Ingenio 2010 CSD2007–00006) and the Junta de Andalucía (grant FQM-119 and project P09-FQM-4832) is acknowledged.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LH5594).

References

- Allen, F. H. (2002). *Acta Cryst.* **B58**, 380–388.
- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Bruker (2003). *SMART*, *SAINTE* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Conejero, S., Paneque, M., Poveda, M. L., Santos, L. L. & Carmona, E. (2010). *Acc. Chem. Res.* **43**, 572–580.
- Dyker, G. (1999). *Angew. Chem. Int. Ed.* **38**, 1698–1712.
- Gutiérrez-Puebla, E., Monge, A., Nicasio, M. C., Pérez, P. J., Poveda, M. L. & Carmona, E. (1998). *Chem. Eur. J.* **4**, 2225–2236.
- Labinger, J. A. & Bercaw, J. E. (2002). *Nature*, **417**, 507–514.
- Lara, P., Paneque, M., Poveda, M. L., Santos, L. L., Valpuesta, J. E. V., Carmona, E., Moncho, S., Ujaque, G., Lledós, A., Álvarez, E. & Mereiter, K. (2009). *Chem. Eur. J.* **15**, 9034–9045.
- Lin, Y.-S. & Yamamoto, A. (1999). In *Activation of Unreactive Bonds and Organic Synthesis*, edited by S. Murai. Berlin: Springer.
- Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.
- Santos, L. L., Mereiter, K. & Paneque, M. (2013). *Organometallics*, **32**, 565–569.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supplementary materials

Acta Cryst. (2013). E69, m224–m225 [doi:10.1107/S1600536813007344]

Chlorido[1-(2-oxidophenyl)ethylidene][tris(3,5-dimethylpyrazol-1-yl)hydroborato]iridium(III) chloroform monosolvate

Laura L. Santos, Margarita Paneque and Kurt Mereiter

Comment

Transition metal compounds capable of inducing C—H bond activation and subsequent C—C bond formation have important applications in the synthesis of complex organic molecules from simple, commonly available substrates (Lin & Yamamoto, 1999; Dyker, 1999; Labinger & Bercaw, 2002). Iridium complexes with hydrogen-tris(pyrazolyl)borate as a stabilizing ligand and labile coordination sites have been found to show an outstanding potential at this respect (Conejero *et al.*, 2010). Part of our work in this field has derived from the study of reactions of ethers with reactive Ir complexes coordinated by the hydrogen-tris(3,5-dimethylpyrazolyl)borate ligand (Tp^{Me2}) (Lara *et al.*, 2009; Conejero *et al.*, 2010; Santos *et al.*, 2013). When the complex [(Tp^{Me2})Ir(C₆H₅)₂(*k*¹-N₂)] (C₆H₅ = phenyl, N₂ = dinitrogen; Gutiérrez-Puebla *et al.*, 1998) and C₆H₅OCH₂CH₃ (ethoxybenzene) are heated to 333K in cyclohexane a mixture of three compounds is formed (see reaction scheme Fig. 3). The major reaction product is **2**, the precursor of the title complex **1**. Compound **2** is a hydride-alkylidene whose formation requires multiple C—H bond activations, C—O bond cleavage and C—C bond formation. The other reaction products are a heteroatom-stabilized hydride-carbene **3** derived from three C—H activations of the organic product, and the minor reaction product **4** (*ca* 5%), which is the hydride-alkene tautomer of **2**. Compound **2** could be prepared independently in nearly quantitative yield (~95%) by the reaction of [(Tp^{Me2})Ir(C₆H₅)₂(*k*¹-N₂)] with 2-ethylphenol (Lara *et al.*, 2009). Compound **2** is stable at room temperature but at higher temperatures is in equilibrium with compound **4**. On the other hand, if **2** is heated in chloroform at 353K for 4 days, a C—Cl bond activation takes place whereby the hydride is exchanged against a Cl atom under concomitant formation of dichloromethane. The resulting complex **1** (Fig. 3) crystallizes from the excess of CHCl₃ under formation of the title compound, an air-stable solvate 1·CHCl₃, (**I**). In **1** the iridium atom exhibits a relatively regular octahedral coordination by three pyrazole nitrogen atoms, the carbene atom C22, the phenolate oxygen O1, and the chloride ligand Cl1 (Fig. 1). The *cis* bond angles about Ir vary from 82.18 (12)° (C22—Ir1—O1) to 98.38 (13)° (C22—Ir1—N3) and the *trans* bond angles from 173.11 (8)° to 177.55 (9)° (N3—Ir1—O1). The metal-carbene bond Ir1—C22 = 1.937 (3) Å is characteristically short and in good accord with Ir-carbene bonds of well refined crystal structures in the Cambridge Structural Database (version 5.33; Allen, 2002), which gave a mean value of 1.942 (64) Å for 57 crystal structures with 69 bonds. The Ir—N bonds (Table 1) show a typical elongation of *ca* 0.1 Å for the bond Ir1—N5 *trans* to the carbene ligand. The Ir—O and Ir—Cl bonds adopt normal values (Allen *et al.*, 1987). Bond lengths and angles in the Tp^{Me2} ligand compare well with related complexes (*e.g.*: Lara *et al.*, 2009; Santos *et al.*, 2013). The chelate ring formed by the carbene ligand has a flat envelope conformation with O1—C16—C21—C22 perfectly planar (r.m.s. deviation from planarity 0.0002 Å) and Ir1 displaced from this plane by -0.352 (5) Å whereas the terminal methyl carbon C23 is 0.419 (7) Å off from this plane. The carbene atom C22 has a very flat pyramidal coordination and deviates by -0.064 (4) Å from the plane defined by Ir1, C21, and C23. In the crystal structure the Ir complexes **1** and the CHCl₃ molecules are organized in a layer-like fashion

parallel to (010) as shown in Fig. 2. Such layers are centered at $y \approx 1/4$; and $y \approx 3/4$. The CHCl_3 molecule is anchored in the structure *via* a pronounced $\text{C—H}\cdots\text{Cl}$ hydrogen bond ($\text{C}\cdots\text{Cl} = 3.360(4) \text{ \AA}$) to the C11 atom of the Ir complex (Fig. 1 and Table 2). It is moreover fixed by the interaction $\text{C11—H11c}\cdots\text{Cl3}$ [$\text{C11}(x, 1/2 - y, 1/2 + z)\cdots\text{Cl3} = 3.608(4) \text{ \AA}$], by the halogen-halogen contact $\text{Cl3}\cdots\text{Cl1}(x, 1/2 - y, 1/2 + z) = 3.498(2) \text{ \AA}$, and two side-on contacts between the π -orbitals of arene rings and Cl [Cl2 with pyrazole ring 1 and the shortest contact distance $\text{Cl2}\cdots\text{C4}(x - 1, y, z) = 3.360(4) \text{ \AA}$; Cl3 with the phenyl ring and the shortest contact distance $\text{Cl3}\cdots\text{C20} = 3.396(4) \text{ \AA}$]. These interactions are included in Fig. 2. Interactions between the Ir complexes are unremarkable and consist essentially of van der Waals contacts. The interaction $\text{C11—H11a}\cdots\text{O3}$ is intramolecular.

Experimental


A solution of **2** (0.030 g, 0.049 mmol; see Fig. 3; for synthesis see Lara *et al.*, 2009) in CHCl_3 (3 ml) was stirred at 353K for 4 days. After this time the solvent was removed under reduced pressure. NMR spectra of the crude product revealed the presence of complex **1** in 70% spectroscopic yield. Crystallization from pentane/ CH_2Cl_2 / CHCl_3 at 253K gave compound **1** as a dark green microcrystalline solid. ^1H NMR (CDCl_3 , 298 K) δ 7.47, 7.28, 7.19, 6.60 (dd, ddd, d, ddd, 1 H each, $^3J_{\text{HH}} \approx 8.5$, $^4J_{\text{HH}} \approx 1$ Hz, 4 CH_{ar}), 5.89, 5.88, 5.50 (s, 1 H each, 3 CH_{pz}), 3.07 (s, 3 H, $\text{Ir}=\text{CCH}_3$), 2.77, 2.49, 2.40, 2.36, 2.32, 1.32 (s, 3 H each, 6 Me_{pz}). $^{13}\text{C}\{^1\text{H}\}$ NMR (CDCl_3 , 25 °C) δ 273.2 ($\text{Ir}=\text{C}$), 192.5 (Ir—O—C), 154.7, 153.8, 153.3, 152.3, 144.6, 144.2, 144.1 ($\text{Ir}=\text{C—C} + \text{C}_{\text{qpz}}$), 141.0, 124.2, 119.7, 115.7 (CH_{ar}), 108.4, 108.3, 108.2 (CH_{pz}), 34.0 ($\text{Ir}=\text{CCH}_3$), 16.4, 14.1, 13.1, 13.0, 12.4, 12.2 (Me_{pz}). Crystals of **1**. CHCl_3 for X-ray diffraction were obtained by recrystallization from CHCl_3 /pentane.

Refinement


H atoms were placed in calculated positions and thereafter treated as riding, $\text{C—H} = 0.95\text{--}1.00 \text{ \AA}$, $\text{B—H} = 1.00 \text{ \AA}$, $U_{\text{iso}}(\text{H}) = 1.2\text{--}1.5U_{\text{eq}}(\text{C}, \text{B})$, using AFIX 137 of program *SHELXL97* (Sheldrick, 2008) for the methyl groups.

Computing details


Data collection: *SMART* (Bruker, 2003); cell refinement: *SAINTE* (Bruker, 2003); data reduction: *SAINTE* (Bruker, 2003); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: Mercury (Macrae *et al.*, 2006); software used to prepare material for publication: *PLATON* (Spek, 2009) and *pubCIF* (Westrip, 2010).

**Figure 1**

The asymmetric unit of the title structure **1**.CHCl₃, with displacement ellipsoids drawn at the 50% probability level and the hydrogen bond C24—H24...Cl1 as a dashed red line.

**Figure 2**

View of the structure of **1**.CHCl₃ approximately along [010] in the range $0 < y < 1/2$; showing the interactions C—H...O, Cl (red), Cl...Cl (green), and Cl... π (blue) as dashed lines.


Figure 3

 Reaction scheme for the synthesis of **1**.

Chlorido[1-(2-oxidophenyl)ethylidene][tris(3,5-dimethylpyrazol-1-yl)hydroborato]iridium(III) chloroform monosolvate
Crystal data
 $[\text{Ir}(\text{C}_{15}\text{H}_{22}\text{BN}_6)(\text{C}_8\text{H}_7\text{O})\text{Cl}]\cdot\text{CHCl}_3$
 $M_r = 763.35$

 Monoclinic, $P2_1/c$

 Hall symbol: $-P\ 2_1/c$
 $a = 10.1271\ (4)\ \text{\AA}$
 $b = 19.1711\ (8)\ \text{\AA}$
 $c = 14.3154\ (6)\ \text{\AA}$
 $\beta = 91.956\ (2)^\circ$
 $V = 2777.7\ (2)\ \text{\AA}^3$
 $Z = 4$
 $F(000) = 1496$
 $D_x = 1.825\ \text{Mg m}^{-3}$

 Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 8986 reflections

 $\theta = 2.3\text{--}30.0^\circ$
 $\mu = 5.22\ \text{mm}^{-1}$
 $T = 173\ \text{K}$

Irregular, dark green

 $0.32 \times 0.15 \times 0.10\ \text{mm}$
Data collection

 Bruker SMART APEX CCD
 diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

 ω and φ scans

Absorption correction: multi-scan

 (*SADABS*; Bruker, 2003)

 $T_{\text{min}} = 0.343$, $T_{\text{max}} = 0.593$

52411 measured reflections

8053 independent reflections

 6999 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.037$
 $\theta_{\text{max}} = 30.0^\circ$, $\theta_{\text{min}} = 2.6^\circ$
 $h = -14 \rightarrow 14$
 $k = -26 \rightarrow 26$
 $l = -20 \rightarrow 20$
Refinement

 Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.028$
 $wR(F^2) = 0.071$
 $S = 1.02$

8053 reflections

341 parameters

0 restraints

 Primary atom site location: structure-invariant
 direct methods

 Secondary atom site location: difference Fourier
 map

 Hydrogen site location: inferred from
 neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.038P)^2 + 4.5744P]$$

where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$

$$\Delta\rho_{\max} = 1.29 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -1.43 \text{ e } \text{\AA}^{-3}$$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Ir1	0.436291 (10)	0.157920 (6)	0.682819 (7)	0.01911 (4)
Cl1	0.23334 (8)	0.20336 (4)	0.62640 (6)	0.02847 (15)
B1	0.6944 (4)	0.16488 (19)	0.5669 (3)	0.0262 (7)
H0B	0.7782	0.1680	0.5322	0.031*
N1	0.6159 (3)	0.11309 (14)	0.71808 (18)	0.0227 (5)
N2	0.7168 (3)	0.12094 (14)	0.65687 (19)	0.0247 (5)
N3	0.5334 (3)	0.24641 (14)	0.64204 (18)	0.0234 (5)
N4	0.6481 (3)	0.23785 (15)	0.59388 (19)	0.0257 (5)
N5	0.4663 (3)	0.11764 (14)	0.54460 (18)	0.0225 (5)
N6	0.5867 (3)	0.12870 (14)	0.50605 (18)	0.0236 (5)
O1	0.3316 (2)	0.07061 (11)	0.72065 (14)	0.0216 (4)
C1	0.5825 (4)	0.0473 (2)	0.8677 (3)	0.0357 (8)
H1A	0.5529	0.0884	0.9018	0.054*
H1B	0.6378	0.0181	0.9095	0.054*
H1C	0.5055	0.0205	0.8451	0.054*
C2	0.6607 (3)	0.06994 (17)	0.7866 (2)	0.0267 (6)
C3	0.7896 (3)	0.05026 (19)	0.7680 (3)	0.0323 (7)
H3	0.8448	0.0201	0.8048	0.039*
C4	0.8217 (3)	0.08277 (18)	0.6863 (2)	0.0281 (7)
C5	0.9489 (3)	0.0795 (2)	0.6354 (3)	0.0380 (8)
H5A	0.9295	0.0740	0.5683	0.057*
H5B	1.0014	0.0397	0.6583	0.057*
H5C	0.9989	0.1227	0.6464	0.057*
C6	0.3925 (4)	0.34615 (18)	0.6934 (3)	0.0340 (7)
H6A	0.3425	0.3095	0.7241	0.051*
H6B	0.3359	0.3689	0.6456	0.051*
H6C	0.4224	0.3807	0.7399	0.051*
C7	0.5096 (4)	0.31472 (17)	0.6486 (2)	0.0277 (6)
C8	0.6101 (4)	0.35093 (18)	0.6050 (3)	0.0337 (8)
H8	0.6178	0.4001	0.5992	0.040*
C9	0.6957 (4)	0.30165 (18)	0.5719 (2)	0.0308 (7)
C10	0.8185 (4)	0.3117 (2)	0.5176 (3)	0.0454 (10)
H10A	0.8053	0.2917	0.4550	0.068*

H10B	0.8929	0.2884	0.5501	0.068*
H10C	0.8374	0.3617	0.5124	0.068*
C11	0.2581 (3)	0.05407 (19)	0.5004 (3)	0.0308 (7)
H11A	0.2475	0.0463	0.5674	0.046*
H11B	0.2415	0.0104	0.4664	0.046*
H11C	0.1952	0.0897	0.4781	0.046*
C12	0.3956 (3)	0.07824 (16)	0.4843 (2)	0.0232 (6)
C13	0.4714 (4)	0.06377 (17)	0.4063 (2)	0.0280 (6)
H13	0.4455	0.0369	0.3530	0.034*
C14	0.5906 (3)	0.09640 (17)	0.4225 (2)	0.0263 (6)
C15	0.7068 (4)	0.1003 (2)	0.3615 (3)	0.0387 (9)
H15A	0.7034	0.0615	0.3169	0.058*
H15B	0.7884	0.0973	0.4001	0.058*
H15C	0.7049	0.1446	0.3274	0.058*
C16	0.2741 (3)	0.08151 (19)	0.7994 (2)	0.0280 (6)
C17	0.1875 (4)	0.0306 (2)	0.8350 (3)	0.0377 (8)
H17	0.1654	-0.0100	0.7999	0.045*
C18	0.1364 (4)	0.0413 (3)	0.9210 (3)	0.0500 (11)
H18	0.0767	0.0077	0.9443	0.060*
C19	0.1686 (4)	0.1001 (3)	0.9767 (3)	0.0493 (11)
H19	0.1324	0.1054	1.0366	0.059*
C20	0.2535 (4)	0.1500 (2)	0.9430 (3)	0.0384 (8)
H20	0.2762	0.1896	0.9799	0.046*
C21	0.3070 (3)	0.14175 (19)	0.8528 (2)	0.0282 (7)
C22	0.4044 (3)	0.18475 (18)	0.8105 (2)	0.0263 (6)
C23	0.4790 (4)	0.2367 (2)	0.8696 (3)	0.0356 (8)
H23A	0.4192	0.2741	0.8877	0.053*
H23B	0.5510	0.2563	0.8340	0.053*
H23C	0.5156	0.2136	0.9258	0.053*
C24	0.0420 (5)	0.2910 (2)	0.7869 (4)	0.0508 (11)
H24	0.1171	0.2666	0.7573	0.061*
Cl2	-0.10099 (13)	0.24331 (7)	0.76482 (9)	0.0592 (3)
Cl3	0.07774 (17)	0.29868 (9)	0.90450 (11)	0.0756 (4)
Cl4	0.02838 (17)	0.37582 (8)	0.73962 (13)	0.0804 (5)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Ir1	0.01915 (6)	0.01930 (6)	0.01887 (6)	-0.00133 (4)	0.00052 (4)	-0.00196 (4)
Cl1	0.0237 (3)	0.0308 (4)	0.0307 (4)	0.0028 (3)	-0.0012 (3)	-0.0030 (3)
B1	0.0226 (16)	0.0281 (18)	0.0280 (17)	-0.0040 (13)	0.0031 (13)	0.0005 (14)
N1	0.0205 (12)	0.0230 (12)	0.0247 (12)	-0.0017 (10)	0.0005 (9)	0.0001 (10)
N2	0.0187 (12)	0.0266 (13)	0.0285 (13)	-0.0012 (10)	-0.0002 (10)	0.0004 (10)
N3	0.0240 (12)	0.0219 (12)	0.0245 (12)	-0.0032 (10)	0.0014 (10)	0.0012 (10)
N4	0.0244 (13)	0.0256 (13)	0.0273 (13)	-0.0041 (10)	0.0022 (10)	0.0008 (10)
N5	0.0227 (12)	0.0223 (12)	0.0225 (12)	-0.0025 (10)	0.0020 (9)	-0.0005 (10)
N6	0.0229 (12)	0.0247 (13)	0.0234 (12)	0.0006 (10)	0.0048 (10)	-0.0001 (10)
O1	0.0229 (10)	0.0216 (10)	0.0207 (10)	-0.0041 (8)	0.0053 (8)	-0.0022 (8)
C1	0.0391 (19)	0.0357 (19)	0.0325 (18)	0.0050 (15)	0.0022 (14)	0.0104 (15)
C2	0.0289 (16)	0.0243 (15)	0.0264 (15)	0.0011 (12)	-0.0046 (12)	0.0003 (12)

C3	0.0267 (16)	0.0341 (18)	0.0355 (18)	0.0043 (13)	-0.0072 (13)	0.0021 (14)
C4	0.0199 (14)	0.0273 (16)	0.0370 (17)	-0.0013 (12)	-0.0032 (12)	-0.0029 (13)
C5	0.0191 (15)	0.046 (2)	0.049 (2)	0.0000 (14)	-0.0007 (14)	-0.0016 (17)
C6	0.046 (2)	0.0254 (17)	0.0305 (17)	0.0046 (14)	-0.0010 (15)	-0.0034 (13)
C7	0.0384 (18)	0.0210 (14)	0.0234 (15)	-0.0032 (13)	-0.0034 (12)	-0.0021 (12)
C8	0.048 (2)	0.0218 (16)	0.0310 (17)	-0.0099 (14)	-0.0032 (15)	0.0013 (13)
C9	0.0359 (18)	0.0272 (16)	0.0291 (16)	-0.0121 (14)	-0.0015 (13)	0.0022 (13)
C10	0.044 (2)	0.042 (2)	0.051 (2)	-0.0161 (18)	0.0114 (19)	0.0053 (19)
C11	0.0297 (16)	0.0307 (17)	0.0316 (17)	-0.0065 (13)	-0.0040 (13)	-0.0030 (13)
C12	0.0288 (15)	0.0188 (13)	0.0217 (14)	-0.0005 (11)	-0.0032 (11)	0.0002 (11)
C13	0.0404 (18)	0.0228 (15)	0.0208 (14)	0.0020 (13)	0.0007 (12)	-0.0010 (11)
C14	0.0353 (17)	0.0212 (14)	0.0226 (14)	0.0042 (12)	0.0051 (12)	0.0015 (11)
C15	0.050 (2)	0.037 (2)	0.0302 (17)	0.0049 (17)	0.0179 (16)	0.0016 (15)
C16	0.0252 (15)	0.0328 (17)	0.0261 (15)	-0.0011 (13)	0.0027 (12)	0.0016 (13)
C17	0.0366 (19)	0.042 (2)	0.0348 (19)	-0.0111 (16)	0.0075 (15)	0.0016 (16)
C18	0.042 (2)	0.069 (3)	0.041 (2)	-0.014 (2)	0.0164 (18)	0.005 (2)
C19	0.042 (2)	0.065 (3)	0.042 (2)	-0.001 (2)	0.0193 (18)	0.004 (2)
C20	0.039 (2)	0.049 (2)	0.0279 (17)	0.0034 (17)	0.0046 (14)	-0.0053 (16)
C21	0.0296 (16)	0.0322 (17)	0.0227 (15)	0.0043 (13)	0.0003 (12)	-0.0021 (12)
C22	0.0260 (15)	0.0267 (15)	0.0261 (15)	0.0035 (12)	-0.0019 (12)	-0.0018 (12)
C23	0.047 (2)	0.0311 (18)	0.0280 (17)	-0.0063 (15)	-0.0033 (15)	-0.0064 (14)
C24	0.044 (2)	0.038 (2)	0.070 (3)	0.0005 (18)	0.014 (2)	-0.014 (2)
Cl2	0.0559 (7)	0.0624 (7)	0.0594 (7)	-0.0145 (6)	0.0056 (5)	-0.0200 (6)
Cl3	0.0868 (10)	0.0707 (9)	0.0678 (9)	-0.0084 (8)	-0.0188 (8)	-0.0070 (7)
Cl4	0.0887 (11)	0.0508 (8)	0.1044 (12)	0.0061 (7)	0.0424 (9)	0.0156 (7)

Geometric parameters (Å, °)

Ir1—C22	1.937 (3)	C8—C9	1.377 (6)
Ir1—N3	2.056 (3)	C8—H8	0.9500
Ir1—N1	2.059 (3)	C9—C10	1.500 (5)
Ir1—O1	2.063 (2)	C10—H10A	0.9800
Ir1—N5	2.155 (3)	C10—H10B	0.9800
Ir1—C11	2.3500 (8)	C10—H10C	0.9800
B1—N4	1.529 (5)	C11—C12	1.492 (5)
B1—N6	1.538 (5)	C11—H11A	0.9800
B1—N2	1.549 (5)	C11—H11B	0.9800
B1—H0B	1.0000	C11—H11C	0.9800
N1—C2	1.350 (4)	C12—C13	1.405 (4)
N1—N2	1.376 (4)	C13—C14	1.373 (5)
N2—C4	1.346 (4)	C13—H13	0.9500
N3—C7	1.336 (4)	C14—C15	1.490 (5)
N3—N4	1.380 (4)	C15—H15A	0.9800
N4—C9	1.356 (4)	C15—H15B	0.9800
N5—C12	1.337 (4)	C15—H15C	0.9800
N5—N6	1.372 (3)	C16—C21	1.418 (5)
N6—C14	1.348 (4)	C16—C17	1.419 (5)
O1—C16	1.303 (4)	C17—C18	1.366 (5)
C1—C2	1.492 (5)	C17—H17	0.9500
C1—H1A	0.9800	C18—C19	1.414 (7)

C1—H1B	0.9800	C18—H18	0.9500
C1—H1C	0.9800	C19—C20	1.383 (6)
C2—C3	1.393 (5)	C19—H19	0.9500
C3—C4	1.374 (5)	C20—C21	1.425 (5)
C3—H3	0.9500	C20—H20	0.9500
C4—C5	1.503 (5)	C21—C22	1.435 (5)
C5—H5A	0.9800	C22—C23	1.494 (5)
C5—H5B	0.9800	C23—H23A	0.9800
C5—H5C	0.9800	C23—H23B	0.9800
C6—C7	1.494 (5)	C23—H23C	0.9800
C6—H6A	0.9800	C24—C13	1.716 (6)
C6—H6B	0.9800	C24—C12	1.732 (5)
C6—H6C	0.9800	C24—C14	1.765 (5)
C7—C8	1.397 (5)	C24—H24	1.0000
C22—Ir1—N3	98.38 (13)	N3—C7—C6	125.0 (3)
C22—Ir1—N1	93.09 (12)	C8—C7—C6	126.3 (3)
N3—Ir1—N1	89.24 (10)	C9—C8—C7	106.8 (3)
C22—Ir1—O1	82.18 (12)	C9—C8—H8	126.6
N3—Ir1—O1	177.55 (9)	C7—C8—H8	126.6
N1—Ir1—O1	93.12 (9)	N4—C9—C8	107.8 (3)
C22—Ir1—N5	174.27 (12)	N4—C9—C10	122.9 (3)
N3—Ir1—N5	87.20 (10)	C8—C9—C10	129.2 (3)
N1—Ir1—N5	85.68 (10)	C9—C10—H10A	109.5
O1—Ir1—N5	92.29 (9)	C9—C10—H10B	109.5
C22—Ir1—C11	93.09 (10)	H10A—C10—H10B	109.5
N3—Ir1—C11	91.03 (8)	C9—C10—H10C	109.5
N1—Ir1—C11	173.71 (8)	H10A—C10—H10C	109.5
O1—Ir1—C11	86.56 (6)	H10B—C10—H10C	109.5
N5—Ir1—C11	88.06 (7)	C12—C11—H11A	109.5
N4—B1—N6	109.7 (3)	C12—C11—H11B	109.5
N4—B1—N2	108.9 (3)	H11A—C11—H11B	109.5
N6—B1—N2	107.8 (3)	C12—C11—H11C	109.5
N4—B1—H0B	110.1	H11A—C11—H11C	109.5
N6—B1—H0B	110.1	H11B—C11—H11C	109.5
N2—B1—H0B	110.1	N5—C12—C13	109.3 (3)
C2—N1—N2	107.0 (3)	N5—C12—C11	123.9 (3)
C2—N1—Ir1	134.9 (2)	C13—C12—C11	126.9 (3)
N2—N1—Ir1	117.83 (19)	C14—C13—C12	106.1 (3)
C4—N2—N1	109.6 (3)	C14—C13—H13	127.0
C4—N2—B1	130.4 (3)	C12—C13—H13	127.0
N1—N2—B1	119.9 (3)	N6—C14—C13	107.9 (3)
C7—N3—N4	108.0 (3)	N6—C14—C15	123.0 (3)
C7—N3—Ir1	134.4 (2)	C13—C14—C15	129.1 (3)
N4—N3—Ir1	117.6 (2)	C14—C15—H15A	109.5
C9—N4—N3	108.7 (3)	C14—C15—H15B	109.5
C9—N4—B1	130.7 (3)	H15A—C15—H15B	109.5
N3—N4—B1	120.4 (3)	C14—C15—H15C	109.5
C12—N5—N6	106.9 (2)	H15A—C15—H15C	109.5

C12—N5—Ir1	135.0 (2)	H15B—C15—H15C	109.5
N6—N5—Ir1	117.95 (19)	O1—C16—C21	119.6 (3)
C14—N6—N5	109.8 (3)	O1—C16—C17	120.1 (3)
C14—N6—B1	131.9 (3)	C21—C16—C17	120.2 (3)
N5—N6—B1	117.7 (2)	C18—C17—C16	118.6 (4)
C16—O1—Ir1	110.2 (2)	C18—C17—H17	120.7
C2—C1—H1A	109.5	C16—C17—H17	120.7
C2—C1—H1B	109.5	C17—C18—C19	122.7 (4)
H1A—C1—H1B	109.5	C17—C18—H18	118.7
C2—C1—H1C	109.5	C19—C18—H18	118.7
H1A—C1—H1C	109.5	C20—C19—C18	119.3 (4)
H1B—C1—H1C	109.5	C20—C19—H19	120.4
N1—C2—C3	108.7 (3)	C18—C19—H19	120.4
N1—C2—C1	124.7 (3)	C19—C20—C21	120.0 (4)
C3—C2—C1	126.5 (3)	C19—C20—H20	120.0
C4—C3—C2	106.9 (3)	C21—C20—H20	120.0
C4—C3—H3	126.6	C16—C21—C20	119.3 (3)
C2—C3—H3	126.6	C16—C21—C22	113.1 (3)
N2—C4—C3	107.9 (3)	C20—C21—C22	127.3 (3)
N2—C4—C5	123.4 (3)	C21—C22—C23	119.1 (3)
C3—C4—C5	128.7 (3)	C21—C22—Ir1	112.6 (2)
C4—C5—H5A	109.5	C23—C22—Ir1	127.8 (3)
C4—C5—H5B	109.5	C22—C23—H23A	109.5
H5A—C5—H5B	109.5	C22—C23—H23B	109.5
C4—C5—H5C	109.5	H23A—C23—H23B	109.5
H5A—C5—H5C	109.5	C22—C23—H23C	109.5
H5B—C5—H5C	109.5	H23A—C23—H23C	109.5
C7—C6—H6A	109.5	H23B—C23—H23C	109.5
C7—C6—H6B	109.5	C13—C24—C12	111.8 (3)
H6A—C6—H6B	109.5	C13—C24—C14	108.0 (3)
C7—C6—H6C	109.5	C12—C24—C14	111.2 (3)
H6A—C6—H6C	109.5	C13—C24—H24	108.6
H6B—C6—H6C	109.5	C12—C24—H24	108.6
N3—C7—C8	108.6 (3)	C14—C24—H24	108.6
C22—Ir1—N1—C2	45.2 (3)	C1—C2—C3—C4	179.5 (3)
N3—Ir1—N1—C2	143.5 (3)	N1—N2—C4—C3	-0.6 (4)
O1—Ir1—N1—C2	-37.2 (3)	B1—N2—C4—C3	-176.4 (3)
N5—Ir1—N1—C2	-129.2 (3)	N1—N2—C4—C5	179.9 (3)
C22—Ir1—N1—N2	-142.2 (2)	B1—N2—C4—C5	4.2 (5)
N3—Ir1—N1—N2	-43.9 (2)	C2—C3—C4—N2	0.3 (4)
O1—Ir1—N1—N2	135.5 (2)	C2—C3—C4—C5	179.7 (3)
N5—Ir1—N1—N2	43.4 (2)	N4—N3—C7—C8	-0.4 (4)
C2—N1—N2—C4	0.6 (3)	Ir1—N3—C7—C8	-178.1 (2)
Ir1—N1—N2—C4	-173.9 (2)	N4—N3—C7—C6	178.2 (3)
C2—N1—N2—B1	176.9 (3)	Ir1—N3—C7—C6	0.5 (5)
Ir1—N1—N2—B1	2.4 (4)	N3—C7—C8—C9	-0.1 (4)
N4—B1—N2—C4	-129.0 (3)	C6—C7—C8—C9	-178.7 (3)
N6—B1—N2—C4	111.9 (4)	N3—N4—C9—C8	-0.8 (4)

N4—B1—N2—N1	55.6 (4)	B1—N4—C9—C8	173.4 (3)
N6—B1—N2—N1	-63.4 (4)	N3—N4—C9—C10	-178.9 (3)
C22—Ir1—N3—C7	-48.8 (3)	B1—N4—C9—C10	-4.6 (6)
N1—Ir1—N3—C7	-141.8 (3)	C7—C8—C9—N4	0.6 (4)
N5—Ir1—N3—C7	132.5 (3)	C7—C8—C9—C10	178.4 (4)
Cl1—Ir1—N3—C7	44.5 (3)	N6—N5—C12—C13	0.1 (3)
C22—Ir1—N3—N4	133.7 (2)	Ir1—N5—C12—C13	-175.2 (2)
N1—Ir1—N3—N4	40.7 (2)	N6—N5—C12—C11	179.6 (3)
N5—Ir1—N3—N4	-45.0 (2)	Ir1—N5—C12—C11	4.4 (5)
Cl1—Ir1—N3—N4	-133.0 (2)	N5—C12—C13—C14	-0.1 (4)
C7—N3—N4—C9	0.8 (4)	C11—C12—C13—C14	-179.6 (3)
Ir1—N3—N4—C9	178.9 (2)	N5—N6—C14—C13	-0.1 (4)
C7—N3—N4—B1	-174.2 (3)	B1—N6—C14—C13	170.8 (3)
Ir1—N3—N4—B1	4.0 (4)	N5—N6—C14—C15	178.1 (3)
N6—B1—N4—C9	-115.4 (4)	B1—N6—C14—C15	-11.1 (5)
N2—B1—N4—C9	126.7 (3)	C12—C13—C14—N6	0.1 (4)
N6—B1—N4—N3	58.3 (4)	C12—C13—C14—C15	-177.9 (3)
N2—B1—N4—N3	-59.6 (4)	Ir1—O1—C16—C21	-10.4 (4)
N3—Ir1—N5—C12	-143.5 (3)	Ir1—O1—C16—C17	174.7 (3)
N1—Ir1—N5—C12	127.0 (3)	O1—C16—C17—C18	175.3 (4)
O1—Ir1—N5—C12	34.1 (3)	C21—C16—C17—C18	0.4 (6)
Cl1—Ir1—N5—C12	-52.4 (3)	C16—C17—C18—C19	-1.4 (7)
N3—Ir1—N5—N6	41.6 (2)	C17—C18—C19—C20	1.1 (7)
N1—Ir1—N5—N6	-47.8 (2)	C18—C19—C20—C21	0.3 (7)
O1—Ir1—N5—N6	-140.8 (2)	O1—C16—C21—C20	-174.0 (3)
Cl1—Ir1—N5—N6	132.8 (2)	C17—C16—C21—C20	0.9 (5)
C12—N5—N6—C14	0.0 (3)	O1—C16—C21—C22	0.0 (5)
Ir1—N5—N6—C14	176.2 (2)	C17—C16—C21—C22	174.8 (3)
C12—N5—N6—B1	-172.3 (3)	C19—C20—C21—C16	-1.2 (6)
Ir1—N5—N6—B1	3.9 (3)	C19—C20—C21—C22	-174.2 (4)
N4—B1—N6—C14	128.6 (3)	C16—C21—C22—C23	-161.3 (3)
N2—B1—N6—C14	-112.9 (4)	C20—C21—C22—C23	12.0 (5)
N4—B1—N6—N5	-61.1 (4)	C16—C21—C22—Ir1	11.4 (4)
N2—B1—N6—N5	57.4 (4)	C20—C21—C22—Ir1	-175.3 (3)
C22—Ir1—O1—C16	13.0 (2)	N3—Ir1—C22—C21	164.6 (2)
N1—Ir1—O1—C16	105.7 (2)	N1—Ir1—C22—C21	-105.8 (2)
N5—Ir1—O1—C16	-168.5 (2)	O1—Ir1—C22—C21	-13.0 (2)
Cl1—Ir1—O1—C16	-80.6 (2)	Cl1—Ir1—C22—C21	73.1 (2)
N2—N1—C2—C3	-0.4 (4)	N3—Ir1—C22—C23	-23.6 (3)
Ir1—N1—C2—C3	172.8 (2)	N1—Ir1—C22—C23	66.1 (3)
N2—N1—C2—C1	-179.9 (3)	O1—Ir1—C22—C23	158.9 (3)
Ir1—N1—C2—C1	-6.7 (5)	Cl1—Ir1—C22—C23	-115.1 (3)
N1—C2—C3—C4	0.1 (4)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
C24—H24...Cl1	1.00	2.55	3.488 (5)	156

supplementary materials

C11—H11A [⋯] O1	0.98	2.37	3.230 (4)	146
C11—H11C [⋯] C13 ⁱ	0.98	2.65	3.609 (4)	166

Symmetry code: (i) $x, -y+1/2, z-1/2$.