


RESEARCH PAPER

OPEN ACCESS 

## Hsa\_circ\_0000418 promotes the progression of glioma by regulating microRNA-409-3p / pyruvate dehydrogenase kinase 1 axis

Zhihui Ma<sup>a</sup>, Zhen Chen<sup>b</sup>, Ying Zhou<sup>c</sup>, Yanping Li<sup>d</sup>, Shuyang Li<sup>d</sup>, Hongxia Wang<sup>e</sup>, and Ji Feng<sup>a</sup>

<sup>a</sup>Department of Neurosurgery, The First Hospital of Qinhuangdao, the First Hospital of Qinhuangdao, Qinhuangdao, Hebei, China;

<sup>b</sup>Department of Anesthesiology, Qinglong Manchu Autonomous County Hospital, Qinhuangdao, Hebei, China; <sup>c</sup>Department of Anesthesiology, The Third Hospital of Qinhuangdao, Qinhuangdao, Hebei, China; <sup>d</sup>Supply Department, Qinglong Manchu Autonomous County Hospital, Qinhuangdao, Hebei, China; <sup>e</sup>Operating Room, Qinglong Manchu Autonomous County Hospital, Qinhuangdao, Hebei, China

### ABSTRACT


Glioma is the commonest intracranial malignancy, and circRNAs are important regulatory factors which are implicated in the development of glioma. Nonetheless, the role of circRNAs in glioma is largely unknown. The research is performed to elaborate on the biological role of has\_circ\_0000418 (circ\_0000418) in glioma progression and its potential molecular mechanism. The differentially expressed circRNAs in glioblastoma patient derived cells and neural progenitor cells were analyzed based on the microarray data of GSE146463. Additionally, qRT-PCR and Western blot experiments were conducted to measure the expression of circ\_0000418, microRNA-409-3p (miR-409-3p) and pyruvate dehydrogenase kinase 1 (*PDK1*) in glioma tissues/cells. Cell growth and cell cycle distribution were monitored using CCK-8 assay, BrdU assay and flow cytometry. Bioinformatics prediction, dual-luciferase reporter gene experiment and RIP assay were conducted to verify the targeting relationship between circ\_0000418 and miR-409-3p, miR-409-3p and PDK1 3'UTR. In this work, we observed that, circ\_0000418 expression level was significantly up-regulated in glioma tissues and cell lines. Circ\_0000418 overexpression facilitated glioma cell growth and accelerated cell cycle progression, while knockdown of circ\_0000418 produced the opposite effects. Circ\_0000418 specifically combined with miR-409-3p, and circ\_0000418 negatively modulated the expression of miR-409-3p. PDK1 acted as a target gene of miR-409-3p, and PDK1 could be positively and indirectly modulated by circ\_0000418 in glioma cells. In summary, circ\_0000418 enhances glioma cell growth and accelerates cell cycle progression by regulating miR-409-3p/PDK1 axis.

### ARTICLE HISTORY

Received 25 January 2022  
Revised 25 February 2022  
Accepted 26 February 2022

### KEYWORDS

Circ\_0000418; glioma; miR-409-3p; PDK1



## 1 Introduction

Glioma is a malignant tumor which originates from neuroepithelial cells, taking up 40–50% of all intracranial malignancies, and is characterized by high morbidity, high relapse rate and high

mortality [1]. The current treatments for glioma include surgical resection, adjuvant radiotherapy and chemotherapy, and targeted therapies, however, the average overall survival time of patients was only 12–18 months [2]. Therefore, the

**CONTACT** Ying Zhou  [helloying.86@163.com](mailto:helloying.86@163.com)  Department of Anesthesiology, The Third Hospital of Qinhuangdao, Qinhuangdao, Hebei 066000, China

© 2022 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

identification of new therapeutic targets for glioma is urgent.

Circular RNAs (circRNAs) are ncRNAs that have a closed-loop structure [2]. They are widely present and stably expressed in eukaryotic transcriptome [3]. Accumulating studies demonstrate that circRNAs are closely correlated with glioma tumorigenesis and development. For instance, circ-AKT3 is under-expressed in glioma tissues; circ-AKT3 overexpression restrains the growth, radioresistance and tumorigenicity of glioma cells [4]. Circ\_0043278 is significantly overexpressed in glioma tissues and cells, and its high expression correlates with unfavorable prognosis of the patients; knockdown of circ\_0043278 inhibits glioma cell growth, migration and invasion, and suppresses *in vivo* tumorigenesis [5]. Circ\_0000418, a member of circRNAs, is involved in mediating the anti-depressant effect of *adenosine deaminase acting on RNA 1* [6]. In the present work, our bioinformatics analysis implies that circ\_0000418 expression is up-regulated in glioblastoma patient derived cells. Nevertheless, the detailed expression pattern and biological function of circ\_0000418 in glioma are rarely reported.

MiR-409-3p is down-regulated in different malignancies, and it can regulate cellular biological processes like growth, apoptosis, differentiation and migration by targeting different genes. For instance, miR-409-3p inhibits the aggressiveness of glioma cells by targeting *high mobility group nucleosome binding domain 5 (HMGN5)* [7]. Pyruvate dehydrogenase kinase 1 (*PDK1*) belongs to the serine/threonine protein kinase family; PDK1 is significantly overexpressed in glioma tissues, and the knockdown of PDK1 restrains the colony formation ability of glioma cells [8,9]. The competitive endogenous RNA (ceRNA) theory states that circRNAs work as molecular sponge that regulate downstream target genes' expression by sponging miRNAs [10]. Our bioinformatics analysis reveals that both of circ\_0000418 and PDK1 3' UTR contain complementary binding sites with miR-409-3p. However, whether circ\_0000418/miR-409-3p/PDK1 forms a ceRNA network in the tumorigenesis of glioma needs to be verified.

As mentioned above, in this work, we supposed that there was a ceRNA regulatory mechanism

among circ\_0000418, miR-409-3p and PDK1 in glioma progression. The goal of this work is to validate this hypothesis. This work showed that circ\_0000418 was up-regulated in glioma tissues and cells, and circ\_0000418 facilitated the growth and cell cycle progression of glioma cells by decoying miR-409-3p and up-regulating PDK1. This study hopefully provides an innovative molecular marker/target for the diagnosis and therapy of glioma.

## 2 Materials and methods

### 2.1 Sample collection

A total of 47 subjects who were diagnosed with glioma from June 2018 to December 2020 were selected. The surgically removed malignant tissues and matching adjacent tissues were collected and instantly preserved in liquid nitrogen. All subjects did not undergo chemotherapy, radiotherapy or other anti-cancer treatments before the surgery. This work was executed under the guidance of the Ethics Committee of the First Hospital of Qinhuangdao (Approval No. 20180305). All subjects signed an informed consent form before surgery.

### 2.2 Cell culture

Astrocyte cell line NHA and human glioma cell lines (U-87 MG, U-138 MG, U-118 MG, T98-G, LN-229, LN-18) were procured from ScienCell (San Diego, CA, USA) or American Type Culture Collection (ATCC) (Manassas, VA, USA), respectively. All cells were cultured in Roswell's Park Memorial Institute 1640 (RPMI-1640) medium (Gibco, Carlsbad, CA, USA) containing 10% fetal bovine serum (FBS) (Gibco, Grand Island, NY, USA), 100 U/mL penicillin (Mediatech, Manassas, VA, USA) and 0.1 mg/mL streptomycin (Mediatech, Manassas, VA, USA) at 37°C with 5% CO<sub>2</sub>.

### 2.3 Cell transfection

Empty plasmid (negative control plasmid, NC), circ\_0000418 overexpression plasmid (circ\_0000418), si-NC, two circ\_0000418 siRNAs (si-circ

\_0000418-1 and si-circ\_0000418-2), miR-negative control (NC), miR-409 mimics, inhibitor-negative control (NC), miR-409-3p inhibitors (miR inhibitors) were procured from GenePharma Co., Ltd. (Shanghai, China). LN-229 cells and T98-G cells were planted into 24-well plates at  $3 \times 10^5$  cells/well and incubated at 37°C with 5% CO<sub>2</sub> for 24 h before cell transfection. The above oligonucleotides/vectors were transfected into LN-229 and T98-G cells using Lipofectamine® 3000 (Invitrogen, Carlsbad, CA, USA) according to the protocols provided by the manufacturer. Transfection efficiency was measured by quantitative real-time PCR (qPCR) 48 h after the transfection.

## 2.4 qPCR

The RNA was extracted from tissues and cell lines by TRIzol reagent (Invitrogen, Shanghai, China), and after determination of concentration and purity, the total RNA was reverse transcribed into cDNA with the RevertAid RT kit (Thermo Fisher Science, Waltham, MA, USA). For the reverse transcription of miRNA, miRNA First Strand cDNA Synthesis Kit (Sigma-Aldrich, Louis, MO, USA) was used. Subsequently, the cDNA (1ng/μL) was used as the template for DNA amplification with the qRT-PCR assay kit (Stratagene, La Jolla, CA, USA). Reaction system (20 μL): 2 μL of cDNA, 10 μL SYBR Green Mix (Takara, TX, USA), 0.4 μL of ROX Reference Dye, 0.8 μL of forward primer, 0.8 μL of reverse primer, 6 μL of ddH<sub>2</sub>O. The DNA amplification was performed on Applied Biosystems 7500 Real-time system (Applied Biosystems; Thermo Fisher Scientific, Inc., Foster City, CA, USA). PCR thermal cycling parameters were as follows: 95°C for 5 min, with a 3-step reaction: 94°C denaturation at 94°C for 30s and annealing at 60°C for 30s. 45 cycles were performed. Circ\_0000418, miR-409-3p and PDK1 mRNA relative expressions were calculated using the  $2^{-\Delta\Delta C_t}$  method using GAPDH and U6 as the internal references. The primer sequences are as follows:

hsa\_circ\_0000418 Forward: 5'-TGG AGTAAATCAACCAAACGA-3' and reverse 5'-TGTGCCCGCAATATTCATTA-3'.

miR-409-3p Forward: 5'-GAATGTTGCTCG GTGA-3' and reverse 5'-GTGCAGGGTCCG AGGT-3'.

PDK1 mRNA Forward: 5'-CTGGCTGGAT TTGGTTACGG-3' and reverse 5'-ACTCCGTTG ACAGAGCCTTAAT-3'

GAPDH Forward: 5'-AATCCCATCACCAT CTTCC-3' and reverse 5'-CATCAGCCACAGTTTCC-3'.

U6 forward: 5'-CGCTTCGGCAGCACAT ATAC-3' and reverse 5'-AACGCTTCACGAATT TGCCT-3'.

## 2.5 Cell counting kit-8 (CCK-8) experiment

The transfected cells were inoculated into 96-well plates ( $2 \times 10^3$  per well). 10 μL of CCK-8 reagent (Dojindo, Rockville, MD, USA) was then supplemented to each well at 24 h, 48 h, and 72 h and incubated for 2 h. The absorbance values of each well were appraised at 450 nm using a Multiskan™ spectrophotometer (Thermo Fisher Scientific, Grand Island, NY, USA).

## 2.6 BrdU assay

Briefly, LN-229 and T98-G cells were plated in 24-well plates at  $2.5 \times 10^4$  cells/well, and cultured for 24 h, followed by the addition of BrdU solution (Beyotime Biotechnology, Shanghai, China) to incubate for another 4 h. Then, the cells were fixed, and incubated with anti-BrdU antibody for 2 h, and then incubated with secondary antibodies for 1 h. The nucleus was marked by 10 μmol/L DAPI for 5 min. Ultimately, an upturned fluorescence microscope was utilized for observing the cells, and the percentage of BrdU-positive cells was calculated.

## 2.7 Flow cytometry

The transfected LN-229 and T98-G cells were fixed in 70% ethanol overnight at 4°C overnight. On the following day, the glioma cells were washed with phosphate buffer saline (PBS) and resuspended in 100 μL of PBS, and 5 μL of propidium iodide solution (PI, BD Bioscience, Franklin Lakes, NJ, USA) was added to stain the cells for 30 min. After the cells were washed by PBS again, a flow cytometer

was employed to detect the cell cycles. FSC v.s. SSC figure was used to exclude the cell debris (R1); FSC v.s. PI figure was used to exclude the dead cells (R2). Then the cell cycle distribution was analyzed, with X-axis indicating the DNA content, and the Y-axis indicating the number of the cells.

## 2.8 Western blot assay

LN-229 and T98-G cells were lysed using RIPA buffer (Beyotime, Shanghai, China). A BCA Protein Assay Kit (Beyotime, Shanghai, China) was used to quantify the protein concentrations. 25  $\mu$ g of protein sample in each group was separated by SDS-PAGE and then transferred into the polyvinylidene fluoride (PVDF) membrane. Afterward, the membranes were incubated with primary antibodies anti-PDK1 (ab202468, 1:1000, Abcam, Cambridge, UK) and anti-GAPDH (ab9485, 1:1000, Abcam, Cambridge, UK) overnight at 4°C. The membrane was cleaned 3 times for 5 min each time with tris buffered saline tween, and then incubated with the secondary antibody Goat Anti-Rabbit IgG H&L (ab150077, 1:2000, Abcam, Cambridge, UK) for 1 h at 37°C. Finally, the protein bands were visualized using the enhanced chemiluminescence (ECL) kit (Beyotime, Shanghai, China).

## 2.9 Dual-luciferase reporter gene experiment

The sequences of circ\_0000418 or 3'UTR of PDK1 including the wild type (WT) or mutated type (MUT) binding sites of miR-409-3p were amplified and inserted into the luciferase reporter plasmid psiCHECK2 (Promega, Madison, WI, USA) to construct wild-type circ\_0000418 (WT circ\_0000418), wild-type PDK1 (WT PDK1), mutant-type circ\_0000418 (MUT circ\_0000418) and mutant-type PDK1 (MUT PDK1) reporting vectors. The above vectors were co-transfected with miR-409-3p mimics or miR-409-3p inhibitors into LN-229 and T98-G cells, respectively. After 48 h of transfection, the luciferase activity was determined using the dual-luciferase Reporter Assay Kit (Promega, Fitchburg, WI, USA).

## 2.10 RNA immunoprecipitation (RIP) experiment

Following the protocols provided by the manufacturer, a Magna RIP Kit (Millipore, Billerica, MA, USA) was applied in the RIP experiment to validate the interaction between circ\_0000418 and miR-409-3p. Briefly, the transfected LN-229 and T98-G cells were lysed in RIP lysis buffer, and the cell extracts were incubated with magnetic beads coated with anti-IgG or anti-Ago2. Then the immunoprecipitate was incubated with Proteinase K. The purified RNA was analyzed using qPCR to detect the enrichment of circ\_0000418 and miR-409-3p in the immunoprecipitate.

## 2.11 Bioinformatics analysis


GSE146463 was downloaded from GEO database (<https://www.ncbi.nlm.nih.gov/gds>). This dataset includes the circRNA microarray data of glioblastoma cells (n = 8) and neural progenitor cells (n = 3). The data of GSE146463 was analyzed by GEO2R. The circRNAs with  $P < 0.05$  and  $|\log_2 \text{fold change}| > 1$  were regarded as the differentially expressed circRNAs. Additionally, Circular RNA Interactome database (<https://circinteractome.nia.nih.gov>) [11], StarBase (<https://starbase.sysu.edu.cn>) [12], TartgetScan 7.1 database ([http://www.tartgetscan.org/vert\\_71](http://www.tartgetscan.org/vert_71)) [13] and miRWalk database (<http://mirwalk.umm.uni-heidelberg.de>) [14] were searched for the targeting relationship among circ\_0000418, miR-409-3p and PDK1.

## 2.12 Statistical analysis

All statistical analysis was performed using GraphPad Prism Ver 8.0. D'Agostino & Pearson omnibus normality test was used to analyze the normality of the data. All of the data were presented as mean  $\pm$  standard deviation (SD), and differences analysis was conducted using Student's *t*-test or one-way ANOVA. Correlations between the data were analyzed by Pearson's correlation analysis.  $P < 0.05$  denoted statistical significance.

## 3 Results

The goal of this work was to explore the differentially expressed circRNAs in glioma progression.



**Figure 1.** Circ\_0000418 expression is significantly up-regulated in glioma a. The microarray dataset GSE146463 was downloaded from GEO database to analyze the differentially expressed circRNAs between glioblastoma patient derived cells and neural progenitor cells, and the volcano plot showed these circRNAs. The yellow indicated the down-regulated circRNAs, and the blue indicated the up-regulated circRNAs. b. Heat map was generated to show 8 representative up-regulated circRNAs in glioma tissues. c. Circ\_0000418 expression in 47 cases of glioma tissues and normal tissues was measured by qPCR. d. Circ\_0000418 expression levels in NHA cells and glioma cells (U-87 MG, U-138 MG, U-118 MG, T98-G, LN-229, LN-18) were detected by qPCR. \*\*  $P < 0.01$ , \*\*\*  $P < 0.001$ .

Bioinformatics helps us obtain a up-regulated circRNA in glioma, circ\_0000418. Then we used a series of experiments to investigate its expression characteristics in glioma tissues and cell lines, its function in the growth and cell cycle of glioma cells, and its downstream mechanism.

### 3.1 Circ\_0000418 is highly expressed in glioma tissues and cell lines

To probe circ\_0000418 expression in glioma, the microarray dataset GSE146463 was accessed from the GEO database to examine the differentially expressed circRNAs between glioblastoma patient derived cells and neural progenitor cells. The


results suggested that 540 circRNAs were down-regulated and 823 circRNAs were up-regulated in glioma (Figure 1(a)), and a heat map of 8 up-regulated circRNAs was generated (Figure 1(b)). Among these, circ\_0000418 was chosen for the subsequent investigation. qPCR was conducted to detect circ\_0000418 expression in tumor tissues and non-cancerous tissues of 47 glioma patients. The data implied that circ\_0000418 expression was remarkably up-regulated in glioma tissues ( $P < 0.001$ ) (Figure 1(c)). Also, circ\_0000418 expression in glioma cells (U-87 MG, U-138 MG, U-118 MG, T98-G, LN-229, LN-18) was also significantly up-regulated compared to that in NHA ( $P < 0.01$ ) (Figure 1(d)).

### 3.2 Circ\_0000418 promotes glioma cell growth, and cell cycle progression


Among the six glioma cells, circ\_0000418 expression level was the lowest in LN-229 cells and the highest in T98-G cells. Hence, circ\_0000418 overexpression plasmid and circ\_0000418 siRNAs were transfected into LN-229 cells and T98-G cells, and we successfully constructed circ\_0000418 overexpression and knockdown models (Figure 2(a,b)). CCK-8 experiment, BrdU experiment and flow cytometry showed that circ\_0000418 overexpression promoted LN-229 cell growth and accelerates the cell cycle as opposed to the NC group ( $P < 0.05$ ) (Figure 2(c-e)). Contrariwise, in T98-G cells, the knockdown of circ\_0000418 suppressed glioma cell growth and cell cycle progression (Figure 2(c-e)).

### 3.3 Circ\_0000418 directly targets miR-409-3p

To validate the targeting correlation between circ\_0000418 and miR-409-3p, the subcellular localization of circ\_0000418 in glioma cells was identified using qPCR. The data implied that circ\_0000098 was primarily found in the cytoplasm ( $P < 0.001$ ) (Figure 3(a)). MiRNAs that can interact with circ\_0000418 were predicted by Circular RNA Interactome, a possible binding site between circ\_0000418 and miR-409-3p was discovered (Figure 3(b)). To validate the targeting relationship between the two, the dual-luciferase reporter gene assays were performed. Transfection with miR-409-3p mimics significantly inhibited the luciferase activity of wild-type circ\_0000418 reporter ( $P < 0.01$ ), while there was no remarkable alteration in the luciferase activity for mutated circ\_0000418 reporter ( $P > 0.05$ ) (Figure 3(c)).



**Figure 2.** The biological function of circ\_0000418 in glioma a-b. qPCR was carried out to detect the transfection efficiency of circ\_0000418 overexpression plasmid and two circ\_0000418 siRNAs. c-d. The effects of overexpression or knockdown of circ\_0000418 on the growth of LN-229 and T98-G cells were analyzed using CCK-8 assay (c) and BrdU assay (d). e. Flow cytometry was executed to detect the effect of overexpression or knockdown of circ\_0000418 on the cell cycle progression of LN-229 and T98-G cells. \*  $P < 0.05$ , \*\*  $P < 0.01$ , \*\*\*  $P < 0.001$ .



**Figure 3.** Circ\_0000418 directly targets miR-409-3p. **a.** The subcellular location of circ\_0000418 in LN-229 and T98-G cells was evaluated by qPCR after that the nuclear and cytoplasmic fractions of glioma cells were isolated. **b.** The Circular RNA Interactome online database was employed to predict the binding site of circ\_0000418 to miR-409-3p. **c.** WT circ\_0000418 and MUT circ\_0000418 were co-transfected with miR mimics or miR inhibitors into LN-229 and T98-G cells, respectively, and the dual-luciferase reporter gene system was employed to measure luciferase activity. **d.** The interaction between circ\_0000418 and miR-409-3p was further validated using RIP assay. **e.** The effect of overexpression/knockdown of circ\_0000418 on miR-409-3p expression in glioma cells was examined by qPCR. **f.** MiR-409-3p expression in 47 cases of glioma tissues and normal tissues was examined by qPCR. **g.** Pearson correlation analysis was performed to analyze the correlation between miR-409-3p expression and circ\_0000418 expression in glioma tissues. \*\*  $P < 0.01$ , \*\*\*  $P < 0.001$ .

Furthermore, RIP assays validated that circ\_0000418 and miR-409-3p were enriched by the Ago2-antibody compared with control IgG ( $P < 0.001$ ) (Figure 3(d)). Furthermore, circ\_0000418 overexpression was unveiled to substantially impede miR-409-3p expression in LN-229 cells, while in T98-G cells ( $P < 0.01$ ), knock-down of circ\_0000418 caused an increase in miR-

409-3p expression ( $P < 0.001$ ) (Figure 3(e)). Notably, miR-409-3p expression was markedly down-regulated in glioma tissues compared with that in non-cancerous brain tissues ( $P < 0.01$ ) (Figure 3(f)), and circ\_0000418 expression was negatively correlated with miR-409-3p expression in glioma tissues ( $r = -0.6199$ ,  $P < 0.001$ ) (Figure 3(g)).


### 3.4 Circ\_0000418 participates in regulating the biological behaviors of glioma cells by sponging miR-409-3p

To probe the role of miR-409-3p in the progression of glioma, circ\_0000418 overexpression plasmid+miR-409-3p mimic and si-circ\_0000418-2 + miR-409-3p inhibitors were co-transfected into LN-229 cells and T98-G cells, respectively. qPCR manifested that the transfection was successful (Figure 4(a)). Subsequently, the CCK-8 experiment, BrdU experiment and flow cytometry showed that compared with the circ\_0000418 group, co-transfection of miR-409-3p mimics into LN-229 cells significantly inhibited cell

growth and blocked the cell cycle progression ( $P < 0.05$ ) (Figure 4(b-e)). On the other hand, compared with si-circ\_0000418-2 alone, co-transfection of si-circ\_0000418-2 and miR-409-3p inhibitors into T98-G cells promoted cell growth and facilitated the cell cycle progression ( $P < 0.05$ ) (Figure 4(b-e)).

### 3.5 Circ\_0000418 up-regulates PDK1 expression through repressing miR-409-3p

To look for the downstream targets of miR-409-3p, StarBase, TartgetScan 7.1 and miRWalk databases were utilized to screen the candidate targets




**Figure 4.** Effects of Circ\_0000418 and miR-409-3p on the biological functions of glioma cells a. The transfection efficiency of circ\_0000418 overexpression plasmid+miR mimics and si-circ\_0000418-2+ miR inhibitors in LN-229 and T98-G cells was examined by qPCR. B-C. CCK-8 assay (b) and BrdU experiment (c) were applied to detect the effects of circ\_0000418+ miR-409-3p mimics or si-circ\_0000418+ miR-409-3p inhibitor on cell growth. d. The effect of circ\_0000418+ miR-409-3p mimics or si-circ\_0000418+ miR-409-3p inhibitor on cell cycle was examined by flow cytometry. \*  $P < 0.05$ , \*\*  $P < 0.01$ , \*\*\*  $P < 0.001$ .



of miR-409-3p, and 162 genes including PDK1 were unearthed to be the candidate targets of miR-409-3p (Figure 5(a)). Additionally, the KEGG database was utilized to conduct pathway enrichment analysis on the potential downstream target genes of miR-409-3p, and the results

suggested that the above target genes were correlated with the triggering of the Wnt signaling pathway (Figure 5(b)). To validate whether miR-409-3p could bind with PDK1 mRNA 3'-UTR, a luciferase reporter experiment was implemented. The data showed that transfection of miR-409-3p



**Figure 5.** Circ\_0000418 up-regulates PDK1 expression through adsorption of miR-409-3p a. The downstream targets of miR-409-3p were predicted through bioinformatics websites StarBase, TargetScan and miRWalk. b. Enrichment analysis of downstream target genes of miR-409-3p was conducted using the KEGG database. c. The binding site between miR-409-3p and PDK1 mRNA 3' UTR. d. WT PDK1 and MUT PDK1 were co-transfected with miR-409-3p mimics and miR-409-3p inhibitors into 293 T cells, and then the luciferase activity was evaluated. e. Western blot was applied to detect the effects of circ\_0000418 and miR-409-3p on PDK1 expression. f. qPCR was implemented to detect PDK1 mRNA expression in 47 pairs of glioma tissues and non-cancerous brain tissues. G-H. Pearson correlation analysis was performed to analyze the correlation between circ\_0000418, miR-409-3p and PDK1 mRNA expression in glioma tissues. \*  $P < 0.05$ , \*\*  $P < 0.01$ , \*\*\*  $P < 0.001$ .

mimics remarkably inhibited the luciferase activity of the 3'-UTR of wild-type PDK1, and transfection of miR-409-3p inhibitors remarkably increased the luciferase activity of the 3'-UTR of wild-type PDK1 ( $P < 0.05$ ), while the two had no significant effects on the luciferase activity of MUT-PDK1 ( $P > 0.05$ ) (Figure 5(c,d)). Additionally, the data of Western blot also manifested that circ\_0000418 overexpression significantly promoted PDK1 protein expression ( $P < 0.001$ ), while co-transfection of miR-409-3p mimics weakened this effect ( $P < 0.01$ ); knockdown of circ\_0000418 significantly inhibited PDK1 protein expression ( $P < 0.01$ ), while suppression of miR-409-3p reversed this effect ( $P < 0.01$ ) (Figure 5(e)). Meanwhile, qPCR presented that PDK1 mRNA was substantially overexpressed in tumor tissues ( $P < 0.001$ ) (Figure 5(f)). In addition, miR-409-3p and PDK1 mRNA expressions were negatively correlated ( $r = -0.6608$ ,  $P < 0.001$ ), while circ\_0000418 and PDK1 mRNA expressions were positively associated ( $r = 0.6509$ ,  $P < 0.001$ ) (Figure 5(g,h)).

#### 4 Discussion

CircRNAs are widely present in mammalian cells [3], affecting different biological activities, including cell growth and differentiation. Accumulating studies have shown that circRNA is differentially expressed between tumor tissues and corresponding paracancerous tissues, and can be employed as novel biomarkers and potential therapeutic targets for diverse malignancies [15,16]. In glioma. Circ-FBXW7 is downregulated in gliomas and correlated with shorter overall survival time of the patients [17]. Circ\_01844 expression is down-regulated in glioma tissues and cells, and circ\_01844 overexpression significantly impedes glioma cell growth and migration and induces cell apoptosis [18]. Circ-EPB41L5 is significantly under-expressed in glioma tissues and cells, and is closely associated with poor prognosis of the patients; functionally, circ-EPB41L5 inhibits glioma cell growth, colony formation, migration and invasion [19]. Circ\_0000418 is much less investigated. A previous study reports that circ\_0000418 is associated with the pathogenesis of depression [6]. In this research, for the first

time, the expression feature and biological function of circ\_0000418 in cancer biology is investigated. Circ\_0000418 expression was revealed to be up-regulated in glioma tissues, and functional studies unveiled that circ\_0000418 remarkably enhanced cell growth and facilitated the cell cycle progression of glioma cells, implying that circ\_0000418 is a potential molecular marker for glioma and a possible therapeutic target.

MiRNA is non-coding small molecule RNA of 18–25 nucleotides in length [20]. MiRNAs regulate gene expression via binding to the 3'-UTR of target mRNAs, and they modulate about 30% of human protein-coding genes [21]. MiRNAs are implicated in diverse biological activities, and they are also involved in the tumorigenesis of human malignancies [22–26]. Reportedly, miR-409-3p is remarkably under-expressed in cervical cancer tissues and cells, and miR-409-3p overexpression specifically down-regulated the expression level of *activating transcription factor 1*, inhibiting cell growth, migration, invasion and glycolysis [27]. MiR-409-3p is also under-expressed in breast cancer tissues and cells and correlates with TNM stage, lymph node metastasis and short patient survival [28]. In addition, the role and mechanism of miR-409-3p in glioma have been reported in some previous studies. For example, in glioma, miR-409-3p under-expression is associated with poor prognosis of the patients [29]. Additionally, miR-409-3p is significantly under-expressed in glioma and miR-409-3p impedes cell growth and invasion by negatively regulating *HBGNBD5* [7]. Similarly, in the current work, it was found that miR-409-3p was significantly under-expressed in glioma tissues and cells, and we also demonstrated that circ\_0000418 promoted glioma cell growth and promoted the cell cycle progression by specifically down-regulating miR-409-3p.

PDK1 is a serine/threonine-protein kinase with a size of 67 KD, which induces the phosphorylation of pyruvate dehydrogenase to regulate the homeostasis of carbohydrate fuels; PDK1 is vital in regulating a lot of biological processes such as cell growth, differentiation and apoptosis [9]. Importantly, PDK1 is highly expressed in cancers including breast cancer, non-small cell lung cancer and pancreatic cancer, and it functions as an

oncogene [30–32]. Notably, PDK1 promotes glioma cell migration and invasion *in vitro* and glioma xenograft growth *in vivo* through up-regulation of c-Jun protein and induction of epithelial-mesenchymal transition [33]. Another study reports that PDK1 expression is up-regulated in glioma tissues and cells; silencing of *PDK1* expression led to reduced level of lactate and ATP, accumulation of ROS, mitochondrial damage, reduced cell growth, and cell apoptosis [34]. PDK1 has been reported to be modulated by several miRNAs. A previous study identifies *PDK1* as a downstream target of miR-128-3p in glioma cells [34]. Additionally, *PDK1* is validated to be a target gene of miR-454 in glioma, and miR-454 overexpression significantly inhibits PDK1 expression, thereby suppressing cell growth and blocking the cell cycle in G0/G1 phase [35]. In the research, we found that circ\_0000418 could decoy miR-409-3p to up-regulate PDK1 expression, which provides a novel explanation of PDK1 dysregulation in glioma cells.

## 5 Conclusion

Taken together, this work found that circ\_0000418 expression was up-regulated in glioma tissues and cells. Circ\_0000418 facilitated glioma cell growth, and accelerated the cell cycle by targeting the miR-409-3p/PDK1 axis, indicating that circ\_0000418 may be a potential diagnostic marker and a new therapeutic target for glioma, which provides a theoretical basis for the clinical therapy of glioma.

## Ethics statement

Our study was approved by the Ethics Review Board of the First Hospital of Qinhuangdao.

## Disclosure statement

No potential conflict of interest was reported by the author(s).

## Funding

The author(s) reported there is no funding associated with the work featured in this article.

## ORCID

Zhihui Ma  <http://orcid.org/0000-0003-0779-3069>

## References

- [1] Chen R, Smith-Cohn M, Cohen AL, et al. Glioma Subclassifications and Their Clinical Significance. *Neurotherapeutics*. 2017;14(2):284–297.
- [2] Zhou D, Lin X, Wang P, et al. Circular RNA circ\_0001162 promotes cell proliferation and invasion of glioma via the miR-936/ERBB4 axis. *Bioengineered*. 2021;12(1):2106–2118.
- [3] Deng T, Liu Y, Yang Y, et al. Regulation of microRNA miR-197-3p/CDC28 protein kinase regulatory subunit 1B (CKS1B) axis by Circular RNA hsa\_circ\_0000285 promotes glioma progression. *Bioengineered*. 2022;13(3):4757–4772.
- [4] Xia X, Li X, Li F, et al. A novel tumor suppressor protein encoded by circular AKT3 RNA inhibits glioblastoma tumorigenicity by competing with active phosphoinositide-dependent Kinase-1. *Mol Cancer*. 2019;18(1):131.
- [5] Wu Z, Zheng M, Zhang Y, et al. Hsa\_circ\_0043278 functions as competitive endogenous RNA to enhance glioblastoma multiforme progression by sponging miR-638. *Aging (Albany NY)*. 2020;12(21):21114–21128.
- [6] Zhang X, Yan W, Xue Y, et al. Roles of miR-432 and circ\_0000418 in mediating the anti-depressant action of ADAR1. *Neurobiol Stress*. 2021;15:100396.
- [7] Cao Y, Zhang, L, Wei, M, et al. MicroRNA-409-3p represses glioma cell invasion and proliferation by targeting high-mobility group nucleosome-binding domain 5. *Oncol Res*. 2017;25(7):1097–1107.
- [8] Wang Z, Xu X, Liu N, et al. SOX9-PDK1 axis is essential for glioma stem cell self-renewal and temozolomide resistance. *Oncotarget*. 2018;9(1):192–204.
- [9] Gagliardi PA, Puliafito A, Primo L. PDK1: at the crossroad of cancer signaling pathways. *Semin Cancer Biol*. 2018;48:27–35.
- [10] Salmena L, Poliseno L, Tay Y, et al. A ceRNA hypothesis: the Rosetta Stone of a hidden RNA language? *Cell*. 2011;146(3):353–358.
- [11] Dudekula DB, Panda AC, Grammatikakis I, et al. CircInteractome: a web tool for exploring circular RNAs and their interacting proteins and microRNAs. *RNA Biol*. 2016;13(1):34–42.
- [12] Li J, Liu S, Zhou H, et al. starBase v2.0: decoding miRNA-ceRNA, miRNA-ncRNA and protein-RNA interaction networks from large-scale CLIP-Seq data. *Nucleic Acids Res*. 2014;42(D1):D92–7.
- [13] Agarwal V, Bell GW, Nam J-W, et al. Predicting effective microRNA target sites in mammalian mRNAs. *Elife*. 2015;4:e05005.

- [14] Sticht C, De La Torre C, Parveen A, et al. miRWalk: an online resource for prediction of microRNA binding sites. *PLoS One*. 2018;13(10):e0206239.
- [15] Yang Y, Yujiao W, Fang W, et al. The roles of miRNA, lncRNA and circRNA in the development of osteoporosis. *Biol Res*. 2020;53(1):40.
- [16] Xu X, Zhang J, Tian Y, et al. CircRNA inhibits DNA damage repair by interacting with host gene. *Mol Cancer*. 2020;19(1):128.
- [17] Yang Y, Gao X, Zhang M, et al. Novel Role of FBXW7 Circular RNA in Repressing Glioma Tumorigenesis. *J Natl Cancer Inst*. 2018;110(3):304–315.
- [18] Zhou JX, Chen K-F, Hu S, et al. Up-regulation of circular RNA hsa\_circ\_01844 induces apoptosis and suppresses proliferation and migration of glioblastoma cells. *Chin Med J (Engl)*. 2020;134(1):81–87.
- [19] Lv T, Miao Y, Xu T, et al. Circ-EPB41L5 regulates the host gene EPB41L5 via sponging miR-19a to repress glioblastoma tumorigenesis. *Aging (Albany NY)*. 2020;12(1):318–339.
- [20] Lu TX, Rothenberg ME. MicroRNA. *J Allergy Clin Immunol*. 2018;141(4):1202–1207.
- [21] Griffiths-Jones S, Grocock R, van Dongen S, et al. miRBase: microRNA sequences, targets and gene nomenclature. *Nucleic Acids Res*. 2006;34(90001):D140–4.
- [22] Bushati N, Cohen SM. microRNA functions. *Annu Rev Cell Dev Biol*. 2007;23(1):175–205.
- [23] Zhang L, Kong L, Yang Y. miR-18a Inhibitor Suppresses Leukemia Cell Proliferation by Upregulation of PTEN Expression. *Med Sci Monit*. 2020;26:e921288.
- [24] Yu T, Gong L, Li W, et al. MiR-30a suppresses metastasis of gastric adenocarcinoma via targeting FAP $\alpha$ . *Cancer Biomark*. 2020;27(4):471–484.
- [25] Yang Y, Tai W, Lu N, et al. lncRNA ZFAS1 promotes lung fibroblast-to-myofibroblast transition and ferroptosis via functioning as a ceRNA through miR-150-5p/SLC38A1 axis. *Aging (Albany NY)*. 2020;12(10):9085–9102.
- [26] Liu ZZ, Tian YF, Wu H, et al. LncRNA H19 promotes glioma angiogenesis through miR-138/HIF-1 $\alpha$ /VEGF axis. *Neoplasma*. 2020;67(1):111–118.
- [27] Cui X, Chen, J, Zheng, Y, et al. Circ\_0000745 Promotes the Progression of Cervical Cancer by Regulating miR-409-3p/ATF1 Axis. *Cancer Biother Radiopharm*. 2020; DOI:10.1089/cbr.2019.3392.
- [28] Cao G-H, Sun X-L, Wu F, et al. Low expression of miR-409-3p is a prognostic marker for breast cancer. *Eur Rev Med Pharmacol Sci*. 2016;20(18):3825–3829.
- [29] Kumar A, Nayak, S, Pathak, P, et al. Identification of miR-379/miR-656 (C14MC) cluster downregulation and associated epigenetic and transcription regulatory mechanism in oligodendrogliomas. *J Neurooncol*. 2018;139(1):23–31.
- [30] Kim S, Lee E, Jung J, et al. microRNA-155 positively regulates glucose metabolism via PIK3R1-FOXO3a-cMYC axis in breast cancer. *Oncogene*. 2018;37(22):2982–2991.
- [31] Huang T, Wen, Y, Peng, B, et al. Upregulated lncRNA H19 promotes non-small cell lung cancer cell proliferation through miR-138/PDK1 axis. *Int J Clin Exp Pathol*. 2017;10(8):9012–9020.
- [32] Xie F, Huang Q, Liu C-H, et al. MiR-1271 negatively regulates AKT/MTOR signaling and promotes apoptosis via targeting PDK1 in pancreatic cancer. *Eur Rev Med Pharmacol Sci*. 2018;22(3):678–686.
- [33] Luo D, Xu X, Li J, et al. The PDK1/c-Jun pathway activated by TGF- $\beta$  induces EMT and promotes proliferation and invasion in human glioblastoma. *Int J Oncol*. 2018;53(5):2067–2080.
- [34] Qu C, Yan C, Cao W, et al. miR-128-3p contributes to mitochondrial dysfunction and induces apoptosis in glioma cells via targeting pyruvate dehydrogenase kinase 1. *IUBMB Life*. 2020;72(3):465–475.
- [35] Fang B, Zhu J, Wang Y, et al. MiR-454 inhibited cell proliferation of human glioblastoma cells by suppressing PDK1 expression. *Biomed Pharmacother*. 2015;75:148–152.