



Investigation of the cytotoxicity of bioinspired coumarin analogues towards human breast cancer cells

Leonidas Gkionis¹ · Eleni Kavetsou² · Alexandros Kalospyros² · Dimitris Manousakis² · Miguel Garzon Sanz¹ · Sam Butterworth^{1,3} · Anastasia Detsi² · Annalisa Tirella^{1,3}

Received: 7 February 2020 / Accepted: 1 April 2020 / Published online: 23 April 2020
© The Author(s) 2020

Abstract

Coumarins possess a wide array of therapeutic capabilities, but often with unclear mechanism of action. We tested a small library of 18 coumarin derivatives against human invasive breast ductal carcinoma cells with the capacity of each compound to inhibit cell proliferation scored, and the most potent coumarin analogues selected for further studies. Interestingly, the presence of two prenyloxy groups (5,7-diprenyloxy-4-methyl-coumarin, **4g**) or the presence of octyloxy substituent (coumarin **4d**) was found to increase the potency of compounds in breast cancer cells, but not against healthy human fibroblasts. The activity of potent compounds on breast cancer cells cultured more similarly to the conditions of the tumour microenvironment was also investigated, and increased toxicity was observed. Results suggest that tested coumarin derivatives could potentially reduce the growth of tumour mass. Moreover, their use as (combination) therapy in cancer treatment might have the potential of causing limited side effects.

Leonidas Gkionis and Eleni Kavetsou: Co-first authors.

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s11030-020-10082-6>) contains supplementary material, which is available to authorized users.

✉ Anastasia Detsi
adetsi@chemeng.ntua.gr


✉ Annalisa Tirella
annalisa.tirella@manchester.ac.uk

¹ Division of Pharmacy and Optometry, Faculty of Biology, Medicine and Health, Manchester Academic Health Science Centre, University of Manchester, Oxford Road, Manchester M13 9PL, UK

² Laboratory of Organic Chemistry, School of Chemical Engineering, National Technical University of Athens, Heroon Polytechniou 9, Zografou Campus, 15780 Athens, Greece

³ NorthWest Centre for Advanced Drug Delivery (NoWCADD), Faculty of Biology, Medicine and Health, University of Manchester, Oxford Road, Manchester M13 9PT, UK

Graphic abstract



Keywords Alkoxy-coumarins · Auraptene · Umbelliprenin · Breast cancer

Introduction

Breast carcinoma is considered the predominant and more common malignancy in women worldwide, with one in eight women potentially developing breast cancer during their lifetime [1] and predictions of 3.2 million newly diagnosed cases per year by 2050 [2]. Early detection and intervention are essential to increase patients' survival rate, yet the treatment of advanced cancer remains an issue. While many biological and physicochemical factors have been identified in cancer development, there is an increasing interest in the role of inflammation and involvement of stromal component of the tumour microenvironment [3]. The challenge of treating breast cancer resides not only in the identification of active compounds capable of targeting the cancer, but mainly in identifying potent therapies with low side effects [4]. In this perspective, natural compounds like coumarins have gained significant interest in the recent years for their numerous pharmacological activities including chemopreventive and antiproliferative properties against various cancer types [5–8].

Coumarins are heterocyclic organic compounds that are widely distributed in nature [4, 9]. Coumarin derivatives have gained high scientific interest as promising drug candidates since they possess multiple pharmacological

properties [10–13], such as antioxidant [14–16], antibacterial [17, 18], antimicrobial [18], antiviral [13], hepatoprotective [19] and anti-inflammatory effects [20–22]. Natural and synthetic coumarins have been also reported as effective chemopreventive and anticancer agents *in vitro* [23–26] and *in vivo* [27].

Chemical modification such as alkylation (the addition of unsaturated or saturated chain to the coumarin scaffold) has been shown to enhance the pharmacological profile of several coumarins, especially their anticancer activity [25]. In particular, the insertion of an unsaturated chain (prenyl, geranyl or farnesyl side chain) is known as prenylation and constitutes a metabolic pathway in nature (including plant kingdom and microorganism such as fungi and bacteria [28]). The process of prenylation is considered to further enhance the pharmacological activity of these metabolites mostly because it strengthens the lipophilicity of the molecules [29]. Recently, natural oxyprenylated coumarins (isopentenylxy (C5), geranylxy (C10) and farnesylxy (C15) compounds and their biosynthetic derivatives) have been studied for their pharmacological properties [28], mainly as potential anticancer agents [30]. Auraptene (7-geranylxy coumarin) and umbelliprenin are the most common plant-derived oxyprenylated coumarins, first isolated from citrus fruits and *Ferula* plant species, respectively, and present a wide range of bioactivities [25, 31–35]. The addition of an

aliphatic chain to the coumarin scaffold is another modification shown to have anticancer effects as reported by Farley et al. [25], who reported that octyloxy-coumarins possess cytotoxicity against pancreatic cancer cells with concentrations in the order of tens of nM. As a continuation to our previous work concerning the biological evaluation of structurally modified coumarin analogues [14, 15], a series of bioinspired synthetic alkoxy coumarin derivatives (bearing saturated and unsaturated chains) were synthesized, structurally characterized and evaluated for their cytotoxicity against breast cancer cell lines (MCF-7 and MDA-MB-231) and fibroblasts. Interestingly we found that the more potent coumarin compounds have no effect on fibroblast (off-target control) and increase their potency on breast cancer cells cultured under nutrient-deprived conditions similar to the tumour microenvironment.

Materials and methods

Coumarins analogues

Synthesis

The chemicals used for synthesis and analysis were purchased from Sigma-Aldrich or Alfa Aesar (7-hydroxycoumarin, 98%) and used without further purification. NMR spectra were recorded on a Varian 300 MHz and 600 MHz spectrometer at the Institute of Chemical Biology of the National Hellenic Research Foundation. The HRMS spectra were obtained using a UHPLC-MSn Orbitrap Velos-Thermo mass spectrometer. Melting points were determined on a Gallenkamp MFB-595 melting point apparatus and are uncorrected.

General procedure for the synthesis of hydroxy or dihydroxy-4-substituted coumarin analogues The desired compounds **3a** and **3b** were synthesized according to the method of Prousis et al. [36].

7-Hydroxy-4-propyl-2H-chromen-2-one (3a) Beige solid; yield 80% (735.4 mg, 3.60 mmol); mp 130 °C (lit. m.p. 127–128 °C) [36]. ¹H NMR (600 MHz, DMSO-*d*₆): δ(ppm) 10.51 (s, 1H, 7-OH), 7.64 (d, *J*=9.0 Hz, 1H, H-5), 6.79 (d, *J*=8.4 Hz, 1H, H-6), 6.71 (s, 1H, H-8), 6.08 (s, 1H, H-3), 2.70 (t, *J*=7.5 Hz, 2H, 4-CH₂CH₂CH₃), 1.62 (m, 2H, 4-CH₂CH₂CH₃), 0.96 (t, *J*=7.2 Hz, 3H, 4-CH₂CH₂CH₃).

5,7-Dihydroxy-4-methyl-2H-chromen-2-one (3b) Beige solid; yield 93% (707.2 mg, 3.68 mmol); mp 288–289 °C (lit. m.p. 289–290 °C) [36]. ¹H NMR (600 MHz, DMSO-*d*₆): δ(ppm) 10.51 (s, 1H, 7-OH), 10.29 (s, 1H, 5-OH), 6.25 (s, 1H, H-8), 6.16 (s, 1H, H-6), 5.84 (s, 1H, H-3), 2.48 (d, *J*=9.6 Hz, 3H, 4-CH₃).

Synthesis of geranylgeranyl iodide The following method was adapted from Alvarez-Manzaneda et al. [37]; briefly, 1170.0 mg (1.72 mmol, 1 eq.) of imidazole and 450.0 mg (1.72 mmol, 1 eq.) of triphenylphosphine were dissolved in 10 mL of anhydrous dichloromethane (DCM) in a round-bottom flask. 435.0 mg (1.72 mol, 1 eq.) of iodine was added slowly, and the mixture was stirred for 30 min. Then, the flask was covered with aluminium foil and placed in an ice bath, followed by slow addition of 0.57 mL (1.72 mol, 1 eq.) of geranylgeranyl. The mixture was stirred for approximately 2 h. After the reaction was complete (monitored by TLC in pure hexane), the mixture was filtered through a plug of silica, which was then washed with pure hexane. The solvent was evaporated in *vacuo*, resulting in a dark oily film. Yield 52% (51.9 mg).

General procedure for the synthesis of alkoxy-coumarins 4a–4m One eq. of the hydroxy- or dihydroxy-4-substituted coumarins, **3a–3c**, and 1 eq. of potassium carbonate (K₂CO₃) were dissolved in dry acetone. Then, 1.2 eq. of the appropriate alkoxy-bromide or geranylgeranyl iodide was added dropwise at room temperature, and the mixture was refluxed for 6 h. After the completion of the reaction, K₂CO₃ was filtrated, the precipitate was washed with acetone and the solvent was removed in *vacuo*. The desired products were purified via silica gel column chromatography in a solvent system of petroleum ether/ethyl acetate (9:1). Diprenyloxy coumarins were obtained in high purity after two steps of silica gel chromatography.

7-Prenyloxy-4-methyl-2H-chromen-2-one (4a) White solid; yield 70% (61.1 mg, 0.25 mmol); mp 84 °C (lit. m.p. 84–86 °C) [38]. ¹H NMR (300 MHz, CDCl₃): δ(ppm) 7.48 (d, *J*=8.7 Hz, 1H, H-5), 6.86 (dd, *J*=8.7 Hz & *J*=2.4 Hz, 1H, H-6), 6.82 (d, *J*=2.4 Hz, 1H, H-8), 5.48 (t, *J*=6.9 Hz, 1H, H-2'), 4.59 (d, *J*=6.9 Hz, 2H, H-1'), 2.42 (s, 3H, 4-CH₃), 1.83 (s, 3H, 3'-CH₃), 1.79 (s, 3H, 4'-CH₃).

7-Geranyloxy-4-methyl-2H-chromen-2-one (4b) [14] Brown gummy solid; yield 68% (296.8 mg, 0.95 mmol). ¹H NMR (300 MHz, DMSO-*d*₆): δ(ppm) 7.64 (d, *J*=8.4 Hz,

1H, H-5), 6.94–6.91 (m, 2H, H-6 & H-8), 6.18 (s, 1H, H-3), 5.42 (t, $J=6.0$ Hz, 1H, H-2'), 5.03 (br, 1H, H-6'), 4.65 (d, $J=6.6$ Hz, 2H, H-1'), 2.39 (s, 3H, 4-CH₃), 2.08–2.06 (m, 4H, H-4' & H-5'), 1.73 (s, 3H, 3'-CH₃), 1.61 (s, 3H, 8'-CH₃), 1.56 (s, 3H, 7'-CH₃).

7-Farnesyloxy-4-methyl-2H-chromen-2-one (4c) Yellow gummy solid; yield 60% (319.6 mg, 0.84 mmol). ¹H NMR (300 MHz, CDCl₃): δ (ppm) 7.48 (d, $J=8.7$ Hz, 1H, H-5), 6.86 (dd, $J=8.7$ Hz & $J=1.8$ Hz, 1H, H-6), 6.82 (d, $J=1.8$ Hz, 1H, H-8), 6.13 (s, 1H, H-3), 5.47 (t, $J=6.9$ Hz, 1H, H-2'), 5.10–5.07 (m, 1H, H-6'), 4.60 (d, $J=6.6$ Hz, 1H, H-1'), 2.39 (s, 3H, 4-CH₃), 2.12–1.95 (m, 8H, H-4' & H-5' & H-8' & H-9'), 1.76 (s, 3H, 3'-CH₃), 1.67 (s, 3H, 7'-CH₃), 1.59 (s, 6H, 11'-CH₃ & 12'-CH₃). ¹³C NMR (75 MHz, DMSO-*d*₆): δ (ppm) 162.0, 160.6, 155.1, 153.9, 141.5, 135.1, 131.1, 126.8, 124.5, 123.9, 119.5, 113.5, 113.1, 111.5, 101.8, 65.6, 39.7, 39.3, 26.6, 26.0, 25.9, 18.6, 17.9, 16.9, 16.3. HRMS (ESI) calcd for C₂₅H₃₂O₃Na: m/z : 403.2244, found: 403.2245.

4-Methyl-7-octyloxy-2H-chromen-2-one (4d) Pale yellow solid; yield 55% (180 mg, 0.62 mmol); mp 51 °C (lit. m.p. 48–50 °C) [24]. ¹H NMR (600 MHz, DMSO): δ (ppm) 8.58 (d, $J=8.4$ Hz, 1H, H-5), 6.78 (dd, $J=9$ Hz & $J=2.4$ Hz, 1H, H-6), 6.68 (d, $J=2.4$ Hz, 1H, H-8), 6.15 (s, 1H, H-3), 4.06 (t, $J=6.6$ Hz, 2H, H-1'), 2.39 (s, 3H, 4-CH₃), 1.72 (m, 2H, H-2'), 1.40 (m, 2H, H-7'), 1.29 (m, 8H, H-3' & H-5' & H-4' & H-6'), 0.86 (t, $J=6.9$ Hz, 3H, 7'-CH₃).

7-Prenyloxy-4-propyl-2H-chromen-2-one (4e) Yellow solid; yield 60% (135.0 mg, 0.49 mmol); mp 89 °C. ¹H NMR (600 MHz, CDCl₃): δ (ppm) 7.51 (d, $J=8.4$ Hz, 1H, H-5), 6.85 (dd, $J=9.0$ Hz & $J=2.4$ Hz, 1H, H-6), 6.82 (d, $J=2.4$ Hz, 1H, H-8), 6.11 (s, 1H, H-3), 5.47 (t, $J=6.9$ Hz, 1H, H-2'), 4.57 (d, $J=7.2$ Hz, 2H, H-1'), 2.69 (t, $J=7.5$ Hz, 2H, 4-CH₂CH₂CH₃), 1.80 (s, 3H, 3'-CH₃), 1.76 (s, 3H, 4'-CH₃), 1.72–1.69 (m, 2H, 4-CH₂CH₂CH₃), 1.04 (t, $J=7.5$ Hz, 3H, 4-CH₂CH₂CH₃). ¹³C NMR (150 MHz, CDCl₃): δ (ppm) 161.9, 161.7, 156.5, 155.6, 139.3, 125.4, 118.8, 113.0, 112.9, 110.9, 101.8, 77.6, 77.2, 76.7, 65.5, 33.9, 25.9, 21.6, 18.4, 14.1. HRMS (ESI) calcd for C₁₇H₂₁O₃ (M+H)⁺: m/z : 273.1485, found: 273.1485.

7-Octyloxy-4-propyl-2H-chromen-2-one (4f) Beige solid; yield 60% (350.0 mg, 1.11 mmol); mp 47 °C. ¹H NMR (600 MHz, CDCl₃): δ (ppm) 7.51 (d, $J=9.0$ Hz, 1H, H-5), 6.84 (dd, $J=8.4$ Hz & $J=1.8$ Hz, 1H, H-6), 6.81 (d, $J=2.4$ Hz, 1H, H-8), 6.12 (s, 1H, H-3), 4.01 (t, $J=6.6$ Hz, 2H, H-1'), 2.71 (t, $J=7.5$ Hz, 2H, 4-CH₂CH₂CH₃), 1.80 (m, 2H, H-2'), 1.73 (m, 2H, 4-CH₂CH₂CH₃), 1.45 (m, 2H, H-7'), 1.31 (m, 8H, H-3' & H-5' & H-4' & H-6'), 1.04 (t, $J=7.2$ Hz,

3H, 4-CH₂CH₂CH₃), 0.88 (t, $J=6.9$ Hz, 3H, 7'-CH₃). ¹³C NMR (150 MHz, CDCl₃): δ (ppm) 162.2, 161.8, 156.6, 155.7, 125.4, 112.9, 110.8, 101.6, 68.8, 33.9, 31.9, 29.4, 29.3, 29.1, 26.1, 22.8, 21.7, 14.2, 14.1. HRMS (APCI) calcd for C₂₀H₂₉O₃(M+H)⁺: m/z : 317.2116, found: 317.2105.

5,7-Diprenyloxy-4-methyl-2H-chromen-2-one (4g) Green solid; yield 40% (272.6 mg, 0.83 mmol); mp 90 °C. ¹H NMR (600 MHz, CDCl₃): δ (ppm) 6.43 (d, $J=2.4$ Hz, 1H, H-6), 6.31 (d, $J=1.8$ Hz, 1H, H-8), 5.93 (d, $J=0.6$ Hz, 1H, H-3), 5.47 (pseudotriplet, 2H, H-2' & H-2''), 4.53 (dd, $J=7.2$ Hz & $J=9.6$ Hz, 4H, H-1' & H-1''), 2.53 (s, 3H, 4-CH₃), 1.81 (s, 3H, 3'-CH₃), 1.80 (s, 3H, 4'-CH₃), 1.76 (s, 3H, 3''-CH₃), 1.73 (s, 3H, 4''-CH₃). ¹³C NMR (150 MHz, CDCl₃): δ (ppm) 162.1, 161.4, 158.4, 157.1, 154.8, 139.4, 138.7, 118.8, 111.3, 105.1, 96.9, 94.1, 65.9, 65.4, 25.9, 25.8, 24.5, 18.4, 18.3. HRMS (ESI) calcd for C₂₀H₂₄O₄ (M+H)⁺: m/z : 351.1567, found: 351.1561.

5,7-Diprenyloxy-4-propyl-2H-chromen-2-one (4h) White solid; yield 42% (272.5 mg, 0.76 mmol); mp 71–72 °C. ¹H NMR (600 MHz, CDCl₃): δ (ppm) 6.44 (d, $J=2.4$ Hz, 1H, H-6), 6.32 (d, $J=1.8$ Hz, 1H, H-8), 5.94 (s, 1H, H-3), 5.50 (t, $J=7.2$ Hz, 1H, H-2''), 5.47 (t, $J=6.0$ Hz, 1H, H-2'), 4.54 (d, $J=6.6$ Hz, 2H, H-1'), 4.51 (d, $J=6.6$ Hz, 2H, H-1''), 2.84 (t, $J=7.8$ Hz, 2H, 4-CH₂CH₂CH₃), 1.81 (s, 6H, 3'-CH₃ & 4'-CH₃), 1.76 (s, 3H, 3''-CH₃), 1.74 (s, 3H, 4''-CH₃), 1.61–1.57 (m, 2H, 4-CH₂CH₂CH₃), 0.97 (t, $J=7.8$ Hz, 3H, 4-CH₂CH₂CH₃). ¹³C NMR (150 MHz, CDCl₃): δ (ppm) 161.9, 161.5, 158.6, 158.0, 157.4, 139.4, 139.3, 118.8, 118.6, 110.8, 104.4, 96.9, 94.3, 65.7, 65.4, 38.8, 25.9, 25.8, 23.4, 18.4, 18.3, 13.9. HRMS (ESI) calcd for C₂₂H₂₈O₄ (M+H)⁺: m/z : 379.1880, found: 379.1869.

5,7-Digeranyloxy-4-propyl-2H-chromen-2-one (4i) White solid; yield 41% (262.6 mg, 0.53 mmol); mp 79 °C. ¹H NMR (600 MHz, CDCl₃): δ (ppm) 6.45 (d, $J=1.8$ Hz, 1H, H-6), 6.33 (d, $J=1.8$ Hz, 1H, H-8), 5.95 (s, 1H, H-3), 5.51 (t, $J=6.6$ Hz, 1H, H-1'), 5.46 (t, $J=6.6$ Hz, 1H, H-1''), 5.09 (pseudotriplet, 2H, H-6' & H-6''), 4.57 (d, $J=6.6$ Hz, 2H, H-2'), 4.54 (d, $J=6.6$ Hz, 2H, H-2''), 2.85 (t, $J=7.8$ Hz, 2H, 4-CH₂CH₂CH₃), 2.13–2.09 (m, 8H, H-4', H-5' & H-4'', H-5''), 1.76 (s, 3H, 3'-CH₃), 1.73 (s, 3H, 3''-CH₃), 1.68 (s, 3H, 7'-CH₃), 1.67 (s, 3H, 8'-CH₃), 1.61 (s, 6H, 7''-CH₃ & 8''-CH₃), 0.98 (t, $J=7.8$ Hz, 3H, 4-CH₂CH₂CH₃). ¹³C NMR (150 MHz, CDCl₃): δ (ppm) 162.0, 161.6, 158.6, 158.0, 157.4, 142.5, 142.4, 132.1, 132.0, 123.8, 123.7, 118.6, 118.4, 110.8, 104.4, 96.9, 94.3, 65.8, 65.4, 39.7, 39.6, 38.8, 31.0, 26.4, 26.3, 25.8, 25.7, 23.3, 17.9, 17.8, 16.9, 16.8, 14.1. HRMS (APCI) calcd for C₃₂H₄₅O₄ (M+H)⁺: m/z : 493.3312, found: 493.3302.

7-Prenyloxy-2H-chromen-2-one (4j) White solid; yield 65% (59.9 mg, 0.26 mmol); mp 77 °C (lit. m.p. 77–78 °C) [39]. ¹H NMR (300 MHz, CDCl₃): δ(ppm) 7.63 (d, *J*=9.6 Hz, 1H, H-4), 7.36 (d, *J*=8.4 Hz, 1H, H-5), 6.86–6.82 (m, 2H, H-6 & H-8), 6.25 (d, *J*=9.3 Hz, 1H, H-6), 5.48 (t, *J*=8.1 Hz, 1H, H-2'), 4.59 (d, *J*=6.9 Hz, 2H, H-1'), 1.84 (s, 3H, 3'-CH₃), 1.79 (s, 3H, 4'-CH₃).

7-Geranyloxy-2H-chromen-2-one (auraptene) (4k) White solid; yield 60% (268.5 mg, 0.90 mmol); mp 63 °C (lit. m.p. 62–63 °C) [40]. ¹H NMR (300 MHz, CDCl₃): δ(ppm) 7.63 (d, *J*=9.3 Hz, 1H, H-4), 7.36 (d, *J*=8.4 Hz, 1H, H-5), 6.86–6.82 (m, 2H, H-6 & H-8), 6.24 (d, *J*=9.3 Hz, 1H, H-3), 5.47 (t, *J*=6 Hz, 1H, H-2'), 5.09 (t, *J*=5.7 Hz, 1H, H-6'), 4.62 (d, *J*=6.6 Hz, 2H, H-1'), 2.15–2.11 (m, 4H, H-4' & H-5'), 1.79 (s, 3H, 3'-CH₃), 1.69 (s, 3H, 8'-CH₃), 1.63 (s, 3H, 7'-CH₃).

7-Farnesyloxy-coumarin (umbelliprenin) (4l) Yellowish solid; yield 80% (439.8 mg, 1.20 mmol); mp 61 °C (lit. m.p. 58–60 °C) [40]. ¹H NMR (300 MHz, CDCl₃): δ(ppm) 7.62 (d, *J*=9.3 Hz, 1H, H-4), 7.35 (d, *J*=8.1 Hz, 1H, H-5), 6.86–6.82 (m, 2H, H-6 & H-8), 6.24 (d, *J*=9.6 Hz, 1H, H-3), 5.48 (t, *J*=6.6 Hz, 1H, H-2'), 4.62 (d, *J*=6.6 Hz, 1H, H-1'), 2.18–2.01 (m, 8H, H-4' & H-5' & H-8' & H-9'), 1.79 (s, 3H, 3'-CH₃), 1.70 (s, 3H, 8'-CH₃), 1.63 (s, 6H, 11'-CH₃ & 12'-CH₃).

7-Geranylgeranyloxy-coumarin (4m) [41] Yellow solid; yield 64% (365.0 mg, 0.94 mmol); mp 72–73 °C. ¹H NMR (400 MHz, CDCl₃): δ(ppm) 7.63 (d, 1H), 7.36 (d, 1H), 6.86–6.82 (m, 2H), 6.24 (d, 1H), 5.47 (td, 1H), 5.10–5.07 (m, 3H), 4.60 (d, 2H), 2.15–2.03 (m, 8H), 1.99–1.95 (m, 4H), 1.76 (s, 3H), 1.67 (s, 3H), 1.60 (s, 3H), 1.59 (s, 3H), 1.58 (s, 3H).

5,7-Diacetyloxy-4-methyl-2H-chromen-2-one (5) 1 eq of the coumarin **3b** and 2 eq of acetic anhydride were added to the appropriate amount of pyridine, and the mixture was heated at 80 °C. After the completion of the reaction, pyridine was removed in vacuo and a solid product was obtained. The final pure coumarin was selected after recrystallization. Green solid; yield 72% (273.5 mg, 0.99 mmol); mp 143–144 °C (lit. m.p. 150–151 °C) [42]. ¹H NMR (600 MHz, CDCl₃): δ(ppm) 7.06 (s, 1H, H-8), 6.87 (s, 1H, H-6), 6.20 (s, 1H, H-3), 2.50 (s, 3H, 4-CH₃), 2.37 (s, 3H, 7-CH₃CO), 2.32 (s, 3H, 5-CH₃CO).

Cell culture

Dulbecco's modified Eagle's medium (DMEM, D6429), foetal bovine serum (FBS, F9665), trypsin (T3924), L-glutamine (G7513), antibiotics (penicillin – streptomycin,

P0781) and (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) (MTT, M2128) were purchased from Sigma-Aldrich (Gillingham, UK). Dulbecco's modified Eagle's medium with no glucose (DMEM, A1443001) was purchased from Gibco Thermo Fisher Scientific, UK. Human breast adenocarcinoma cell lines MCF-7 (HTB-22™) and MDA-MB-231 (HTB-26™) were kindly donated from Manchester Cancer Research Labs (University of Manchester, UK). Human colon fibroblasts 18-Co (CRL-1459™) were purchased from ATCC.

General cell culture

Unless otherwise specified, all cell culture experiments were performed in a humidified 5% (v/v) CO₂ air atmosphere at 37 °C in complete medium, and cell culture growth media were supplemented with 10% (v/v) foetal bovine serum and 2 mM L-glutamine. Human breast adenocarcinoma cell lines were cultured, maintained at densities lower than 1 × 10⁶ cells/cm² and discarded upon reaching passage number 60. Stromal healthy cells (human colorectal fibroblasts, 18Co) were cultured using complete DMEM medium supplemented also with 1% (v/v) penicillin–streptomycin. Cells were maintained at densities less than 1 × 10⁶ cells/cm² and discarded upon reaching passage number 12.

Nutrient-deprived conditions

Cells were culture using nutrient-deprived cell culture conditions (i.e. cell culture media with no glucose, L-glutamine, HEPES and sodium pyruvate) to mimic conditions similar to the tumour microenvironment. Note that these experiments were performed only using breast cancer cells. MCF-7 and MDA-MB-231 cells were seeded in 96-well plates (Corning Inc., NY, USA) at a density of 1 × 10⁴ and 6.7 × 10³ cells/cm², respectively. Cells were incubated with coumarin derivatives at concentrations of 0.1, 1, 10, 50, 100 and 250 μM up to 48 h. Untreated cells (negative) and cells incubated with 0.5% (v/v) DMSO in complete media (positive) were used as controls.

Coumarins cytotoxicity

The full library of coumarins was tested in both breast cancer cell lines, i.e. MCF-7 and MDA-MB-231. Fibroblast were used as “healthy” control. Stocks of compounds **3a–3c**, **4a–4m** and **5** were dissolved in pure DMSO and then diluted in complete media. (Note that DMSO concentration was kept lower than 0.5% (v/v).) Briefly, MCF-7 and MDA-MB-231 cells were seeded in 96-well plates (3799, Corning Inc., NY, USA) at a density of 1 × 10⁴ and 6.7 × 10³ cells/cm², respectively, whereas 18Co fibroblasts were seeded at a density of 1 × 10⁴ cells/cm². Cells were incubated with coumarin

derivatives at concentrations 0.1, 1, 10, 50, 100 and 250 μM for 48 h. Untreated cells (negative) and cells incubated with 0.5% (v/v) DMSO in complete media (positive) were used as controls.

Cell metabolism assay

For each treatment, cell viability was measured via MTT assay after 48-h incubation as following described. Cell culture medium was replaced with 150 μL of fresh medium and 30 μL of MTT solution, and cells were incubated for 4 h (37 $^{\circ}\text{C}$, 5% CO_2). After the formazan crystal formation, cell culture medium was removed from each well and replaced with 200 μL of DMSO. The absorbance was measured at 540-nm wavelength using a plate reader (Synergy 2 Biotek plate reader, Gen5 software).

Toxicity (IC_{50}) and identification of potent coumarins

IC_{50} values were calculated (nonlinear regression, normalized response–variable slope) with GraphPad Prism (version 7.04). Values were ranked and classified as: high toxicity, moderate toxicity, poor toxicity and no toxicity. Threshold were set as 60, 80 and > 100 μM , respectively. The selection was used in order to test only the most toxic coumarins under cell culture condition more relevant to the tumour microenvironment, i.e. deprived cell culture media (Sect. 3.3).


Results and discussion

Design and synthesis of coumarin analogues

Our previous results concerning the cytotoxic activity evaluation of natural oxyprenylated coumarins [14] in combination with the latest literature data led us to design, a new series of diprenyloxy as well as dialkyloxy coumarins. The new series were designed in order to evaluate the influence of the disubstitution as well as the length of the lipophilic chain in the cytotoxicity against breast cancer cell lines.

In order to efficiently synthesize the desired coumarin analogues and the naturally occurring oxyprenylated coumarins, the appropriate hydroxy-4-substituted coumarins **3a** and **3b** were firstly synthesized via Pechmann reaction using iron (III) chloride (FeCl_3) as the catalyst [36]. Compounds **3a** and **3b** as well as the commercially available 7-hydroxy-coumarin (umbelliferone, **3c**) were subsequently alkylated with the appropriate commercially available alkyl bromide using potassium carbonate (K_2CO_3) in acetone (Scheme 1). For the preparation of **4m**, the required geranylgeranyl iodide was prepared according to the method of Alvarez-Manzaneda et al. [37].

All the coumarin derivatives were purified using flash column chromatography and were structurally identified using ^1H and ^{13}C NMR spectroscopy and HRMS spectrometry.



Scheme 1 Synthesis of **3a**, **3b**, **4a–4m** and **5**; Reagents and conditions: **a** $\text{FeCl}_3/70$ $^{\circ}\text{C}$, 80–90%, **b** R_2Br or geranylgeranyl iodide, K_2CO_3 , acetone, 50–60%, **c** Ac_2O , pyridine, 80 $^{\circ}\text{C}$

Cytotoxicity towards breast cancer cells

The systematic variations on the alkyl chain as well as the position of substitution at the coumarin scaffold were examined as potential factors which could affect anticancer activity. The first set of experiments identified the most potent candidates among the coumarin derivatives herein synthesized. Cytotoxicity was firstly evaluated on two breast cancer cell lines: MCF-7 and MDA-MB-231. MCF-7 cells were selected as they retain several characteristics of differentiated mammary epithelium proliferation, as well as expressing oestrogen receptor, whereas MDA-MB-231 was selected as expressing a more aggressive and metastatic cells that do not express high levels of the oestrogen, progesterone or HER2 receptors (i.e. triple negative).

Coumarins were classified as possessing high ($IC_{50} < 80 \mu\text{M}$), moderate ($80 \mu\text{M} < IC_{50} < 100 \mu\text{M}$) and poor toxicity ($100 \mu\text{M} < IC_{50} < 250 \mu\text{M}$); compounds with IC_{50} values $> 250 \mu\text{M}$ were classified as non-toxic. Coumarin derivatives **4c**, **4d**, **4g**, **4k** and **4l** were identified as the top-five most potent compounds tested in this study (Table 1) and were selected for further investigation.

One of the main drawbacks of cytotoxic compounds is the poor selectivity towards cancer cells, with undesired effects on healthy cells, e.g. fibroblasts and epithelia. In an effort to understand whether coumarins have any effect on ‘healthy’ cells, human fibroblasts (18-Co) were treated with the most potent coumarin derivatives. As control, fibroblasts were also treated with a non-toxic coumarin (i.e. umbelliferone, **3c**). Interestingly, cytotoxicity data evidenced no effect of the selected coumarins on fibroblasts (shown in Table 2) with the exception of coumarin **4k** that showed some toxicity towards ‘healthy’ cells. The tested compounds were not as toxic as typical chemotherapeutics with IC_{50} s at the scale of few hundred nM such as doxorubicin [43–46] or gemcitabine [47–49], but they did appear to have no significant effects on ‘healthy’ fibroblasts as compared to the aforementioned chemo-agents. This is a very positive result in view of development of (nano)formulations and further translation of such compounds.

Auraptene (**4j**) was found to be the most potent compound among the library of coumarins tested in this study, confirming what has been already observed in other *in vitro* studies and in various *in vivo* animal models [50]. Its effect on cancer cells is still not clear and could be associated with induction of carcinogen-detoxifying enzymes, inhibition of free radical generation or metalloproteinase production [51]. The length of the prenyl chain seems to affect the activity of the compounds: auraptene (**4k**, 10 carbons) is more potent

compared to its prenyloxy analogue (**4j**, 5 carbons) against the tested breast cancer cells. However, umbelliprenin (**4l**, 15 carbons) and coumarin **4m** (20 carbons) exhibited lower antitumour potency, with umbelliprenin being more toxic than coumarin **4m** in both cancer cell lines (Fig. 1).

The number of substituents on the aromatic ring could also play a role in the activity: 5,7-diprenyloxy-4-methylcoumarin (**4g**) is approximately 2.5 times more cytotoxic compared to its monosubstituted analogue (compound **4a**) against MCF-7 cells. Increasing the chain length of the substituents, as in 7-geranylgeranyloxy-coumarin (**4n**), results in complete loss of activity against both cell lines (Fig. 1, Table 1).

In an effort to better investigate the role of unsaturation on cytotoxicity, the coumarin analogues **4d** and **4f**, which possess a saturated alkyloxy substituent, were synthesized. Only coumarin **4d** exhibited a moderate potency against MCF7 cells, whereas only a slight toxic effect was observed on MDA-MB-231 (Table 1); moreover, no toxicity was observed on fibroblasts (Table 2). These results suggest that this specific modification can participate in different biochemical pathways compared to unsaturated substituents; however, further research is necessary to confirm this and identify specific pathways.

Finally, we investigated the activity of derivatives as function of lipophilicity through the introduction of a methyl group at position 4 of the coumarin scaffold. The methylated coumarin derivatives **4b**, **4a** and **4c** exhibited a rather moderate cytotoxic effect compared to their corresponded non-methylated analogues **4k**, **4j** and **4l**. This observation suggests that substitution at position 4 might not directly link to increased toxicity. Furthermore, comparing the compounds with different substitutions at position 4 (e.g. coumarins **4d** vs **4f**, coumarins **4g** vs **4h**) increased toxicity was observed in both cancer cell lines for compounds with methyl substitution (Fig. 1).

Cytotoxicity in tumour relevant *in vitro* models

In accordance with the work of Jun et al. [52] and Devji et al. [41], we were motivated to assess the activity of some selected derivatives under nutrient-deprived conditions (NDCs). Cancer cells are programmed in a non-ordinary way to exhibit high glycolytic activity even under sufficient aerobic conditions [53]. In hypoxic tumour conditions, when oxygen depletion and low vascularization take place, cancer cells often find the way to proliferate rapidly by foregoing oxidative phosphorylation and instead ferment large

Table 1 IC₅₀ values (μM) of coumarin derivatives (18 compounds) on MCF-7 and MDA-MB-231 breast cancer cells obtained after 48-h incubation

Coumarin derivative	Chemical structure	MCF-7 IC ₅₀ values (μM)	MDA-MB-231 IC ₅₀ values (μM)
3a		225 ± 19	206 ± 9
3b		> 250	> 250
3c		> 250	> 250
4a		179 ± 35	169 ± 23
4b		129 ± 13	137 ± 9
4c		105 ± 3	110 ± 29
4d		82 ± 19	113 ± 7

Table 1 (continued)











Coumarin derivative	Chemical structure	MCF-7 IC ₅₀ values (μM)	MDA-MB-231 IC ₅₀ values (μM)
4e		> 250	> 250
4f		124 ± 39	143 ± 34
4g		69 ± 14	144 ± 16
4h		234 ± 76	159 ± 61
4i		> 250	133 ± 34
4j		143 ± 11	175 ± 33
4k		70 ± 5	60 ± 4

Table 1 (continued)

Coumarin derivative	Chemical structure	MCF-7 IC ₅₀ values (μM)	MDA-MB-231 IC ₅₀ values (μM)
4l		98 ± 9	88 ± 3
4m		> 250	187 ± 66
5		> 250	> 250







Data are expressed as mean ± SD. of two independent experiments ($n=3$ samples for each experiment). As control, MCF-7 and MDA-231 were incubated with [0.001–50] μg/mL of doxorubicin for 48 h, reporting IC₅₀ values of (0.97 ± 0.60) μM and (0.48 ± 0.14) μM, respectively

amounts of glucose into lactate under aerobic glycolysis, known as the Warburg effect [54, 55]. Moreover, hypoxia tends to boost this phenomenon by up-regulating the HIF-1α factor that “switches on” glycolytic and glucose transporter gene expression [56]. Breast carcinoma cell lines behave in a glucose-dependent manner and derive the majority of energy needed from high-throughput glycolysis [56, 57]. Hyperglycaemic systemic conditions, i.e. diabetes, have been proved to further promote the migratory invasiveness of breast malignancies in patients [55].

On that basis, we exposed the breast cancer cells to nutrient-deprived conditions where culture media were supplemented only with 2.5% v/v FBS, but not additional glucose, L-glutamine, sodium pyruvate. The coumarins tested were auraptene (**4k**), umbelliprenin (**4l**) and analogues **4d**, **4c**

and **4g**. As shown in Table 3, the tested compounds showed selective preferential cytotoxicity under nutrient-deprived conditions with umbelliprenin (**4l**) to be the most potent candidate as its pharmacological activity was remarkably enhanced by 15 times (IC₅₀ = 9.0 and 7.0 for MCF7 and MDA-MB231 cells, respectively). In similar studies, Zhang et al. and Jun et al. reported the high preferential cytotoxicity of umbelliprenin (**4l**) and its C6 analogue under NDC against pancreatic cancer cells [52, 58]. It should be therefore noted that these derivatives could represent a potential new tool for treating aggressively metastatic hypoxic tumours. The exact mechanism of action, though, should be further investigated.



Table 2 IC₅₀ values (μM) of the most potent coumarin derivatives (five compounds: **4c**, **4d**, **4g**, **4k** and **4l**) on 18Co colon fibroblasts (healthy control)

Coumarin derivative	Chemical structure	18-Co IC ₅₀ values (μM)
3c		> 250
4c		> 250
4d		206 ± 17
4g		216 ± 23
4k		176 ± 62
4l		> 250


Data are expressed as mean ± SD and are obtained from $n = 2$ independent experiments

Fig. 1 Effect of substituents on coumarins on MCF-7 cells: cell viability was measured after 48-h incubation with different concentrations of coumarins and IC_{50} values determined using nonlinear regression (GraphPad Prism, v7). Coumarins were compared on the basis of the following structural features: **a** length of lipophilic chain, coumarin **4k** versus **4j** (10 vs 5 carbons) and coumarin **4l** versus **4m** (15 vs 20 carbons); **b** position of the substituent on the coumarin scaffold, coumarin **4g** vs **4a**; **c** effect of the saturation degree (substitution at position C7), coumarin **4d** vs **4f**; **d** the presence of methyl group (substitution at position C4), coumarin **4b** versus **4k** and coumarin **4a** versus **4j** (data not shown), coumarin **4c** versus **4l** (data not shown); **e** the presence of propyl group (substitution at position C4), coumarins **4g** versus **4h** and coumarins **4d** versus **4f** (data not shown)


A | Length of lipophilic chain




B | Group substitution



C | Saturation degree in 7' carbon



D | Substitution in 3' carbon



E | Alkylation in 3' carbon









Table 3 IC₅₀ values (μM) of the tested coumarin derivatives **4c**, **4d**, **4g**, **4k** and **4l**. Values were obtained for both MCF-7 and MDA-MB-231 breast cancer cells cultured under nutrient-deprived conditions and incubated with coumarin derivatives up to 48 h

Coumarin derivative	Chemical structure	MCF-7 IC ₅₀ values (μM)	MDA-MB-231 IC ₅₀ values (μM)
4c		19 ± 3	51 ± 9
4d		29 ± 3	63 ± 5
4g		35 ± 9	11 ± 2
4k		14 ± 5	12 ± 8
4l		9 ± 1	7 ± 2

Data are expressed as average ± SD of $n = 3$ independent experiments

Conclusions

A series of novel alkoxy-coumarin derivatives were synthesized and tested for their cytotoxicity against the MCF7 and MDA-MB-231 breast cancer cells. The results of this study indicate that alkylation modification induces noticeable differentiation in pharmacological activity of coumarins. Auraptene (**4k**) was found to possess the most potent cytotoxic activity among the tested derivatives followed by compounds **4c**, **4d**, **4g** and **4l**. The tested compounds seemed not to affect the cell viability of the healthy 18Co fibroblasts but for the highest dose only. The amplification of the cytotoxic effect of the above pharmacophores under nutrient-deprived conditions, with umbelliprenin (**4l**) being the lead compound, indicates that these compounds could lead to potential new therapeutics for highly metastatic hypoxic tumours once their mechanisms are fully understood.

Acknowledgements Eleni Kavetsou gratefully acknowledges financial support from State Scholarships Foundation (IKY): The realization of the doctoral dissertation was co-funded by the “Grant Scheme for Post-graduate Studies in Secondary Education” of the Operational Program “Human Resource Development, Education and Lifelong Learning” of the NSRF 2014–2020 with the co-financing of the European Social Fund. Leonidas Gkionis is indebted to EPSRC for a Ph.D. studentship as part of the Graphene NOWNANO Doctoral Training Centre.

Funding Not applicable.

Availability of data and material (data transparency) The raw data supporting the conclusions of this manuscript will be made available by the authors, without undue reservation, to any qualified researcher.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

Ethics approval (include appropriate approvals or waivers) Not applicable.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Marcom PK (2017) Breast cancer. In: Ginsburg G, Willard H, David S (eds) *Genomic and precision medicine: primary care*, 3rd edn. Elsevier Ltd, pp 181–194
- Tao ZQ, Shi A, Lu C et al (2015) Breast cancer: epidemiology and etiology. *Cell Biochem Biophys* 72:333–338. <https://doi.org/10.1007/s12013-014-0459-6>
- Hanahan D, Weinberg RA (2011) Hallmarks of cancer: the next generation. *Cell* 144:646–674
- Guerra AR, Duarte MF, Duarte IF (2018) Targeting tumor metabolism with plant-derived natural products: emerging trends in cancer therapy. *J Agric Food Chem* 66:10663–10685. <https://doi.org/10.1021/acs.jafc.8b04104>
- Lee KW, Bode AM, Dong Z (2011) Molecular targets of phytochemicals for cancer prevention. *Nat Rev Cancer* 11:211–218
- Zhang L, Xu Z (2019) Coumarin-containing hybrids and their anticancer activities. *Eur J Med Chem* 181:111587
- Fayed EA, Sabour R, Harras MF, Mehany ABM (2019) Design, synthesis, biological evaluation and molecular modeling of new coumarin derivatives as potent anticancer agents. *Med Chem Res* 28:1284–1297. <https://doi.org/10.1007/s00044-019-02373-x>
- Kaur M, Kohli S, Sandhu S et al (2015) Coumarin: a promising scaffold for anticancer agents. *Anticancer Agents Med Chem* 15:1032–1048. <https://doi.org/10.2174/1871520615666150101125503>
- O'Kennedy R, Thornes DR (1997) *Coumarins: biology, applications and mode of action*. Wiley, Chichester
- Detsi A, Kontogiorgis C, Hadjipavlou-Litina D (2017) Coumarin derivatives: an updated patent review (2015–2016). *Expert Opin Ther Pat* 27:1201–1226
- Stefanachi A, Leonetti F, Pisani L et al (2018) Coumarin: a natural, privileged and versatile scaffold for bioactive compounds. *Molecules* 23:1–34
- Barot KP, Jain SV, Kremer L et al (2015) Recent advances and therapeutic journey of coumarins: current status and perspectives. *Med Chem Res* 24:2771–2798
- Venugopala KN, Rashmi V, Odhav B (2013) Review on natural coumarin lead compounds for their pharmacological activity. *Biomed Res Int* 2013:1–14
- Kavetsou E, Gkionis L, Galani G et al (2017) Synthesis of prenyloxy coumarin analogues and evaluation of their antioxidant, lipoxygenase (LOX) inhibitory and cytotoxic activity. *Med Chem Res* 26:856–866. <https://doi.org/10.1007/s00044-017-1800-6>
- Roussaki M, Zelianaios K, Kavetsou E et al (2014) Structural modifications of coumarin derivatives: determination of antioxidant and lipoxygenase (LOX) inhibitory activity. *Bioorganic Med Chem* 22:6586–6594. <https://doi.org/10.1016/j.bmc.2014.10.008>
- Al-Majedy YK, Al-Duhaidhawi DL, Al-Azawi KF et al (2016) Coumarins as potential antioxidant agents complemented with suggested mechanisms and approved by molecular modeling studies. *Molecules* 21:2–11. <https://doi.org/10.3390/molecules21020135>
- Tamene D, Endale M (2019) Antibacterial activity of coumarins and carbazole alkaloid from roots of *clausena anisata*. *Adv Pharmacol Sci* 2019:1–8. <https://doi.org/10.1155/2019/5419854>
- Yang L, Wu L, Yao X et al (2018) Hydroxycoumarins: new, effective plant-derived compounds reduce *Ralstonia pseudosolanacearum* populations and control tobacco bacterial wilt. *Microbiol Res* 215:15–21. <https://doi.org/10.1016/j.micres.2018.05.011>
- Tian D, Wang F, Duan M et al (2019) Coumarin analogues from the *Citrus grandis* (L.) osbeck and their hepatoprotective activity. *J Agric Food Chem* 67:1937–1947. <https://doi.org/10.1021/acs.jafc.8b06489>
- Berrino E, Milazzo L, Micheli L et al (2019) Synthesis and evaluation of carbonic anhydrase inhibitors with carbon monoxide releasing properties for the management of rheumatoid arthritis. *J Med Chem* 62:7233–7249. <https://doi.org/10.1021/acs.jmedchem.9b00845>
- Zhao YL, Yang XW, Wu BF et al (2019) Anti-inflammatory effect of Pomelo peel and its bioactive coumarins. *J Agric Food Chem* 67:8810–8818. <https://doi.org/10.1021/acs.jafc.9b02511>
- Peperidou A, Bua S, Bozdog M et al (2018) Novel 6- and 7-substituted coumarins with inhibitory action against lipoxygenase and tumor-associated carbonic anhydrase IX. *Molecules* 23:13–17. <https://doi.org/10.3390/molecules23010153>
- Zhao H, Donnelly AC, Kusuma BR et al (2011) Engineering an antibiotic to fight cancer: optimization of the novobiocin scaffold to produce anti-proliferative agents. *J Med Chem* 54:3839–3853. <https://doi.org/10.1021/jm200148p>
- Herrera-R A, Castrillón W, Otero E et al (2018) Synthesis and antiproliferative activity of 3- and 7-styrylcoumarins. *Med Chem Res* 27:1893–1905. <https://doi.org/10.1007/s00044-018-2202-0>
- Farley CM, Dibwe DF, Ueda JY et al (2016) Evaluation of synthetic coumarins for antiausterity cytotoxicity against pancreatic cancers. *Bioorganic Med Chem Lett* 26:1471–1474. <https://doi.org/10.1016/j.bmcl.2016.01.054>
- Ricci F, Carrassa L, Christodoulou MS et al (2018) A high-throughput screening of a chemical compound library in ovarian cancer stem cells. *Comb Chem High Throughput Screen* 21:50–56. <https://doi.org/10.2174/1386207321666180124093406>
- Cao D, Liu Y, Yan W et al (2016) Design, synthesis, and evaluation of in vitro and in vivo anticancer activity of 4-substituted coumarins: a novel class of potent tubulin polymerization inhibitors. *J Med Chem* 59:5721–5739. <https://doi.org/10.1021/acs.jmedchem.6b00158>
- Fiorito S, Epifano F, Taddeo VA, Genovese S (2018) Recent acquisitions on oxyprenylated secondary metabolites as anti-inflammatory agents. *Eur J Med Chem* 153:116–122. <https://doi.org/10.1016/j.ejmech.2017.08.038>
- Alhassan AM, Abdullahi MI, Uba A, Umar A (2014) Prenylation of aromatic secondary metabolites: a new frontier for development of novel drugs. *Trop J Pharm Res* 13:307–314
- Hasan M, Genovese S, Fiorito S et al (2017) Oxyprenylated phenylpropanoids bind to MT1 melatonin receptors and inhibit breast cancer cell proliferation and migration. *J Nat Prod* 80:3324–3329. <https://doi.org/10.1021/acs.jnatprod.7b00853>
- Okuyama S, Minami S, Shimada N et al (2013) Anti-inflammatory and neuroprotective effects of auraptene, a citrus coumarin, following cerebral global ischemia in mice. *Eur J Pharmacol* 699:118–123. <https://doi.org/10.1016/j.ejphar.2012.11.043>
- Okuyama S, Semba T, Toyoda N et al (2016) Auraptene and other prenyloxyphenylpropanoids suppress microglial activation and dopaminergic neuronal cell death in a

- lipopolysaccharide-induced model of Parkinson's disease. *Int J Mol Sci* 17:1–11. <https://doi.org/10.3390/ijms17101716>
33. Iranshahi M, Askari M, Sahebkar A, Hadjipavlou-Litina D (2009) Evaluation of antioxidant, anti-inflammatory and lipoxigenase inhibitory activities of the prenylated coumarin umbelliprenin. *Daru* 17:99–103
 34. Yan H, Ma Z, Peng S, Deng X (2013) Anti-inflammatory effect of auraptene extracted from trifoliolate orange (*Poncirus trifoliolate*) on LPS-stimulated RAW 264.7 cells. *Inflammation* 36:1525–1532. <https://doi.org/10.1007/s10753-013-9695-y>
 35. La VD, Zhao L, Epifano F et al (2013) Anti-inflammatory and wound healing potential of citrus auraptene. *J Med Food* 16:961–964. <https://doi.org/10.1089/jmf.2013.0029>
 36. Prousis KC, Avlonitis N, Heropoulos GA, Calogeropoulou T (2014) FeCl₃-catalysed ultrasonic-assisted, solvent-free synthesis of 4-substituted coumarins. A useful complement to the Pechmann reaction. *Ultrason Sonochem* 21:937–942. <https://doi.org/10.1016/j.ultsonch.2013.10.018>
 37. Alvarez-Manzaneda EJ, Chahboun R, Cabrera Torres E et al (2005) Reaction of allylic and benzylic alcohols and esters with PPh₃/I₂: one-pot synthesis of β , γ -unsaturated compounds. *Tetrahedron Lett* 46:3755–3759. <https://doi.org/10.1016/j.tetlet.2005.03.132>
 38. Takeuch N, Kasama T, Aida Y et al (1991) Pharmacological activities of the prenylcoumarins, developed from folk usage as a medicine of *Peucedanum japonicum* thunb. *Chem Pharm Bull* 39:1415–1421. <https://doi.org/10.1248/cpb.39.1415>
 39. Row EC, Brown SA, Stachulski AV, Lennard MS (2006) Design, synthesis and evaluation of furanocoumarin monomers as inhibitors of CYP3A4. *Org Biomol Chem* 4:1604–1610. <https://doi.org/10.1039/b601096b>
 40. Askari M, Sahebkar A, Iranshahi M (2009) Synthesis and purification of 7-prenyloxycoumarins and herniarin as bioactive natural coumarins. *Iran J Basic Med Sci* 12:63–69. <https://doi.org/10.22038/ijbms.2009.5145>
 41. Devji T, Reddy C, Woo C et al (2011) Pancreatic anticancer activity of a novel geranylgeranylated coumarin derivative. *Bioorganic Med Chem Lett* 21:5770–5773. <https://doi.org/10.1016/j.bmcl.2011.08.005>
 42. Canter FW, Curd FH, Robertson A (1931) Hydroxy-carbonyl compounds. Part III. The preparation of coumarins and 1:4-benzopyrones from phloroglucinol and resorcinol. *J Chem Soc*. <https://doi.org/10.1039/jr9310001255>
 43. Wen Sh, Su Sc, Liou Bh et al (2018) Sulbactam-enhanced cytotoxicity of doxorubicin in breast cancer cells. *Cancer Cell Int* 18:1–18. <https://doi.org/10.1186/s12935-018-0625-9>
 44. Arif IS, Hooper CL, Greco F et al (2013) Increasing doxorubicin activity against breast cancer cells using PPAR γ -ligands and by exploiting circadian rhythms. *Br J Pharmacol* 169:1178–1188. <https://doi.org/10.1111/bph.12202>
 45. Fang XJ, Jiang H, Zhu YQ et al (2014) Doxorubicin induces drug resistance and expression of the novel CD44st via NF- κ B in human breast cancer MCF-7 cells. *Oncol Rep* 31:2735–2742. <https://doi.org/10.3892/or.2014.3131>
 46. You Y, Xu Z, Chen Y (2018) Doxorubicin conjugated with a trastuzumab epitope and an MMP-2 sensitive peptide linker for the treatment of HER2-positive breast cancer. *Drug Deliv* 25:448–460. <https://doi.org/10.1080/10717544.2018.1435746>
 47. Tao XM, Wang JC, Wang JB et al (2012) Enhanced anticancer activity of gemcitabine coupling with conjugated linoleic acid against human breast cancer in vitro and in vivo. *Eur J Pharm Biopharm* 82:401–409. <https://doi.org/10.1016/j.ejpb.2012.06.007>
 48. Li A, Fridley B, Kalari K et al (2008) Gemcitabine and cytosine arabinoside cytotoxicity: association with lymphoblastoid cell expression. *Cancer Res* 68:7050–7058. <https://doi.org/10.1158/0008-5472.CAN-08-0405>
 49. Hernández-Vargas H, Rodríguez-Pinilla SM, Julián-Tendero M et al (2007) Gene expression profiling of breast cancer cells in response to gemcitabine: NF- κ B pathway activation as a potential mechanism of resistance. *Breast Cancer Res Treat* 102:157–172. <https://doi.org/10.1007/s10549-006-9322-9>
 50. Krishnan P, Yan KJ, Windler D et al (2009) Citrus auraptene suppresses cyclin D1 and significantly delays N-methyl nitrosourea induced mammary carcinogenesis in female Sprague-Dawley rats. *BMC Cancer* 9:1–12. <https://doi.org/10.1186/1471-2407-9-259>
 51. Krishnan P, Kleiner-Hancock H (2012) Effects of Auraptene on IGF-1 stimulated cell cycle progression in the human breast cancer cell line, MCF-7. *Int J Breast Cancer* 2012:1–8. <https://doi.org/10.1155/2012/502092>
 52. Jun M, Bacay AF, Moyer J et al (2014) Synthesis and biological evaluation of isoprenylated coumarins as potential anti-pancreatic cancer agents. *Bioorganic Med Chem Lett* 24:4654–4658. <https://doi.org/10.1016/j.bmcl.2014.08.038>
 53. Izuishi K, Kato K, Ogura T et al (2000) Remarkable tolerance of tumor cells to nutrient deprivation: possible new biochemical target for cancer therapy. *Cancer Res* 60:6201–6207
 54. Zhuang Y, Chan DK, Haugrud AB, Miskimins WK (2014) Mechanisms by which low glucose enhances the cytotoxicity of metformin to cancer cells both in vitro and in vivo. *PLoS ONE*. <https://doi.org/10.1371/journal.pone.0108444>
 55. Takatani-Nakase T, Matsui C, Maeda S et al (2014) High glucose level promotes migration behavior of breast cancer cells through zinc and its transporters. *PLoS ONE* 9:e90136. <https://doi.org/10.1371/journal.pone.0090136>
 56. O'Mahony F, Razandi M, Pedram A et al (2012) Estrogen modulates metabolic pathway adaptation to available glucose in breast cancer cells. *Mol Endocrinol* 26:2058–2070. <https://doi.org/10.1210/me.2012-1191>
 57. Simões RV, Serganova IS, Kruchevsky N et al (2015) Metabolic plasticity of metastatic breast cancer cells: adaptation to changes in the microenvironment. *Neoplasia* 17:671–684. <https://doi.org/10.1016/j.neo.2015.08.005>
 58. Zhang H, Zhou R, Jun M et al (2016) Identification of the factors responsible for the selective in vitro cytotoxic activity of isoprenylated coumarin derivatives under nutrient-deprived conditions. *J Cancer* 7:160–166. <https://doi.org/10.7150/jca.13243>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.