

MINI-REVIEW

Open Access



Methanogens: biochemical background and biotechnological applications

Franziska Enzmann¹, Florian Mayer¹ , Michael Rother² and Dirk Holtmann^{1*} 

Abstract

Since fossil sources for fuel and platform chemicals will become limited in the near future, it is important to develop new concepts for energy supply and production of basic reagents for chemical industry. One alternative to crude oil and fossil natural gas could be the biological conversion of CO₂ or small organic molecules to methane via methanogenic archaea. This process has been known from biogas plants, but recently, new insights into the methanogenic metabolism, technical optimizations and new technology combinations were gained, which would allow moving beyond the mere conversion of biomass. In biogas plants, steps have been undertaken to increase yield and purity of the biogas, such as addition of hydrogen or metal granulate. Furthermore, the integration of electrodes led to the development of microbial electrosynthesis (MES). The idea behind this technique is to use CO₂ and electrical power to generate methane via the microbial metabolism. This review summarizes the biochemical and metabolic background of methanogenesis as well as the latest technical applications of methanogens. As a result, it shall give a sufficient overview over the topic to both, biologists and engineers handling biological or bioelectrochemical methanogenesis.

Keywords: Methanogens, Genetic tools, Biogas, Microbial electrosynthesis, Electroactivity

Introduction

Methanogens are biocatalysts, which have the potential to contribute to a solution for future energy problems by producing methane as storable energy carrier. The very diverse archaeal group of methanogens is characterized by the ability of methane production (Balch et al. 1979). The flammable gas methane is considered to be a suitable future replacement for fossil oil, which is about to be depleted during the next decades (Ren et al. 2008). Methane can be used as a storable energy carrier, as fuel for vehicles, for the production of electricity, or as base chemical for synthesis and many countries do already have well developed natural gas grids (Ren et al. 2008). In terms of the necessary transition from chemical to biological processes, methanotrophic bacteria can use methane as a carbon and energy source to produce biomass, enzymes, PHB or methanol (Strong et al. 2015; Ge et al. 2014). The biological methanation is the main industrial

process involving methanogens. These archaea use CO₂ and H₂ and/or small organic molecules, such as acetate, formate, and methylamine and convert it to methane. Although the electrochemical production of methane is still more energy efficient than the biological production [below 0.3 kWh/cubic meter of methane (0.16 MPa, Bär et al. 2015)], the biological conversion may be advantageous due to its higher tolerance against impurities (H₂S and NH₃) within the educt streams, especially if CO₂ rich waste gas streams shall be used (Bär et al. 2015). Apart from that, research is going on to increase the energy efficiency of the biological process, so that it might be the preferred way of methane production in the future (Bär et al. 2015). Biological methanation occurs naturally in swamps, digestive systems of animals, oil fields and other environments (Garcia et al. 2000) and is already commonly used in sewage water plants and biogas plants. New applications for methanogens such as electromethanogenesis are on the rise, and yet, there is still a lot of basic research, such as strain characterization and development of basic genetic tools, going on about the very diverse, unique group of methanogens (Blasco-Gómez et al. 2017). This review will summarize important

*Correspondence: Holtmann@dechema.de

¹ DECHEMA Research Institute, Industrial Biotechnology, Theodor-Heuss-Allee 25, 60486 Frankfurt am Main, Germany
Full list of author information is available at the end of the article

facts about the biological properties and possibilities of genetic modification of methanogenic organisms as well as the latest technical applications. It shall therefore give an overview over the applicability of methanogens and serve as a start-up point for new technical developments.

Biochemical and microbial background

Methanogens are the only group of microorganisms on earth producing significant amounts of methane. They are unique in terms of metabolism and energy conservation, are widespread in different habitats and show a high diversity in morphology and physiological parameters.

Phylogeny and habitats of methanogens

For decades known methanogenic archaea belonged exclusively to the phylum *Euryarchaeota*. There, methanogens were classified first into five orders, namely *Methanococcales*, *Methanobacteriales*, *Methanosarcinales*, *Methanomicrobiales* and *Methanopyrales* (Balch et al. 1979; Stadtman and Barker 1951; Kurr et al. 1991). Between the years 2008 and 2012 another two orders of methanogens, namely *Methanocellales* (Sakai et al. 2008) and *Methanomassiliicoccales* (Dridi et al. 2012; Iino et al. 2013), were added to the phylum *Euryarchaeota*. Hydrogenotrophic methanogenesis from H₂ and CO₂ is found in almost all methanogenic orders with the exception of the *Methanomassiliicoccales*. Due to its broad distribution it is postulated that this type of methanogenesis is the ancestral form of methane production (Bapteste et al. 2005). Methane formation from acetate, called acetoclastic methanogenesis, can be found only in the order *Methanosarcinales*. In contrast to that, methylotrophic methanogenesis, which is the methane formation from different methylated compounds such as methanol, methylamines or methylated thiols, is found in the orders *Methanomassiliicoccales*, *Methanobacteriales* and *Methanosarcinales*. Extensive recent metagenomic analyses suggested that methanogens may no longer be restricted to the *Euryarchaeota*. Two new phyla, namely the *Bathyarchaeota* (Evans et al. 2015) and the *Verstraetearchaeota* (Vanwonterghem et al. 2016) were postulated. Genome sequences from both phyla indicate a methylotrophic methane metabolism in these -as of yet uncultivated- potential methanogens.

Methanogens are a relatively diverse group of archaea and can be found in various anoxic habitats (Garcia et al. 2000). For example, they can be cultured from extreme environments such as hydrothermal vents or saline lakes. *Methanocaldococcus jannaschii* was isolated from a white smoker chimney of the East Pacific Rise at a depth of 2600 m (Jones et al. 1983) and *Methanopyrus kandleri* from a black smoker chimney

from the Gulf of California in a depth of 2000 m (Kurr et al. 1991). From a saline lake in Egypt the halophilic methanogen *Methanohalophilus zhilinae* was cultured (Mathrani et al. 1988). But methanogens also colonize non-extreme environments. They can be isolated from anoxic soil sediments such as rice fields, peat bogs, marshland or wet lands. For example, *Methanoregula boonei* was obtained from an acidic peat bog (Bräuer et al. 2006, 2011) and several strains of *Methanobacterium* as well as *Methanosarcina mazei* TMA and *Methanobrevibacter arboriphilus* were isolated from rice fields (Asakawa et al. 1995).

Some methanogens can also associate with plants, animals and could be found in the human body. *Methanobacterium arboriphilicum* could be isolated from a tree wetwood tissue and uses the H₂ resulting from pectin and cellulose degradation by *Clostridium butyricum* for methanogenesis (Schink et al. 1981; Zeikus and Henning 1975). From the feces of cattle, horse, sheep and goose the methanogens *Methanobrevibacter thaueri*, *Methanobrevibacter gottschalkii*, *Methanobrevibacter wolinii* and *Methanobrevibacter woesei* have been isolated, respectively (Miller and Lin 2002). In addition, different *Methanobrevibacter* species could be found in the intestinal tract of insects such as termites (Leadbetter and Breznak 1996). Beside the intestinal tract of herbivorous mammals also the rumen contains methanogens. One of the major species here is *Methanobrevibacter ruminantium* (Hook et al. 2010). Methanogenic archaea are also present in the human body. *Methanobrevibacter smithii* and *Methanosphaera stadtmanae* as well as *Methanomassiliicoccus luminyensis* could be detected in human feces (Dridi et al. 2009, 2012; Miller et al. 1982). Further *Methanosarcina* sp., *Methanosphaera* sp. and *Methanobrevibacter oralis* were discovered in human dental plaque (Belay et al. 1988; Ferrari et al. 1994; Robichaux et al. 2003).

Methanogens can be also found in non-natural habitats such as landfills, digesters or biogas plants. There, the microbial community varies with the substrate. In biogas plants, due to hydrolysis of complex polymers to sugars and amino acids, followed by fermentation and acetogenesis, acetate, H₂ and CO₂ is produced as substrates for methanogenesis. Therefore, hydrogenotrophic and acetoclastic methanogens are prevalent in mesophilic biogas plants, often dominated by species of *Methanosarcina* (*Methanohrix* at low acetate concentrations) or *Methanoculleus* (Kern et al. 2016b; Karakashev et al. 2005; Lucas et al. 2015; Sundberg et al. 2013). However, under certain conditions syntrophic acetate oxidation may be the dominant path towards methane (Schnürer and Nordberg 2008; Westerholm et al. 2016).

Diversity of methanogens in morphology and physiological parameters

Methanogens show not only a wide diversity in regard to their habitats but are also highly diverse in terms of morphology, temperature optimum, pH and osmolarity. The shapes of methanogens (only some typical methanogens are mentioned here) can be coccoid as for *Methanococcus*, *Methanosphaera* or *Methanococcoides*, long or short rods as for *Methanobacterium* or *Methanobrevibacter*, or rods in chains as for *Methanopyrus* (Kurr et al. 1991). *Methanoplanus* (Ollivier 1997) has a plate-shaped morphology and *Methanospirillum* (Zeikus and Bowen 1975), as the name says, a spirally shape. *Methanosarcina* (Balch et al. 1979; Bryant and Boone 1987; Kern et al. 2016a; Mah 1980) are irregularly shaped cocci, most often arranged to sarcina cell packages. In addition long filaments formed with rods were observed by species of *Methanotrix* [formerly designated *Methanosaeta* (Kamagata et al. 1992)]. The formation of multicellular aggregates irrespective of the individual cell shape can also occur, like for species of *Methanolobus* (Mochimaru et al. 2009), *Methanosarcina* (Kern et al. 2016a), or *Methanobacterium* (Kern et al. 2015).

The diversity of methanogens is also reflected in the different growth conditions. Many methanogens have a mesophilic temperature spectrum, as, e.g. *Methanosarcina*, *Methanobacterium*, or most *Methanococcus*. However, thermophilic and even hyperthermophilic methanogens are known, like *Methanothermobacter thermautotrophicus* or *M. jannaschii* which grow at temperatures of up to 75 and 86 °C, respectively. Even growth up to 110 °C is possible in hot environments as shown for the hyperthermophilic strain *M. kandleri* (Kurr et al. 1991). In contrast, also cold-loving methanogenic strains could be isolated. One example is the methanol-converting archaeon *Methanolobus psychrophilus*, which grows optimally at 18 °C and shows still metabolic activity at 0 °C (Zhang et al. 2008).

Beside the temperature, salt concentration may also be an important physiological parameter for methanogens. A few methanogens have colonized niches such as saline lakes, which are extreme environments for microorganisms because of their high salinity. Microorganisms living under such salty conditions have to protect themselves from losing water and “salting-out”. Due to the fact that biological membranes are permeable to water, a higher solute concentration outside the cell, as in the case of environments with a high salinity, would drag water out of the cell and would lead to cell death. To prevent the loss of water, and as a countermeasure, microbes increase the cytoplasmatic osmolarity to survive in such salty environments. This can be done in two ways. The first is the synthesis and accumulation of osmoprotectants,

also known as compatible solutes, which have a small molecular mass and a high solubility. This has been shown for example for *M. mazei*. At a NaCl concentration of 400 mM the methanogen synthesizes glutamate in response to hypersalinity. At higher salt concentration (800 mM NaCl) *N*-acetyl- β -lysine is synthesized in addition to glutamate (Pflüger et al. 2005, 2003). But *N*-acetyl- β -lysine is not essential for growth and can be also substituted by glutamate and alanine at high salinity (Saum et al. 2009). Moreover it has been also shown that *M. mazei* can take up the osmoprotectant glycine betaine from its environment (Roessler et al. 2002). The second way to protect the cell from losing water, and to balance the cytoplasm osmotically with the high salinity of the environment, is an influx of potassium and chloride into the cytoplasm (Oren 2008). This as “high-salt-in strategy” known way may be also used by the recently discovered “Methanonatronarchaea” (Sorokin et al. 2017). They appear to be extremely halophilic, methyl-reducing methanogens related to the haloarchaea.

Although most (by far) methanogens grow optimally around neutral pH, some, which are halophilic or halotolerant, show also an adaptation to alkaline pH. *Methanocalculus alkaliphilus* grows alkaliphilically with an optimum at pH 9.5 and a moderate salinity up to 2 M of total Na⁺, whereas *Methanosalsum natronophilum* can even tolerate higher salinities, up to 3.5 M of total Na⁺, at the same alkaline pH (Sorokin et al. 2015). Moderately acidic environments can also be inhabited by methanogens as, for example, *Methanoregula booneii*, which was isolated from an acidic peat bog and has a pH optimum for growth of 5.1 (Bräuer et al. 2006, 2011).

Substrates and metabolism of methanogens

Methanogens use the substrate CO₂ and the electron donor H₂ during hydrogenotrophic methanogenesis. In the first step, CO₂ is reduced and activated to formylmethanofuran (Wagner et al. 2016) in which reduced ferredoxin (Fd_{red}) is the electron donor for this reaction (Fig. 1).

In the second step the formyl group is transferred to tetrahydromethanopterin (H₄MTP) obtaining formyl-H₄MTP. Then the formyl group is dehydrated and reduced to methylene-H₄MTP and subsequently to methyl-H₄MTP with reduced F₄₂₀ (F₄₂₀H₂) as electron donor. The methyl group is then transferred to coenzyme M (HS-CoM). Finally, methyl-CoM is reduced to methane with coenzyme B (HS-CoB) as electron donor. The resulting heterodisulfide (CoM-S-S-CoB) is reduced with H₂ to recycle the coenzymes (Liu and Whitman 2008; Thauer et al. 2008). It is also important to note that several methanogens can use formate instead of H₂ as electron source for CO₂ reduction. There, four formate




Fig. 1 Schematic overview of hydrogenotrophic (a), aceticlastic (b) and methylotrophic (c) methanogenesis. Hydrogenotrophic methanogenesis for Ech-containing methanogens is shown. The methylotrophic methanogenesis from methanol is displayed. Abbreviations are mentioned in the text (Adapted from (Thauer et al. 2008; Welte and Deppenmeier 2014; Welander and Metcalf, 2005))

molecules are first oxidized to CO_2 by formate dehydrogenase (Fdh) followed by the reduction of one molecule of CO_2 to methane (Liu and Whitman 2008). Instead of H_2 , a few methanogens can also use alcohols like ethanol or 2-propanol as electron donors (Frimmer and Widdel 1989; Widdel 1986).

Some methanogens can also use carbon monoxide (CO) for methanogenesis. In *Methanosarcina barkeri* and *M. thermautotrophicus* four molecules of CO are oxidized to CO_2 by CO dehydrogenase (CODH) followed by the reduction of one molecule of CO_2 to methane with H_2 as electron donor (Daniels et al. 1977; O'Brien et al. 1984). Thus, growth on H_2 and CO_2 is still possible with both methanogens. In contrast, CO metabolism of *Methanosarcina acetivorans* seems to be different. It can also use CO, but is unable to grow on H_2 and CO_2 due to the lack of a functioning hydrogenase system. Further, the organism produces high amounts of acetate and formate from CO during methanogenesis (Rother and Metcalf

2004). The genera *Methanosarcina* and *Methanotrix* can use acetate for methane production. In this aceticlastic methanogenesis acetate has to be activated first. It is converted with ATP and coenzyme A (CoA) to acetyl-CoA, which is then split by the CODH/acetyl-CoA synthase complex. The methyl group is transferred to H_4MTP [which is tetrahydrosarcinapterin (H_4SPT) in *Methanosarcina*] and further converted to methane like in the CO_2 reduction pathway. The carbonyl group is oxidized to CO_2 , thus providing the electrons for the methyl group reduction (Welte and Deppenmeier 2014).

The third way of biological methanation is methylotrophic methanogenesis in which methylated substrates as methanol, methylamines or methylated sulfur compounds like methanethiol or dimethyl sulfide, are utilized. Most methylotrophic methanogens belong to the *Methanosarcinales*. In the first step the methyl-group from the methylated substrate is transferred to a corrinoid protein by a substrate-specific methyltransferase

(MT1) and subsequently to HS-CoM by another methyltransferase (MT2), thus forming methyl-CoM (Burke and Krzycki 1997). One methyl-CoM is oxidized to CO₂ (via the hydrogenotrophic pathway in reverse) generating the reducing equivalents to reduce three methyl-CoM to methane and also generating a proton motive force (Timmers et al. 2017; Welte and Deppenmeier 2014).

Energy conservation in methanogens

In general, methanogens can be divided into two groups according to their mode of energy conservation: methanogens without and with cytochromes (Mayer and Müller 2014; Thauer et al. 2008). Most of the methanogenic archaea do not contain cytochromes. They have a methyl-H₄MPT:coenzyme M methyltransferase (Mtr) which couples the methyl group transfer to a primary, electrochemical Na⁺ gradient over the membrane (Becher et al. 1992; Gottschalk and Thauer 2001). Furthermore, the H₂-dependent reduction of CoM-S-S-CoB in cytochrome-free methanogens is catalyzed by a complex consisting of a (methyl viologen-reducing) hydrogenase and heterodisulfide reductase (Mvh-Hdr), which also couples this exergonic process to the concomitant endergonic reduction of oxidized ferredoxin (Fd_{ox}) via flavin-based electron bifurcation (Buckel and Thauer 2013). Due to the existence of a Na⁺ binding motif in the *c* subunits of A₁A_O ATP synthases of almost all non-cytochrome containing methanogens (one exception is *Methanosalsum zhilinae*), the established Na⁺ gradient can be used for ATP synthesis (Mayer and Müller 2014; Grüber et al. 2014).

Cytochrome-containing methanogens such as *M. mazei* or *M. barkeri*, also employ Mtr, thus, generating a Na⁺ gradient over the membrane. However, reduction of CoM-S-S-CoB is catalyzed by a membrane-bound heterodisulfide reductase (HdrED), which obtains electrons from reduced methanophenazine (MPhH₂, functionally analogous to quinones) via its cytochrome *b* subunit, which is coupled to the generation of a proton motive force. During hydrogenotrophic methanogenesis, a membrane-bound (F₄₂₀ non-reducing) hydrogenase (Vho) oxidizes H₂ and transfers electrons via cytochrome *b* to oxidized methanophenazine (MPh), again generating a proton motive force. Further, another membranous energy converting hydrogenase, Ech (which is similar to complex I) couples the endergonic reduction of Fd_{ox} with H₂ to the intrusion of H⁺, i.e., uses the proton motive force (Mayer and Müller 2014; Thauer et al. 2008; Welte and Deppenmeier 2014). Under environmental conditions, e.g. as in a biogas plants, cytochrome-containing *Methanosarcina* are outcompeted by “true” hydrogenotrophic methanogens, which produce methane from CO₂ and H₂ exclusively.

Methanosarcina acetivorans lacks both Vho and Ech. Instead it employs an Rnf complex which is thought to establish a Na⁺ gradient over the membrane by transferring electrons from Fd_{red} (accrued from, e.g., oxidation of CO or oxidation of the carbonyl-group from acetyl-CoA) to MPh. Subsequent electron transport from MPhH₂ to HdrED again generates a H⁺ gradient (Mayer and Müller 2014; Schlegel et al. 2012b; Welte and Deppenmeier 2014).

The fact that methanogenesis in cytochrome-containing methanogens is coupled to the generation of both a H⁺ and a Na⁺ gradient (Schlegel and Müller 2013) may be also reflected by the ion dependence of their A₁A_O ATP synthases. It has been shown that the A₁A_O ATP synthase from *M. acetivorans* can use both ion gradients (Schlegel et al. 2012a).

During methylotrophic growth of cytochrome-containing methanogens oxidation to CO₂ involves reduction of cofactor F₄₂₀, which is a 5-deazaflavin derivative. F₄₂₀H₂ is re-oxidized by F₄₂₀H₂ dehydrogenase (Fpo), which is a membrane-bound complex (similar to Nuo of *E. coli*) and transfers electrons to MPh, thereby establishing a H⁺ gradient over the membrane in addition to the H⁺ gradients at Hdr and Ech, and the use of the Na⁺ gradient at Mtr (Welte and Deppenmeier 2014).

Analyses of genomes from *Bathyarchaeota* (Evans et al. 2015) and *Verstraetearchaeota* (Vanwonterghem et al. 2016) suggest a methylotrophic methane metabolism for members of these two new phyla. Reduction of the CoM-S-S-CoB in the *Verstraetearchaeota* might be accomplished by the Mvh-Hdr complex which might be coupled to re-oxidation of Fd_{red} by an Ehb or and Fpo-like complex. However, what type of ion gradient (H⁺ and/or Na⁺) might be established over the membrane, is unclear, although H⁺ are predicted to be the coupling ion of the respective A₁A_O ATP synthase (Vanwonterghem et al. 2016). It is obvious that pure culture isolation of *Verstraetearchaeota* is required in order to address the physiology and energy conservation in these potential methanogens.

In the *Bathyarchaeota* energy conservation is even more of a mystery. Two available metagenomes, BA1 and BA2 (proposed to be 91.6 and 93.8% complete, respectively), are missing most of the genes encoding for methanogenic energy conservation. Mtr is incomplete, Fpo as well as an energy-converting hydrogenase (like EhaB, establishing a H⁺ or Na⁺ gradient over the membrane), are missing. In the genome of BA1 only an Ech hydrogenase is encoded. Also, genes encoding for an A₁A_O ATP synthase are absent, which would restrict the organism to ATP synthesis by substrate level phosphorylation (SLP) (Evans et al. 2015).

Electroactivity of methanogens

Electron transfer

When electrodes are inserted into a reactor with methanogens, these electrodes can eventually be used by the organisms to produce methane. An external potential leads to the electrolysis of water at the anode; oxygen and protons are produced, electrons are transferred to the anode. Otherwise, excess electrons out of metabolic reactions can be transferred to the anode, like it would happen in a microbial fuel cell. The electrons migrate to the cathode through an external circuit. At the cathode surface, the electrons are transferred to the methanogens, which can use them to produce methane. The complete mechanism is not yet elucidated, but mainly, three possibilities are suggested (Fig. 2) (Sydow et al. 2014; Geppert et al. 2016). Probably, more than one of these mechanisms contributes to the electron transfer (Zhen et al. 2015).

One possible way would be the transfer of electrons from the cathode to protons, which have been

produced at the anode and migrated through the membrane between anodic and cathodic chamber. Thereby, hydrogen is produced at the cathode, which is then consumed by the methanogens. This indirect electron transfer (IET) would allow the production of methane out of hydrogen and CO₂ (Villano et al. 2010). As an example, IET was observed in *M. thermotrophicus* (Hara et al. 2013). It has also been shown that some *Methanococcus maripaludis* secrete hydrogenases and probably formate dehydrogenases which catalyze the formation of hydrogen and formate directly at the electrode surface; the produced hydrogen and formate is then metabolized by the cells (Deutzmann et al. 2015). This has to be seen as an indirect electron transfer, since the cells were not directly attached to the electrode; from the experimental results, it may be mistaken for a direct electron transport, since the abiotically (without catalyzing hydrogenases) produced amounts of hydrogen and formate cannot explain the amount of methane produced (Deutzmann et al. 2015).




Fig. 2 Extracellular electron transfer. Means of electron transfer within a separated, electromethanogenic system at the cathode: indirect electron transfer (IET), mediated electron transfer (MET) and direct electron transfer (DET)

Another possibility suggests that mediator molecules could accept the electrons at the cathode surface, shuttle it to the methanogens and donate it to the microorganisms. This mediated electron transfer (MET) would imply that the methanogens take up electrons, protons and CO₂ to form methane (Choi and Sang 2016). Flavins, phenazines or quinones can serve as mediator, either naturally secreted by the organisms or added to the reaction medium (Sydow et al. 2014; Patil et al. 2012). A natural secretion of mediators with a redox potential suitable for microbial electrosynthesis (should be < - 0.4 V vs. SHE) has not been observed yet (Sydow et al. 2014). In methanogens, MET could be performed by using neutral red as an electron shuttle (Park et al. 1999).

The third option would be the direct electron transfer (DET) from the cathode surface to the methanogens, e.g. via surface proteins or conductive filaments (so-called nanowires). To generate methane, the microorganisms would use electrons, protons and CO₂ (Cheng et al. 2009). Several studies suggest that direct electron transfer indeed occurs in methanogens (Zhen et al. 2016; Lohner et al. 2014). For a hydrogenase-deficient strain of *M. maripaludis* hydrogenase-independent electron uptake was demonstrated (Lohner et al. 2014), ruling out IET.

In a mixed microbial consortium, direct interspecies electron transfer (DIET) is another possible way of electron transfer. There, one microbial strain takes up electrons at the cathode surface and transfers it to another strain. This may happen, e.g. via conductive filaments (Gorby et al. 2006). It has been reported that this (syntrophic) electron transfer can be very specific between two species, e.g. based on conductive filaments between *M. thermautotrophicus* and *Pelotomaculum thermopropionicum* (Gorby et al. 2006) or between *M. barkeri* and *Geobacter metallireducens* (Rotaru et al. 2014). Apart from this direct interspecies electron transfer, an interspecies hydrogen transfer can occur. Here, one organisms takes up electrons, produces hydrogen as an intermediate and transfers them to a second organism that forms another product. An example is the defined co-culture between the iron-corroding, sulfate-reducing bacterium 'Desulfopila corrodens' IS4 (former name: *Desulfobacterium corrodens*) for electron uptake and *M. maripaludis* for methane production (Deutzmann and Spormann 2017).

Electroactive methanogens

Up to date, most investigations on electromethanogenesis have been carried out with mixed cultures, e.g. from wastewater treatment plants, biogas plants or microbial fuel cells. In technical applications, mixed cultures might be more resistant against environmental stress (Babanova et al. 2017), but it is hard to conclude how many

and which methanogenic strains are electroactive by themselves. From analysis of the mixed cultures studied, it can be concluded which methanogens are enriched and are therefore likely to be electroactive, although mixed culture experiments cannot replace pure culture studies to prove electroactivity. These are for example *Methanobacterium palustre* (Cheng et al. 2009; Battle-Vilanova et al. 2015; Jiang et al. 2014), *Methanosarcina thermophila* (Sasaki et al. 2013), *M. thermautotrophicus* (Sasaki et al. 2013; Fu et al. 2015), *Methanoculleus thermophilus* (Sasaki et al. 2013), *Methanobacterium formicicum* (Sasaki et al. 2013), *M. maripaludis* (Deutzmann and Spormann 2017), *Methanococcus aeolicus* (Feng et al. 2015), *M. mazei* (Feng et al. 2015), *M. arboriphilus* (Jiang et al. 2014), *Methanocorpusculum parvum* (Jiang et al. 2014) and *Methanocorpusculum bavaricum* (Kobayashi et al. 2013). In other studies, the dominant methanogenic organism has not been defined exactly or not explicitly mentioned (Battle-Vilanova et al. 2015; Bo et al. 2014; Zhen et al. 2015). Only few studies have been carried out with pure cultures instead of mixed cultures, so these methanogens are the only ones that are certainly electroactive. To mention are *M. thermautotrophicus* (Hara et al. 2013), and a *Methanobacterium*-like strain IM1 (Beese-Vasbender et al. 2015).

Yet, just a minority of methanogenic strains has been tested for electroactivity, mostly under similar growth conditions. Unfortunately, no specific marker for electroactivity has been found yet (Koch and Harnisch 2016). It is therefore possible that more electroactive methanogens, active even under more extreme conditions, exist.

Genetic tools for methanogens

Many properties of a (model) organisms are unraveled by biochemical and physiological analysis; however where neither of the two lead to satisfactory insight, genetic analysis is often desirable. Furthermore, the accessibility of an organism relevant for applied purposes to genetic manipulating opens the possibility for targeted engineering by removal of -or amendment with- metabolic or regulatory functions. The principal requirements for such a system are sufficiently efficient methods to (a) isolate clonal populations (e.g., via plating on solid media), to (b) transfer genetic material (i.e., protocols for transformation, transduction, or conjugation), and to (c) link the transfer of the genetic material to an identifiable (i.e., screenable or selectable) phenotype (e.g., conferred by marker genes).

The biochemistry of the methanogenic pathway, the trace elements required, as well as the nature and structure of unusual (C1-carrying) cofactors involved has been elucidated using various *Methanobacterium* strains (some of them now reclassified as *Methanothermobacter*).

Therefore, it was a logical next step to develop genetic systems for these models. Plating of *Methanothermobacter* on solid media could be achieved which allowed isolation (and consequently characterization) of randomly induced mutations (Harris and Pinn 1985; Hummel and Böck 1985). However, this species could not be developed into model organisms for genetic analysis because the transfer of genetic material is too inefficient (Worrell et al. 1988). Furthermore, the use of selectable phenotypes was (and still is) restricted because antibiotics (in conjunction with the respective genes conferring resistance) commonly used in bacterial genetics are ineffective in archaea due to the differences in the target structures (e.g., cell wall, ribosomes). Therefore, the establishing of an antibiotic selectable marker (the *pac* gene from *Streptomyces alboniger* conferring resistance to puromycin) in *Methanococcus voltae* (Gernhardt et al. 1990) was key to the development of gene exchange systems in methanogens. Another feature of *Methanococcus*, which facilitated method development, is absence of pseudomurein from its cell wall; instead, the organism is surrounded by a proteinaceous surface (S-) layer that can be removed with polyethylene glycol (PEG), resulting in protoplasts, which apparently can take up DNA. Combined with its comparably robust and fast growth on $H_2 + CO_2$ *Methanococcus* species, most prominently *M. maripaludis*, prevailed as the genetic model for hydrogenotrophic methanogens, for which many useful genetic tools have been developed (Table 1, and see Sarmiento et al. 2011 for a review).

For methylotrophic methanogens containing cytochromes (*Methanosarcina* species) genetic methodology was initially developed on existing tools. PEG-mediated transformation was reported to be ineffective [but later shown to require only modest modifications of the existing protocol (Oelgeschläger and Rother 2009)], but cationic liposomes could be used to transform *Methanosarcina* species (Metcalf et al. 1997). Like in *Methanococcus*, presence of an S-layer and availability of an autonomously replicating cryptic plasmid [pC2A in *Methanosarcina* (Metcalf et al. 1997) and pURB500 in *Methanococcus* (Tumbula et al. 1997)], which could be engineered into shuttle vectors also replicating in *E. coli*, made (heterologous) gene expression comparably easy. Markerless insertion and/or deletion of genes was achieved by establishing counter-selective markers, which are used to remove “unwanted” DNA from the chromosome (Pritchett et al. 2004; Moore and Leigh 2005).

Chromosomal integration and deletion of DNA in methanogens, which can be rather inefficient, mostly relies on homologous recombination requiring sequences of substantial length (500–1000 bp) to be cloned. Thus,

establishing site-specific recombination by engineering a *Streptomyces* phage recombination system (Φ C31) to integrate DNA into (Guss et al. 2008) -and the yeast FLP/FRT system to remove DNA from- the chromosome (Welander and Metcalf 2008), was a major progress for the genetic manipulation of *Methanosarcina*. The recent successful -and highly efficient- application of the CRISPR/Cas9-system from *Streptococcus pyogenes* (Doudna and Charpentier 2014) for gene deletion and insertion in *M. acetivorans* (Nayak and Metcalf 2017) holds the promise of an even easier way to genetically manipulate these important organisms. Most tools (Table 1) developed for one methanoarchaeal model organism can usually be adapted for use in another, as exemplified by exploiting the insect transposable element *Himar1* together with its transposase for random mutagenesis in *Methanosarcina* (Zhang et al. 2000) and, later, in *Methanococcus* (Sattler et al. 2013). Thus, any progress made will likely be useful for all other model systems.

Although it might not be possible to use genetically modified methanogens in the established methanogenic processes like biogasproduction or wastewater treatment, new genetic tools are necessary to guarantee the progress in methanogenic research. It will get clear in the next sections that modified methanogens can be used for bioproduction.

Applications of methanogens

Methanogenic archaea are a very diverse group and some strains can grow under extreme conditions, like extremely high or low temperatures, high osmolarities or pH values. Therefore, the development and optimization of industrially applicable processes making use of methanogens is desirable. This is not only true in terms of methane production as a technical relevant fuel (Ravichandran et al. 2015), but also for other products and applications.

Hydrogen production

It has been observed that several methanogenic strains can also produce hydrogen (Valentine et al. 2000; Goyal et al. 2016). This can happen if the amount of available hydrogen is limited (sub-nanomolar), so that the methanogens seem to start metabolic hydrogen production instead of hydrogen consumption; it has turned out that not methane, but formate and possibly other metabolites can be the source of H_2 ; this cannot be seen as reverse methanogenesis (Valentine et al. 2000; Lupa et al. 2008). The hydrogen production observed by Valentine et al. reached 0.25 $\mu\text{mol}/\text{mg}$ cell dry mass for *Methanothermobacter marburgensis*, 0.23 $\mu\text{mol}/\text{mg}$ cell dry mass for *Methanosaeta thermophila* strain CALS-1

Table 1 Genetic elements used for manipulation of methanogens

Element(s)	Type	Use/phenotype	Organism ^a	References
(Counter-) selectable markers				
<i>pac</i>	Resistance gene	Puromycin resistance	<i>Methanococcus, Methanosarcina</i>	Gernhardt et al. (1990) and Metcalf et al. (1997)
APH3II	Resistance gene	Neomycin resistance	<i>M. maripaludis</i>	Argyle et al. (1996)
<i>aphH-2b</i>	Resistance gene	Neomycin resistance	<i>M. mazei</i> ^b	Mondorf et al. (2012)
<i>ileS3</i>	Resistance gene	Pseudomonic acid resistance	<i>M. acetivorans</i>	Bocazzi et al. (2000)
<i>serC</i>	Biosynthesis gene	Auxotrophic marker	<i>M. barkeri</i>	Metcalf et al. (1996)
<i>proC</i>	Biosynthesis gene	Auxotrophic marker	<i>M. acetivorans</i>	Pritchett et al. (2004)
<i>hpt</i>	Purine salvage gene	Counter-selectable marker (8-aza-2,6-diaminopurine)	<i>Methanosarcina</i>	Pritchett et al. (2004)
<i>upt</i>	Purine salvage gene	Counter-selectable marker (6-aza-uracil)	<i>M. maripaludis</i>	Moore and Leigh (2005)
Controlling gene expression				
<i>PhmVA</i>	Promoter	Constitutive gene expression	<i>Methanococcus</i>	Beneke et al. (1995)
<i>Psi</i>	Promoter	Constitutive gene expression	<i>Methanococcus</i>	Sun and Klein (2004)
<i>PmcrB/Tmcr</i>	Promoter/terminator	Constitutive gene expression	<i>Methanococcus, Methanosarcina</i>	Gernhardt et al. (1990) and Müller et al. (1985)
<i>PmcrB(tetO)/TetR</i>	Promoter/repressor	Tetracycline-dependent gene de-repression	<i>Methanosarcina</i>	Guss et al. (2008)
<i>PmtaCB1</i>	Promoter	Methanol-dependent gene induction	<i>M. acetivorans</i>	Rother et al. (2005)
<i>PmtbC</i>	Promoter	Methylamine-dependent gene induction	<i>M. mazei</i>	Mondorf et al. (2012)
<i>Pmif/NrpR</i>	Promoter/repressor	N-dependent gene repression	<i>M. maripaludis</i>	Cohen-Kupiec et al. (1997) and Lie and Leigh (2003)
tc-RS4	Riboswitch	Tetracycline-dependent gene repression	<i>M. acetivorans</i>	Demolli et al. (2014)
Random mutagenesis				
<i>Himar1/Tnp</i>	Insect transposon/transposase	Random transposon mutagenesis	<i>M. acetivorans, M. maripaludis</i>	Zhang et al. (2000) and Sattler et al. (2013)
<i>Tn5/Tnp</i>	Transposon/transposase	Random transposon mutagenesis	<i>M. maripaludis</i>	Porat and Whitman (2009)
Reporter genes				
<i>lacZ</i>	Reporter gene	Quantification of gene expression/promoter function	<i>M. maripaludis</i>	Gardner and Whitman (1999)
<i>uidA</i>	Reporter gene	Quantification of gene expression/promoter function	<i>M. voltae, Methanosarcina</i>	Beneke et al. (1995) and Pritchett et al. (2004)
<i>bla</i>	Reporter gene	Quantification of gene expression/promoter function	<i>M. acetivorans</i>	Demolli et al. (2014)
Facilitating recombination/mutagenesis				
Φ C31	Phage recombination system	Site specific chromosomal integration	<i>Methanosarcina</i>	Guss et al. (2008)
<i>int</i>				
<i>attB/P</i>				
<i>flp/FRT</i>	Yeast recombination system	Marker rescue	<i>Methanosarcina, Methanococcus</i>	Welander and Metcalf (2008) and Hohn et al. (2011)
Cas9	Nuclease	Targeted genome editing	<i>M. acetivorans</i>	Nayak and Metcalf (2017)

^a If genus is given, the element is effective in various model species

^b The construct used does not confer resistance to neomycin in *M. acetivorans* (Matschiavelli and Rother, unpublished)

and 0.21 $\mu\text{mol}/\text{mg}$ cell dry mass for *M. barkeri* strain 227 (Valentine et al. 2000). Several strains of *M. maripaludis* produced 1.4 $\mu\text{mol}/\text{mg}$ of hydrogen per milligram of cell dry mass, out of formate (Lupa et al. 2008). This application is still restricted to the lab scale (Valentine et al. 2000) and to create a reasonable process, genetic engineering would have to be done to increase the hydrogen yield (Goyal et al. 2016). It is assumed that the hydrogenases present in methanogens are the enzymes catalyzing the hydrogen production (Valentine et al. 2000). A possible way to increase the hydrogen yield could therefore be the detection of the relevant hydrogenase and afterwards overexpressing it.

Biotechnological production by genetically modified methanogens

During recent years, genetic tools for methanogens have been improved, opening a new field of research on these important microorganisms. As a first step, the product spectra of methanogens could be increased. For example, it has been possible to modify *M. maripaludis* to produce geraniol instead of methane from $\text{CO}_2 + \text{H}_2$ or from formate (Lyu et al. 2016). Apart from allowing different products, it has also been possible to broaden the substrate range. As an example, the introduction of a bacterial esterase allowed *M. acetivorans* to grow on methyl-esters (like methyl acetate and methyl propionate, Lessner et al. 2010). In wild type methanogens, “trace methane oxidation” (i.e., “reverse methanogenesis”) has been reported to occur during net methane production (Timmers et al. 2017). It has been possible to use this effect for acetate production: Heterologous expression in *M. acetivorans* of genes encoding methyl-CoM reductase from anaerobic methanotrophic archaea (ANME-1) resulted in a strain that converted methane to acetate three times faster than the parental strain (Soo et al. 2016). Also, additional expression of the gene encoding 3-hydroxybutyryl-CoA dehydrogenase (Hbd) from *Clostridium acetobutylicum* resulted in formation of L-lactate (0.59 g/g methane) from methane with acetate as intermediate, possibly by Hbd exhibiting lactate dehydrogenase activity in the heterologous host (McAnulty et al. 2017). Thus, the principal possibility might exist to engineer *M. acetivorans* for industrial production. However, as both conversion rates and product yields were low and for neither case the conversion stoichiometries reported, the applicability of such a system remains in question. The same holds true for the production of other high value products like amino acids or vitamins with methanogens, and due to their slow growth, a technical application is not yet developed (Schiraldi et al. 2002). But since there is continuous progress in the development of genetic tools for methanogens, as described

above, it is thinkable that new processes with heterologous methanogens will emerge during the next years.

Methane from oil and coal beds

Nearly two-thirds of the fossil oil remains within the oil fields if using conventional production methods (Gieg et al. 2008). It was observed that the residual oil can be converted to natural gas by a methanogenic consortium, which was added to the oil field (Gieg et al. 2008). The consortium used was gained from subsurface sediments and could be enriched with crude oil. *Methanosaeta* spec. was the dominant archaeon in the enrichment, which also contained syntrophic sulfate-reducing bacteria, *Clostridiales*, *Bacteroidetes* and *Chloroflexi*. The consortium was added to samples of petroliferous cores from different oilfields, with residual oil saturation of the sandstone grains of approximately 30–40%. Methane could be produced with yields of up to 3.14 mmol/g crude oil (Gieg et al. 2008). Apart from oil fields, also oil sands tailing ponds or other oil–water emulsions could be treated that way (Voordouw 2011). But since costs for natural gas remain relatively low, whilst those for crude oil are significantly higher, this approach remains experimental due to lack of benefit (Voordouw 2011). A natural source of methane is coal bed methane. It has been discovered that about 40% of this methane are produced by microbial consortia containing methanogens (e.g. *Methanosarcinales*); the substrates for this production are methoxylated aromatic compounds within the coal beds (Mayumi et al. 2016). It was recently discovered that pure cultures of *Methermicoccus shengliensis* can produce up to 10.8 $\mu\text{M}/(\text{g coal})$ methane (Mayumi et al. 2016). Coal bed methane is already industrially used; it might be possible to use *M. shengliensis* for methane production from other sedimentary organic material (Mayumi et al. 2016).

Biogas production from organic matter

The main technical application of methanogens is the production of biogas by digestion of organic substrates. It is estimated that up to 25% of the bioenergy used in Europe could be produced using the biogas process until 2020 (Holm-Nielsen et al. 2009). Digestion of organic matter can be seen as a four-stage process. During the first step (hydrolysis), complex organic matter (proteins, polysaccharides, lipids) is hydrolyzed by exo-enzymes to oligo- and monomers (amino acids, sugars, long chain fatty acids), which can be taken up by microorganisms (Vavilin et al. 2008). The second step, fermentation or acidogenesis, leads to an oxidation of the compounds formed during hydrolysis to typical fermentation products like butyrate, propionate, acetate, formate, ethanol, H_2 and CO_2 . Acetogenesis represent the third step, where the fermentation products are oxidized, mostly to acetate

and CO₂ with the concomitant formation of H₂ (Batstone et al. 2002). However, this process is only sufficiently exergonic for the organisms if the H₂ partial pressure is kept very low (McInerney et al. 2008). This requires the fourth step, methanogenesis, where acetate (and methylated compounds) and CO₂ and H₂ is converted to methane by the methanogens. This implicates that a syntrophic consortium of microorganisms is always needed, whereas the exact composition of this consortium can not only change over time, but also vary between different reactors (Solli et al. 2014). Depending on the microbial community and the type of methanogens within, this process can be carried out in psychrophilic, mesophilic or thermophilic temperature range (Vanegas and Bartlett 2013). For stable biogas production, hydrolysis, acidogenesis, acetogenesis and methanogenesis have to run within the digester in balanced reaction rates to prevent the overacidification of the reactor by surplus protons. However, the microorganisms responsible for these different steps often have different optimal growth conditions, so it is crucial that conditions are maintained, which favor all steps (Niu et al. 2015). Therefore, careful control of process parameters like temperature (Vanegas and Bartlett 2013), hydraulic retention time (Rincón et al. 2008), pH (Lay et al. 1997) and ammonia concentration (Karakashev et al. 2005) are necessary. Apart from that, the biogas yield and the process operation and conditions strongly depend on the type of substrate used (Niu et al. 2015). It was for example observed that the methanogenic consortium, which is strongly depending on the substrate type, is usually dominated by *Methanosaetaeaceae* in digesters with sludge as substrate, while solid waste digesters operated with manure explained in the following section usually host a majority of *Methanosarcinaceae* (Karakashev et al. 2005). In both cases, methanogens that can metabolize acetate (see also “[Substrates and metabolism of methanogens](#)” section) are preferred in biogas systems, compared to those feeding on hydrogen and CO₂. Apart from the substrate type itself, a differentiation is made between wet and dry fermentation, whereby the more common wet fermentation includes up to 10% of solids in the substrate, and the dry fermentation between 15 and 35% (Stolze et al. 2015).

Treatment of sewage water

The treatment of sewage water by anaerobic digestion does not only lead to biogas production but also to clean water. Using a methanogenic process to convert the organic matter within wastewater to biogas reduces the amount of sludge to be disposed, lowers its pathogenic potential and usually needs less additional energy than aerobic processes, since biogas as energy fuel is produced and no energy intense aeration is necessary (Martin et al.

2011). Apart from that, the greenhouse gas emission of the anaerobic process is lower when treating high strength waste waters, although no greenhouse gas savings could be detected for low strength sewage water (Cakir and Stenstrom 2005).

A commonly used system for the anaerobic treatment of wastewater is the upflow anaerobic sludge blanket (UASB) reactor; wastewater enters the reactor from the bottom and flows to an outlet in the upper part of the reactor. Sludge particles out of the waste water agglomerate and form a sludge blanket, which has to be passed by the incoming wastewater. In this zone, the methanogenic consortium digests organic material and produces biogas, which leaves the reactor at its top. Since the solubility of methane in water is low compared to that of CO₂, the holdup of methane within the water is negligible (Sander 2015). The contact between organic material and microorganisms is sufficient for efficient methane production due to the sludge blanket, thus allowing higher loading rates than in other reactor types. The system only requires a low energy input, but needs a long start-up phase of several months, until the sludge blanket has fully established (Rajeshwari et al. 2000). To overcome long start up phases, continuously stirred tank reactors can be used, but here, organic loading rates are about tenfold lower than in the UASB reactor (Rajeshwari et al. 2000). It is important to consider the type of sewage water (e.g. from breweries, paper mills, oil mills, dairy production or other) when estimating the biogas yield of digestion. Different organic loads or different substrate composition lead not only to fluctuating amounts of biogas, but also to changes of the biogas composition (reviewed by Tabatabaei et al. 2010). Instead of treating sewage water itself via anaerobic digestion, it is also possible to purify the water by aerobic processes and anaerobically digest the remaining sewage sludge (Van Lier et al. 2008). Usually, a pretreatment of the sludge can increase the biogas yield. This can for example, but not only, be an alkaline pretreatment, ozonation, ultrasonic pretreatment or electric pulses to increase the biodegradability of the sludge (Wonglertarak and Wichitsathian 2014; Bougrier et al. 2007; Rittmann et al. 2008; for a recent review see: Neumann et al. 2016).

Treatment of solids

The largest amount of biodegradable waste for biogas production can be obtained from the agricultural sector. This includes animal manure and slurry from the production of pig, poultry, fish and cattle (Holm-Nielsen et al. 2009). The treatment of agricultural wastes like animal manure with methanogenic consortia is not only beneficial in terms of the biogas produced. It also reduces odors and pathogens and is therefore increasing the fertilizer

qualities of the manure (Sahlström 2003). The process of biogas formation does not necessarily have to be coupled to waste treatment. Biogas plants can also be operated with energy crops cultured for the biogas production, like sugar beet or maize silage (Demirel and Scherer 2008; Lebuhn et al. 2008). Another possibility is the anaerobic digestion of microalgae, which lowers the necessary cultivation area (Mussnug et al. 2010). Especially if energy crops without addition of manure are digested, it can be necessary to add micronutrients to ensure optimum growth conditions (Choong et al. 2016). It is also important to consider that lignocellulosic materials are not fully convertible without pretreatment, which leads to lower methane yields (Zheng et al. 2014). Table 2 shows production yields for different solid substrates.

A crucial aspect of the biogas process is the design of the anaerobic digester (Nizami and Murphy 2010). There are several digester types for the anaerobic digestion of wastewater. For the digestion of solids, biogas plants

are usually designed as continuously stirred tank reactors (CSTRs). Even though this might be the easiest and cheapest way of biogas production, it turned out that the efficiency can be increased by using a serial system. Here, two CSTRs are used; biogas yield was increased by a longer overall retention time (Boe and Angelidaki 2009; Kaparaju et al. 2009). Instead of CSTRs, plug-flow systems have been invented by different companies to perform continuous processes (Fig. 3); in a serial digestion, they would usually be taken for the first stage (Weiland 2010). Another possibility is the use of a batch process, especially for substrates with low water contents, for example in a garage type fermenter (Li et al. 2011; Nizami and Murphy 2010).

Micro biogas systems

An interesting application of the biogas process is the use of micro biogas plants in developing countries. These plants of up to 10 m³ can be operated using

Table 2 Biogas production from organic wastes

Substrate	Biogas (ml/gVS)	Methane (ml/gVS)	Methane content (%)	References
Food waste	784	518	66.1	Liu et al. (2009)
Green waste	631	357	56.5	Liu et al. (2009)
Bovine manure	150	40	46.5	Fantozzi and Buratti (2009)
Chicken manure	220	110	66.6	Fantozzi and Buratti (2009)
Pig manure	412	216	52	Amon et al. (2006)
Sugar beet	730	387	53	Weiland (2010)
Grass	211	150	71	Yu et al. (2002)
Maize	560	291	52	Weiland (2010)
Microalgae (<i>Chlamydomonas reinhardtii</i>)	784	518	66.1	Mussnug et al. (2010)
Microalgae (<i>Arthrospira platensis</i>)	631	357	56.5	Mussnug et al. (2010)

VS volatile solids




Fig. 3 Plug flow digesters for biogas production. **a** "Kompogas" reactor. Horizontal plug flow reactor. Additional mixing by axial mixer. Increased process condition stability by partial effluent recycling. Gas outlet on top of the outlet side. 23–28% total solids. **b** Valorga reactor. Substrate entry at the bottom; plug flow over a vertical barrier to the outlet. Additional mixing by biogas injection at the bottom. 25–35% total solid content. **c** Dranco reactor. Substrate entry with partial effluent recycling at the bottom, upward flow through substrate pipes. Downward plug flow to outlet. 30–40% total solids (Li et al. 2011; Nizami and Murphy 2010)

domestic organic waste or feces, while the produced gas can be used directly for heating and cooking. There are also attempts to convert the biogas out of those digesters with volumes of up to 10 m³ to electricity, which might be valuable in rural areas (Plöchl and Heiermann 2006). These reactors are particularly popular in China and India and programs to equip households with biogas energy are supported by the government (Bond and Templeton 2011). Domestic biogas plants are especially beneficial in warm regions (e.g. Africa around the equator, South-East Asia) with sufficient water available. In general, 3 types of digesters are used, which are the fixed dome, the floating cover digester, which was further developed to the ARTI biogas system, and the plug flow (or tube) system (Fig. 4).

Although micro biogas systems might not solve the energy problems in developing countries, and the investment costs may not be covered without governmental subsidy, some positive impacts of this technology can be observed. The deforestation in rural areas decreases since wood is not needed for heating, at the same time risks caused by open fire in closed buildings

are minimized by the use of a biogas driven stove. The amount of pathogens in the substrate (waste and feces) is decreased, so that it can be reused as fertilizer (Bond and Templeton 2011). Therefore, microbiogas systems are an important contribution to the development of third world countries and a use of the biogas process not standing in conflict to food-production, since organic waste is the main substrate.

Biogas composition and process optimizations

The composition of biogas does not only include methane, but also up to 40% CO₂, water, hydrogen sulfide and other trace gases. Biogas is usually flammable due to the high yield of methane (40–75%), but for the use in engines or for injection into the natural gas grid it has to be purified and upgraded in methane content. This leads to higher calorific values of the biogas and avoids the presence of corrosive gases like hydrogen sulfide, which could cause damages to engines and pipes if remaining in the biogas (Ryckebosch et al. 2011). There are several upgrading techniques, which take place after digestion (extensively reviewed by Bauer et al. 2013).




Fig. 4 Micro biogas systems. **a** Arti biogas (India). Material two plastic water tanks (working volume of 1 m³). Substrate mainly kitchen waste. Disadvantage of gas losses of up to 20% (Voegeli et al. 2009). **b** Floating cover (India). Material bricks and metal cover. Top rises when gas is produced. Substrate mainly pig and cow manure (Bond and Templeton 2011). **c** Fixed dome (China). Material bricks and clay. Substrate mainly pig and cow manure (Plöchl and Heiermann 2006). **d** Plug flow. Material affordable plastic foils (Bond and Templeton 2011)


Process optimization can influence the biogas composition already during the process, lowering the costs of after-process purification. Numerous investigations on improvement of the biogas process have been undertaken, either to increase the overall amount of biogas, or to increase the methane content of the biogas.

It has turned out that careful pretreatment of the organic substrates leads to higher percentages of methane in the biogas. Several pretreatment methods such as chopping, alkali treatment and thermal treatment are reviewed in Andriani et al. (2014). From a biotechnological point of view, biological pretreatment of substrate is especially interesting. Biological pretreatment can increase the biogas production; this method was described by Zhong et al. (2011) which led to a 33% increase of biogas production (Zhong et al. 2011). The substrates were exposed to a microbial agent including yeasts, cellulolytic bacteria and lactic acid bacteria, which degraded the substrate before the actual start of the anaerobic digestion. A reduction in lignin, cellulose and hemicelluloses content could be observed after 15 days of pretreatment. The following anaerobic digestion showed an increase of biogas yield and methane content (Zhong et al. 2011). Apart from the pretreatment of the single substrates, a mixture of different substrates (co-digestion) or a backmixing of digester effluent can lead to a better performance of the system (Weiland 2010; Sosnowski et al. 2003). Co-digestions can be carried out with mixtures of manure and energy plants or sewage slug and solid wastes and increase the methane production because of stabilizing the C:N ratio within the digester (Ward et al. 2008; Sosnowski et al. 2003). Another optimization method is addition of inorganic particles to the fermentation medium. Addition of nanoparticles of zero-valent iron could enhance the methane production by 28% (Carpenter et al. 2015). An increase in biogas formation could also be observed with magnetic iron oxide particles (Abdelsalam et al. 2017). Other particles include

charcoal, silica and mineral salts were investigated (reviewed by Yadvika et al. 2004). The improvement in biogas yield could be due to aggregation of bacteria and methanogens around the particles, leading to a lower washout and higher culture densities; it is also possible that metal particles release electrons to the surrounding medium, which can be used for methane formation, but the exact mechanism remains unclear (Yadvika et al. 2004; Carpenter et al. 2015).

One promising method for biological biogas upgrading in methane content is the conversion of the residual CO₂ to additional methane using hydrogenotrophic methanogens, which are capable of producing methane solely out of CO₂ and H₂ (Bassani et al. 2015). H₂ can either be injected into the anaerobic digester (Luo et al. 2012), or H₂ and biogas can be mixed in a second reactor containing methanogens (Bassani et al. 2015; Luo and Angelidaki 2012) (Fig. 5). If introducing hydrogen to the anaerobic digester, there may be a shift within the methanogenic community: acetoclastic methanogens decrease, while hydrogenotrophic methanogens (especially *Methanoculleus*) are enriched; also, hydrolyzing and acidifying bacteria decrease, while syntrophic bacteria producing acetate increase (see also “Substrates and metabolism of methanogens” section; Bassani et al. 2015). Technical concepts for the integration of H₂ into existing biogas plants and effective new means of process control are necessary to make this process commercially attractive. Therefore, experiments have to be carried out under industrial conditions, i.e. under fluctuating substrate compositions, in reactors with zones of different substrate concentrations, changing microbial consortium and different pressure zones according to a larger reactor height; these conditions will usually not appear in lab scale, unless they are particularly tested.

H₂ is usually produced by water electrolysis, a process in which electricity is used to split water and generate oxygen and hydrogen. To couple water electrolysis to



anaerobic methanogenesis and provide a constant level of H₂ within the digester, methanogenic bioelectrochemical systems were invented.

Methanogenic bioelectrochemical systems

The successful increase of methane production by iron addition leads to the conclusion that methanogens may use inorganic surfaces to boost their metabolism by exchange of electrons with the inorganic material (Carpenter et al. 2015). On the other hand, hydrogen addition could also increase the methane output of a biogas plant. A methanogenic bioelectrochemical system (BES) combines these two improvements for increased methane production (Koch et al. 2015). Here, electrodes are introduced into the reaction medium and an external potential is applied. Methanogens can now either interact directly with the electrode surface to gain electrons (Cheng et al. 2009), and/or hydrogen can be produced at the cathode, which can then be consumed by the methanogens to produce methane (Geppert et al. 2016). The whole process belongs to the field of microbial electrosynthesis (MES), which includes processes that convert a substrate into a desired organic product by using microorganisms and electrical current (Schröder et al. 2015; Lovley 2012; Holtmann et al. 2014). The advantage of a BES system

compared to the external production of hydrogen is that short time storage and gassing in of the hardly soluble hydrogen can be avoided (Butler and Lovley 2016).

Notably, the electrode material and size, the membrane material and size and the applied voltage strongly influences the performance of electromethanogenesis, (see Babanova et al. 2017; Krieg et al. 2014; Ribot-Llobet et al. 2013; Siegert et al. 2014 for reviews), but “optimal” conditions for microbial growth and production have not yet been found (Blasco-Gómez et al. 2017). Investigations of this (relatively new) technology have been mostly carried out in lab scale so far, with very few pilot scale approaches (for hydrogen production with methane as side product, see Cusick et al. 2011). Yet, no scale up concept or even well characterized reactor concept exists for electromethanogenesis, whereas various types of bioelectrochemical reactors have been designed (reviewed in Geppert et al. 2016; Krieg et al. 2014; Kadier et al. 2016). Two general modes of integrating electrochemistry into the methanogenic process can be distinguished: first, the electrodes can be integrated into the anaerobic digestion of sewage water or other organic wastes, and secondly, the methanogenic BES can be placed into a second reactor as a stand-alone-process, fed with CO₂, but without additional organic substrates (Fig. 6).




Fig. 6 Increasing methane yield by electrode integration. Top: integration of electrodes into the anaerobic digester; bottom: biogas upgrading in an external, separated MES system fed with CO₂ and electricity

Integration of electrodes into waste and wastewater treatment

To enhance the production of biogas and increase biogas purity, electrodes can be inserted into the anaerobic digester for in situ biogas upgrading. CO₂, which is produced during the digestion of organic matter, can be converted to methane at the electrodes without an additional reactor (Bo et al. 2014). Therefore, the biogas production can be performed during wastewater treatment (Guo et al. 2017) or sewage sludge treatment (Guo et al. 2013) as well as in a mere biogas producing process (Gajaraj et al. 2017). The methane content within the biogas reached up to 98.1% during the digestion of activated sludge and acetate (Bo et al. 2014). It has been shown that the integration of electrodes alters the microbial consortium within the plant, while it is also possible to use adapted consortia, e.g. for psychrophilic temperature ranges (Koch et al. 2015; Bo et al. 2014; Liu et al. 2016). To achieve a reasonable process, ways of electrode integration into existing treatment plants need to be established.

Bioelectrochemical systems fed with CO₂

Methanogenic microbial electrosynthesis can also be carried out in a second reactor, which is equipped with electrodes and fed with CO₂ or a gas containing CO₂. Gas streams rich in CO₂ can be biogas, syngas, or industrial flue gas. The CO₂ contained is often considered a waste component of these gas streams, and since it is also a greenhouse gas, the conversion of CO₂ to more useful chemicals is desirable (Dürre and Eikmanns 2015; Geppert et al. 2016). The conversion of CO₂ by methanogens takes place at the cathode of the system. Since anodic processes like oxygen generation or acid production could inhibit the methanogens, the process can be carried out in a two-chamber system, where anode and cathode chamber are separated by a proton-exchange membrane, which allows the transfer of protons from anode to cathode chamber; this is necessary to allow electrical current in the system and maintain the pH within the cathode chamber (Dykstra and Pavlostathis 2017; Cheng et al. 2009). In this system, it is possible to use a pure methanogenic culture (Beese-Vasbender et al. 2015) or an enriched methanogenic consortium (Dykstra and Pavlostathis 2017) at the cathode, while the anode chamber can be abiotic (water electrolysis) or biotic (degradation of organic matter) (Dykstra and Pavlostathis 2017).

As mentioned, the bioelectrochemical methanogenesis is currently still a lab-scale application. To gain an economical technical process, concepts for process characterization and control, reactor balancing, and scale

up of reactors have to be developed. To create further progress in this field and also in bioelectrochemical applications, genetic tools might be necessary to create methanogens with higher electron uptake rates, e.g. via the integration of (more) cytochromes into the membrane or the heterologous secretion of electron shuttles.

Conclusions

Methanogens are interesting organisms, both from a biological, as well as for a technological, point of view. Research of the last years made it clear that this unique group of microbes is far from being fully understood. During the last years, several reviews on biological aspects of methanogens (Borrel et al. 2016; Goyal et al. 2016), on natural methanogenesis (e.g. Park and Liang 2016; Bao et al. 2016) or on single technical applications, eventually in combination with the very specific biology within the process (Biogas: Wang et al. 2017; Braguglia et al. 2017; Koo et al. 2017; Biogas upgrading and optimization: Neumann et al. 2016; Choong et al. 2016; Romero-Güiza et al. 2016; Bioelectromethanation: Blasco-Gómez et al. 2017; Geppert et al. 2016) have been published. All these review articles are rather specialized to one single aspect of methanogens. This review combines all these aspects, including a review of recently developed tools, to give an overview over the whole field of methanogenic research. Therefore, it makes it possible to understand challenges in industrial applications by giving the biological basics and helps to imagine applications for results from basic research in industry. Industry mainly focused on the production of biogas with methanogens, but other applications, especially when considering electroactivity of methanogens, seem feasible. Newly developed genetic tools for methanogens are useful to design a wider product spectrum, which raises the technical relevance of methanogens. However, most processes possible with methanogens are still not economically feasible, since their strict requirement for anaerobic conditions raises the investment costs and their slow growth leads to long process times. It would be desirable to have further comparable knowledge of the efficiency of different methanogenic strains in terms of space time yield and conversion rates under industrially relevant conditions, for example by performing pure culture studies with fluctuating substrate composition, fluctuating pH and under different substrate concentrations. A major problem here remains the comparability of published data about methanogenic performance in biogas plants as well as in electrochemical systems, since studies have been carried out under various conditions. For some applications, especially microbial electrosynthesis,

more research of the methanogenic community and comparisons between pure and mixed cultures have to be done to increase methane yields. Still, process optimization, like the use of CO₂-rich waste gas streams as substrates and intelligent process integration will favor methanogenic processes beyond waste treatment in the future. Scale-up of reactors, e.g., for electromethanogenesis or biogas-upgrading, are a major task for process engineers, while genetic engineering may pave the way to produce higher value products from waste CO₂ employing methanogens.

Abbreviations

BES: bioelectrochemical system; CoA: coenzyme A; CoB: coenzyme B; CoM: coenzyme M; CODH: CO dehydrogenase; CSTRs: continuously stirred tank reactors; DET: direct electron transfer; DIET: direct interspecies electron transfer; Ech: energy converting hydrogenase; Fd: ferredoxin; Fd_{ox}: oxidized ferredoxin; Fd_{red}: reduced ferredoxin; Fdh: formate dehydrogenase; Fpo: F₄₂₀H₂ dehydrogenase; Hbd: 3-hydroxybutyryl-CoA dehydrogenase; Hdr: heterodisulfide reductase; H₄MTP: tetrahydromethanopterin; H₄SPT: tetrahydrosarcinapterin; IET: indirect electron transfer; MES: microbial electrosynthesis; MET: mediated electron transfer; MFR: methanofuran; MPH: methanophenazine; MPhH₂: reduced methanophenazine; Mtr: methyl-H₄MPT:coenzyme M methyltransferase; Mvh-Hdr: (methyl viologen-reducing) hydrogenase and heterodisulfide reductase; PHB: polyhydroxybutyric acid; SHE: standard hydrogen electrode; SLP: substrate level phosphorylation; UASB: upflow anaerobic sludge blanket; Vho: (F₄₂₀ non-reducing) hydrogenase.

Authors' contributions

All authors contributed equally in writing this review article. All authors read and approved the final manuscript.

Acknowledgements

The authors thank the DECHEMA Research Institute and the Technical University Dresden for their help in accomplishing this work.

Competing interests

The authors declare that they have no competing interests.

Availability of data and materials

Not applicable.

Consent for publication

Not applicable.

Ethics approval and consent to participate

Not applicable.

Funding

FE, FM and DH thank the BMBF for funding the projects "Microbial electrosynthesis—an integrated research project for the utilization of electric energy in microbial production processes" (FKZ 031A226) and MIKE-Methanation of CO₂ from biogas by microbial electrosynthesis (FKZ: 033RC013A). MR is grateful for funding from BMBF and from the Deutsche Forschungsgemeinschaft.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Author details

¹ DECHEMA Research Institute, Industrial Biotechnology, Theodor-Heuss-Allee 25, 60486 Frankfurt am Main, Germany. ² Technische Universität Dresden, Institut für Mikrobiologie, Zellescher Weg 20b, 01217 Dresden, Germany.

Received: 5 December 2017 Accepted: 19 December 2017
Published online: 04 January 2018

References

- Abdelsalam E, Samer M, Attia YA, Abdel-Hadi MA, Hassan HE, Badr Y (2017) Influence of zero valent iron nanoparticles and magnetic iron oxide nanoparticles on biogas and methane production from anaerobic digestion of manure. *Energy* 120:842–853
- Amon T, Amon B, Kryvoruchko V, Bodirosa V, Pötsch E, Zollitsch W (2006) Optimising methane yield from anaerobic digestion of manure: effects of dairy systems and of glycerine supplementation. *Int Congr Ser* 1293:217–220
- Andriani D, Wresta A, Atmaja TD, Saepudin A (2014) A review on optimization production and upgrading biogas through CO₂ removal using various techniques. *Appl Biochem Biotechnol* 172:1909–1928
- Argyle JL, Tumbula DL, Leigh JA (1996) Neomycin resistance as a selectable marker in *Methanococcus maripaludis*. *Appl Environ Microbiol* 62:4233–4237
- Asakawa S, Akagawa-Matsushita M, Morii H, Koga Y, Hayano K (1995) Characterization of *Methanosarcina mazei* TMA isolated from a paddy field soil. *Curr Microbiol* 31:34–38
- Babanova S, Carpenter K, Phadke S, Suzuki S, Ishii S, Phan T, Grossi-Soyster E, Flynn M, Hogan J, Bretschger O (2017) The effect of membrane type on the performance of microbial electrosynthesis cells for methane production. *J Electrochem Soc* 164:H3015–H3023
- Balch WE, Fox GE, Magrum LJ, Woese CR, Wolfe RS (1979) Methanogens: reevaluation of a unique biological group. *Microbiol Rev* 43:260–296
- Bao Y, Huang H, He D, Ju Y, Qi Y (2016) Microbial enhancing coal-bed methane generation potential, constraints and mechanism - a mini-review. *J Nat Gas Sci Eng* 35:68–78
- Bapteste E, Brochier C, Boucher Y (2005) Higher-level classification of the Archaea: evolution of methanogenesis and methanogens. *Archaea* 1:353–363
- Bär K, Mörs F, Götz M, Graf F (2015) Vergleich der biologischen und katalytischen Methanisierung für den Einsatz bei PtG-Konzepten. *gwf-Gas* 7:1–8
- Bassani I, Kougiaris PG, Treu L, Angelidaki I (2015) Biogas upgrading via hydrogenotrophic methanogenesis in two-stage continuous stirred tank reactors at mesophilic and thermophilic conditions. *Environ Sci Technol* 49:12585–12593
- Battle-Vilanova P, Puig S, Gonzalez-Olmos R, Vilajeliu-Pons A, Balaguer MD, Colprim J (2015) Deciphering the electron transfer mechanisms for biogas upgrading to biomethane within a mixed culture biocathode. *RSC Adv* 5:52243–52251
- Batstone DJ, Keller J, Angelidaki I, Kalyuzhnyi SV, Pavlostathis SG, Rozzi A, Sanders WT, Siegrist H, Vavilin VA (2002) The IWA anaerobic digestion model no 1 (ADM1). *Water Sci Technol* 45:65–73
- Bauer F, Persson T, Hultberg C, Tamm D (2013) Biogas upgrading - technology overview, comparison and perspectives for the future. *Biofuels, Bioprod Biorefin* 7:499–511
- Becher B, Müller V, Gottschalk G (1992) The methyl-tetrahydromethanopterin: coenzyme M methyltransferase of *Methanosarcina* strain Gö1 is a primary sodium pump. *FEMS Microbiol Lett* 91:239–243
- Beese-Vasbender PF, Grote J-P, Garrelfs J, Stratmann M, Mayrhofer KJJ (2015) Selective microbial electrosynthesis of methane by a pure culture of a marine lithoautotrophic archaeon. *Bioelectrochemistry* 102:50–55
- Belay N, Johnson R, Rajagopal BS, Conway de Macario E, Daniels L (1988) Methanogenic bacteria from human dental plaque. *Appl Environ Microbiol* 54:600–603
- Beneke S, Bestgen H, Klein A (1995) Use of the *Escherichia coli uidA* gene as a reporter in *Methanococcus voltae* for the analysis of the regulatory function of the intergenic region between the operons encoding selenium-free hydrogenases. *Mol Gen Genet* 248:225–228
- Blasco-Gómez R, Battle-Vilanova P, Villano M, Balaguer M, Colprim J, Puig S (2017) On the edge of research and technological application: a critical review of electromethanogenesis. *Int J Mol Sci* 18:874
- Bo T, Zhu X, Zhang L, Tao Y, He X, Li D, Yan Z (2014) A new upgraded biogas production process: coupling microbial electrolysis cell and anaerobic

- digestion in single-chamber, barrel-shape stainless steel reactor. *Electrochim Commun* 45:67–70
- Boccazzi P, Zhang JK, Metcalf WW (2000) Generation of dominant selectable markers for resistance to pseudomonadic acid by cloning and mutagenesis of the *ileS* gene from the archaeon *Methanosarcina barkeri* fusaro. *J Bacteriol* 182:2611–2618
- Boe K, Angelidaki I (2009) Serial CSTR digester configuration for improving biogas production from manure. *Water Res* 43:166–172
- Bond T, Templeton MR (2011) History and future of domestic biogas plants in the developing world. *Energy Sustain Dev* 15:347–354
- Borrel G, Adam PS, Gribaldo S (2016) Methanogenesis and the Wood–Ljungdahl pathway: an ancient, versatile, and fragile association. *Genome Biol Evol* 8:1706–1711
- Bougrier C, Battimelli A, Delgenes JP, Carrere H (2007) Combined ozone pretreatment and anaerobic digestion for the reduction of biological sludge production in wastewater treatment. *Ozone Sci Eng* 29:201–206
- Braguglia CM, Gallipoli A, Gianico A, Pagliaccia P (2017) Anaerobic bioconversion of food waste into energy: a critical review. *Bioresour Technol*. 248(Pt A):37–56
- Bräuer SL, Cadillo-Quiroz H, Yashiro E, Yavitt JB, Zinder SH (2006) Isolation of a novel acidiphilic methanogen from an acidic peat bog. *Nature* 442:192–194
- Bräuer SL, Cadillo-Quiroz H, Ward RJ, Yavitt JB, Zinder SH (2011) *Methanoregula boonei* gen. nov., sp. nov., an acidiphilic methanogen isolated from an acidic peat bog. *Int J Syst Evol Microbiol* 61:45–52
- Bryant MP, Boone DR (1987) Emended description of strain MST (DSM 800T), the type strain of *Methanosarcina barkeri*. *Int J Syst Bacteriol* 37:169–170
- Buckel W, Thauer RK (2013) Energy conservation via electron bifurcating ferredoxin reduction and proton/Na⁺ translocating ferredoxin oxidation. *Biochim Biophys Acta Bioenerg* 1827:94–113
- Burke SA, Krzycki JA (1997) Reconstitution of monomethylamine: coenzyme M methyl transfer with a corrinoid protein and two methyltransferases purified from *Methanosarcina barkeri*. *J Biol Chem* 272:16570–16577
- Butler CS, Lovley DR (2016) How to sustainably feed a microbe: strategies for biological production of carbon-based commodities with renewable electricity. *Front Microbiol* 7:1879
- Cakir FY, Stenstrom MK (2005) Greenhouse gas production: a comparison between aerobic and anaerobic wastewater treatment technology. *Water Res* 39:4197–4203
- Carpenter AW, Loughton SN, Wiesner MR (2015) Enhanced biogas production from nanoscale zero valent iron-amended anaerobic bioreactors. *Environ Eng Sci* 32:647–655
- Cheng S, Xing D, Call DF, Logan BE (2009) Direct biological conversion of electrical current into methane by electromethanogenesis. *Environ Sci Technol* 43:3953–3958
- Choi O, Sang B-I (2016) Extracellular electron transfer from cathode to microbes: application for biofuel production. *Biotechnol Biofuels* 9:11
- Choong YY, Norli I, Abdullah AZ, Yahya MF (2016) Impacts of trace element supplementation on the performance of anaerobic digestion process: a critical review. *Bioresour Technol* 209:369–379
- Cohen-Kupiec R, Blank C, Leigh JA (1997) Transcriptional regulation in Archaea: in vivo demonstration of a repressor binding site in a methanogen. *Proc Natl Acad Sci USA* 94:1316–1320
- Cusick RD, Bryan B, Parker DS, Merrill MD, Mehanna M, Kiely PD, Liu G, Logan BE (2011) Performance of a pilot-scale continuous flow microbial electrolysis cell fed winery wastewater. *Appl Microbiol Biotechnol* 89:2053–2063
- Daniels L, Fuchs G, Thauer RK, Zeikus JG (1977) Carbon monoxide oxidation by methanogenic bacteria. *J Bacteriol* 132:118–126
- Demirel B, Scherer P (2008) Production of methane from sugar beet silage without manure addition by a single-stage anaerobic digestion process. *Biomass Bioenergy* 32:203–209
- Demolli S, Geist MM, Weigand JE, Matschiavelli N, Suess B, Rother M (2014) Development of β -lactamase as a tool for monitoring conditional gene expression by a tetracycline-riboswitch in *Methanosarcina acetivorans*. *Archaea* 2014:1–10
- Deutzmann JS, Spormann AM (2017) Enhanced microbial electrosynthesis by using defined co-cultures. *ISME J* 11:704–714
- Deutzmann JS, Sahin M, Spormann AM (2015) Extracellular enzymes facilitate electron uptake in biocorrosion and bioelectrosynthesis. *mBio* 6(2):e00496–15
- Doudna JA, Charpentier E (2014) The new frontier of genome engineering with CRISPR-Cas9. *Science* 346:1258096
- Dridi B, Henry M, El Khéchine A, Raoult D, Drancourt M (2009) High prevalence of *Methanobrevibacter smithii* and *Methanosphaera stadtmanae* detected in the human gut using an improved DNA detection protocol. *PLoS ONE* 4:e7063
- Dridi B, Fardeau M-L, Ollivier B, Raoult D, Drancourt M (2012) *Methanomassiliicoccus luminyensis* gen. nov., sp. nov., a methanogenic archaeon isolated from human faeces. *Int J Syst Evol Microbiol* 62:1902–1907
- Dürre P, Eikmanns BJ (2015) C1-carbon sources for chemical and fuel production by microbial gas fermentation. *Curr Opin Biotechnol* 35:63–72
- Dykstra CM, Pavlostathis SG (2017) Evaluation of gas and carbon transport in a methanogenic bioelectrochemical system (BES). *Biotechnol Bioeng* 114:961–969
- Evans PN, Parks DH, Chadwick GL, Robbins SJ, Orphan VJ, Golding SD, Tyson GW (2015) Methane metabolism in the archaeal phylum *Bathyarchaeota* revealed by genome-centric metagenomics. *Science* 350:434–438
- Fantozzi F, Buratti C (2009) Biogas production from different substrates in an experimental Continuously Stirred Tank Reactor anaerobic digester. *Bioresour Technol* 100:5783–5789
- Feng Y, Zhang Y, Chen S, Quan X (2015) Enhanced production of methane from waste activated sludge by the combination of high-solid anaerobic digestion and microbial electrolysis cell with iron-graphite electrode. *Chem Eng J* 259:787–794
- Ferrari A, Brusa T, Rutili A, Canzi E, Biavati B (1994) Isolation and characterization of *Methanobrevibacter oralis* sp. nov. *Curr Microbiol* 29:7–12
- Frimmer U, Widdel F (1989) Oxidation of ethanol by methanogenic bacteria. *Arch Microbiol* 152:479–483
- Fu Q, Kuramochi Y, Fukushima N, Maeda H, Sato K, Kobayashi H (2015) Bioelectrochemical analyses of the development of a thermophilic biocathode catalyzing electromethanogenesis. *Environ Sci Technol* 49:1225–1232
- Gajaraj S, Huang Y, Zheng P, Hu Z (2017) Methane production improvement and associated methanogenic assemblages in bioelectrochemically assisted anaerobic digestion. *Biochem Eng J* 117:105–112
- Garcia J-L, Patel BK, Ollivier B (2000) Taxonomic, phylogenetic, and ecological diversity of methanogenic Archaea. *Anaerobe* 6:205–226
- Gardner WL, Whitman WB (1999) Expression vectors for *Methanococcus maripaludis*: overexpression of acetohydroxyacid synthase and beta-galactosidase. *Genetics* 152:1439–1447
- Ge X, Yang L, Sheets JP, Yu Z, Li Y (2014) Biological conversion of methane to liquid fuels: status and opportunities. *Biotechnol Adv* 32:1460–1475
- Geppert F, Liu D, van Eerten-Jansen M, Weidner E, Buisman C, ter Heijne A (2016) Bioelectrochemical power-to-gas: state of the art and future perspectives. *Trends Biotechnol* 34:879–894
- Gernhardt P, Possot O, Foglino M, Sibold L, Klein A (1990) Construction of an integration vector for use in the archaeobacterium *Methanococcus voltae* and expression of a eubacterial resistance gene. *Mol Gen Genet* 221:273–279
- Gieg LM, Duncan KE, Sulflita JM (2008) Bioenergy production via microbial conversion of residual oil to natural gas. *Appl Environ Microbiol* 74:3022–3029
- Gorby YA, Yanina S, McLean JS, Rosso KM, Moyses D, Dohnalkova A, Beveridge TJ, Chang IS, Kim BH, Kim KS, Cullley DE, Reed SB, Romine MF, Saffarini DA, Hill EA, Shi L, Elias DA, Kennedy DW, Pinchuk G, Watanabe K, Ishii S, Logan B, Nealon KH, Fredrickson JK (2006) Electrically conductive bacterial nanowires produced by *Shewanella oneidensis* strain MR-1 and other microorganisms. *Proc Natl Acad Sci* 103:11358–11363
- Gottschalk G, Thauer RK (2001) The Na(+)-translocating methyltransferase complex from methanogenic archaea. *Biochim Biophys Acta* 1505:28–36
- Goyal N, Zhou Z, Karimi IA (2016) Metabolic processes of *Methanococcus maripaludis* and potential applications. *Microb Cell Fact* 15:107
- Grüber G, Manimekalai MSS, Mayer F, Müller V (2014) ATP synthases from archaea: the beauty of a molecular motor. *Biochim Biophys Acta Bioenerg* 1837:940–952
- Guo X, Liu J, Xiao B (2013) Bioelectrochemical enhancement of hydrogen and methane production from the anaerobic digestion of sewage sludge in single-chamber membrane-free microbial electrolysis cells. *Int J Hydrogen Energy* 38:1342–1347
- Guo Z, Thangavel S, Wang L, He Z, Cai W, Wang A, Liu W (2017) Efficient methane production from beer wastewater in a membraneless microbial

- electrolysis cell with a stacked cathode: the effect of the cathode/anode ratio on bioenergy recovery. *Energy Fuels* 31:615–620
- Guss AM, Rother M, Zhang JK, Kulkarni G, Metcalf WW (2008) New methods for tightly regulated gene expression and highly efficient chromosomal integration of cloned genes for *Methanosarcina* species. *Archaea* 2:193–203
- Hara M, Onaka Y, Kobayashi H, Fu Q, Kawaguchi H, Vilcaez J, Sato K (2013) Mechanism of electromethanogenic reduction of CO₂ by a thermophilic methanogen. *Energy Procedia* 37:7021–7028
- Harris JE, Pinn PA (1985) Bacitracin-resistant mutants of a mesophilic *Methanobacterium* species. *Arch Microbiol* 143:151–153
- Hohn MJ, Palioura S, Su D, Yuan J, Söll D (2011) Genetic analysis of selenocysteine biosynthesis in the archaeon *Methanococcus maripaludis*. *Mol Microbiol* 81:249–258
- Holm-Nielsen JB, Al Seadi T, Oleskowicz-Popiel P (2009) The future of anaerobic digestion and biogas utilization. *Bioresour Technol* 100:5478–5484
- Holtmann D, Hannappel A, Schrader J (2014) Microbial electrosynthesis. *Encyclopedia of applied electrochemistry*. Springer, New York, pp 1268–1275
- Hook SE, Wright A-DG, McBride BW (2010) Methanogens: methane producers of the rumen and mitigation strategies. *Archaea* 2010:1–11
- Hummel H, Böck A (1985) Mutations in *Methanobacterium formicicum* conferring resistance to anti-80S ribosome-targeted antibiotics. *MGG Mol Gen Genet* 198:529–533
- Iino T, Tamaki H, Tamazawa S, Ueno Y, Ohkuma M, Suzuki K-I, Igarashi Y, Haruta S (2013) Candidatus *Methanogranum caenicola*: a novel methanogen from the anaerobic digested sludge, and proposal of *Methanomassiliicoccaceae* fam. nov. and *Methanomassiliicoccales* ord. nov., for a methanogenic lineage of the class *Thermoplasmata*. *Microbes Environ* 28:244–250
- Jiang Y, Su M, Li D (2014) Removal of sulfide and production of methane from carbon dioxide in microbial fuel cells–microbial electrolysis cell (MFCs–MEC) coupled system. *Appl Biochem Biotechnol* 172:2720–2731
- Jones WJ, Leigh JA, Mayer F, Woese CR, Wolfe RS (1983) *Methanococcus jannaschii* sp. nov., an extremely thermophilic methanogen from a submarine hydrothermal vent. *Arch Microbiol* 136:254–261
- Kadier A, Simayi Y, Abdeshahian P, Azman NF, Chandrasekhar K, Kalil MS (2016) A comprehensive review of microbial electrolysis cells (MEC) reactor designs and configurations for sustainable hydrogen gas production. *Alexandria Eng J* 55:427–443
- Kamagata Y, Kawasaki H, Oyaizu H, Nakamura K, Mikami E, Endo G, Koga Y, Yamasato K (1992) Characterization of three thermophilic strains of *Methanotherix* (“*Methanoseta*”) *thermophila* sp. nov. and rejection of *Methanotherix* (“*Methanoseta*”) *thermoacetophila*. *Int J Syst Bacteriol* 42:463–468
- Kaparaju P, Ellegaard L, Angelidaki I (2009) Optimisation of biogas production from manure through serial digestion: lab-scale and pilot-scale studies. *Bioresour Technol* 100:701–709
- Karakashev D, Batstone DJ, Angelidaki I (2005) Influence of environmental conditions on methanogenic compositions in anaerobic biogas reactors. *Appl Environ Microbiol* 71:331–338
- Kern T, Linge M, Rother M (2015) *Methanobacterium aggregans* sp. nov., a hydrogenotrophic methanogenic archaeon isolated from an anaerobic digester. *Int J Syst Evol Microbiol* 65:1975–1980
- Kern T, Fischer Martin A, Deppenmeier U, Schmitz RA, Rother M (2016a) *Methanosarcina flavescens* sp. nov., a methanogenic archaeon isolated from a full-scale anaerobic digester. *Int J Syst Evol Microbiol* 66:1533–1538
- Kern T, Theiss J, Röske K, Rother M (2016b) Assessment of hydrogen metabolism in commercial anaerobic digesters. *Appl Microbiol Biotechnol* 100:4699–4710
- Kobayashi H, Saito N, Fu Q, Kawaguchi H, Vilcaez J, Wakayama T, Maeda H, Sato K (2013) Bio-electrochemical property and phylogenetic diversity of microbial communities associated with bioelectrodes of an electromethanogenic reactor. *J Biosci Bioeng* 116:114–117
- Koch C, Harnisch F (2016) What is the essence of microbial electroactivity? *Front Microbiol* 7:1890
- Koch C, Kuchenbuch A, Kretzschmar J, Wedwitschka H, Liebetau J, Müller S, Harnisch F (2015) Coupling electric energy and biogas production in anaerobic digesters - impacts on the microbiome. *RSC Adv* 5:31329–31340
- Koo T, Shin SG, Lee J, Han G, Kim W, Cho K, Hwang S (2017) Identifying methanogen community structures and their correlations with performance parameters in four full-scale anaerobic sludge digesters. *Bioresour Technol* 228:368–373
- Krieg T, Sydow A, Schröder U, Schrader J, Holtmann D (2014) Reactor concepts for bioelectrochemical syntheses and energy conversion. *Trends Biotechnol* 32:645–655
- Kurr M, Huber R, König H, Jannasch HW, Fricke H, Trincone A, Kristjansson JK, Stetter KO (1991) *Methanopyrus kandleri*, gen. and sp. nov. represents a novel group of hyperthermophilic methanogens, growing at 110 °C. *Arch Microbiol* 156:239–247
- Lay J-J, Li Y-Y, Noike T (1997) Influences of pH and moisture content on the methane production in high-solids sludge digestion. *Water Res* 31:1518–1524
- Leadbetter JR, Breznak JA (1996) Physiological ecology of *Methanobrevibacter cuticularis* sp. nov. and *Methanobrevibacter curvatus* sp. nov., isolated from the hindgut of the termite *Reticulitermes flavipes*. *Appl Environ Microbiol* 62:3620–3631
- Lebuhn M, Liu F, Heuwinkel H, Gronauer A (2008) Biogas production from mono-digestion of maize silage-long-term process stability and requirements. *Water Sci Technol* 58:1645–1651
- Lessner DJ, Lhu L, Wahal CS, Ferry JG (2010) An engineered methanogenic pathway derived from the domains *Bacteria* and *Archaea*. *mBio* 1(5):e00243–10.
- Li Y, Park SY, Zhu J (2011) Solid-state anaerobic digestion for methane production from organic waste. *Renew Sustain Energy Rev* 15:821–826
- Lie TJ, Leigh JA (2003) A novel repressor of *nif* and *glnA* expression in the methanogenic archaeon *Methanococcus maripaludis*. *Mol Microbiol* 47:235–246
- Liu Y, Whitman WB (2008) Metabolic, phylogenetic, and ecological diversity of the methanogenic archaea. *Ann NY Acad Sci* 1125:171–189
- Liu G, Zhang R, El-Mashad HM, Dong R (2009) Effect of feed to inoculum ratios on biogas yields of food and green wastes. *Bioresour Technol* 100:5103–5108
- Liu D, Zhang L, Chen S, Buisman C, ter Heijne A (2016) Bioelectrochemical enhancement of methane production in low temperature anaerobic digestion at 10 °C. *Water Res* 99:281–287
- Lohner ST, Deutzmann JS, Logan BE, Leigh J, Spormann AM (2014) Hydrogenase-independent uptake and metabolism of electrons by the archaeon *Methanococcus maripaludis*. *ISME J* 8:1673–1681
- Lovley DR (2012) Electromicrobiology. *Annu Rev Microbiol* 66:391–409
- Lucas R, Kuchenbuch A, Fetzer I, Harms H, Kleinstüber S (2015) Long-term monitoring reveals stable and remarkably similar microbial communities in parallel full-scale biogas reactors digesting energy crops. *FEMS Microbiol Ecol* 91:3
- Luo G, Angelidaki I (2012) Integrated biogas upgrading and hydrogen utilization in an anaerobic reactor containing enriched hydrogenotrophic methanogenic culture. *Biotechnol Bioeng* 109:2729–2736
- Luo G, Johansson S, Boe K, Xie L, Zhou Q, Angelidaki I (2012) Simultaneous hydrogen utilization and in situ biogas upgrading in an anaerobic reactor. *Biotechnol Bioeng* 109:1088–1094
- Lupa B, Hendrickson EL, Leigh JA, Whitman WB (2008) Formate-dependent H₂ production by the mesophilic methanogen *Methanococcus maripaludis*. *Appl Environ Microbiol* 74:6584–6590
- Lyu Z, Jain R, Smith P, Fetchko T, Yan Y, Whitman WB (2016) Engineering the autotroph *Methanococcus maripaludis* for geraniol production. *ACS Synth Biol* 5:577–581
- Mah RA (1980) Isolation and characterization of *Methanococcus mazei*. *Curr Microbiol* 3:321–325
- Martin I, Pidou M, Soares A, Judd S, Jefferson B (2011) Modelling the energy demands of aerobic and anaerobic membrane bioreactors for wastewater treatment. *Environ Technol* 32:921–932
- Mathrani IM, Boone DR, Mah RA, Fox GE, Lau PP (1988) *Methanohalophilus zhilinae* sp. nov., an alkaliphilic, halophilic, methylotrophic methanogen. *Int J Syst Bacteriol* 38:139–142
- Mayer F, Müller V (2014) Adaptations of anaerobic archaea to life under extreme energy limitation. *FEMS Microbiol Rev* 38:449–472
- Mayumi D, Mochimaru H, Tamaki H, Yamamoto T, Yoshioka H, Suzuki Y, Kamagata Y, Sakata S (2016) Methane production from coal by a single methanogen. *Science* 354:222–225

- McAnulty MJ, Poosarla VG, Li J, Soo WVC, Zhu F, Wood TK (2017) Metabolic engineering of *Methanosarcina acetivorans* for lactate production from methane. *Biotechnol Bioeng* 114:852–861
- McInerney MJ, Struchtemeyer CG, Sieber J, Mouttaki H, Stams AJ, Schink B, Rohlin L, Gunsalus RP (2008) Physiology, ecology, phylogeny, and genomics of microorganisms capable of syntrophic metabolism. *Ann NY Acad Sci* 1125:58–72
- Metcalf WW, Zhang JK, Shi X, Wolfe RS (1996) Molecular, genetic, and biochemical characterization of the *serC* gene of *Methanosarcina barkeri* Fusaro. *J Bacteriol* 178:5797–5802
- Metcalf WW, Zhang JK, Apolinario E, Sowers KR, Wolfe RS (1997) A genetic system for Archaea of the genus *Methanosarcina*: liposome-mediated transformation and construction of shuttle vectors. *Proc Natl Acad Sci USA* 94:2626–2631
- Miller TL, Lin C (2002) Description of *Methanobrevibacter gottschalkii* sp. nov., *Methanobrevibacter thaueri* sp. nov., *Methanobrevibacter woesei* sp. nov. and *Methanobrevibacter wolinii* sp. nov. *Int J Syst Evol Microbiol* 52:819–822
- Miller TL, Wolin MJ, Conway de Macario E, Macario AJ (1982) Isolation of *Methanobrevibacter smithii* from human feces. *Appl Environ Microbiol* 43:227–232
- Mochimaru H, Tamaki H, Hanada S, Imachi H, Nakamura K, Sakata S, Kamagata Y (2009) *Methanobolus profundus* sp. nov., a methylophilic methanogen isolated from deep subsurface sediments in a natural gas field. *Int J Syst Evol Microbiol* 59:714–718
- Mondorf S, Deppenmeier U, Welte C (2012) A novel inducible protein production system and neomycin resistance as selection marker for *Methanosarcina mazei*. *Archaea* 2012:1–8
- Moore BC, Leigh JA (2005) Markerless mutagenesis in *Methanococcus maripaludis* demonstrates roles for alanine dehydrogenase, alanine racemase, and alanine permease. *J Bacteriol* 187:972–979
- Müller B, Allmansberger R, Klein A (1985) Termination of a transcription unit comprising highly expressed genes in the archaeobacterium *Methanococcus voltae*. *Nucleic Acids Research* 13(18):6439–6445
- Mussgnug JH, Klassen V, Schlüter A, Kruse O (2010) Microalgae as substrates for fermentative biogas production in a combined biorefinery concept. *J Biotechnol* 150:51–56
- Nayak DD, Metcalf WW (2017) Cas9-mediated genome editing in the methanogenic archaeon *Methanosarcina acetivorans*. *Proc Natl Acad Sci* 114:2976–2981
- Neumann P, Pesante S, Venegas M, Vidal G (2016) Developments in pre-treatment methods to improve anaerobic digestion of sewage sludge. *Rev Environ Sci Biotechnol* 15:173–211
- Niu Q, Kobayashi T, Takemura Y, Kubota K, Li Y-Y (2015) Evaluation of functional microbial community's difference in full-scale and lab-scale anaerobic digesters feeding with different organic solid waste: effects of substrate and operation factors. *Bioresour Technol* 193:110–118
- Nizami A-S, Murphy JD (2010) What type of digester configurations should be employed to produce biomethane from grass silage? *Renew Sustain Energy Rev* 14:1558–1568
- O'Brien JM, Wolkin RH, Moench TT, Morgan JB, Zeikus JG (1984) Association of hydrogen metabolism with unitrophic or mixotrophic growth of *Methanosarcina barkeri* on carbon monoxide. *J Bacteriol* 158:373–375
- Oelgeschläger E, Rother M (2009) In vivo role of three fused corrinoid/methyl transfer proteins in *Methanosarcina acetivorans*. *Mol Microbiol* 72:1260–1272
- Ollivier B (1997) *Methanoplanus petrolearius* sp. nov., a novel methanogenic bacterium from an oil-producing well. *FEMS Microbiol Lett* 147:51–56
- Oren A (2008) Microbial life at high salt concentrations: phylogenetic and metabolic diversity. *Saline Syst* 4:2
- Park SY, Liang Y (2016) Biogenic methane production from coal: a review on recent research and development on microbially enhanced coalbed methane (MECBM). *Fuel* 166:258–267
- Park DH, Laivenieks M, Guettler MV, Jain MK, Zeikus JG (1999) Microbial utilization of electrically reduced neutral red as the sole electron donor for growth and metabolite production. *Appl Environ Microbiol* 65:2912–2917
- Patil SA, Hägerhäll C, Gorton L (2012) Electron transfer mechanisms between microorganisms and electrodes in bioelectrochemical systems. *Bioanal Rev* 4:159–192
- Pflüger K, Baumann S, Gottschalk G, Lin W, Santos H, Müller V (2003) Lysine-2,3-aminomutase and beta-lysine acetyltransferase genes of methanogenic archaea are salt induced and are essential for the biosynthesis of N(epsilon)-acetyl-beta-lysine and growth at high salinity. *Appl Environ Microbiol* 69:6047–6055
- Pflüger K, Wieland H, Müller V (2005) Osmoadaptation in methanogenic archaea: recent insights from a genomic perspective. In: Gundel-Cimerman N, Oren A, Plemenitas A (eds) *Adaptation of life at high salt concentrations in Archaea, Bacteria, and Eukarya*. Springer, Dordrecht, pp 241–251
- Plöchl M, Heiermann M (2006) Biogas farming in central and northern Europe: a strategy for developing countries? *Agric Eng Int CIGR Ejournal* VIII:1–15
- Porat I, Whitman WB (2009) Tryptophan auxotrophs were obtained by random transposon insertions in the *Methanococcus maripaludis* tryptophan operon. *FEMS Microbiol Lett* 297:250–254
- Pritchett MA, Zhang JK, Metcalf WW (2004) Development of a markerless genetic exchange method for *Methanosarcina acetivorans* C2A and its use in construction of new genetic tools for methanogenic archaea. *Appl Environ Microbiol* 70:1425–1433
- Rajeshwari K, Balakrishnan M, Kansal A, Lata K, Kishore VV (2000) State-of-the-art of anaerobic digestion technology for industrial wastewater treatment. *Renew Sustain Energy Rev* 4:135–156
- Ravichandran M, Munisamy P, Varadharaju C, Natarajan S (2015) Methanogenic archaea: a multipotent biological candidate focusing toward realizing future global energy. *Int J Curr Microbiol Appl Sci* 4:785–793
- Ren T, Patel M, Blok K (2008) Steam cracking and methane to olefins: energy use, CO₂ emissions and production costs. *Energy* 33:817–833
- Ribot-Llobet E, Nam J-Y, Tokash JC, Guisasola A, Logan BE (2013) Assessment of four different cathode materials at different initial pHs using unbuffered catholytes in microbial electrolysis cells. *Int J Hydrog Energy* 38:2951–2956
- Rincón B, Borja R, González JM, Portillo MC, Sáiz-Jiménez C (2008) Influence of organic loading rate and hydraulic retention time on the performance, stability and microbial communities of one-stage anaerobic digestion of two-phase olive mill solid residue. *Biochem Eng J* 40:253–261
- Rittmann BE, Lee H, Zhang H, Alder J, Banaszak JE, Lopez R (2008) Full-scale application of focused-pulsed pre-treatment for improving biosolids digestion and conversion to methane. *Water Sci Technol* 58:1895
- Robichaux M, Howell M, Boopathy R (2003) Methanogenic activity in human periodontal pocket. *Curr Microbiol* 46:53–58
- Roeßler M, Pflüger K, Flach H, Lienard T, Gottschalk G, Müller V (2002) Identification of a salt-induced primary transporter for glycine betaine in the methanogen *Methanosarcina mazei* Gö1. *Appl Environ Microbiol* 68:2133–2139
- Romero-Güiza MS, Vila J, Mata-Alvarez J, Chimenos JM, Astals S (2016) The role of additives on anaerobic digestion: a review. *Renew Sustain Energy Rev* 58:1486–1499
- Rotaru A-E, Shrestha PM, Liu F, Markovaite B, Chen S, Nevin KP, Lovley DR (2014) Direct interspecies electron transfer between *Geobacter metallireducens* and *Methanosarcina barkeri*. *Appl Environ Microbiol* 80:4599–4605
- Rother M, Metcalf WW (2004) Anaerobic growth of *Methanosarcina acetivorans* C2A on carbon monoxide: an unusual way of life for a methanogenic archaeon. *Proc Natl Acad Sci* 101:16929–16934
- Rother M, Boccazzi P, Bose A, Pritchett MA, Metcalf WW (2005) Methanol-dependent gene expression demonstrates that methyl-coenzyme M reductase is essential in *Methanosarcina acetivorans* C2A and allows isolation of mutants with defects in regulation of the methanol utilization pathway. *J Bacteriol* 187:5552–5559
- Ryckebosch E, Drouillon M, Vervaeren H (2011) Techniques for transformation of biogas to biomethane. *Biomass Bioenergy* 35:1633–1645
- Sahlström L (2003) A review of survival of pathogenic bacteria in organic waste used in biogas plants. *Bioresour Technol* 87:161–166
- Sakai S, Imachi H, Hanada S, Ohashi A, Harada H, Kamagata Y (2008) *Methanocella paludicola* gen. nov., sp. nov., a methane-producing archaeon, the first isolate of the lineage “Rice Cluster I”, and proposal of the new archaeal order *Methanocellales* ord. nov. *Int J Syst Evol Microbiol* 58:929–936
- Sander R (2015) Compilation of Henry's law constants (version 4.0) for water as solvent. *Atmos Chem Phys* 15:4399–4981

- Sarmiento FB, Leigh JA, Whitman WB (2011) Genetic systems for hydrogenotrophic methanogens. pp 43–73
- Sasaki D, Sasaki K, Watanabe A, Morita M, Igarashi Y, Ohmura N (2013) Efficient production of methane from artificial garbage waste by a cylindrical bioelectrochemical reactor containing carbon fiber textiles. *AMB Express* 3:17
- Sattler C, Wolf S, Fersch J, Goetz S, Rother M (2013) Random mutagenesis identifies factors involved in formate-dependent growth of the methanogenic archaeon *Methanococcus maripaludis*. *Mol Genet Genomics* 288:413–424
- Saum R, Mingote A, Santos H, Müller V (2009) A novel limb in the osmoregulatory network of *Methanosarcina mazei* Gö1: N(epsilon)-acetyl-beta-lysine can be substituted by glutamate and alanine. *Environ Microbiol* 11:1056–1065
- Schink B, Ward JC, Zeikus JG (1981) Microbiology of wetwood: importance of pectin degradation and clostridium species in living trees. *Appl Environ Microbiol* 42:526–532
- Schiraldi C, Giuliano M, De Rosa M (2002) Perspectives on biotechnological applications of archaea. *Archaea* 1:75–86
- Schlegel K, Müller V (2013) Evolution of Na⁺ and H⁺ bioenergetics in methanogenic archaea. *Biochem Soc Trans* 41:421–426
- Schlegel K, Leone V, Faraldo-Gomez JD, Müller V (2012a) Promiscuous archaeal ATP synthase concurrently coupled to Na⁺ and H⁺ translocation. *Proc Natl Acad Sci* 109:947–952
- Schlegel K, Welte C, Deppenmeier U, Müller V (2012b) Electron transport during aceticlastic methanogenesis by *Methanosarcina acetivorans* involves a sodium-translocating Rnf complex. *FEBS J* 279:4444–4452
- Schnürer A, Nordberg A (2008) Ammonia, a selective agent for methane production by syntrophic acetate oxidation at mesophilic temperature. *Water Sci Technol* 57:735–740
- Schröder U, Harnisch F, Angenent LT (2015) Microbial electrochemistry and technology: terminology and classification. *Energy Environ Sci* 8:513–519
- Siegert M, Yates MD, Call DF, Zhu X, Spormann A, Logan BE (2014) Comparison of nonprecious metal cathode materials for methane production by electromethanogenesis. *ACS Sustain Chem Eng* 2:910–917
- Solli L, Hävelsrud OE, Horn SJ, Rike AG (2014) A metagenomic study of the microbial communities in four parallel biogas reactors. *Biotechnol Biofuels* 7:146
- Soo VWC, McNulty MJ, Tripathi A, Zhu F, Zhang L, Hatzakis E, Smith PB, Agrawal S, Nazem-Bokaei H, Gopalakrishnan S, Salis HM, Ferry JG, Maranas CD, Patterson AD, Wood TK (2016) Reversing methanogenesis to capture methane for liquid biofuel precursors. *Microb Cell Fact* 15:11
- Sorokin DY, Abbas B, Merkel AY, Rijpstra WIC, Damste JSS, Sukhacheva MV, van Loosdrecht MCM (2015) *Methanosalsum natronophilum* sp. nov., and *Methanocalculus alkaliphilus* sp. nov., haloalkaliphilic methanogens from hypersaline soda lakes. *Int J Syst Evol Microbiol* 65:3739–3745
- Sorokin DY, Makarova KS, Abbas B, Ferrer M, Golyshin PN, Galinski EA, Ciordia S, Mena MC, Merkel AY, Wolf YI, van Loosdrecht MCM, Koonin EV (2017) Discovery of extremely halophilic, methyl-reducing euryarchaea provides insights into the evolutionary origin of methanogenesis. *Nat Microbiol* 2:17081
- Sosnowski P, Wieczorek A, Ledakowicz S (2003) Anaerobic co-digestion of sewage sludge and organic fraction of municipal solid wastes. *Adv Environ Res* 7:609–616
- Stadtman TC, Barker HA (1951) Studies on the methane fermentation. X. A new formate-decomposing bacterium, *Methanococcus vannielii*. *J Bacteriol* 62:269–280
- Stolze Y, Zakrzewski M, Maus I, Eikmeyer F, Jaenicke S, Rottmann N, Siebner C, Pühler A, Schlüter A (2015) Comparative metagenomics of biogas-producing microbial communities from production-scale biogas plants operating under wet or dry fermentation conditions. *Biotechnol Biofuels* 8:14
- Strong PJ, Xie S, Clarke WP (2015) Methane as a resource: can the methanotrophs add value? *Environ Sci Technol* 49:4001–4018
- Sun J, Klein A (2004) A lysR-type regulator is involved in the negative regulation of genes encoding selenium-free hydrogenases in the archaeon *Methanococcus voltae*. *Mol Microbiol* 52:563–571
- Sundberg C, Al-Soud WA, Larsson M, Alm E, Yekta SS, Svensson BH, Sorensen SJ, Karlsson A (2013) 454 pyrosequencing analyses of bacterial and archaeal richness in 21 full-scale biogas digesters. *FEMS Microbiol Ecol* 85:612–626
- Sydow A, Krieg T, Mayer F, Schrader J, Holtmann D (2014) Electroactive bacteria - molecular mechanisms and genetic tools. *Appl Microbiol Biotechnol* 98:8481–8495
- Tabatabaei M, Rahim RA, Abdullah N, Wright AG, Shirai Y, Sakai K, Sulaiman A, Hassan MA (2010) Importance of the methanogenic archaea populations in anaerobic wastewater treatments. *Process Biochem* 45:1214–1225
- Thauer RK, Kaster A-K, Seedorf H, Buckel W, Hedderich R (2008) Methanogenic archaea: ecologically relevant differences in energy conservation. *Nat Rev Microbiol* 6:579–591
- Timmers PHA, Welte CU, Koehorst JJ, Plugge CM, Jetten MSM, Stams AJM (2017) Reverse methanogenesis and respiration in methanotrophic archaea. *Archaea* 2017:1–22
- Tumbula DL, Bowen TL, Whitman WB (1997) Characterization of pURB500 from the archaeon *Methanococcus maripaludis* and construction of a shuttle vector. *J Bacteriol* 179:2976–2986
- Valentine DL, Blanton DC, Reeburgh WS (2000) Hydrogen production by methanogens under low-hydrogen conditions. *Arch Microbiol* 174:415–421
- Van Lier JB, Mahmoud N, Zeeman G (2008) Anaerobic wastewater treatment. In: Henze M, van Loosdrecht MCM, Ekama GA, Brdjanovic D (eds) *Anaerobic wastewater treatment: principles modelling and design*. IWA Publishing, London, pp 401–442
- Vanegas C, Bartlett J (2013) Anaerobic digestion of laminaria digitata: the effect of temperature on biogas production and composition. *Waste Biomass Valorization* 4:509–515
- Vanwonterghem I, Evans PN, Parks DH, Jensen PD, Woodcroft BJ, Hugenholtz P, Tyson GW (2016) Methylophilic methanogenesis discovered in the archaeal phylum Verstraetearchaeota. *Nat Microbiol* 1:16170
- Vavilin VA, Fernandez B, Palatsi J, Flotats X (2008) Hydrolysis kinetics in anaerobic degradation of particulate organic material: an overview. *Waste Manag* 28:939–951
- Villano M, Aulenta F, Ciucci C, Ferri T, Giuliano A, Majone M (2010) Bioelectrochemical reduction of CO₂ to CH₄ via direct and indirect extracellular electron transfer by a hydrogenophilic methanogenic culture. *Bioresour Technol* 101:3085–3090
- Voegeli Y, Lohri C, Kassenga G, Baier U, Zurbrugg C (2009) Technical and biological performance of the Arti compact biogas plant for kitchen waste - case study from Tanzania. In: *Proceedings Sardinia, 2009, twelfth international waste management and landfill symposium*. CISA Publisher, Italy, S. Margherita die Pula, Cagliari
- Voordouw G (2011) Production-related petroleum microbiology: progress and prospects. *Curr Opin Biotechnol* 22:401–405
- Wagner T, Ermler U, Shima S (2016) The methanogenic CO₂ reducing-and-fixing enzyme is bifunctional and contains 46 [4Fe–4S] clusters. *Science* 354:114–117
- Wang P, Wang H, Qiu Y, Ren L, Jiang B (2017) Microbial characteristics in anaerobic digestion process of food waste for methane production - a review. *Bioresour Technol* 17:31056–31058
- Ward AJ, Hobbs PJ, Holliman PJ, Jones DL (2008) Optimisation of the anaerobic digestion of agricultural resources. *Bioresour Technol* 99:7928–7940
- Weiland P (2010) Biogas production: current state and perspectives. *Appl Microbiol Biotechnol* 85:849–860
- Welander PV, Metcalf WW (2005) Loss of the *mtr* operon in *Methanosarcina* blocks growth on methanol, but not methanogenesis, and reveals an unknown methanogenic pathway. *PNAS* 102:10664–10669
- Welander PV, Metcalf WW (2008) Mutagenesis of the C1 oxidation pathway in *Methanosarcina barkeri*: new insights into the Mtr/Mer bypass pathway. *J Bacteriol* 190:1928–1936
- Welte C, Deppenmeier U (2014) Bioenergetics and anaerobic respiratory chains of aceticlastic methanogens. *Biochim Biophys Acta Bioenerg* 1837:1130–1147
- Westerholm M, Moestedt J, Schnürer A (2016) Biogas production through syntrophic acetate oxidation and deliberate operating strategies for improved digester performance. *Appl Energ* 179:124–135
- Widdel F (1986) Growth of methanogenic bacteria in pure culture with 2-propanol and other alcohols as hydrogen donors. *Appl Environ Microbiol* 51:1056–1062

- Wonglertarak W, Wichitsathian B (2014) Alkaline pretreatment of waste activated sludge in anaerobic digestion. *J Clean Energy Technol* 2:118–121
- Worrell VE, Nagle DP, McCarthy D, Eisenbraun A (1988) Genetic transformation system in the archaeobacterium *Methanobacterium thermoautotrophicum* Marburg. *J Bacteriol* 170:653–656
- Yadvika Santosh, Sreekrishnan TR, Kohli S, Rana V (2004) Enhancement of biogas production from solid substrates using different techniques - a review. *Bioresour Technol* 95:1–10
- Yu HW, Samani Z, Hanson A, Smith G (2002) Energy recovery from grass using two-phase anaerobic digestion. *Waste Manag* 22:1–5
- Zeikus JG, Bowen VG (1975) Fine structure of *Methanospirillum hungatii*. *J Bacteriol* 121:373–380
- Zeikus JG, Henning DL (1975) *Methanobacterium arbophilicum* sp. nov. An obligate anaerobe isolated from wetwood of living trees. *Antonie Van Leeuwenhoek* 41:543–552
- Zhang JK, Pritchett MA, Lampe DJ, Robertson HM, Metcalf WW (2000) In vivo transposon mutagenesis of the methanogenic archaeon *Methanosarcina acetivorans* C2A using a modified version of the insect mariner-family transposable element *Himar1*. *Proc Natl Acad Sci USA* 97:9665–9670
- Zhang G, Jiang N, Liu X, Dong X (2008) Methanogenesis from methanol at low temperatures by a novel psychrophilic Methanogen, "*Methanobus psychrophilus*" sp. nov., prevalent in Zoige Wetland of the Tibetan plateau. *Appl Environ Microbiol* 74:6114–6120
- Zhen G, Kobayashi T, Lu X, Xu K (2015) Understanding methane bioelectrosynthesis from carbon dioxide in a two-chamber microbial electrolysis cells (MECs) containing a carbon biocathode. *Bioresour Technol* 186:141–148
- Zhen G, Lu X, Kobayashi T, Kumar G, Xu K (2016) Promoted electromethanogenesis in a two-chamber microbial electrolysis cells (MECs) containing a hybrid biocathode covered with graphite felt (GF). *Chem Eng J* 284:1146–1155
- Zheng Y, Zhao J, Xu F, Li Y (2014) Pretreatment of lignocellulosic biomass for enhanced biogas production. *PECS* 42:35–53
- Zhong W, Zhang Z, Luo Y, Sun S, Qiao W, Xiao M (2011) Effect of biological pretreatments in enhancing corn straw biogas production. *Bioresour Technol* 102:11177–11182

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com
