



# Complete Genome Sequence of *Clostridium perfringens* LLY\_N11, a Necrotic Enteritis-Inducing Strain Isolated from a Healthy Chicken Intestine

Charles Li,<sup>a</sup> Xianghe Yan,<sup>b</sup> Hyun S. Lillehoj<sup>a</sup>

Animal Biosciences and Biotechnology Laboratory, Beltsville Agricultural Research Center, Agricultural Research Service, U.S. Department of Agriculture, Beltsville, Maryland, USA<sup>a</sup>; Environmental Microbial and Food Safety Laboratory, Beltsville Agricultural Research Center, Agricultural Research Service, U.S. Department of Agriculture, Beltsville, Maryland, USA<sup>b</sup>

**ABSTRACT** *Clostridium perfringens* strain LLY\_N11, a commensal bacterium, which previously induced necrotic enteritis in an experimental study, was isolated from the intestine of a young healthy chicken. Here, we present the complete genome sequence of this strain, which may provide a better understanding of the molecular mechanisms involved in necrotic enteritis pathogenesis.

*Clostridium perfringens* is a Gram-positive, spore-forming, and anaerobic bacterium responsible for a variety of diseases, such as gas gangrene, bacteremia, and food poisoning in humans and necrotic enteritis (NE) in food animals (1–4). Commensal *C. perfringens* strains are widely distributed in nature, especially in the soil and in the intestines of humans and animals (1, 5). Annual economic loss caused by NE is estimated to be over \$6 billion globally in the poultry industry (6). In our previous report, strain LLY\_N11, isolated from the intestine of a healthy chicken, was found to belong to *C. perfringens* and induced NE in an experimental model (7). In this study, the whole-genome sequencing of *C. perfringens* strain LLY\_N11 was conducted to characterize potential virulence factors involved in molecular pathogenesis.

A sample was prepared for genome sequencing by culturing strain CP15 anaerobically overnight at 37°C in brain heart infusion nutrient broth (Becton, Dickinson and Company, Sparks, MD, USA). The genomic DNA was then extracted from the cultures using a cetyltrimethylammonium bromide method (8). The complete genome sequence of *C. perfringens* LLY\_N11 was determined with the PacBio RS II platform (Pacific Biosciences, Menlo Park, CA, USA) at the Institute for Genomic Sciences, University of Maryland at Baltimore (Baltimore, MD, USA). The *de novo*-assembled whole-genome shotgun sequence was verified with the Illumina HiSeq 4000 platform (Illumina, Inc., San Diego, CA, USA) by Novogene, Inc. (Sacramento, CA, USA). Gene prediction and functional analysis were carried out using EDGE bioinformatics tools (9) and the NCBI Prokaryotic Genome Annotation Pipeline ([http://www.ncbi.nlm.nih.gov/genome/annotation\\_prok](http://www.ncbi.nlm.nih.gov/genome/annotation_prok)). Multiple-genome alignment showed that strain LLY\_N11 has an average nucleotide identity value of 99.1% to *C. perfringens* reference strain ATCC 13124 and a 99.9% identity score to *C. perfringens* strains Del1 and JP55 (7).

The complete genome sequence of strain LLY\_N11 contained 3,346,739 bp in one chromosome with a G+C content of 28.5%, 3,031 coding sequences (total), 30 rRNAs, 90 tRNAs, and 4 noncoding RNAs. Three plasmids, designated pLLY\_N11\_1, pLLY\_N11\_2, and pLLY\_N11\_3, comprised 13,363 bp, 14,754 bp, and 72,060 bp, respectively. The genome of strain LLY\_N11 was analyzed for the presence of antibiotic resistance genes

Received 28 September 2017 Accepted 2 October 2017 Published 2 November 2017

**Citation** Li C, Yan X, Lillehoj HS. 2017. Complete genome sequence of *Clostridium perfringens* LLY\_N11, a necrotic enteritis-inducing strain isolated from a healthy chicken intestine. *Genome Announc* 5:e01225-17. <https://doi.org/10.1128/genomeA.01225-17>.

This is a work of the U.S. Government and is not subject to copyright protection in the United States. Foreign copyrights may apply.

Address correspondence to Charles Li, [charles.li@ars.usda.gov](mailto:charles.li@ars.usda.gov), or Xianghe Yan, [xianghe.yan@ars.usda.gov](mailto:xianghe.yan@ars.usda.gov).

(10) and virulence factors (Virulence Factor Database, <http://www.mgc.ac.cn/VFs>). *C. perfringens* strain LLY\_N11 possessed *mepA*, *tet38*, *tetA(P)*, and *tetB(P)*, all of which are associated with tetracycline resistance; it also contained the antibiotic resistance gene *mprF* and the rifampin resistance gene *rpoB*. Computational analysis revealed that strain LLY\_N11 contained 20 toxin genes, including alpha-toxin, alpha-clostripain, enterotoxins (EntA, EntB, and EntD), hemolysin, kappa-toxin, mu-toxin, and beta2-toxin. Alpha-toxin (encoded by *cpa*) was the most toxic extracellular enzyme that hydrolyzed important constituents of eukaryotic cell membranes (11, 12), while beta2-toxin was found to be associated with intestinal disorders in chickens and other food animals (13, 14). Based on the toxin gene types (1), *C. perfringens* strain LLY\_N11 is a new strain of *C. perfringens* type A due to absence of beta, iota, and epsilon toxin.

**Accession number(s).** This whole-genome sequence has been deposited at GenBank under the accession numbers [CP023410](#) (chromosome), [CP023411](#) (plasmid pLLY\_N11\_1), [CP023412](#) (plasmid pLLY\_N11\_2), and [CP023413](#) (plasmid pLLY\_N11\_3).

## ACKNOWLEDGMENTS

This work was supported by funding from ARS/USDA.

The PacBio sequencing was completed using the genomic platform at the Institute for Genome Sciences, University of Maryland at Baltimore (Baltimore, MD, USA). The Illumina sequencing was performed by Novogene, Inc. (Sacramento, CA, USA). We also thank Ransom Baldwin for critically reading the manuscript.

## REFERENCES

- Li J, Adams V, Bannam TL, Miyamoto K, Garcia JP, Uzal FA, Rood JI, McClane BA. 2013. Toxin plasmids of *Clostridium perfringens*. *Microbiol Mol Biol Rev* 77:208–233. <https://doi.org/10.1128/MMBR.00062-12>.
- Genigeorgis CA, Riemann H. 1973. Food safety and food poisoning. *World Rev Nutr Diet* 16:363–397. <https://doi.org/10.1159/000393598>.
- Rogstad B, Ritland S, Lunde S, Hagen AG. 1993. *Clostridium perfringens* septicemia with massive hemolysis. *Infection* 21:54–56. <https://doi.org/10.1007/BF01739316>.
- Brook I. 2002. Meningitis and shunt infection caused by anaerobic bacteria in children. *Pediatr Neurol* 26:99–105. [https://doi.org/10.1016/S0887-8994\(01\)00330-7](https://doi.org/10.1016/S0887-8994(01)00330-7).
- Brynstad S, Granum PE. 2002. *Clostridium perfringens* and foodborne infections. *Int J Food Microbiol* 74:195–202. [https://doi.org/10.1016/S0168-1605\(01\)00680-8](https://doi.org/10.1016/S0168-1605(01)00680-8).
- Wade B, Keyburn A. 9 October 2015. The true cost of necrotic enteritis. *World Poultry*. <http://www.poultryworld.net/Meat/Articles/2015/10/The-true-cost-of-necrotic-enteritis-2699819W/>.
- Li C, Lillehoj HS, Gadde UD, Ritter D, Oh S. 2017. Characterization of *Clostridium perfringens* strains isolated from healthy and necrotic enteritis-afflicted broiler chickens. *Avian Dis* 61:178–185. <https://doi.org/10.1637/11507-093016-Reg.1>.
- Wilson K. 2001. Preparation of genomic DNA from bacteria. *Curr Protoc Mol Biol* Chapter 2:Unit 2.4.
- Li PE, Lo CC, Anderson JJ, Davenport KW, Bishop-Lilly KA, Xu Y, Ahmed S, Feng S, Mokashi VP, Chain PS. 2017. Enabling the democratization of the genomics revolution with a fully integrated Web-based bioinformatics platform. *Nucleic Acids Res* 45:67–80. <https://doi.org/10.1093/nar/gkw1027>.
- Turnidge J, Christiansen K. 2005. Antibiotic use and resistance—proving the obvious. *Lancet* 365:548–549.
- Awad MM, Bryant AE, Stevens DL, Rood JI. 1995. Virulence studies on chromosomal  $\alpha$ -toxin and  $\Theta$ -toxin mutants constructed by allelic exchange provide genetic evidence for the essential role of  $\alpha$ -toxin in *Clostridium perfringens*-mediated gas gangrene. *Mol Microbiol* 15:191–202. <https://doi.org/10.1111/j.1365-2958.1995.tb02234.x>.
- Rood JI. 1998. Virulence genes of *Clostridium perfringens*. *Annu Rev Microbiol* 52:333–360. <https://doi.org/10.1146/annurev.micro.52.1.333>.
- Gibert M, Jolivet-Reynaud C, Popoff MR, Jolivet-Renaud C. 1997. Beta2 toxin, a novel toxin produced by *Clostridium perfringens*. *Gene* 203:65–73. [https://doi.org/10.1016/S0378-1119\(97\)00493-9](https://doi.org/10.1016/S0378-1119(97)00493-9).
- França M, Barrios MA, Stabler L, Zavala G, Shivaprasad HL, Lee MD, Villegas AM, Uzal FA. 2016. Association of beta2-positive *Clostridium perfringens* type A with focal duodenal necrosis in egg-laying chickens in the United States. *Avian Dis* 60:43–49. <https://doi.org/10.1637/11263-081915-Reg.1>.