





The patency of graft and anastomoses in sequential and individual coronary artery bypass grafting: A meta-analysis

 Yongxing Li*,  Baotang Liu*,  Caifei Li**,  Yunchen Yu*,  Xiaobo Liu*,  Lei Li*,
 Zijun Li*,  Chenxi Duan*,  Sheng Luo***,  Wenming Hou*

Departments of *Cardiovascular Surgery, and **Operating Room, ***Public Health Management, Affiliated Hospital of Weifang Medical University; Weifang-China

ABSTRACT

Objective: To compare the patency of graft and anastomoses in sequential and individual coronary artery bypass grafting (CABG).

Methods: Our study used the Cochrane Library database, Excerpta Medica database, Web of Science, and PubMed. Studies comparing the outcomes of graft or anastomosis patency were assessed independently by two reviewers to identify the literature of satisfaction. We used Review Manager and STATA software for statistical analysis.

Results: Fifteen cohort studies were analyzed, including 10681 patients, 12957 grafts, and 4341 anastomoses, under sequential and individual CABG. Compared with the sequential group, the individual one is statistically significant in the graft patency [risk ratio (RR)=1.07, 95% confidence interval (CI) 1.01–1.13; p=0.02] and anastomosis patency (RR=1.06, 95% CI 1.01–1.12; p=0.005).

Conclusion: Our study suggested that the patency of the individual group, in terms of graft and anastomosis patency, is better than that of the sequential one. (*Anatol J Cardiol* 2020; 24: 235-43)

Keywords: coronary artery bypass grafting, sequential graft, meta-analysis

Introduction

Coronary heart disease (CHD) is currently the primary cause of death worldwide, with the percentage getting higher and higher. Coronary artery bypass grafting (CABG) is one of the common treatments for CHD patients. Flemma et al. (1) introduced the sequential grafting technique in 1971 for the first time. Despite different operations as sequential and individual veins have been applied, controversy still exists about the graft and anastomosis patency of these methods.

We aimed to conduct a meta-analysis of cohort studies for the comparison of the patency of graft and anastomoses in sequential and individual CABG.

Methods

Data sources and search strategy

Relevant trials that were included in this meta-analysis were searched in PubMed, Cochrane Library database, Excerpta Med-

ica database, and Web of Science, using the keywords “coronary artery bypass graft,” “sequential,” and “individual.” The search was limited to trials with humans and without publication date, language, and imposed publication status restrictions. The information on each study was chosen for the abstract. Two investigators reviewed the titles, abstracts, and studies independently to determine whether or not the inclusion criteria were met. The conflict between investigators was solved by consensus. The protocol of the meta-analysis was not registered.

Inclusion selection

The following criteria must be met by literature for it to be included: (1) patients must have undergone CABG; (2) the study must compare the patency of sequential and individual coronary artery bypass; and (3) graft and anastomosis patency denouements of the study were evaluated by ultrafast computed tomography (CT) or angiography. The Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) (2) was followed in our meta-analysis.

Address for correspondence: Wenming Hou, MD, Department of Cardiovascular Surgery, Affiliated Hospital of Weifang Medical University, 2428 Yuhe Road, 261031, Weifang-China

Phone: 1525 366 81 78 E-mail: houdoc815@126.com

Accepted Date: 20.08.2020 **Available Online Date:** 30.09.2020

©Copyright 2020 by Turkish Society of Cardiology - Available online at www.anatoljcardiol.com
DOI:10.14744/AnatolJCardiol.2020.10406



Outcome measures

Our study focused on the occurrence rate of graft and anastomosis patency and graft conduit with the artery or vein, on-pump or off-pump CABG performance, and assessment by ultrafast CT or angiography in patients during follow-up. Each segment was assessed as a separate graft in sequential grafts.

Data extraction

Relevant data were extracted independently and evaluated by two reviewers according to the pre-stipulated search strategy. The baseline demographic and quality characteristics from each study, author names, year of publication, number of patients, patient characteristics (age, percentages of women, comorbidities), follow-up time and rates, assessment method, operation, and conduit style, were extracted. The number of unobstructed graft and anastomoses in sequential and individual groups was recorded. Any disagreements were discussed between the two reviewers.

Quality assessment

The Newcastle-Ottawa Scale (NOS), a cohort study evaluation criteria suggested by the Cochrane Non-Randomized Studies Methods Group (3), was used to assess the methodological quality of this meta-analysis, which was evaluated using the following aspects: study selection, comparability between groups, and outcome determination.

Statistical analysis

The study effect of the destinations was measured using risk ratio (RR) as the pooled estimate, and the results were analyzed based on 95% confidence intervals (CIs). Two-sided p -values <0.05 were considered statistically significant. The chi-square test was conducted to examine the heterogeneity among the studies, and I^2 was also estimated. $I^2 <40\%$ might not be important, 30%–60% represent moderate heterogeneity, 50%–90% represent substantial heterogeneity, and $<70\%$ represent considerable heterogeneity (4). The fixed effects model was employed when I^2 was $<40\%$, while the random-effects model was employed when I^2 was $\geq 40\%$. The source of heterogeneity was explained using subgroup analyses. When the number of studies was >10 , we conducted Egger's test to evaluate the potential of publication bias. Review Manager (Nordic Cochrane Centre, Collaboration) and STATA (Stata Corp, College Station, Texas, USA) software were used for statistical analysis.

Results

Study selection

The search strategy brought out 701 literature, of which 15 (5-19) conformed to our inclusion criteria, and the selection process is shown in Figure 1. The study characteristics and NOS-dependent quality assessment are outlined in Table 1. All

manuscripts were middle- to high-quality cohort studies. A total of 10681 patients were included, and 8407 grafts and 2648 anastomoses in the sequential group and 4550 grafts and 1693 anastomoses in the individual group, respectively, were included.

Graft patency

Ten of the 15 studies reported information about graft patency, showing that the individual group had better patency than the sequential group (RR=1.07; 95% CI, 1.01–1.13) (Fig. 2). A considerable statistical heterogeneity ($I^2=93\%$, $p<0.00001$) was noted among the studies, so a random-effects model was used. However, there was no potential for significant publication bias after Egger's test ($p=0.27$).

Significant inconsistencies were noted in the follow-up rate, patency evaluation, graft selection (divided into arterial and venous), and operations. To exclude these possible confounding factors, four subgroup analyses were performed (i.e., the studies with a follow-up rate of $>70\%$, the studies of angiography to evaluate graft patency, the studies of grafts by the saphenous vein, and the studies of surgery method by off-pump).

Four studies had follow-up rates of $>70\%$, the results of which showed that the individual group was better than the sequential group (RR=1.09; 95% CI, 1.01–1.18; $p=0.02$) (Fig. 3a) and that statistical heterogeneity ($I^2=96\%$, $p<0.00001$) was noted among the studies, but the limited number of subgroups did not allow the assessment of the publication bias.

Seven studies reported on angiography to evaluate graft patency. No statistically significant difference between the two groups (RR=1.07; 95% CI, 0.98–1.18; $p=0.15$) was noted (Fig. 3b), and the results showed statistical heterogeneity ($I^2=92\%$, $p<0.00001$) among the studies, but the limited number of subgroups did not allow the assessment of publication bias.




Figure 1. PRISMA flow chart of date selection

Table 1. Study characteristics and quality assessment

| Study | Comparability | | | Follow-up | Rate method | Assessment | Patency measured on | Definition of occlusion | Conduit style | Surgical method | Quality assessment (NOS) | | | | |
|------------------------------|---------------|----------------------|---------|------------------------------------|---|------------|---------------------|-------------------------|---------------|-----------------|--------------------------|---------------|-------------|------------|---------------------|
| | Country | Age | Female | | | | | | | | Duration | Other factors | Rate method | Assessment | Patency measured on |
| Wendt et al. (5) 2010 | USA | 64±2/61±1 | 7/13 | Hypertension, diabetes mellitus | 1842±32/2070±33 days | >70 | CT | Anastomoses | Unclear | LITA | Unclear | 4 | 1 | 2 | 7 |
| Vural et al. (6) 2001 | Turkey | 49±8 | 11 | Atherosclerotic risk factor | 5.8±3 year | - | Angiography | Graft, anastomoses | Stenoses ≥50% | SVG | Unclear | 4 | 2 | 2 | 8 |
| Schwann et al. (7) 2009 | USA | 63 (30-90) | 17.7 | Dyslipidemia diabetes mellitus | 970±911 days | <70 | Angiography | Graft | Stenoses ≥75% | RA | On-pump | 3 | 2 | 2 | 7 |
| Park et al. (8) 2020 | South Korea | 64.6±8.7/63.6±9.2 | 25/23 | Body mass index, diabetes mellitus | 88.0 (46.3-119.2) months | >70 | CT | Graft | Stenoses ≥70% | SVG | Off-pump, on-pump | 4 | 2 | 2 | 8 |
| Oz et al. (9) 2006 | Turkey | 51.1±15.5/54.14±12.2 | 37 | Diabetes, smoking | 49.4±13.2 months | >70 | Angiography | Graft | Stenoses ≥50% | RA, SVG | Unclear | 4 | 2 | 2 | 8 |
| Meurala et al. (10) 1982 | Finland | 48±6.2 | 10 | Acute myocardial infarction | 26 (10-62) months | >70 | Angiography | Anastomoses | Unclear | SVG | Unclear | 4 | 2 | 2 | 8 |
| Kim et al. (11) 2011 | South Korea | 63.5±8.3 | 30.5 | Hypertension, diabetes mellitus | 14.8 (1-70.2) months | >70 | CT | Anastomoses | Stenoses ≥50% | SVG | Off-pump, on-pump | 4 | 2 | 3 | 9 |
| Gao et al. (12) 2010 | China | 63.6±10.3 | 11 | - | 26.4±23.6 month | - | CT | Graft, anastomoses | Stenoses ≥50% | SVG | Off-pump | 4 | 2 | 1 | 7 |
| Fukui et al. (13) 2012 | Japan | 67.2±10.4 | 19.5 | Previous myocardial infarction | 12.1 (2-21) months | <70 | Angiography | Graft, anastomoses | Stenoses ≥90% | RA | Off-pump | 4 | 2 | 2 | 8 |
| Farsak et al. (14) 2003 | Turkey | 55.2±9.3 | 13 | Atherosclerotic risk factor | 55.4±17.6 months | - | Angiography | Graft, anastomoses | Stenoses ≥50% | SVG | Unclear | 4 | 2 | 2 | 8 |
| Ji et al. (15) 2017 | China | 62.9±9.4/63.6±8.5 | 10/12.5 | Diabetes mellitus, smoking | 27.0±7.3 months, 27.2±7.2 months, unclear | >70 | CT | Graft | Stenoses ≥50% | LITA | Off-pump | 4 | 2 | 3 | 9 |
| Takazawa et al. (16) 2015 | Japan | 71±8 | 28.7 | Diabetes, hypertension | Unclear | <70 | Angiography | Graft | Unclear | SVG | Off-pump | 4 | 2 | 1 | 7 |
| Ohira et al. (17) 2016 | Japan | 65.7±9.3/65.8±8.4 | 16.6 | Body mass index, ejection fraction | Unclear | >70 | Angiography, CT | Anastomoses | Unclear | LITA | Off-pump | 4 | 2 | 2 | 8 |
| Brower et al. (18) 1981 | Netherlands | 52 (37-65) | 5 | - | 1 year | <70 | Unclear | Anastomoses | Unclear | - | Unclear | 3 | 1 | 2 | 6 |
| Christenson et al. (19) 1998 | Switzerland | 58.2±9.2 | 19 | Hypertension, smoking | 76 months | >70 | Angiography | Graft | Unclear | SVG | On-pump | 3 | 2 | 3 | 8 |

CT - computed tomography; LITA - left internal thoracic artery; RA - radial artery; SVG - saphenous vein graft; NOS - Newcastle-Ottawa scale




Figure 2. Forest plot of graft patency

M-H - Mantel-Haenszel; CI - confidence interval

Seven studies reported on grafts by a saphenous vein, the results of which showed that the individual group was better than the sequential group (RR=1.11; 95% CI, 1.03–1.21; $p=0.01$) (Fig. 3c) and that considerable statistical heterogeneity ($I^2=95%$, $p<0.00001$) was noted among the studies, but the limited number of subgroups did not allow the assessment of the publication bias.

Four studies with surgery methods by off-pump were noted, the results of which demonstrated no statistically significant difference among the two groups (RR=1.00; 95% CI, 0.94–1.05; $p=0.87$) (Fig. 3d) and substantial statistical heterogeneity ($I^2=72%$, $p<0.00001$) among the studies, but the limited number of subgroups did not allow the assessment of publication bias.

Anastomosis patency

Nine studies described the information about anastomosis patency, the results of which showed that the individual group also had better patency than the sequential group (RR=1.06; 95% CI, 1.02–1.11) (Fig. 4) and that considerable statistical heterogeneity ($I^2=93%$, $p<0.00001$) was noted among the studies, but the limited number of studies did not allow the assessment of the publication bias. Four subgroup analyses were also performed (i.e., the studies with a follow-up rate of >70%, the studies of angiography to evaluate graft patency, the studies of grafts by saphenous vein, the studies of surgery method by off-pump) to exclude these possible confounding factors.

Four studies had follow-up rates of >70%, the results of which demonstrated no statistically significant difference among the two groups (RR=1.06; 95% CI, 1.00–1.13; $p=0.04$) (Fig. 5a) and substantial statistical heterogeneity ($I^2=78%$, $p=0.003$) among the studies, but the limited number of subgroups did not allow the assessment of the publication bias.

Four studies with angiography evaluated graft patency, the results of which showed that the individual group was better than the sequential group (RR=1.11; 95% CI, 1.03–1.19; $p=0.004$) (Fig. 5b) and that substantial statistical heterogeneity ($I^2=60%$, $p=0.06$) was noted among the studies, but the limited number of subgroups did not allow the assessment of the publication bias.

Seven studies with grafts by a saphenous vein were noted, the results of which showed that the individual group was better than the sequential group (RR=1.10; 95% CI, 1.06–1.15; $p<0.00001$) (Fig. 5c) and that moderate heterogeneity ($I^2=47%$, $p=0.11$) was noted among the studies, but the limited number of subgroups did not allow the assessment of the publication bias.

Three studies with surgery methods by off-pump were noted, the results of which demonstrated no statistically significant difference among the two groups (RR=1.03; 95% CI, 1.00–1.07; $p=0.05$) (Fig. 5d) and no statistical heterogeneity ($I^2=38%$, $p=0.20$) among the studies.

Sensitivity analyses

By excluding individual studies one by one, we performed two sensitivity analyses on graft and anastomosis patency, respectively (Fig. 6a and 6b). No statistical significance was noted, suggesting that our results were stable and the comprehensive results were not influenced by this heterogeneity.

Discussion

Main finding

To our knowledge, this meta-analysis is the most comprehensive one, including 15 studies and 10681 patients in total, comparing the patency of graft and anastomoses in patients who underwent sequential and individual CABG, which indi-




Figure 3. Forest plot for subgroup analysis of (a) follow-up rates >70%; (b) angiography to evaluate graft patency; (c) grafts by saphenous vein; and (d) surgery method by off-pump
 M-H - Mantel-Haenszel; CI - confidence interval




Figure 4. Forest plot of anastomosis patency

M-H - Mantel-Haenszel; CI - confidence interval

cated that the individual group in graft and anastomoses has a higher patency than that of the sequential one.

Compared with prior studies

Our studies suggested that the individual group has better patency than the sequential group in graft and anastomoses, the result of which is not consistent with previous studies by Li et al. (20) in 2011 and Li and Liu (21) in 2019. The earliest article indicated that the patency in sequential grafts was greater than in the individual group (RR=0.67; 95% CI, 0.60–0.74) and the rate of patency in side-to-side anastomoses was significantly greater than that of end-to-side anastomoses. This difference is first due to the surgical level not being mature enough at that time and people preferring sequential anastomoses and, second, due to the high proportion of SVG in their articles, up to 75%. The article published in 2019 suggested no significant statistical differences between these groups on the patency of grafts (RR=0.96; 95% CI, 0.91–1.02) and anastomoses (RR=0.95; 95% CI, 0.91–1.00), which is different from our research results because we have included more subjects and a larger sample size. Secondly, it may be related to Park et al.'s (8) study, which reported that the patency rate of a single branch is higher than that of the sequential.

The type of graft may be associated with the long-term patency rate. As the gold standard for CABG grafts (22), the internal thoracic artery has reached international consensus with clinical benefits of improving survival and reducing cardiovascular events. Raza et al. (23) found the sequential bridges to have the same long-term patency rate as single bridges and to be higher than Y-type grafts in a study comparing the long-term patency rates of individual segments of different internal thoracic artery grafts. The radial artery, as an alternative to the recommended internal thoracic artery graft (22), has a better long-term patency rate than the great saphenous vein (24) but is not routinely used.

If a radial artery graft is used, it should be anastomosed to a highly stenotic (>90%) target vessel for maximum clinical benefit (25). Mehta et al.'s trial (26), a phase 3, randomized, double-blind, placebo-controlled, and multicenter trial, indicated that the vein graft occlusion rate was higher for the sequential group than for the individual one (adjusted odds ratio 1.24; 95% CI, 1.03–1.48). Park et al. (8), in a 10-year long-term follow-up, found that sequential bridges have a higher rate of venous graft patency than single branch bridges (HR 0.61; 95% CI, 0.45–0.82; $p < 0.001$), which are very safe and effective. The difference in the proportion of cardiopulmonary bypass used between the two groups may be associated with the difference in outcomes.

Different surgical methods may affect graft patency. ROOBY trial (27), a large prospective randomized controlled clinical, was followed up by angiography 1 year later, which showed a significantly lower rate of the graft patency in the off-pump coronary artery bypass graft group than that in the on-pump coronary artery bypass graft group (82.6% vs. 87.8%). However, both the CORONARY (28) and the GOPCABE (29) trials were followed up for 5 years, the results of which showed no significant difference in the rate of revascularization between both bypass graft groups. Zhang et al. (30) showed that the operation of on-pump CABG significantly reduced the risk of saphenous vein graft occlusion than off-pump CABG (RR=1.41; 95% CI, 1.24–1.60), which is not identical with the result of our subgroup analysis.

Heterogeneity

In our study, the heterogeneities of graft and anastomosis patency were considerably high, which may be due to the provision of occlusion or patency of some studies. Errors, causing heterogeneity between studies, are due to the conversion of this information on occlusion into patency data and differences in age, sex, ethnicity, disease severity, and primary disease. The




Figure 5. Forest plot for subgroup analysis of (a) follow-up rates >70%; (b) angiography to evaluate anastomosis patency; (c) grafts by saphenous vein; and (d) surgery method by off-pump
 M-H - Mantel-Haenszel; CI - confidence interval




Figure 6. Sensitivity analysis of (a) graft patency and (b) anastomosis patency

high heterogeneities may also be ascribed to differences in time background, the level of surgeons' expertise, economy levels, and regional and cultural differences. However, these were not analyzed in our study because the data was not enough. Our meta-analysis used a random-effects model to explain these heterogeneities. In research evaluating graft patency, four subgroup analyses were conducted to explain the high heterogeneity, but the results did not obtain the source of heterogeneity. So, Egger's test was performed, the results of which indicated no significant publication bias. We also implemented sensitivity analysis, which proved that our result is stable and this heterogeneity is not influenced by the merged result. In research evaluating anastomosis patency, four subgroup analyses were also conducted to explain the heterogeneity, which was reduced into two subgroups (i.e., the studies of grafts by saphenous vein, the studies of surgery method by off-pump). We also implemented sensitivity analysis, which testified that our result is stable and this heterogeneity is not influenced by the merged result.

Study limitations

However, our meta-analysis also has some inherent limitations. First, some studies only provided the rate of occlusion. Data differences may be noted after conversion to the patency

rate, which may have influenced the dependability of the final results. Second, the coronary artery bypass anastomoses are not uniform, since both side-to-side and end-to-side anastomoses are found in our meta-analysis, which may influence the stability of the final results. Third, in our study, national research in Asia accounts for the majority, which may have some bias and is not representative of the whole world. In addition, our study included standard cohort studies rather than randomized trials, lowering the level of evidence. Therefore, it is necessary to conduct a larger, multicenter, prospective, and randomized international trials to further confirm these results.

Conclusion

Despite the inherent limitations of this meta-analysis, our findings show that the patency of the individual graft was better than that of the sequential group and that the patency of individual anastomoses was better than that of the sequential one.

Conflict of interest: None declared.

Peer-review: Externally peer-reviewed.

Authorship contributions: Concept – Y. L., B.L.; Design – X.L., Z.L.; Supervision – X.L., C.D.; Fundings – W. H.; Materials – None; Data collection &/or processing – Y.L., C.L., Y.Y., L.L., S.L.; Analysis &/or interpretation – Y.L., Y.Y.; Literature search – Y.L., B.L.; Writing – Y.L.; Critical review – Y. L., B.L., W.H.

References

1. Flemma RJ, Johnson WD, Lepley D Jr. Triple aorto-coronary vein bypass as treatment for coronary insufficiency. *Arch Surg* 1971; 103: 82-3.
2. Walther S, Schuetz GM, Hamm B, Dewey M. Quality of reporting of systematic reviews and meta-analyses: PRISMA (Preferred Reporting Items for Systematic reviews and Meta-Analyses). *Rofu* 2011; 183: 1106-10.
3. Wells GA, Shea B, O'Connell, Peterson J, Welch V, Losos M, et al. The Newcastle-Ottawa Scale (NOS) for assessing the quality of non-randomised studies in meta-analysis. Ottawa (CA): Ottawa Hospital Research Institute; c2020 [cited 2020. May. 18]. Available from: URL; http://www.ohri.ca/programs/clinical_epidemiology/oxford.asp
4. Deeks JJ, Higgins JPT, Altman DG. (editors). Chapter 9: Analysing data and undertaking meta-analyses. In: Higgins JPT, Green S (editors). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 (updated March 2011). The Cochrane Collaboration, 2011. Available from: URL; <https://handbook-5-1.cochrane.org/>
5. Wendt D, Schmidt D, Wasserfuhr D, Osswald B, Thielmann M, Tossios P, et al. Comparison of sequential left internal thoracic artery grafting and separate left internal thoracic artery and venous grafting: A 5-year follow-up. *Herz* 2010; 35: 397-402.
6. Vural KM, Sener E, Taşdemir O. Long-term patency of sequential and individual saphenous vein coronary bypass grafts. *Eur J Cardiothorac Surg* 2001; 19: 140-4.

7. Schwann TA, Zacharias A, Riordan CJ, Durham SJ, Shah AS, Habib RH. Sequential radial artery grafts for multivessel coronary artery bypass graft surgery: 10-year survival and angiography results. *Ann Thorac Surg* 2009; 88: 31-9.
8. Park SJ, Kim HJ, Kim JB, Jung SH, Choo SJ, Lee JW, et al. Sequential Versus Individual Saphenous Vein Grafting During Coronary Arterial Bypass Surgery. *Ann Thorac Surg* 2020; 109: 1165-73.
9. Oz BS, Iyem H, Akay HT, Bolcal C, Yokusoglu M, Kuralay E, et al. Mid-term angiographic comparison of sequential and individual anastomosis techniques for diagonal artery. *J Cardiac Surg* 2006; 21: 471-4.
10. Meurala H, Valle M, Hekali P, Somer K, Frick MH, Harjola PT. Patency of sequential versus single vein grafts in coronary bypass surgery. *Thorac Cardiovasc Surg* 1982; 30: 147-51.
11. Kim HJ, Lee TY, Kim JB, Cho WC, Jung SH, Chung CH, et al. The impact of sequential versus single anastomoses on flow characteristics and mid-term patency of saphenous vein grafts in coronary bypass grafting. *J Thorac Cardiovasc Surg* 2011; 141: 750-4.
12. Gao C, Wang M, Wang G, Xiao C, Wu Y, Li B, et al. Patency of sequential and individual saphenous vein grafts after off-pump coronary artery bypass grafting. *J Card Surg* 2010; 25: 633-7.
13. Fukui T, Tabata M, Morita S, Takanashi S. Sequential free right internal thoracic artery grafting for multivessel coronary artery bypass grafting. *J Thorac Cardiovasc Surg* 2012; 144: 824-9.
14. Farsak B, Tokmakoglu H, Kandemir O, Günaydin S, Aydin H, Yorgancıoğlu C, et al. Angiographic assessment of sequential and individual coronary artery bypass grafting. *J Card Surg* 2003; 18: 524-9.
15. Ji Q, Shi Y, Xia L, Ma R, Shen J, Lai H, et al. Revascularization of Left Coronary System Using a Skeletonized Left Internal Mammary Artery- Sequential vs. Separate Grafting. *Circ J* 2017; 82: 102-9.
16. Takazawa A, Nakajima H, Iguchi A, Tabata M, Morita K, Koike H, et al. Impacts of intraoperative flow on graft patency of sequential and individual saphenous vein grafts. *Innovations (Phila)* 2015; 10: 85-9.
17. Ohira S, Doi K, Okawa K, Dohi M, Yamamoto T, Kawajiri H, et al. Safety and Efficacy of Sequential Left Internal Thoracic Artery Grafting to Left Circumflex Area. *Ann Thorac Surg* 2016; 102: 766-73.
18. Brower RW, van Eijk KF, Spek J, Bos E. Sequential versus conventional coronary artery bypass graft surgery in matched patient groups. *Thorac Cardiovasc Surg* 1981; 29: 158-62.
19. Christenson JT, Simonet F, Schmuziger M. Sequential vein bypass grafting: tactics and long-term results. *Cardiovasc Surg* 1998; 6: 389-97.
20. Li J, Liu Y, Zheng J, Bai T, Liu Y, Wang X, et al. The patency of sequential and individual vein coronary bypass grafts: a systematic review. *Ann Thorac Surg* 2011; 92: 1292-8.
21. Li Z, Liu L. Patency of Individual and Sequential Coronary Artery Bypass in Patients with Ischemic Heart Disease: A Meta-analysis. *Braz J Cardiovasc Surg* 2019; 34: 420-7.
22. Sousa-Uva M, Neumann FJ, Ahlsson A, Alfonso F, Banning AP, Benedetto U, et al.; ESC Scientific Document Group. 2018 ESC/EACTS Guidelines on myocardial revascularization. *Eur J Cardiothorac Surg* 2019; 55: 4-90.
23. Raza S, Blackstone EH, Bakaeen FG, Ravichandren K, Tappuni B, Ahmad MA, et al. Long-Term Patency of Individual Segments of Different Internal Thoracic Artery Graft Configurations. *Ann Thorac Surg* 2019; 107: 740-6.
24. Gaudino M, Benedetto U, Fremes S, Biondi-Zoccai G, Sedrakyan A, Puskas JD, et al.; RADIAL Investigators. Radial-Artery or Saphenous-Vein Grafts in Coronary-Artery Bypass Surgery. *N Engl J Med* 2018; 378: 2069-77.
25. Desai ND, Cohen EA, Naylor CD, Fremes SE; Radial Artery Patency Study Investigators. A randomized comparison of radial-artery and saphenous-vein coronary bypass grafts. *N Engl J Med* 2004; 351: 2302-9.
26. Mehta RH, Ferguson TB, Lopes RD, Hafley GE, Mack MJ, Kouchoukos NT, et al.; Project of Ex-vivo Vein Graft Engineering via Transfection (PREVENT) IV Investigators. Saphenous vein grafts with multiple versus single distal targets in patients undergoing coronary artery bypass surgery: one-year graft failure and five-year outcomes from the Project of Ex-Vivo Vein Graft Engineering via Transfection (PREVENT) IV trial. *Circulation* 2011; 124: 280-8.
27. Shroyer AL, Grover FL, Hattler B, Collins JF, McDonald GO, Kozora E, et al.; Veterans Affairs Randomized On/Off Bypass (ROOBY) Study Group. On-pump versus off-pump coronary-artery bypass surgery. *N Engl J Med* 2009; 361: 1827-37.
28. Lamy A, Devereaux PJ, Prabhakaran D, Taggart DP, Hu S, Straka Z, et al.; CORONARY Investigators. Five-Year Outcomes after Off-Pump or On-Pump Coronary-Artery Bypass Grafting. *N Engl J Med* 2016; 375: 2359-68.
29. Diegeler A, Börgermann J, Kappert U, Hilker M, Doenst T, Böning A, et al. Five-Year Outcome After Off-Pump or On-Pump Coronary Artery Bypass Grafting in Elderly Patients. *Circulation* 2019; 139: 1865-71.
30. Zhang B, Zhou J, Li H, Liu Z, Chen A, Zhao Q. Comparison of graft patency between off-pump and on-pump coronary artery bypass grafting: an updated meta-analysis. *Ann Thorac Surg* 2014; 97: 1335-41.