

Hazards and risks in oncology: radiation oncology

Abstract

Adverse effects and hazards which have their origin from radiation using conventional techniques like 3-D conformal radiotherapy and total radiation doses are well known. However little is known about the spectrum of especially late toxicity after radiation using new technologies like intensity modulated radiotherapy (IMRT) combined with novel target volume and dose concepts. Since IMRT allows for selective protection of the large salivary glands this technique improves the intermediate term quality of life and is the standard of care despite many details need further prospective evaluation. Combining cytotoxic drugs and radiotherapy yield improved survival in well-defined high risk patients. However morbidity and mortality of these protocols are high and deserve special expertise and supportive therapy. EGF-receptor antibodies have gained well defined indications, albeit specific toxicities in combination with irradiation deserve prospective studies and special attention.

Keywords: intensity modulated radiotherapy, chemoradiation, adverse events, epidermal growth factor receptor, head and neck cancer

Thomas G. Wendt¹

1 Department of Radiation Oncology, University Hospital Jena, Germany

Introduction

Radiation oncology is a mainstay in treating extracranial head and neck cancers. Its impact both as a single modality but also in combination with surgery and/or systemic therapy has been elucidated in numerous studies of different levels of evidence. A longstanding tradition over ten decades has led to an accumulated knowledge upon therapeutic and adverse effects of radiotherapy performed in conventional 2 dimensional and recently 3 dimensional radiation technique. Initially conventional xrays and telecobalt gamma rays were used and with the advent of electron accelerators in the second half of last century with photons and electrons. These developments led to a large body of knowledge upon the volumes to be treated and their total and single radiation doses to be given to obtain high local control rates and keep acute and chronic adverse effects at a reasonable level. During last three decades unconventional fractionation schedules have been studied in prospective randomized studies in order to optimize tumor control but also to clinical course of acute and chronic adverse effects at normal tissue inevitably exposed to ionizing radiation dose causing treatment associated acute and late morbidity.

Shortly after computer tomography and magnetic resonance tomography revolutionized diagnostics of tumors both became the basis of individualized anatomy based radiation dose distribution planning. Another step forward was computerization including digital processing of linac accelerators used in medicine. New algorithms were established for individual dose calculation including Monte Carlo calculation which allowed much more sophisticated dose distribution in tumor and surrounding normal tissues and organs. This led to the clinical implementation of in-

tensity modulated radiotherapy allowing different radiation doses within one target volume, e.g. higher doses to hypoxic subvolumes of a tumor (dose painting) [1]. Positron emission tomography has recently been integrated not only in the staging process [2] but also in the radiation therapy planning process. Different radiopharmaceuticals e.g. deoxyglucose, fmisonidazol were used to detect tumor subvolumes with different radiobiologically relevant characteristics e.g. proliferation or tumor hypoxia. These characteristics may be exploited in order to optimize an individual dose distribution plan and thus increase local tumor control probability (biologic radiation planning) [3]. However correlation of tumor size in excised specimen and metabolic seems loose [4]. New dose distribution concepts and fractionation schedules including integrated boost harbour which has an at least in part unknown toxicity profile which may impact negatively on chronic morbidity particularly in long term surviving patients.

Another new therapeutic avenue became clinically important with increasing knowledge upon interaction between cytotoxic drugs and more recently target drugs and ionizing radiation on a cellular basis (e.g. radiosensitization of cells). Much experience with regard to tumor control and adverse effects has been accumulated with simultaneous and sequential chemoradiation after extensive clinical use in numerous studies. However this is not true at the same degree for drugs used in target therapy.

Intensity modulated radiotherapy (IMRT)

Radiation induced xerostomia is a major source for chronic morbidity after primary or postoperative radiother-

apy. Conventional 3D conformal radiotherapy does not allow sparing of the major salivary glands except for few cases. Acute and chronic xerostomia causes a couple of adverse effects which impairs the quality of life in particular in long term survivors. Among others the most important are malnutrition and subsequent loss of weight, dental decay and chronic infection of the oral cavity. The technologically challenging intensity modulated radiotherapy allows for dedicated sparing of one or both parotid glands, which consequently results in lower probability of stimulated xerostomia provided radiation dose constraints, e.g. median organ dose of 26 Gy or less have been fulfilled [5], [6]. IMRT is associated with higher global quality of life compared to conventional 3D conformal radiotherapy [7], [8] which is particularly observed with increasing length of survival. The retrospective pattern of care data of the SEER showed equivalent tumor control and survival figures at 5-years obtained in 1,600 patients treated with both methods [9]. For the individual not being treated in a radiation oncology department not experienced with IMRT eventually means an adverse factor with respect of his further quality of life [10]. However with increasing experience with IMRT it became clear that the departure from conventional radiation techniques may carry risks hitherto not known. With 3D conformal radiotherapy in primaries other than larynx and hypopharynx the lower neck nodes have been irradiated by a direct ventral portal and by shielding the spine the lower part of the pharyngeal muscles have been shielded too. At the beginning of the IMRT era sparing of these muscles have not been recognized as important for preventing from dysphagia and aspiration and were initially not considered organs at risk. Not surprising that in a series of studies an increase of dysphagia after primary high dose chemoradiation using IMRT technique has been described [11]. Recent concepts focus on sparing of pharyngeal musculature, esophageal sphincter and supraglottic larynx. Total doses of less than 60 Gy should reduce the risk of dysphagia and aspiration [12], [13].

Chemoradiation and radioimmunotherapy

Advantages and risks of chemoradiation

Conventionally used high total radiation doses (applied with 3D conformal technique), e.g. 70 Gy to advanced primaries and/or neck nodes result in local control rates of around 30%. Since these results cannot be improved by increased total radiation doses, cytotoxic drugs have been integrated in protocols giving chemotherapy simultaneously or sequentially to irradiation or more recently both.

When 5-fluorouracil, mitomycin C and cis-platinum are given simultaneously to radiotherapy, not only the cytotoxic effect contributes to tumor cell kill but a radiosensitizing effect of drugs is exploited.

Simultaneous chemoradiation using 5-fluorouracil and cis-platinum or mitomycin C or carboplatinum augmented loco-regional control rates in locally advanced cancer often considered inoperable due to local extension compared to radical radiotherapy alone. Since not only 3–5 years tumor control rates but also overall survival has been increased simultaneous chemoradiation has become the standard of care [14], [15], [16], [17], [18], [19].

Induction chemotherapy preceding definitive radiotherapy results in an increased overall treatment duration of typically 12 to 15 weeks (in which tumor surviving stem cells may proliferate) and higher cumulated drug doses. The two metaanalyses however showed only a smaller increase of tumor control rates compared to simultaneous chemoradiation [Budach W, Pignon]. Noteworthy recent protocols including docetaxel have not been considered. When triplets consisting of taxane containing induction chemotherapy followed by radical radiotherapy and simultaneous taxane containing chemotherapy are given, increased median progression and overall survival span are reported compared to 5-fluorouracil based induction chemotherapy at the expense of more frequent hematologic $^{\circ}3/^{\circ}4$ toxicity [20], [21], [22]. In these studies increase progression free survival was essentially gained by better loco-regional tumor control. Although total drug amount delivered increased the risk of distant metastases has not been reduced [20], [21] pointing to distant metastases being the achilles' heel of new concepts. In a randomized comparison of immediate chemoradiation compared to induction chemotherapy followed by chemoradiation the latter resulted in a marginal increase in progression free survival but no increase in overall survival at 3 years [20]. Acute toxicity of induction chemotherapy can jeopardize chemoradiation [23].

Postoperative radiotherapy is a standard of care in patients with loco-regionally advanced tumors with distinct pathohistological risk factors. In two randomized studies of the RTOG and EORTC patients with risk factors R1-status, extracapsular spread and more than 3 involved lymph nodes benefitted from the additional simultaneously applied chemotherapy [24], [25]. However a subgroup analysis of the RTOG study showed increased survival only in patients with R1-status and extracapsular spread of involved lymph nodes [26]. Since chemoradiation exposes patients with a treatment associated lethality of 1–4%, it is crucial to restrict chemotherapy to those who are likely to benefit.

An organ sparing strategy is pursued particularly in patients who would need a total laryngectomy for cure of their laryngeal or hypopharyngeal cancer. Radiotherapy can contribute to preserve organ and function. Historical data showed a concordant responsiveness of cancers to chemo- and radiotherapy. Therefore in actual protocols a short induction chemotherapy select those cancers suitable for further chemoradiation and sort out those which need early salvage surgery. Also for smaller tumors simultaneous, sequential chemoradiation and radiotherapy alone was studied in a RTOG protocol. Like in other

primary tumor sites simultaneous chemoradiation was superior [27]. This protocol however carries a higher risk of late laryngeal toxicity compared to sequential therapy (15 vs. 10%).

Therapy associated morbidity of simultaneous chemoradiation

In loco-regionally advanced cancers simultaneous chemoradiation results in a 6–8% gain in 5-year survival compared to irradiation alone at the expense of increased acute morbidity. In these protocols the clinically leading toxicity is (often reversible) hematologic. When the treatment is intensified beside hematologic toxicity other not dominating adverse effects like nausea and emesis, alopecia, dysphagia, stomatitis, neuropathy and others also increase in frequency and severity. However they are less likely assessed. This “underreporting” leads to an unjustified favourable estimation of intensified protocols [28]. Not only acute morbidity is augmented but also subacute adverse effects are reported to be more frequent 3 months after end of chemoradiation compared to radiation alone [29]. Beside acute morbidity late adverse effects, e.g. 5 years after end of chemoradiation gain increasing interest. However data are more scarce compared to those of acute morbidity [24], [30]. Among chronic symptoms aspiration and dysphagia, both vital risks, are increasingly investigated.

Among all cytotoxic drugs cis-platinum, carboplatin, 5-fluorouracil and taxanes are the drugs most used in modern protocols. When radiotherapy and cis-platinum are combined, acute adverse effects increase. Given in postoperative protocols $\geq 3+$ (WHO) adverse effects increase from 34% to 77% compared to irradiation alone. Hence only 60% of all patients receive all 3 scheduled courses [25]. The toxicity profile change when cytotoxic drugs are administered concomitantly. Neuropathy and nephropathy are emerging risks which forces on intensified clinical monitoring but also intensified supportive care including optimized enteral feeding [31].

Cis-platinum is administered in anglo-saxonian protocols often on day 1 and 29 at a dosage of 100mg/kg body weight. In central europe the application of the same dose over 5 days (5 times 20 mg/kg body weight day 1–5, 29–33) has become standard in protocols. With the latter dosage the same tumor control rates are achieved with lower hematologic side effects [32].

In patients with preexisting diabetes mellitus more variation of their blood glucose profile have been observed when treated with cis-platinum for head and neck cancer. This may be due to an altered glucose metabolism. An intensive blood glucose monitoring is mandatory and existing medical antidiabetic therapy has to be adapted or initiated if no medication was applied before [33]. A cis-platinum associated hyperglycemic coma has been described [34].

The large salivary glands are highly radiosensitive. Due to their proximity to the target volume (tumor or tumorbed and safety margins) it is often impossible to spare them

from high radiation dose applied to the target volume. If one parotid gland is exposed to a median cumulated dose of 26 Gy or more irreversible subacute and chronic xerostomia must be anticipated [6]. Even lower median doses have been advocated [5]. If cis-platinum is given simultaneously to irradiation the tolerance of salivary glands and residual salivation further decreases [35]. If cis-platinum is used in a simultaneous chemoradiation protocol, a radiation technique should be applied which allows an optimal sparing of at least one parotid gland. In these cases the target median dose should probably be less than 26 Gy in order to reduce the risk of permanent xerostomia. This is best accomplished by sophisticated intensity modulated radiotherapy which should be regarded as a standard of care in chemoradiation protocols. Further improvement of parotid gland sparing is achieved by radiotherapy with protons [36], which however is not available for a larger population. Moreover chemoradiation protocols are associated with a treatment related mortality of 1–4% [14], [19], [26], [27], [37]. Particularly in polymorbid patients these hazards need to be counterbalanced critically against the possible gain in survival achieved by additional chemotherapy.

New concepts by antibodies and tyrosinkinase inhibitors

Considerable improvements have been achieved by radical and postoperative chemoradiation protocols. Despite intensification of treatment no further gain can be expected due to concomitant morbidity of most patients, which limits their tolerance and compliance. A particular issue are distant metastases, which are experienced more often as loco-regional control increases. As a consequence new drugs with more specific tumoricidal activity and less hematological and mucosal toxicity are investigated. Among the vast body of molecular strategies up to now only a few gained clinical significance or are studied in phase III trials: 1. monoclonal antibodies against the extracellular domain of the epidermal growth factor receptor (EGFR inhibitors), 2. the inhibition of intra- and extracellular domains with tyrosinkinase inhibitors with low molecular weight (“small molecules”) (EGFR tyrosinkinase inhibitors) and 3. antiangiogenic drugs against vascular endothelial growth factor (VEGFR inhibitors) blocking the neoangiogenesis in tumors [38]. Currently in Europe only Cetuximab for EGFR inhibition has been approved in combination with radiotherapy of squamous cell head and neck cancers.

Cetuximab

More than 90% of all squamous cell carcinoma of the head and neck show an overexpression of the epidermal growth factor receptor, which is associated with lower survival [39]. Recent data show versatile activities of cetuximab at different cellular levels and clinical response does not closely correlate with the degree of cellular

overexpression of EGFR. The exact mode of action of cetuximab remains unclear [40]. Despite that cetuximab serves as an alternative to cis-platinum in clinical chemoradiation protocols for head and neck cancers. In the TREMPLIN protocol [41], a larynx preservation protocol, cis-platinum was randomized against cetuximab. Equivalent 2-year survival data without laryngectomy were reported (cis-platinum: 79%, cetuximab: 71%, n.s.). However cetuximab was associated with less protocol violation and less late morbidity.

Low acute toxicity is a characteristic of treatment with cetuximab. However the toxicity profile is different for that of cis-platinum. One typical side effect is the higher risk of radiation dermatitis °3–4 including skin necrosis, which is confined to the irradiated skin and differs from acneiform rash. Various studies observed radiation dermatitis up to 23% of patients [42], [43], [44]. For management of cetuximab associated radiation dermatitis recently multidisciplinary guidelines have been published [45], [46]. When antibodies against EGFR cetuximab or panitumumab are infused acute bronchospasm is occasionally observed and therefore monitoring is mandatory. Case reports of bronchiolitis and lung fibrosis are published [47]. Cetuximab may rarely cause hypocalcemia and hypomagnesemia. When misdiagnosed the latter may proceed and cause convulsions and arrhythmia [48].

Cardiac toxicity of new drugs

A large hospital based survey admission diagnosis and prevalent comorbidity was analysed. Patients admitted with head and neck cancer had a higher incidence of cardiovascular (41% vs. 27%) and obstructive lung (12% vs. 5%) disease compared to patients admitted due to a non-cancer diagnosis. During a 12 months follow-up patients treated with chemotherapy for their cancer experienced 1.7 fold chronic anemia and 2.6 fold a second cancer compared to the non-cancer patients [49]. For cetuximab no cardiac or hematopoietic adverse effects are documented. Therefore cetuximab may be the preferred drug for in comorbid head and neck cancer patients [50].

Hypothyroidism

Due to proximity of the thyroid gland and the target volume of radiotherapy of most tumors a large part of the gland inevitably will be exposed to high radiation doses. 6 months after exposure of the gland thyroid stimulation hormone will rise [51]. The incidence of a subclinical or clinical hypothyroidism will increase over years and has been observed even 20 years after radiotherapy. However with increasing time span, other etiologic factors may contribute [52]. Hemithyroidectomy and radiotherapy were predictive for hypothyroidism in a multivariate analysis [53]. Typically 10 years after radiotherapy a subclinical or clinical hypothyroidism is diagnosed in 20–25% in all irradiated head and neck cancer patients.

Advanced radiation techniques like the IMRT are able to spare at least parts of the thyroid gland. However long term experience upon the incidence of hypothyroidism after IMRT is lacking. Irrespective of radiation technique and total dose used all patients need a yearly follow-up for their functional status. Radiogenic thyroid diseases are treated like spontaneously ones.

Notes

Competing interests

The author received honorarium for oral presentations from Serono-Merck.

References

- Ling CC, Humm J, Larson S, Amols H, Fuks Z, Leibel S, Koutcher JA. Towards multidimensional radiotherapy (MD-CRT): biological imaging and biological conformality. *Int J Radiat Oncol Biol Phys.* 2000 Jun;47(3):551-60. DOI: 10.1016/S0360-3016(00)00467-3
- Abramyuk A, Appold S, Zöphel K, Baumann M, Abolmaali N. Modification of staging and treatment of head and neck cancer by FDG-PET/CT prior to radiotherapy. *Strahlenther Onkol.* 2013 Mar;189(3):197-201. DOI: 10.1007/s00066-012-0283-0
- Madani I, Duthoy W, Derie C, De Gerssem W, Boterberg T, Saerens M, Jacobs F, Grégoire V, Lonneux M, Vakaet L, Vanderstraeten B, Bauters W, Bonte K, Thierens H, De Neve W. Positron emission tomography-guided, focal-dose escalation using intensity-modulated radiotherapy for head and neck cancer. *Int J Radiat Oncol Biol Phys.* 2007 May;68(1):126-35. DOI: 10.1016/j.ijrobp.2006.12.070
- Burri RJ, Rangaswamy B, Kostakoglu L, Hoch B, Genden EM, Som PM, Kao J. Correlation of positron emission tomography standard uptake value and pathologic specimen size in cancer of the head and neck. *Int J Radiat Oncol Biol Phys.* 2008 Jul;71(3):682-8. DOI: 10.1016/j.ijrobp.2007.10.055
- Amosson CM, Teh BS, Van TJ, Uy N, Huang E, Mai WY, Frolov A, Woo SY, Chiu JK, Carpenter LS, Lu HH, Grant WH 3rd, Butler EB. Dosimetric predictors of xerostomia for head-and-neck cancer patients treated with the smart (simultaneous modulated accelerated radiation therapy) boost technique. *Int J Radiat Oncol Biol Phys.* 2003 May;56(1):136-44. DOI: 10.1016/S0360-3016(03)00093-2
- Eisbruch A, Ten Haken RK, Kim HM, Marsh LH, Ship JA. Dose, volume, and function relationships in parotid salivary glands following conformal and intensity-modulated irradiation of head and neck cancer. *Int J Radiat Oncol Biol Phys.* 1999 Oct;45(3):577-87. DOI: 10.1016/S0360-3016(99)00247-3
- Chen AM, Farwell DG, Luu Q, Vazquez EG, Lau DH, Purdy JA. Intensity-modulated radiotherapy is associated with improved global quality of life among long-term survivors of head-and-neck cancer. *Int J Radiat Oncol Biol Phys.* 2012 Sep;84(1):170-5. DOI: 10.1016/j.ijrobp.2011.11.026
- Vergeer MR, Doornaert PA, Rietveld DH, Leemans CR, Slotman BJ, Langendijk JA. Intensity-modulated radiotherapy reduces radiation-induced morbidity and improves health-related quality of life: results of a nonrandomized prospective study using a standardized follow-up program. *Int J Radiat Oncol Biol Phys.* 2009 May;74(1):1-8. DOI: 10.1016/j.ijrobp.2008.07.059

9. Yu JB, Soulos PR, Sharma R, Makarov DV, Decker RH, Smith BD, Desai RA, Cramer LD, Gross CP. Patterns of care and outcomes associated with intensity-modulated radiation therapy versus conventional radiation therapy for older patients with head-and-neck cancer. *Int J Radiat Oncol Biol Phys.* 2012 May;83(1):e101-7. DOI: 10.1016/j.ijrobp.2011.11.067
10. Jabbari S, Kim HM, Feng M, Lin A, Tsien C, Elshaikh M, Terrel JE, Murdoch-Kinch C, Eisbruch A. Matched case-control study of quality of life and xerostomia after intensity-modulated radiotherapy or standard radiotherapy for head-and-neck cancer: initial report. *Int J Radiat Oncol Biol Phys.* 2005 Nov;63(3):725-31. DOI: 10.1016/j.ijrobp.2005.02.045
11. van der Molen L, van Rossum MA, Burkhead LM, Smeele LE, Hilgers FJ. Functional outcomes and rehabilitation strategies in patients treated with chemoradiotherapy for advanced head and neck cancer: a systematic review. *Eur Arch Otorhinolaryngol.* 2009 Jun;266(6):889-900. DOI: 10.1007/s00405-008-0817-3
12. Jensen K, Lambertsen K, Grau C. Late swallowing dysfunction and dysphagia after radiotherapy for pharynx cancer: frequency, intensity and correlation with dose and volume parameters. *Radiother Oncol.* 2007 Oct;85(1):74-82. DOI: 10.1016/j.radonc.2007.06.004
13. Li B, Li D, Lau DH, Farwell DG, Luu Q, Rocke DM, Newman K, Courquin J, Purdy JA, Chen AM. Clinical-dosimetric analysis of measures of dysphagia including gastrostomy-tube dependence among head and neck cancer patients treated definitively by intensity-modulated radiotherapy with concurrent chemotherapy. *Radiat Oncol.* 2009;4:52. DOI: 10.1186/1748-717X-4-52
14. Budach V, Stuschke M, Budach W, Baumann M, Geismar D, Grabenbauer G, Lammert I, Jahnke K, Stueben G, Herrmann T, Bamberg M, Wust P, Hinkelbein W, Wernecke KD. Hyperfractionated accelerated chemoradiation with concurrent fluorouracil-mitomycin is more effective than dose-escalated hyperfractionated accelerated radiation therapy alone in locally advanced head and neck cancer: final results of the radiotherapy cooperative clinical trials group of the German Cancer Society 95-06 Prospective Randomized Trial. *J Clin Oncol.* 2005 Feb;23(6):1125-35. DOI: 10.1200/JCO.2005.07.010
15. Budach W, Hehr T, Budach V, Belka C, Dietz K. A meta-analysis of hyperfractionated and accelerated radiotherapy and combined chemotherapy and radiotherapy regimens in unresected locally advanced squamous cell carcinoma of the head and neck. *BMC Cancer.* 2006;6:28. DOI: 10.1186/1471-2407-6-28
16. Calais G, Alfonsi M, Bardet E, Sire C, Germain T, Bergerot P, Rhein B, Tortochaux J, Oudinot P, Bertrand P. Randomized trial of radiation therapy versus concomitant chemotherapy and radiation therapy for advanced-stage oropharynx carcinoma. *J Natl Cancer Inst.* 1999 Dec;91(24):2081-6. DOI: 10.1093/jnci/91.24.2081
17. Pignon JP, le Maître A, Maillard E, Bourhis J; MACH-NC Collaborative Group. Meta-analysis of chemotherapy in head and neck cancer (MACH-NC): an update on 93 randomised trials and 17,346 patients. *Radiother Oncol.* 2009 Jul;92(1):4-14. DOI: 10.1016/j.radonc.2009.04.014
18. Semrau R, Mueller RP, Stuetzer H, Staar S, Schroeder U, Guntinas-Lichius O, Kocher M, Eich HT, Dietz A, Flentje M, Rudat V, Volling P, Schroeder M, Eckel HE. Efficacy of intensified hyperfractionated and accelerated radiotherapy and concurrent chemotherapy with carboplatin and 5-fluorouracil: updated results of a randomized multicentric trial in advanced head-and-neck cancer. *Int J Radiat Oncol Biol Phys.* 2006 Apr;64(5):1308-16. DOI: 10.1016/j.ijrobp.2005.10.039
19. Wendt TG, Grabenbauer GG, Rödel CM, Thiel HJ, Aydin H, Rohloff R, Wustrow TP, Iro H, Popella C, Schalhorn A. Simultaneous radiochemotherapy versus radiotherapy alone in advanced head and neck cancer: a randomized multicenter study. *J Clin Oncol.* 1998 Apr;16(4):1318-24.
20. Paccagnella A, Ghi MG, Loreggian L, Buffoli A, Koussis H, Mione CA, Bonetti A, Campostrini F, Gardani G, Ardzioia A, Dondi D, Guaraldi M, Cavallo R, Tomio L, Gava A; Gruppo di Studio Tumori della Testa e del Collo XRP 6976 F/2501 Study. Concomitant chemoradiotherapy versus induction docetaxel, cisplatin and 5 fluorouracil (TPF) followed by concomitant chemoradiotherapy in locally advanced head and neck cancer: a phase II randomized study. *Ann Oncol.* 2010 Jul;21(7):1515-22. DOI: 10.1093/annonc/mdp573
21. Posner MR, Hershock DM, Blajman CR, Mickiewicz E, Winquist E, Gorbounova V, Tjulandin S, Shin DM, Cullen K, Ervin TJ, Murphy BA, Raez LE, Cohen RB, Spaulding M, Tishler RB, Roth B, Viroglia Rdel C, Venkatesan V, Romanov I, Agarwala S, Harter KW, Dugan M, Cmelak A, Markoe AM, Read PW, Steinbrenner L, Colevas AD, Norris CM Jr, Haddad RI; TAX 324 Study Group. Cisplatin and fluorouracil alone or with docetaxel in head and neck cancer. *N Engl J Med.* 2007 Oct;357(17):1705-15. DOI: 10.1056/NEJMoa070956
22. Vermorken JB, Remenar E, van Herpen C, Gorlia T, Mesia R, Degardin M, Stewart JS, Jelic S, Betka J, Preiss JH, van den Weyngaert D, Awada A, Cupissol D, Kienzer HR, Rey A, Desanois I, Bernier J, Lefebvre JL; EORTC 24971/TAX 323 Study Group. Cisplatin, fluorouracil, and docetaxel in unresectable head and neck cancer. *N Engl J Med.* 2007 Oct;357(17):1695-704. DOI: 10.1056/NEJMoa071028
23. Ghi MG, Paccagnella A, D'Amanzo P, Mione CA, Fasan S, Paro S, Mastromauro C, Carnuccio R, Turcato G, Gatti C, Pallini A, Nascimben O, Biason R, Oniga F, Medici M, Rossi F, Fila G. Neoadjuvant docetaxel, cisplatin, 5-fluorouracil before concurrent chemoradiotherapy in locally advanced squamous cell carcinoma of the head and neck versus concomitant chemoradiotherapy: a phase II feasibility study. *Int J Radiat Oncol Biol Phys.* 2004 Jun;59(2):481-7. DOI: 10.1016/j.ijrobp.2003.10.055
24. Bernier J, Cooper JS, Pajak TF, van Glabbeke M, Bourhis J, Forastiere A, Ozsahin EM, Jacobs JR, Jassem J, Ang KK, Lefebvre JL. Defining risk levels in locally advanced head and neck cancers: a comparative analysis of concurrent postoperative radiation plus chemotherapy trials of the EORTC (#22931) and RTOG (# 9501). *Head Neck.* 2005 Oct;27(10):843-50. DOI: 10.1002/hed.20279
25. Cooper JS, Pajak TF, Forastiere AA, Jacobs J, Campbell BH, Saxman SB, Kish JA, Kim HE, Cmelak AJ, Rotman M, Machtay M, Ensley JF, Chao KS, Schultz CJ, Lee N, Fu KK; Radiation Therapy Oncology Group 9501/Intergroup. Postoperative concurrent radiotherapy and chemotherapy for high-risk squamous-cell carcinoma of the head and neck. *N Engl J Med.* 2004 May;350(19):1937-44. DOI: 10.1056/NEJMoa032646
26. Cooper JS, Zhang Q, Pajak TF, Forastiere AA, Jacobs J, Saxman SB, Kish JA, Kim HE, Cmelak AJ, Rotman M, Lustig R, Ensley JF, Thorstad W, Schultz CJ, Yom SS, Ang KK. Long-term follow-up of the RTOG 9501/intergroup phase III trial: postoperative concurrent radiation therapy and chemotherapy in high-risk squamous cell carcinoma of the head and neck. *Int J Radiat Oncol Biol Phys.* 2012 Dec;84(5):1198-205. DOI: 10.1016/j.ijrobp.2012.05.008
27. Forastiere AA, Zhang Q, Weber RS, Maor MH, Goepfert H, Pajak TF, Morrison W, Glisson B, Trotti A, Ridge JA, Thorstad W, Wagner H, Ensley JF, Cooper JS. Long-term results of RTOG 91-11: a comparison of three nonsurgical treatment strategies to preserve the larynx in patients with locally advanced larynx cancer. *J Clin Oncol.* 2013 Mar;31(7):845-52. DOI: 10.1200/JCO.2012.43.6097
28. Trotti A, Pajak TF, Gwede CK, Paulus R, Cooper J, Forastiere A, Ridge JA, Watkins-Bruner D, Garden AS, Ang KK, Curran W. TAME: development of a new method for summarising adverse events of cancer treatment by the Radiation Therapy Oncology Group. *Lancet Oncol.* 2007 Jul;8(7):613-24. DOI: 10.1016/S1470-2045(07)70144-4

29. Bieri S, Bentzen SM, Huguenin P, Allal AS, Cozzi L, Landmann C, Monney M, Bernier J. Early morbidity after radiotherapy with or without chemotherapy in advanced head and neck cancer. Experience from four nonrandomized studies. *Strahlenther Onkol*. 2003 Jun;179(6):390-5. DOI: 10.1007/s00066-003-1077-1
30. Machtay M, Moughan J, Trotti A, Garden AS, Weber RS, Cooper JS, Forastiere A, Ang KK. Factors associated with severe late toxicity after concurrent chemoradiation for locally advanced head and neck cancer: an RTOG analysis. *J Clin Oncol*. 2008 Jul;26(21):3582-9. DOI: 10.1200/JCO.2007.14.8841
31. Fietkau R, Lewitzki V, Kuhnt T, Hölscher T, Hess CF, Berger B, Wiegel T, Rödel C, Niewald M, Hermann RM, Lubgan D. A disease-specific enteral nutrition formula improves nutritional status and functional performance in patients with head and neck and esophageal cancer undergoing chemoradiotherapy: Results of a randomized, controlled, multicenter trial. *Cancer*. 2013 Sep;119(18):3343-53. DOI: 10.1002/cncr.28197
32. Rades D, Kronemann S, Meyners T, Bohlen G, Tribius S, Kazic N, Schroeder U, Hakim SG, Schild SE, Dunst J. Comparison of four cisplatin-based radiochemotherapy regimens for nonmetastatic stage III/IV squamous cell carcinoma of the head and neck. *Int J Radiat Oncol Biol Phys*. 2011 Jul;80(4):1037-44. DOI: 10.1016/j.ijrobp.2010.03.033
33. Nguyen NP, Vos P, Vinh-Hung V, Borok TL, Dutta S, Karlsson U, Lee H, Martinez T, Jo BH, Nguyen LM, Nguyen N, Sallah S. Altered glucose metabolism during chemoradiation for head and neck cancer. *Anticancer Res*. 2009 Nov;29(11):4683-7.
34. Komdeur R, Derksen J, Legdeur MC, Hylkema BS. Cisplatin-induced hyperglycemic hyperosmolar coma. *Neth J Med*. 2007 Jan;65(1):36-7.
35. Hey J, Setz J, Gerlach R, Vordermark D, Gernhardt CR, Kuhnt T. Effect of Cisplatin on parotid gland function in concomitant radiochemotherapy. *Int J Radiat Oncol Biol Phys*. 2009 Dec;75(5):1475-80. DOI: 10.1016/j.ijrobp.2008.12.071
36. Simone CB 2nd, Ly D, Dan TD, Ondos J, Ning H, Belard A, O'Connell J, Miller RW, Simone NL. Comparison of intensity-modulated radiotherapy, adaptive radiotherapy, proton radiotherapy, and adaptive proton radiotherapy for treatment of locally advanced head and neck cancer. *Radiother Oncol*. 2011 Dec;101(3):376-82. DOI: 10.1016/j.radonc.2011.05.028
37. Staar S, Rudat V, Stuetzer H, Dietz A, Volling P, Schroeder M, Flentje M, Eckel HE, Mueller RP. Intensified hyperfractionated accelerated radiotherapy limits the additional benefit of simultaneous chemotherapy—results of a multicentric randomized German trial in advanced head-and-neck cancer. *Int J Radiat Oncol Biol Phys*. 2001 Aug;50(5):1161-71. DOI: 10.1016/S0360-3016(01)01544-9
38. Kundu SK, Nestor M. Targeted therapy in head and neck cancer. *Tumour Biol*. 2012 Jun;33(3):707-21. DOI: 10.1007/s13277-012-0350-2
39. Ang KK, Berkey BA, Tu X, Zhang HZ, Katz R, Hammond EH, Fu KK, Milas L. Impact of epidermal growth factor receptor expression on survival and pattern of relapse in patients with advanced head and neck carcinoma. *Cancer Res*. 2002 Dec;62(24):7350-6.
40. Sharafinski ME, Ferris RL, Ferrone S, Grandis JR. Epidermal growth factor receptor targeted therapy of squamous cell carcinoma of the head and neck. *Head Neck*. 2010 Oct;32(10):1412-21. DOI: 10.1002/hed.21365
41. Lefebvre J, Pointreau Y, Rolland F, Alfonsi M, Baudoux A, Sire C, De Raucourt D, Bardet E, Tuchsais C, Garaud P, G. Calais G. Sequential chemoradiotherapy (SCRT) for larynx preservation (LP): Results of the randomized phase II TREMPIN study. *J Clin Oncol*. 2011;29(15 Suppl):abstr 5501.
42. Bonner JA, Harari PM, Giralt J, Azarnia N, Shin DM, Cohen RB, Jones CU, Sur R, Raben D, Jassem J, Ove R, Kies MS, Baselga J, Youssoufian H, Amellal N, Rowinsky EK, Ang KK. Radiotherapy plus cetuximab for squamous-cell carcinoma of the head and neck. *N Engl J Med*. 2006 Feb;354(6):567-78. DOI: 10.1056/NEJMoa053422
43. Budach W, Bölke E, Homey B. Severe cutaneous reaction during radiation therapy with concurrent cetuximab. *N Engl J Med*. 2007 Aug;357(5):514-5. DOI: 10.1056/NEJMco071075
44. Koutcher LD, Wolden S, Lee N. Severe radiation dermatitis in patients with locally advanced head and neck cancer treated with concurrent radiation and cetuximab. *Am J Clin Oncol*. 2009 Oct;32(5):472-6. DOI: 10.1097/COC.0b013e318193125c
45. Potthoff K, Hofheinz R, Hassel JC, Volkenandt M, Lordick F, Hartmann JT, Karthaus M, Riess H, Lipp HP, Hauschild A, Trarbach T, Wollenberg A. Interdisciplinary management of EGFR-inhibitor-induced skin reactions: a German expert opinion. *Ann Oncol*. 2011 Mar;22(3):524-35. DOI: 10.1093/annonc/mdq387
46. Bernier J, Bonner J, Vermorken JB, Bensadoun RJ, Dummer R, Giralt J, Kornek G, Hartley A, Mesia R, Robert C, Segal S, Ang KK. Consensus guidelines for the management of radiation dermatitis and coexisting acne-like rash in patients receiving radiotherapy plus EGFR inhibitors for the treatment of squamous cell carcinoma of the head and neck. *Ann Oncol*. 2008 Jan;19(1):142-9. DOI: 10.1093/annonc/mdm400
47. Barber NA, Ganti AK. Pulmonary toxicities from targeted therapies: a review. *Target Oncol*. 2011 Dec;6(4):235-43. DOI: 10.1007/s11523-011-0199-0
48. Costa A, Tejpar S, Prenen H, Van Cutsem E. Hypomagnesaemia and targeted anti-epidermal growth factor receptor (EGFR) agents. *Target Oncol*. 2011 Dec;6(4):227-33. DOI: 10.1007/s11523-011-0200-y
49. Landis SH, El-Hariry IA, van Herk-Sukel MP, van den Haak P, Janssen-Heijnen ML, Penning-van Beest FJ, Herings RM. Prevalence and incidence of acute and chronic comorbidity in patients with squamous cell carcinoma of the head and neck. *Head Neck*. 2012 Feb;34(2):238-44. DOI: 10.1002/hed.21720
50. Rottlaender D, Reda S, Motloch LJ, Hoppe UC. Neue Tyrosinkinase- und EGFR-Inhibitoren in der Tumorthherapie. Herz und Haut als wichtige Schädigungsorgane. Teil A: Herz [New tyrosine kinase and EGFR inhibitors in cancer therapy. Cardiac and skin toxicity as relevant side effects. Part A: heart]. *Internist (Berl)*. 2011 Oct;52(10):1245-55. DOI: 10.1007/s00108-011-2895-3
51. Nishiyama K, Tanaka E, Tarui Y, Miyauchi K, Okagawa K. A prospective analysis of subacute thyroid dysfunction after neck irradiation. *Int J Radiat Oncol Biol Phys*. 1996 Jan;34(2):439-44. DOI: 10.1016/0360-3016(95)02079-9
52. Jerezek-Fossa BA, Alterio D, Jassem J, Gibelli B, Tradati N, Orecchia R. Radiotherapy-induced thyroid disorders. *Cancer Treat Rev*. 2004 Jun;30(4):369-84. DOI: 10.1016/j.ctrv.2003.12.003
53. Léon X, Gras JR, Pérez A, Rodríguez J, de Andrés L, Orús C, Quer M. Hypothyroidism in patients treated with total laryngectomy. A multivariate study. *Eur Arch Otorhinolaryngol*. 2002 Apr;259(4):193-6. DOI: 10.1007/s00405-001-0418-x

Corresponding author:

Thomas G. Wendt

Department of Radiation Oncology, University Hospital Jena, Bachstraße 18, 07743 Jena, Germany
thomas.wendt@med.uni-jena.de

Please cite as

Wendt TG. Hazards and risks in oncology: radiation oncology. GMS Curr Top Otorhinolaryngol Head Neck Surg. 2013;12:Doc03.
DOI: 10.3205/cto000095, URN: urn:nbn:de:0183-cto0000951

This article is freely available from

<http://www.egms.de/en/journals/cto/2013-12/cto000095.shtml>

Published: 2013-12-13

Copyright

©2013 Wendt. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by-nc-nd/3.0/deed.en>). You are free: to Share – to copy, distribute and transmit the work, provided the original author and source are credited.