

Optimization of hydroxychloroquine dosing scheme based on COVID-19 patients' characteristics: a review of the literature and simulations

Eleni Karatza^a , George Ismailos^b , Markos Marangos^c  and Vangelis Karalis^a 

^aDepartment of Pharmacy, School of Health Sciences, National and Kapodistrian University of Athens, Athens, Greece; ^bExperimental-Research Center ELPEN, ELPEN Pharmaceuticals, Pikermi, Greece; ^cDivision of Infectious Diseases, University Hospital of Patras, Rio, Greece

ABSTRACT

1. During the recent COVID-19 outbreak hydroxychloroquine (HCQ) has been proposed as a safe and effective therapeutic option. However, a wide variety of dosing schemes has been applied in the clinical practice and tested in clinical studies.
2. An extended literature survey was performed investigating the pharmacokinetics, the efficacy and safety of HCQ in COVID-19 treatment. Population pharmacokinetic models were retrieved from the literature and after evaluation and assessment one was selected in order to perform simulations.
3. The most commonly applied dosing schemes were explored for patients with different weights and different levels of HCQ clearance impairment. Model-based simulations of HCQ concentrations revealed that high initial doses followed by low and sparse doses may offer significant benefits to patients by decreasing the viral load without reaching levels considered to produce adverse effects. For instance, the dosing scheme proposed for a 70 kg adult with moderate COVID-19 symptoms would be 600 mg upon diagnosis, 400 mg after 12 h, 300 mg after 24 h, 200 mg after 36 h, followed by 200 mg BID for 4 d, followed by 200 mg OD for 5 d.
4. Based on the results from simulations performed and the currently published knowledge regarding HCQ in COVID-19 treatment, this study provides evidence that a high loading dose followed by sparse doses could offer significant benefits to the patients.

ARTICLE HISTORY

Received 31 July 2020
Revised 11 September 2020
Accepted 12 September 2020

KEYWORDS

COVID-19; hydroxychloroquine; efficacy and safety; modeling and simulation; precision dosing

Introduction

During the recently emerged pandemic of coronavirus disease 2019 (COVID-19) due to SARS-CoV-2, hydroxychloroquine (HCQ) has been proposed as a drug of choice (Alpern & Gertner, 2020). COVID-19 severity has been categorized based on symptoms by the WHO in mild, moderate and severe, while a life-threatening state of critical disease has been noted, as well (WHO, 2020). Most interestingly, it has been shown that mortality, as well as disease severity and immune response are closely related to the viral load of the patient (Casadevall et al., 2020; Liu et al., 2020a; Pujadas et al., 2020; Wang et al., 2020; Zheng et al., 2020).

Up to this point, HCQ has been used for the treatment of malaria or autoimmune diseases such as rheumatoid arthritis, discoid and systemic lupus erythematosus (Plaquenil Label, 2017; Plaquenil SmPC, 2020). However, HCQ seems to offer a dual beneficial action for COVID-19 patients, as it exerts both immunomodulatory and antiviral effects (Drożdżal et al., 2020). In fact, some recent studies have revealed the drug's *in vitro* antiviral activity (Liu et al., 2020b; PAHO/WHO, 2020; Yao et al., 2020). A wide variety of observational and randomized controlled trials (RCT) worldwide are currently

evaluating the efficacy of HCQ for the treatment of COVID-19, implementing many different dosing schemes and patients with different levels of disease severity according to the WHO, (2020) categorization (Alpern & Gertner, 2020; Cortegiani et al., 2020; PAHO/WHO, 2020). Several studies have yielded positive results, indicating that HCQ may accelerate the alleviation of symptoms and diminish the length of hospital stay (Arshad et al 2020; Cavalcanti et al., 2020; Chen et al., 2020a; Gautret et al. 2020a,b; Tang et al 2020; Yu et al., 2020), while others yielded negative results, indicating that HCQ does not show any clinical effects in improving symptoms or was associated with increased length of hospital stay or disease progression (Barbosa et al., 2020; Chen et al., 2020b; Horby et al., 2020; Molina et al 2020). It has to be noted, that in many cases these studies report adverse effects promoted by HCQ, such as prolongation of the QT-interval or diarrhea (Gautret et al 2020b; Horby et al. 2020; Tang et al., 2020).

In view of the high heterogeneity of the so far study outcomes, the WHO recommends not to use HCQ outside the context of clinical trials (WHO, 2020). In any case, due to the lack of a better option, HCQ is currently included in the

therapeutic protocols designed by numerous hospitals and health systems worldwide, for the treatment of COVID-19 patients (Singh et al., 2020). In addition, the FDA that initially approved the use of HCQ for COVID-19 (Alpern & Gertner, 2020) has recently revoked the emergency off-label use of this drug (FDA, 2020). It should be noted that currently there is not substantial evidence to prove the safety and efficacy of HCQ in the treatment of COVID-19, and thus results from well-designed randomized trials are required for this drug's repurposing (Elavarasi et al., 2020; Singh-Uttam et al., 2020).

Indicatively, some dosing schemes that have been applied or evaluated in clinical studies are presented in Table 1.

Considering all the above and based on the reported antiviral activity of the compound (Drożdżal et al., 2020; Liu et al., 2020b; Yao et al., 2020), there is currently a need to rationalize the administered dosing schemes in order to maximize the efficacy and safety of HCQ in patients with COVID-19. Eventually, additional time is needed in order to retrieve concrete results from new *in vitro* and *in vivo* studies specifically designed for HCQ against COVID-19. However, in view of its long-standing clinical use a lot of data and experience has been accumulated for this drug. In this vein, modeling and simulation techniques have been proved very useful as they can combine all the available data from *in vitro*, pre-clinical and clinical studies (Lavé et al., 2007; Lowe et al., 2007). Thus, by implementing all the relevant knowledge that has been gathered so far and using modeling and simulation techniques it is possible to attain a better approximation to the optimum dosing scheme.

The aim of the present study was firstly, to review all the relevant to HCQ literature, focusing on COVID-19 treatment and secondly, through simulations, to investigate the efficacy

and safety of the dosing schemes of HCQ for COVID-19 treatment currently applied and to propose optimized dosing schemes.

Materials and methods

Pharmacokinetics of HCQ

After oral administration HCQ is rapidly and almost completely absorbed with bioavailability ranging from 0.67 to 0.74. Mean peak plasma concentrations (C_{max}) after a single dose of 400 mg was approximately 105 ng/ml, and mean time to peak plasma concentration (T_{max}) was 1.83 h. HCQ was found to present linear pharmacokinetics in a therapeutic dose range (Furst, 1996; Plaquenil SmPC, 2020).

HCQ binds avidly to tissues, leading to large volumes of distribution, significant accumulation of the drug and low clearance. In fact, the increased amount of time needed for this accumulation to occur, accounts for the delayed appearance of its clinical effects but also of its adverse effects. Approximately 30-40% of the administered dose is bound to proteins, both albumin and alpha glycoprotein. HCQ is metabolized by the liver with its main metabolite (desethyl-hydroxychloroquine) presenting some immunomodulatory activity (Furst, 1996; Munster et al., 2002).

The drug is primarily eliminated via the urine, where about 3% of the administered dose is recovered in 24 h. Following a single 200 mg oral dose the half-life of HCQ was found to be 537 h (22.4 d) (Plaquenil Label, 2017; Plaquenil SmPC, 2020).

Blood levels were found to vary significantly among individuals, with a variation in mean maximum/minimum concentration reaching 40%. This variability has been attributed

Table 1. Dosing schemes of hydroxychloroquine applied for the treatment of COVID-19.

Dosing Schemes*	Applied by	Reference
200 mg TID for 10 d	Observational study	Gautret et al., 2020a,b
400 mg at diagnosis, 400 mg 12 h later, followed by 200 mg BID for 5 d	Massachusetts General Hospital (clinical practice)	Massachusetts General Hospital, 2020
200 mg per day for 10 d	Italian Society of Infectious and Tropical Diseases Lombardy Section (clinical practice)	Singh et al., 2020
400 mg orally per day for 7–10 d	Central Clinical Task Force, Korea (clinical practice)	Singh et al., 2020
400 mg BID × 2 doses then 12 h later start 400 mg OD for 5–10 d	Mount Sinai Health System, Canada (clinical practice)	Singh et al., 2020
400 mg BID × 1 day followed by 200 mg BID × 4 d	Clinical guidance for patients with suspected or confirmed COVID-19 in Belgium (clinical practice)	Arnold & Buckner, 2020; Singh et al., 2020; Yao et al., 2020
400 mg twice daily for 2 doses on day 1, followed by 200 mg BID on days 2–5	Multi-center retrospective observational study	Arshad et al., 2020
400 mg BID for 7 d	Randomized controlled trial Mild-to-moderate hospitalized patients	Cavalcanti et al., 2020
A loading dose of 1200 mg daily for 3 d followed by a maintenance dose of 800 mg daily for 2 or 3 weeks	Randomized controlled trial Mild-to-moderate (2 weeks) or severe (3 weeks) disease	Tang et al., 2020
800 mg upon diagnosis and at 6 h, followed by 400 mg starting at 12 after the initial dose and then every 12 h for the next 9 d or until discharge	Randomized controlled trial (RECOVERY)	Horby et al., 2020
200 mg BID for 5 d	Randomized controlled trial	Chen et al., 2020a
200 mg BID for 7–10 d	Observational (retrospective)	Yu et al., 2020
400 mg per day for 5 d	Randomized controlled trial	Chen et al., 2020b
200 mg TID for 10 d	Observational (narrative review)	Molina et al., 2020
400 mg BID for 1–2 d and 3–4 subsequent d of 200 mg to 400 mg OD	Randomized controlled trial	Barbosa et al., 2020

*Doses are expressed in mg of hydroxychloroquine sulfate (200 mg tablet contains 155 mg base equivalent).

to the absorption and distribution rather than clearance characteristics of the drug (Al-Rawi et al., 2018; Miller et al., 1991).

Numerous factors have been shown to affect blood levels of HCQ, such as genetic variants of the CYP enzyme family, age, gender, body weight, gastric emptying, and co-administration with immunosuppressants, such as corticosteroids (Al-Rawi et al., 2018; Lee et al., 2016, 2017). Even though it has been reported that renal clearance of unchanged drug was approximately 16 to 30% and did not correlate with creatinine clearance (Miller et al., 1991), renal insufficiency has been reported to increase the risk of toxicity (Abdulaziz et al., 2018).

Model selection for simulations

A total of eight population pharmacokinetic studies of HCQ were identified in the literature (Balevic et al., 2019; Carmichael et al., 2003; Haas et al., 2019; Lim et al. 2009; Morita et al., 2016; Rangwala et al., 2014; Rosenfeld et al., 2014; Vogl et al., 2014). Population parameter estimates and characteristics of the studies are presented in Table 2.

In the majority of studies assessed, samples were retrieved from patients suffering from severe diseases and receiving co-medications especially immunosuppressants that have been reported to alter HCQ pharmacokinetics (Lee et al., 2017). In addition, given that HCQ binds avidly to tissues, a fact that results in high volumes of distribution and slow clearance (Furst, 1996), a model with two compartments, a central one for plasma and tissues with an instantaneous distribution and a peripheral one standing for tissues where the drug distribution is slower seemed to be more appropriate for the description of HCQ disposition. In this vein, the model of Lim et al. (2009) was developed with samples

retrieved from 91 healthy volunteers and patients with vivax malaria that did not receive any other medications and resulted in a two compartmental model. Also, the study of Lim et al. (2009) that included the highest number of samples, i.e., a total of 431 concentration measurements, from all the studies identified resulting in a two compartmental model, a fact that increases the reliability of the parameter estimates.

In view of the above, the model developed by Lim et al., (2009) was selected in order to perform the simulations. Parameters estimated in this study, were similar to those obtained with other population pharmacokinetic models developed (Rangwala et al., 2014; Rosenfeld et al., 2014), as well as to those retrieved after non-compartmental pharmacokinetic analysis (Furst, 1996; Munster et al., 2002).

Body weight has been reported to affect significantly HCQ *in vivo* concentrations (Plaquenil SmPC, 2020; Lee et al., 2017) and several models have identified it as a statistically significant covariate affecting clearance allometrically (Balevic et al., 2019; Morita et al., 2016; Vogl et al., 2014). Thus, weight has been included in the model used for simulations with an allometric exponent of 0.8, as estimated in these studies.

Toxicity, adverse effects and upper limit of hydroxychloroquine levels

The maximum tolerated dose in adults with rheumatoid arthritis has been reported 1200 mg per day (Munster et al., 2002), while in patients with newly diagnosed glioblastoma multiforme the maximum tolerated dose reported was 600 mg (Rosenfeld et al., 2014).

In order to prevent retinopathy that HCQ induces after long term use, a concentration range of 500–2000 ng/ml has

Table 2. Population pharmacokinetic models of hydroxychloroquine identified in the literature.

	Balevic et al., 2019 Pregnant women with rheumatoid arthritis	Carmichael et al., 2003 Rheumatoid arthritis	Morita et al., 2016 Cutaneous or systemic lupus erythematosus		
Population					
Co-medication	Yes	No	Yes		
Compartments	1	1	1		
<i>t</i> _{lag} (h)	No lag time	0.654	0.389		
<i>k</i> _a (1/h)	1.15	0.91	1.15		
V/F (L)	1850	820	2440		
Cl/F (L/h)	51	9. 13.13	68.2		
Covariate	Weight (allometric exponent = 1)	No covariate	Weight (allometric exponent = 0.844)		
	Haas et al., 2019 Renal carcinoma	Rosenfeld et al., 2014 Glioblastoma multiforme	Rangwala et al., 2014 Advanced solid tumours and melanoma	Vogl et al., 2014 Relapsed/refractory myeloma	Lim et al., 2009 Healthy adults and patients with vivax malaria
Population					
Co-medication	Yes	Yes	Yes	Yes	No
Compartments	2	2	2	2	2
<i>t</i> _{lag} (h)	No lag time	1.06	No lag time	1.74	0.389
<i>k</i> _a (1/h)	0.93	0.49	0.998	1.27	1.15
V/F (L)	599.89	361.28	485.747	243.87	437
V ₂ /F (L)	3604.83	947.26	1406.52	2537.68	1390
Cl/F (L/h)	7.98	11.44	9.97	3	10.9
Q/F (L/h)	14.98	103.9	49.043	15	45.1
Covariate	No	No	No	Weight allometric (value not reported)	No

Tag: absorption lag time; *k*_a: first order absorption rate; V/F: apparent volume of distribution in the central compartment; Cl/F: apparent first order clearance from the central compartment; V₂/F: apparent volume of distribution in the peripheral compartment; Q/F: apparent intercompartmental clearance between the central and the peripheral compartment.

been proposed to be safe and effective in patients with systemic lupus erythematosus under chronic therapeutic treatment (Durcan et al., 2015).

Regarding its acute use, the most significant adverse effects to consider are cardiac, gastrointestinal, extrapyramidal and neuropsychiatric effects (Plaquenil SmPC, 2020; Juurlink, 2020). The cardiotoxicity of chloroquines has been shown to be dose-dependent, with mean increases in QTc of 6.1 ms after a chloroquine dose of 600 mg and 28 ms after a chloroquine dose of 1200 mg. It should be noted that many drugs co-administered with HCQ for the treatment of COVID-19 as azithromycin, ceftriaxone and fluoroquinolones have also been proved to promote prolongation of the QTc interval, increasing the risk of cardiotoxicity (Briasoulis et al., 2011; Juurlink, 2020; Teng et al., 2019).

Gastrointestinal (GI) toxicity of HCQ has been studied in relation to blood levels in 212 patients with rheumatoid arthritis followed for 24 weeks. It was revealed that blood levels of 2250 ng/ml promote GI adverse effects in 30% of patients, while 5250 ng/ml in 50% of patients. This is a rather significant observation in terms of HCQ treatment, as generally GI symptoms are the first and most frequent to occur constituting a warning, even for cardiotoxicity (Munster et al., 2002; Tang et al 2020).

Risk factors related to HCQ toxicity include old age, renal and liver disease, genetic variants, concomitant drug use, high body mass index and, obviously, high dose and long duration of treatment (Marmor et al., 2016).

Symptoms of HCQ overdose manifest rapidly within 30 minutes after administration. They include headache, visual disturbances, cardiovascular collapse, convulsions, and hypokalemia, cardiac rhythm and conduction disorders, including QT prolongation, Torsade de pointes, ventricular tachycardia and ventricular fibrillation, followed by sudden potentially fatal respiratory and cardiac arrest (Plaquenil Label, 2017; de Olano et al., 2019).

In view of the above, HCQ levels of 2250 ng/ml, even though rather conservatory, were used as an upper limit, considering also the cardiotoxic effects of the drug, possible co-administered medications and impaired clearance due to co-morbidities (Abdulaziz et al., 2018).

It should be noted that this concentration is far below the cytotoxic concentration of the drug (CC50 = 249.50 μ M or 83799 ng/ml) in green monkey kidney VeroE6 cells (Liu et al., 2020b). Regarding cytotoxicity, Yang et al., 2020 tested cytotoxicity of HCQ in 8 cell lines retrieved from heart, liver, kidney, retina, intestine and lung. This study identified that the lowest cytotoxic level of HCQ was 15.26 μ M or 5125.40 ng/ml, i.e., a concentration far below the safety threshold selected in the present study (Yang et al., 2020).

Efficacy and lower limit of hydroxychloroquine levels

In the study of Liu et al. (2020b) the 50% maximal effective concentration (EC50) of HCQ against SARS-CoV-2 was estimated using the same cell line for 4 different multiplicities of infections (MOIs) namely (0.01, 0.02, 0.2, and 0.8) after treatment with the drug for 1 h. The EC50s found were 4.51, 4.06, 17.31, 12.96 μ M or 1514.46, 1363.6, 5813.9 and 4352.8 ng/ml,

Table 3. Population parameters used for simulations (Lim et al., 2009).

Parameters	Parameter estimates	Inter-individual variability
tlag (h)	0.389	0.0359
ka (1/h)	1.15	–
V/F (L)	437	0.232
V2/F (L)	1390	0.715
Cl/F (L/h)	10.9	0.161
Weight exponentially on Cl/F	0.8	–
Q/F (L/h)	45.1	–

tlag: absorption lag time; ka: first order absorption rate; V/F: apparent volume of distribution in the central compartment; Cl/F: apparent first order clearance from the central compartment; V2/F: apparent volume of distribution in the peripheral compartment; Q/F: apparent intercompartmental clearance between the central and the peripheral compartment; Weight exponentially on Cl/F: allometric exponent for weight scaling of apparent clearance.

respectively (Liu et al., 2020b). Most interestingly, in another study performed by Yao et al. (2020), also in green monkey kidney VeroE6 cells, a time – dependency of the EC50 was demonstrated, with EC50 values corresponding to 6.14 μ M or 2062.25 ng/ml and 0.72 μ M or 241.82 ng/ml after 24 and 48 h, respectively.

This phenomenon was attributed to the fact that HCQ is accumulated within the cells, and its actions present a delay to be manifested (Yao et al., 2020). Besides its anti-viral activity, HCQ's therapeutic effects against COVID-19 reside also on its immunomodulatory effects which are manifested with blood concentrations above 500 ng/ml (Durcan et al., 2015).

Simulations

Simulations were performed using the R function 'Simulx' included in the 'mlxR 4.0' package (Lavielle, 2019). The model parameters used are summarized in Table 3. Inter-individual variability was also taken into account and thus a population of 500 patients was simulated. The dosing schemes explored were selected from published literature studies and simulations were performed assuming patients with weights of 50, 70 and 90 kg. Then, the dosing scheme providing the most favorable profile was sought.

HCQ is metabolized by the liver, while it is primarily excreted by the kidney. Both renal insufficiency and impaired hepatic clearance, can reduce HCQ's total clearance. It has been reported that renal clearance constitutes the 55% of total HCQ's clearance (White et al., 2020). COVID-19 patients have been proved to present increased risk of renal impairment (Cheng et al., 2020; Naicker et al., 2020; Ronco and Reis 2020), while extracorporeal membrane oxygenation (ECMO) has been showed to impair the drug's clearance (Tukacs, 2018). Therefore, during simulations an intermediate and a high level of renal impairment reducing the total clearance of HCQ by 30 and by 50%, i.e., a population with an apparent clearance of 7.63 and 5.45 L/h, were also considered.

Results and discussion

Simulations were performed in order to explore the expected blood levels of HCQ upon administration of various dosing

schemes currently used in clinical practice, as well as, in order to propose an optimal dosing scheme in relation to patients' body weight, clearance impairment and COVID-19 severity.

The lower and upper bounds, selected herein, are in line with other published studies (Arnold & Buckner, 2020; Perinel et al., 2020; Yao et al., 2020), while concentrations achieved with the simulations are in accordance with the dose-concentrations results presented in previously pharmacokinetic analyses (Morita et al., 2016; Rangwala et al., 2014; Vogl et al., 2014) indicating the good predictability of our simulations.

As it may be noted in Figures 1 and 2, low frequent doses of HCQ result after a period of 5 d in toxic levels even in patients with normal body weight, while high initial bolus doses do not increase blood levels significantly. This phenomenon is due to HCQ pharmacokinetics that imply its accumulation into tissues leading to delayed clearance (Furst, 1996).

In terms of efficacy, three factors should be considered for the determination of the appropriate HCQ blood levels: (a) higher multiplicity of infection *in vitro*, i.e., higher viral load, results in higher EC50, (b) increasing the duration of exposure to HCQ results in lower EC50 that decreases from 2062.25 ng/ml to 241.82 ng/ml in 24 h, and (c) the immunomodulatory effect of the drug is promoted at concentrations over 500 ng/ml. Therefore, blood levels should ideally reach 1500 ng/ml during the first days upon diagnosis, especially in severe cases with high viral load, and be kept above 500 ng/ml for the following days (Durcan et al., 2015; Liu et al., 2020b; Yao et al., 2020).

On the other hand, levels exceeding 2250 ng/ml are considered more probable to promote gastrointestinal adverse effects, which seem to constitute a warning for cardiotoxicity as they occur first and are more common (Munster et al., 2002; Gautret et al. 2020a,b; Plaquenil SmPC, 2020). In addition, this concentration is far below the concentration found to be cardiotoxic *in vitro*, i.e., 5125.40 ng/ml. More interestingly, cardiotoxicity of chloroquines is known to be dose-dependent, with 600 mg of chloroquine, a far more toxic compound than HCQ (Liu et al., 2020b), promoting a prolongation of 6.1 ms, when prolongations less than 10 ms are considered to be of low concern (US FDA E14, 2017). In general, significant prolongations of the QT interval with HCQ have been noted with either high doses or with long term-use and frequent doses (Gautret et al. 2020a,b; Jankelson et al., 2020; Juurlink, 2020; Saleh et al., 2020). As a result, in the present study the dosing schemes proposed, were designed in order to avoid unnecessarily high and frequent doses.

Based on simulations among the currently published dosing schemes, the most adequate dose for a patient of 50 or 70 kg, even with 30% impaired clearance, would be 800 mg once daily on day 1, followed by 200 mg twice daily for 7 d (Scheme_7) or 400 mg upon diagnosis, 400 mg 12 h later, followed by 200 mg BID for 5 d (Scheme_2). For a patient of 90 kg, 400 mg BID for 2 doses followed 12 h later by 400 mg OD for 5–10 d (Scheme_5) seems to be the most adequate

scheme. In this vein, for a patient of 70 kg with 50%, a dose of 200 mg per day for 10 d (Scheme_3) seems to constitute a safe option, even though this dosing scheme is not expected to be effective enough to significantly decrease the viral load, during the first days of the treatment.

Through model-based simulations, optimal dosing schemes were developed (Table 4 and Supplementary material) and the corresponding HCQ blood concentrations versus time were retrieved (Figures 3 and 4). Dosing schemes were designed in order to achieve a fast onset of "high" concentrations during the initial phase of the disease, since an initial higher viral load is anticipated, especially for patients with severe COVID-19, and then keep HCQ blood levels below 2250 ng/ml and over 500 ng/ml, at all times. This can be noted in Figures 3 and 4 where the simulated HCQ concentrations versus time profiles for 500 volunteers are presented as 90% prediction intervals. In Table 4 the dosing scheme designed for a patient of 70 kg with normal clearance is presented, while Tables with the dosing schemes designed for patients of 30 kg, 50 kg, 90 kg, 110 kg, 70 kg with 30% impaired clearance and 70 kg with 50% impaired clearance are included in the Supplementary Material.

It should be noted that some of the dosing schemes applied currently in the clinical setting start with a loading dose of 400 mg. Despite that, the rest of the treatment in most cases differs significantly, as in most cases doses that lead to drug accumulation and thus increase the possibility of adverse effects are widely noted in the literature. In addition, only one dose scheme is used in all cases and all patients, irrespectively of their body weight, HCQ clearance or their condition.

In this study, the need to reach higher concentrations in patients whose immune system is not effectively reducing the viral load, leading to more intense symptoms is addressed. In fact, it has been shown that viral load relates significantly to disease severity and immune response. High viral loads were linked to intense immune response, even in peripheral tissues, complicating the patient's condition and increasing the risk of mortality (Casadevall et al., 2020; Liu et al., 2020a; Pujadas et al., 2020; Wang et al., 2020; Zheng et al., 2020). HCQ in an appropriate dosage may be of benefit thanks to its dual action, i.e., immunomodulatory and antiviral.

In addition, the possibility of impaired clearance is explored and its effect on the drug's levels estimated. Based on the pharmacokinetics of HCQ, the fact that the drug binds avidly to tissues and that patient's weight can significantly affect its clearance was taken under consideration. Thus, depending on the patient and his/her condition, the dosing schemes proposed in this study actually present some significant differences compared to those currently applied (e.g., Supplementary Table S4 and S6), improving the possibility of a safer use of HCQ, depending on patient's characteristics.

A possible disadvantage of the dosing schemes proposed is their complexity, as they involve different doses and varying frequency of administration, especially during the first days upon diagnosis. However, based on the results from




Figure 1. Simulated hydroxychloroquine concentrations (ng/ml) versus time (hours) in 500 patients receiving dosing schemes proposed in the literature for **A:** a population of 50 kg. **B:** a population of 70 kg, **C:** a population of 90 kg. **Scheme_1:** 200 mg TID for 10 d used by Gautret et al France (Gautret et al., 2020a,b). **Scheme_2:** 400 mg at diagnosis, 400 mg 12 h later, followed by 200 mg BID for 5 d used by the Massachusetts General Hospital (Massachusetts General Hospital, 2020). **Scheme_3:** 200 mg per day for 10 d used by the Italian Society of Infectious and Tropical Diseases Lombardy Section (Singh et al., 2020). **Scheme_4:** 400 mg orally per day for 7–10 d used by the Central Clinical Task Force, Korea. (Singh et al., 2020). **Scheme_5:** 400 mg BID \times 2 doses then 12 h later start 400 mg OD for 5–10 d used by the Mount Sinai Health System, Canada (Singh et al., 2020). **Scheme_6:** 400 mg BID for day 1, followed by 200 mg BID for 4 d. (Singh et al., 2020). **Scheme_7:** 800 mg once daily on day 1, followed by 200 mg twice daily for 7 d (Perinel et al., 2020) Red line: 2235 ng/ml, known to promote adverse effects to 30% of patients Blue line: 1500 ng/ml, levels to achieve during the first doses. Green line: 500 ng/ml, minimum levels eliciting immunomodulatory effect.




Figure 2. Simulated hydroxychloroquine concentrations (ng/ml) versus time (hours) in 500 patients receiving dosing schemes proposed in the literature for **A:** a population of 70 kg with 30% impaired clearance and **B:** a population of 70 kg with 50% impaired clearance. **Scheme_1:** 200 mg TID for 10 d used by Gautret et al. France (Gautret et al., 2020a,b). **Scheme_2:** 400 mg at diagnosis, 400 mg 12 h later, followed by 200 mg BID for 5 d used by the Massachusetts General Hospital (Massachusetts General Hospital, 2020). **Scheme_3:** 200 mg per day for 10 d used by the Italian Society of Infectious and Tropical Diseases Lombardy Section. (Singh et al., 2020). **Scheme_4:** 400 mg orally per day for 7–10 d used by the Central Clinical Task Force, Korea. (Singh et al., 2020). **Scheme_5:** 400 mg BID \times 2 doses then 12 h later start 400 mg OD for 5–10 d used by the Mount Sinai Health System, Canada. (Singh et al., 2020). **Scheme_6:** 400 mg BID for day 1, followed by 200 mg BID for 4 d (Singh et al., 2020). **Scheme_7:** 800 mg once daily on day 1, followed by 200 mg twice daily for 7 d (Perinel et al., 2020) Red line: 2235 ng/ml, known to promote adverse effects to 30% of patients. Blue line: 1500 ng/ml, levels to achieve during the first doses. Green line: 500 ng/ml, minimum levels eliciting immunomodulatory effect.

Table 4. Proposed dosing scheme for treatment of COVID-19 with hydroxychloroquine for a 70 kg patient, depending on disease severity.

Patient of 70kg					
Mild		Moderate		Severe	
Time (h)	Doses (mg)	Time (h)	Doses (mg)	Time (h)	Doses (mg)
0	400	0	600	0	800
12	400	12	400	12	400
24	300	24	300	24	400
36	300	36	200	36	300
48	200	48	200	48	200
72	200	60	200	60	200
96	200	72	200	72	200
120	200	84	200	84	200
144	200	96	200	96	200
168	200	108	200	108	200
192	200	120	200	120	200
216	200	144	200	132	200
240	200	168	200	144	200
		192	200	156	200
		216	200	168	200
		240	200	192	200
				216	200
				240	200

*Doses are expressed in mg of hydroxychloroquine sulfate (a 200 mg tablet contains 155 mg base equivalent).

simulations performed and the currently published data regarding hydroxychloroquine in COVID-19 treatment, these schemes could offer significant benefits to the patients, while after the first 3 d, in most cases they require only a “low” dose once or twice daily.

A maximum of 10 d of treatment is proposed in view of the course of COVID-19 (Harapan et al., 2020). In fact, as it has been shown for the 90% of mild cases, viral clearance is achieved within 10 d post-onset (Liu et al., 2020a). However, clinical evaluation of patients has to be performed and decisions should be made on a case-by-case basis. Therefore, continuation of a daily dose may be required for a longer period, keeping HCQ concentrations steady. Despite the high doses proposed upon diagnosis, the MTD of HCQ is not exceeded. It should be noted that continuous ECG monitoring should be conducted, especially during the first-high dose phase, in patients with moderate and severe COVID-19.

HCQ concentrations in the lungs have been reported to be significantly higher than the corresponding ones in plasma (Yao et al., 2020). Therefore, reaching HCQ blood concentrations of 1500 ng/ml assures that the EC50 concentrations are achieved within the main organ affected by the disease (Harapan et al., 2020). In addition, there is some evidence that the virus attacks red blood cells, rendering them incapable of transporting oxygen (Liu & Li, 2020) and T cells, decreasing their number significantly (Qin et al., 2020). Thus, a high loading dose providing an initial phase of high blood levels, followed by sparse small doses is anticipated to be both more efficacious and safer in comparison to a frequent low dose scheme. This can also be supported by both the pharmacokinetics of HCQ and by its time-dependent EC50, which resulted in 9-fold lower values after an incubation of 48 h, as compared to an incubation period of 24 h (Yao et al., 2020). Consequently, in a dosage regimen with a high initial dose followed by small maintenance doses, the viral load is expected to decrease during the initial phase, while the virus

becomes more susceptible to the drug’s concentration, allowing, therefore, for a gradual decrease of the dose to be administered.

Regarding the immunomodulatory effect of the drug, it has been found to be exerted with relatively low HCQ blood-levels (range 500–2000 ng/ml) (Durcan et al., 2015).

It’s worth mentioning that the aim of the present study was not to propose a cure for COVID-19. Instead, given that, HCQ is used both in the clinical setting and in clinical trials as an option for the management of COVID-19 with a ‘one-dose fits all’ approach, this study aimed to provide guidance on the dose to be selected depending on patient’s characteristics.

Other studies, where modeling and simulation approaches have been used in order to identify an optimized dosing scheme for HCQ in COVID-19 treatment suggest, as well, that a higher dose upon diagnosis will significantly benefit the patients (Arnold & Buckner, 2020; Fan et al., 2020; Garcia-Cremades et al 2020; Perinel et al., 2020; White et al., 2020; Yao et al., 2020). However, in these studies the currently applied dosing schemes were not investigated, while they focused primarily on efficacy or on safety aspects. In addition, patients’ characteristics were not taken into consideration and a specific dose was proposed for all cases. Only in the study of White et al., (2020) renal impairment and body weight were taken under consideration. The remarkable effect on HCQ’s blood concentrations was made evident and the investigators addressed the importance of dose adjustment per weight (White et al., 2020). In the present study, an integrated approach was used taking into consideration all the main aspects of treatment with HCQ, while the proposed dosing schemes were designed by taking into consideration patient’s weight, disease severity, and his/her HCQ clearance.

It is worth mentioning, that the significant advantage of a high loading dose has been demonstrated in an open-label, randomized, controlled trial including 150 patients with COVID-19 (Tang et al., 2020), while the safety of a sparse dosing strategy in order to avoid accumulation has also been indicated in a recently published study for chloroquine (Karalis et al., 2020), that exhibits similar chemical structure, mechanism of action and adverse effects with HCQ (Singh et al., 2020).

Conclusion

After an extensive literature survey and simulations performed, several dosing schemes of HCQ have been proposed for the treatment of COVID-19 in relation to patient’s weight and disease severity that could indicate patient’s viral load. A high initial dose followed by lower sparse doses seems to be the most appropriate approach to apply in this case, as it is postulated to be more effective and safer compared to small frequent doses. Indeed, these dosing schemes were designed aiming to lower the viral load both in blood and in the lungs, without allowing for HCQ accumulation that could lead to adverse effects. Despite their complexity in terms of clinical practice we believe that they may offer significant advantages to clinicians coping with COVID-19.




Figure 3. Simulated hydroxychloroquine concentrations (ng/ml) versus time (hours) in 500 patients receiving the proposed dosing schemes depending on disease severity, i.e., mild (I), moderate (II) or severe (III) for A: a patients' population of 30 kg, B: a patients' population of 50 kg, C: a patients' population of 70 kg, D: a patients' population of 90 kg, E: a patients' population of 110 kg. Red line: 2235 ng/ml, known to promote adverse effects to 30% of patients. Blue line: 1500 ng/ml, levels to achieve during the first doses. Green line: 500 ng/ml, minimum levels eliciting immunomodulatory effect.




Figure 4. Simulated hydroxychloroquine concentrations (ng/ml) versus time (hours) in 500 patients receiving the proposed dosing schemes depending on disease severity, i.e., mild (I), moderate (II) or severe (III) for A: a patients' population of 70 kg with 30% impaired clearance, B: a patients' population of 70 kg with 50% impaired clearance. Red line: 2235 ng/ml, known to promote adverse effects to 30% of patients. Blue line: 1500 ng/ml, levels to achieve during the first doses. Green line: 500 ng/ml, minimum levels eliciting immunomodulatory effect.

Disclosure statement

No potential conflict of interest was reported by the author(s).

ORCID

Eleni Karatza  <http://orcid.org/0000-0001-8406-4121>
 George Ismailos  <http://orcid.org/0000-0003-1466-4978>
 Markos Marangos  <http://orcid.org/0000-0001-5030-2398>
 Vangelis Karalis  <http://orcid.org/0000-0003-0492-0712>

Data availability statement

Data sharing is not applicable to this article as no new data were created or analyzed in this study.

References

- Abdulaziz N, Shah AR, McCune WJ. (2018). Hydroxychloroquine: balancing the need to maintain therapeutic levels with ocular safety: an update. *Curr Opin Rheumatol* 30:249–55.
- Al-Rawi H, Meggitt SJ, Williams FM, Wahie S. (2018). Steady-state pharmacokinetics of hydroxychloroquine in patients with cutaneous lupus erythematosus. *Lupus* 27:847–52.
- Alpern JD, Gertner E. (2020). Off-label therapies for COVID-19—are we all in this together? *Clin Pharmacol Ther* 108:182–4.
- Arnold S, Buckner F. (2020). Hydroxychloroquine for treatment of SARS-CoV-2 infection? Improving our confidence in a model-based approach to dose selection. *Clin Transl Sci* 13:642–5.
- Arshad S, Kilgore P, Chaudhry ZS, et al. (2020). Treatment with hydroxychloroquine, azithromycin, and combination in patients hospitalized with COVID-19. *Int J Infect Dis* 97:396–403.
- Balevic SJ, Green TP, Clowse M, et al. (2019). Pharmacokinetics of hydroxychloroquine in pregnancies with rheumatic diseases. *Clin Pharmacokinet* 58:525–33.
- Barbosa J, Kaitis D, Freedman R, Le K, Lin X. (2020) Clinical outcomes of hydroxychloroquine in hospitalized patients with COVID-19: a quasi-randomized comparative study. Submitted to the *New England Journal of Medicine*. (April 4th 2020); https://www.sefq.es/_pdfs/NEJM_Hydroxychloroquine.pdf [last accessed 29 Jul 2020].
- Briasoulis A, Agarwal V, Pierce WJ. (2011). QT prolongation and torsade de pointes induced by fluoroquinolones: infrequent side effects from commonly used medications. *Cardiology* 120:103–10.
- Carmichael SJ, Charles B, Tett SE. (2003). Population pharmacokinetics of hydroxychloroquine in patients with rheumatoid arthritis. *Ther Drug Monitor* 25:671–81.
- Casadevall A, Joyner MJ, Pirofski LA. (2020). SARS-CoV-2 viral load and antibody responses: the case for convalescent plasma therapy. *J Clin Invest* 139760. Advance online publication. doi: 10.1172/JCI139760. [Online ahead of print].
- Cavalcanti AB, Zampieri FG, Rosa RG, et al. (2020). Hydroxychloroquine with or without azithromycin in mild-to-moderate covid-19. *N Engl J Med*. Advance online publication. doi: 10.1056/NEJMoa2019014. [Online ahead of print].
- Chen Z, Hu J, Zhang Z, et al. (2020a) Efficacy of hydroxychloroquine in patients with COVID-19: results of a randomized clinical trial url: medRxiv preprint Posted April 10, 2020. (Pre-publication. Not peer-reviewed). *BMJ*. doi: 10.1101/2020.03.22.20040758. [Online ahead of print].
- Chen J, Liu D, Liu L, et al. (2020b). [A pilot study of hydroxychloroquine in treatment of patients with moderate COVID-19]. *Zhejiang Da Xue Xue Bao Yi Xue Ban* 49:215–9. 9292.2020.03.03.

- Cheng Y, Luo R, Wang K, et al. (2020). Kidney disease is associated with in-hospital death of patients with COVID-19. *Kidney Int* 97:829–38.
- Cortegiani A, Ingoglia G, Ippolito M, et al. (2020). A systematic review on the efficacy and safety of chloroquine for the treatment of COVID-19. *J Crit Care* 57:279–83.
- Olano J, Howland MA, Su MK, et al. (2019). Toxicokinetics of hydroxychloroquine following a massive overdose. *Am J Emerg Med* 37: 2264.e5–e8.
- Drożdżal S, Rosik J, Lechowicz K, et al. (2020). FDA approved drugs with pharmacotherapeutic potential for SARS-CoV-2 (COVID-19) therapy. *Drug Resist Updat* 53:100719. Advance online publication.
- Durcan L, Clarke WA, Magder LS, Petri M. (2015). Hydroxychloroquine blood levels in systemic lupus erythematosus: clarifying dosing controversies and improving adherence. *J Rheumatol* 42:2092–7.
- Elavarasi A, Prasad M, Seth T, et al. (2020). Chloroquine and hydroxychloroquine for the treatment of COVID-19: a systematic review and meta-analysis. *J General Internal Med.* 1–7. Advance online publication. hdoi: 10.1007/s11606-020-06146-w. [Online ahead of print].
- Fan J, Zhang X, Liu J, et al. (2020). Connecting hydroxychloroquine in vitro antiviral activity to in vivo concentration for prediction of antiviral effect: a critical step in treating COVID-19 patients. *Clin Infect Dis:ciaa623*. Advance online publication. doi: 10.1093/cid/ciaa623. [Online ahead of print].
- FDA (2020). U.S. Food and Drug Administration: memorandum explaining basis for revocation of emergency use authorization for emergency use of chloroquine phosphate and hydroxychloroquine sulfate. June 15 2020. Available from: <https://www.fda.gov/media/138945/download> [last accessed 29 Jul 2020].
- Furst DE. (1996). Pharmacokinetics of hydroxychloroquine and chloroquine during treatment of rheumatic diseases. *Lupus* 5(Suppl 1): S11–S15.
- Garcia-Cremades M, Solans BP, Hughes E, et al. (2020). Optimizing hydroxychloroquine dosing for patients with COVID-19: an integrative modeling approach for effective drug repurposing. *Clin Pharmacol Ther* 108:253–63.
- Gautret P, Lagier JC, Parola P, et al. (2020a). Hydroxychloroquine and azithromycin as a treatment of COVID-19: results of an open-label non-randomized clinical trial. *Int J Antimicrob Agents* 56:105949.
- Gautret P, Lagier JC, Parola P, et al. (2020b). Clinical and microbiological effect of a combination of hydroxychloroquine and azithromycin in 80 COVID-19 patients with at least a six-day follow up: a pilot observational study. *Travel Med Infect Dis* 34:101663.
- Haas NB, Appleman LJ, Stein M, et al. (2019). Autophagy inhibition to augment mTOR inhibition: a phase I/II trial of everolimus and hydroxychloroquine in patients with previously treated renal cell carcinoma. *Clin Cancer Res* 25:2080–7.
- Harapan H, Itoh N, Yufika A, et al. (2020). Coronavirus disease 2019 (COVID-19): a literature review. *J Infect Public Health* 13:667–73.
- Horby P, Mafham M, Linsell L, et al. (2020). Effect of hydroxychloroquine in hospitalized patients with COVID-19: preliminary results from a multi-centre, randomized, controlled trial. *medRxiv*. 2020.07.15.20151852; Posted July 15, 2020. (Pre-publication. Not peer-reviewed). *BMJ*. doi: 10.1101/2020.07.15.20151852. [Online ahead of print].
- Jankelson L, Karam G, Becker ML, et al. (2020). QT prolongation, torsades de pointes, and sudden death with short courses of chloroquine or hydroxychloroquine as used in COVID-19: a systematic review. *Heart Rhythm* 17:1472–9.
- Juurlink DN. (2020). Safety considerations with chloroquine, hydroxychloroquine and azithromycin in the management of SARS-CoV-2 infection. *Canad Med Assoc J* 192:E450–E453.
- Karalis V, Ismailos G, Karatza E. (2020). Chloroquine dosage regimens in patients with COVID-19: safety risks and optimization using simulations. *Safety Sci* 129:104842.
- Lavé T, Parrott N, Grimm HP, et al. (2007). Challenges and opportunities with modelling and simulation in drug discovery and drug development. *Xenobiotica* 37:1295–310.
- Lavielle M. (2019). mlxR: simulation of longitudinal data. R package version 4.0.6. <https://CRAN.R-project.org/package=mlxR> [last accessed on 29 Jul 2020].
- Lee JY, Vinayagamoorthy N, Han K, et al. (2016). Association of polymorphisms of cytochrome P450 2D6 with blood hydroxychloroquine levels in patients with systemic lupus erythematosus. *Arthritis Rheumatol* 68:184–90.
- Lee JY, Lee J, Ki Kwok S, et al. (2017). Factors related to blood hydroxychloroquine concentration in patients with systemic lupus erythematosus. *Arthritis Care Res* 69:536–42.
- Lim HS, Im JS, Cho JY, et al. (2009). Pharmacokinetics of hydroxychloroquine and its clinical implications in chemoprophylaxis against malaria caused by *Plasmodium vivax*. *Antimicrob Agents Chemother* 53: 1468–75.
- Liu W, Li H. (2020) COVID-19: attacks the 1-beta chain of hemoglobin and captures the porphyrin to inhibit human heme metabolism. *ChemRxiv*. Available from: https://chemrxiv.org/articles/COVID-19_Disease_ORF8_and_Surface_Glycoprotein_Inhibit_Heme_Metabolism_by_Binding_to_Porphyrin/11938173 [last accessed 29 Jul 2020]
- Liu Y, Yan LM, Wan L, et al. (2020a). Viral dynamics in mild and severe cases of COVID-19. *Lancet Infect Dis* 20:656–7.
- Liu J, Cao R, Xu M, et al. (2020b). Hydroxychloroquine, a less toxic derivative of chloroquine, is effective in inhibiting SARS-CoV-2 infection in vitro. *Cell Discov* 6:16.
- Lowe PJ, Hijazi Y, Luttringer O, et al. (2007). On the anticipation of the human dose in first-in-man trials from preclinical and prior clinical information in early drug development. *Xenobiotica* 37:1331–54.
- Marmor MF, Kellner U, Lai TY, et al. (2016). Recommendations on screening for chloroquine and hydroxychloroquine retinopathy (2016 revision). *Ophthalmology* 123:1386–94.
- Massachusetts General Hospital. 2020: Massachusetts General Hospital COVID-19 treatment guidance. Available from: <https://www.massgeneral.org/news/coronavirus/treatment-guidance> [last accessed 29 Jul 2020]
- Miller DR, Khalil SK, Nygard GA. (1991). Steady-state pharmacokinetics of hydroxychloroquine in rheumatoid arthritis patients. *DICP* 25:1302–5.
- Molina JM, Delaugerre C, Le Goff J, et al. (2020). No evidence of rapid antiviral clearance or clinical benefit with the combination of hydroxychloroquine and azithromycin in patients with severe COVID-19 infection. *Medecine et Maladies Infectieuses* 50:384.
- Morita S, Takahashi T, Yoshida Y, Yokota N. (2016). Population pharmacokinetics of hydroxychloroquine in Japanese patients with cutaneous or systemic lupus erythematosus. *Ther Drug Monitor* 38:259–67.
- Munster T, Gibbs JP, Shen D, et al. (2002). Hydroxychloroquine concentration-response relationships in patients with rheumatoid arthritis. *Arthritis Rheum* 46:1460–9.
- Naicker S, Yang CW, Hwang SJ, et al. (2020). The Novel Coronavirus 2019 epidemic and kidneys. *Kidney Int* 97:824–8.
- PAHO/WHO (2020). RAPID REVIEW – May 8th, 2020.COVID-19: chloroquine and hydroxychloroquine research. Available from: https://iris.paho.org/bitstream/handle/10665.2/52105/RAPID%20UPDATE%20REVIEW_Chloroquine%20May%2020b.pdf?sequence=5&isAllowed=y [last accessed 29 Jul 2020]
- Perinel S, Launay M, Botelho-Nevers É, et al. (2020). Towards optimization of hydroxychloroquine dosing in intensive care unit COVID-19 patients. *Clin Infect Dis* 2020 Apr 7;ciaa394. doi: 10.1093/cid/ciaa394. Online ahead of print.
- Plaquenil Label. (2017) Plaquenil® Hydroxychloroquine Sulfate Tablets, USP, US Food and Drug Administration (USFDA) Label. Revised on January 2017. Barbados (USA): Concordia Pharmaceuticals Inc. Available from: https://www.accessdata.fda.gov/drugsatfda_docs/label/2017/009768s037s045s047lbl.pdf
- Plaquenil Smpc. (2020). Plaquenil® Hydroxychloroquine sulfate 200 mg film-coated Tablets. Summary of Product characteristics (Smpc), revised on March 2020. Surrey (UK): Zentiva Pharma UK Limited. Available from: <https://www.medicines.org.uk/emc/product/1764/smpc>
- Pujadas E, Chaudhry F, McBride R, et al. (2020). SARS-CoV-2 viral load predicts COVID-19 mortality. *Lancet Respir Med* 8:e70.
- Qin C, Zhou L, Hu Z, et al. (2020). Dysregulation of immune response in patients with coronavirus 2019 (COVID-19) in Wuhan, China. *Clin Infect Dis* 71:762–8.

- Rangwala R, Leone R, Chang YC, et al. (2014). Phase I trial of hydroxychloroquine with dose-intense temozolomide in patients with advanced solid tumors and melanoma. *Autophagy* 10:1369–79.
- Ronco C, Reis T. (2020). Kidney involvement in COVID-19 and rationale for extracorporeal therapies. *Nat Rev Nephrol* 16:308–10.
- Rosenfeld MR, Ye X, Supko JG, et al. (2014). A phase I/II trial of hydroxychloroquine in conjunction with radiation therapy and concurrent and adjuvant temozolomide in patients with newly diagnosed glioblastoma multiforme. *Autophagy* 10:1359–68.
- Saleh M, Gabriels J, Chang D, et al. (2020). Effect of chloroquine, hydroxychloroquine, and azithromycin on the corrected QT interval in patients with SARS-CoV-2 infection. *Circ Arrhythm Electrophysiol* 13:e008662.
- Singh-Uttam T, Parida S, Lingaraju MC, et al. (2020). Drug repurposing approach to fight COVID-19. *Pharmacol Rep*;1–30. doi: [10.1007/s43440-020-00155-6](https://doi.org/10.1007/s43440-020-00155-6). [Online ahead of print].
- Singh AK, Singh A, Shaikh A, et al. (2020). Chloroquine and hydroxychloroquine in the treatment of COVID-19 with or without diabetes: a systematic search and a narrative review with a special reference to India and other developing countries. *Diabet Metabol Syndr* 14:241–6.
- Tang W, Cao Z, Han M, et al. (2020). Hydroxychloroquine in patients with mainly mild to moderate coronavirus disease 2019: open label, randomised controlled trial. *BMJ* 369:m1849.
- Teng C, Walter EA, Gaspar D, et al. (2019). Torsades de pointes and QT prolongation Associations with antibiotics: a pharmacovigilance study of the FDA adverse event reporting system. *Int J Med Sci* 16:1018–22.
- Tukacs M. (2018). Pharmacokinetics and extracorporeal membrane oxygenation in adults: a literature review. *AACN Adv Crit Care* 29:246–58.
- U.S. Food & Drug Administration. (2017) E14 clinical evaluation of QT/QTc interval prolongation and proarrhythmic potential for non-antiarrhythmic drugs – questions and answers (R3) guidance for industry. available at: <https://www.fda.gov/regulatory-information/search-fda-guidance-documents/e14-clinical-evaluation-qtqc-interval-prolongation-and-proarrhythmic-potential-non-antiarrhythmic-1>, FDA, Rockville, MD, USA
- Vogl DT, Stadtmauer EA, Tan KS, et al. (2014). Combined autophagy and proteasome inhibition: a phase 1 trial of hydroxychloroquine and bortezomib in patients with relapsed/refractory myeloma. *Autophagy* 10:1380–90.
- Wang Y, Zhang L, Sang L, et al. (2020). Kinetics of viral load and antibody response in relation to COVID-19 severity. *J Clin Invest*, 138759. Advance Online Publication. doi: [10.1172/JCI13875](https://doi.org/10.1172/JCI13875). [Online ahead of print].
- White NJ, Watson JA, Hoglund RM, et al. (2020). COVID-19 prevention and treatment: a critical analysis of chloroquine and hydroxychloroquine clinical pharmacology. *PLoS Med* 17:e1003252.
- WHO (2020), Clinical management of COVID-19. Interim Guidance (27 May 2020). Available from: <https://www.who.int/publications/i/item/clinical-management-of-covid-19> [last accessed 29 Jul 2020]
- Yang J, Wu M, Liu X, et al. (2020). Cytotoxicity evaluation of chloroquine and hydroxychloroquine in multiple cell lines and tissues by dynamic imaging system and PBPK model. (Pre-publication. Not peer-reviewed). *bioRxiv*. 2020.04.22.056762. doi: [10.1101/2020.04.22.056762](https://doi.org/10.1101/2020.04.22.056762). [Online ahead of print].
- Yao X, Ye F, Zhang M, et al. (2020). In vitro antiviral activity and projection of optimized dosing design of hydroxychloroquine for the treatment of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). *Clin Infect Dis* 71:732–9.
- Yu B, Wang DW, Li C. (2020) Hydroxychloroquine application is associated with a decreased mortality in critically ill patients with COVID-19. *medRxiv*. doi: 01 May 2020. Available from: <https://www.medrxiv.org/content/10.1101/2020.04.27.20073379v1> [last accessed August 2020]
- Zheng S, Fan J, Yu F, et al. (2020). Viral load dynamics and disease severity in patients infected with SARS-CoV-2 in Zhejiang province, China, January–March 2020: retrospective cohort study. *BMJ* 369:m1443.