

Thermostable vacuum foam dried Newcastle disease vaccine: Process optimization and pilot-scale study

Fang Lyu ^{1,2,4,5}, Yan-hong Zhao ^{1,5}, Xiao-xin Zuo ^{1,6}, Babalwa Nyide ², Bi-hua Deng ^{1,4,5}, Ming-xu Zhou ^{1, 4}, Jibo Hou ¹, Jia-jie Jiao ^{1,5}, Min-qian Zeng ^{1,5}, Hong-ying Jie¹, Ademola Olaniran ³, Yu Lu ^{1,4,5,7*}, Thandeka Khoza ^{2*}

¹ Institute of Veterinary Immunology & Engineering, National Research Center of Engineering and Technology for Veterinary Biologicals, Jiangsu Academy of Agricultural Sciences, Nanjing 210014, China

² Department of Biochemistry, School of Life Sciences, College of Agriculture, Engineering & Science, University of KwaZulu-Natal, Pietermaritzburg 3209, South Africa

³ Department of Microbiology, School of Life Sciences, College of Agriculture, Engineering & Science, University of KwaZulu-Natal, Durban 4000, South Africa

⁴ GuoTai (Taizhou) Center of Technology Innovation for Veterinary Biologicals, Taizhou 225300, China

⁵ School of Animal Medicine, Nanjing Agricultural University, Nanjing 210095, Jiangsu, China;

⁶ Jiangsu Key Laboratory for Food Quality and Safety-State Key Laboratory Cultivation Base, Ministry of Science and Technology, Nanjing 210014, China

⁷ School of Pharmacy, Jiangsu University, Zhenjiang, 212013, China

*Correspondence:

khozat1@ukzn.ac.za; Tel: +27 33 2606281

luyu@jaas.ac.cn (Y. Lu); Tel: +86-25-8439-2088

Supplementary information




Figure S1 The sampling design showing different positions of the selected NDV-VFD vaccine samples in the shelves. There are 3 shelves in the lab lyophilizer (0.27 m²). The colors represent the location and position of the samples that were selected for analysis.