


RESEARCH ARTICLE

A clinical method for estimating the modulus of elasticity of the human cornea *in vivo*

David C. Pye *

School of Optometry and Vision Science, University of New South Wales, Sydney, Australia

* d.pye@unsw.edu.au

Abstract

Background

To develop a method, using current clinical instrumentation, to estimate the Young's modulus of the human cornea *in vivo*.

Methods

Central corneal curvature (CCC), central corneal thickness (CCT), intraocular pressure (IOP) was measured with the Goldmann tonometer (IOPG) and the Pascal Dynamic Corneal Tonometer (PDCT) in one eye of 100 normal young human subjects (21.07 ± 2.94 years) *in vivo*. The Orssengo and Pye algorithm was used to calculate the Young's modulus of the corneas of these subjects.

Results


The Young's modulus (E) of the corneas of the subjects using the PDCT and IOPG results (E_{calc}) was 0.25 ± 0.10 MPa, and without the PDCT results (E_{iopg}) was 0.29 ± 0.06 MPa. The difference in these results is due to the difference in tonometry results between the two instruments, as the mean PDCT result for the subjects was 16.89 ± 2.49 mmHg and the IOPG result 15.06 ± 2.71 mmHg. E was affected by the CCC of the subjects but more particularly by the CCT and IOP measurements. Corneal stiffness results are also presented.

Conclusion

Two methods have been developed to estimate the Young's modulus of the human cornea *in vivo* using current clinical instrumentation. One method (E_{calc}) is applicable to the general corneal condition, and E_{iopg} to the normal cornea, and these results can be used to calculate corneal stiffness.

Introduction

There has been an increasing interest in the biomechanical behaviour of the eye [1], and especially of the cornea as demonstrated by Dupps and Roberts [2]. The biomechanical behaviour



OPEN ACCESS

Citation: Pye DC (2020) A clinical method for estimating the modulus of elasticity of the human cornea *in vivo*. PLoS ONE 15(1): e0224824. <https://doi.org/10.1371/journal.pone.0224824>

Editor: Sanjoy Bhattacharya, Bascom Palmer Eye Institute, UNITED STATES

Received: April 22, 2019

Accepted: October 22, 2019

Published: January 8, 2020

Copyright: © 2020 David C. Pye. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the manuscript and its Supporting Information files.

Funding: The authors received no specific funding for this work.

Competing interests: The authors have declared that no competing interests exist.

of the cornea may be particularly relevant in clinical practice, as the cornea may be able to be considered as a surrogate for what might be happening elsewhere in the eye[3]. Corneal biomechanics have been reported to be altered in myopic patients, and with the degree of myopia [4], be combined with baseline age to predict the rate of axial elongation in myopic children [5], be altered in diabetic patients[6], in patients using topical prostaglandins[7] and patients with keratoconus[8] and as a sensitive marker of the ocular activity of collagen vascular diseases[9]. As a result, the study of corneal biomechanics has emerged as a very hot topic for research in ophthalmology[10].

One of the key values for the biomechanical behaviour of a tissue is a measurement of the way the tissue behaves when subjected to an applied load (or stress). This gives rise to a stress/strain ratio called Young's modulus, and this is a key value to be incorporated into engineering models of the cornea which include corneal topography, intraocular pressure and other characteristics of the eye to enable the behaviour of the tissue to be better understood.

There have been a number of methods used to try to establish the biomechanical behaviour of the human cornea *ex vivo* including strip extensometry[11] and inflation testing[12,13,14]. More recent *ex vivo* techniques include supersonic shear imaging[15] radial shear speckle pattern interferometry[16], atomic force microscopy[17], corneal optical coherence elastography [18] and a Surface Force Apparatus[19].

There are two clinical instruments currently available to investigate the biomechanical behaviour of the cornea, the Ocular Response Analyzer and the Corvis ST. Both devices employ a puff of air of short duration to investigate corneal behaviour, but these instruments do not currently produce a measurement of Young's modulus of the tissue.

There are, however, techniques under development which may enable a value for corneal tissue elasticity to be obtained for the human cornea *in vivo*. These instruments include Brillouin optical microscopy[20,21], corneal indentation[22], estimation of Young's modulus based on a fluid-filled spherical shell model with Scheimpflug imaging[23] and ultrasound surface wave elastography[24].

However, many of these techniques are still experimental and not available in the clinical environment. There is a need to develop a method to estimate the modulus of elasticity of the cornea using currently available clinical instruments which most ophthalmic practitioners would already have available to them in their practices. Having this knowledge may enable practitioners to better understand corneal behaviour, and how the cornea may be affected by corneal surgical techniques or rigid contact lenses. But the application of this information could be helpful for tissue matching for corneal grafts, modelling of new surgical techniques, interpreting tonometry results and determining whether the cornea may be a surrogate for what may be happening elsewhere in the eye.

In this study it is investigated whether current standard clinical instrumentation may be used to estimate the Young's modulus of the human cornea *in vivo* using two methods: one to provide an accurate estimation in all corneas, and the other to provide an estimation in normal corneas.

Materials and methods

Subjects

One hundred subjects (100 eyes) aged between 17 and 30 years were recruited from the student and staff population of the University of New South Wales School of Optometry and Vision Science over a 3-month period, selected without a preference for gender or race. The subjects in this experiment were not the same subjects as used in the Hamilton and Pye (2008) [25] paper, and a different group of researchers collected the data. Subjects were examined and

excluded from the study if they had ocular or systemic disease, a history of ocular trauma or surgery, were using ocular medications or had corneal astigmatism $>2.00\text{D}$. Contact lens wearers of soft contact lenses were accepted into the study provided they had not worn lenses on the day of participation in the study.

The study was conducted in accordance with the declaration of Helsinki and approved by the Human Research Ethics Committee at the University of New South Wales. Written informed consent was obtained from all subjects after a full explanation of the nature of the study.

Procedure

Central corneal curvature (two readings, EyeChek autokeratometer, Reichert Ophthalmic Instruments, NY), applanation IOP (three readings, slit-lamp based Goldmann tonometer), Pascal DCT IOP (Swiss Microtechnology AG, Ziemer Ophthalmic Systems Group Co, Port Switzerland, three readings of Q3 or above) and ultrasonography CCT (three readings, BV International, Clermont-Ferrand, France) were measured in both eyes of each subject, with a maximum interval of 5 minutes between the use of each instrument. No measurements were taken within the first 2 hours of the subjects awakening to allow for overnight corneal swelling to dissipate, and the mean values for each instrument were used for analysis. Each procedure was conducted by a dedicated experienced independent observer who was masked from the measurements obtained from the other procedures. All results obtained for both eyes were statistically significantly correlated with $p < 0.0001$ using the paired, two tailed t-test. As a result, only the results for one eye (the left eye) are discussed in this paper in order to avoid statistical bias from the use of both eyes of the subjects.

The baseline demographics are summarized in [Table 1](#).

Calculations

The Orssengo-Pye algorithm[25,26] was used to calculate the elastic (or Young's) modulus of the cornea. These equations have been discussed in detail elsewhere, with the key equations being:

1. The Pascal DCT tonometer(PDCT) is a tonometer whose intraocular pressure results in the human eye appear to not be influenced by the material properties of the cornea of the eye being measured[27,28,29]. This is due to the design of the probe tip, and the Pascal DCT tonometry result is thought to be a better indication of the true intraocular pressure, and hence the PDCT result is used in the equations below as the true IOP.
2. B is the coefficient of the Goldmann tonometry result (IOPG) of the calibration cornea (no units) and C is the coefficient of the Pascal DCT tonometry result (IOPT) in the test cornea

Table 1. Baseline demographics.

Parameter	Mean (\pm SD)
No. of subjects	100
Male:Female	42:58
Age (years)	21.07 (\pm 2.94)
Central corneal curvature (mm)	7.75 (\pm 0.26)
Central corneal thickness (μm)	549.9 (\pm 32.8)
Pascal DCT IOP (mmHg)	16.89 (\pm 2.49)
Goldmann IOP (mmHg)	15.06 (\pm 2.71)

<https://doi.org/10.1371/journal.pone.0224824.t001>

(no units). R = average radius of curvature of the anterior cornea (mm); t = central corneal thickness (mm); ν = Poisson's ratio of the cornea = 0.49 (no units) and A = area of appplanation of the cornea (mm^2).

$$B = \frac{0.6\pi R \left(R - \frac{t}{2}\right) \sqrt{1 - \nu^2}}{t^2}$$

$$C = \frac{\pi R \left(R - \frac{t}{2}\right)^2 (1 - \nu)}{At}$$

3. IOPT_{calc} was calculated from the equation

$$IOPT_{calc} = \frac{IOPG}{(B_c - C_c + C)/B}$$

Where IOPT_{calc} is the calculated true IOP (mmHg); IOPG is the Goldmann tonometer intraocular pressure (mmHg); B_c is the coefficient of IOPG of the calibration cornea (no units); C_c is the coefficient of the calibration cornea (no units); B is the coefficient of IOPG in the test cornea (no units); and C is the coefficient of IOPT in the test cornea (no units).

4. Young's modulus (E_{calc} , MPa) was calculated *in vivo* using the Pascal DCT tonometry value (IOPT, mmHg) in the following equation.

$$E_{calc} = \frac{(B \cdot IOPG - C \cdot IOPT)}{7500}$$

5. Eiopg was calculated using the above equation but using the Orssengo and Pye calculated IOPT value (IOPT_{calc}), and hence Step 4 enabled E_{calc} and Eiopg to be calculated for the 100 subjects.
6. Step 3 enabled E_{calc} and Eiopg to be plotted against IOPG and IOPT separately.
7. As the values for E_{calc} and IOPG were normally distributed, a linear regression and Pearson correlation coefficient was determined ($r = 0.749$, $p < 0.0001$). The linear regression equation was used to then determine the error in IOPG at differing CCTs.
8. The Orssengo and Pye algorithm was used to determine the predicted effect of CCT of the subjects on the Eiopg results.

Statistics

GraphPad Prism (Graph Pad) version 7.02 and Excel (Excel for Office 365) was used to perform the statistical analysis. All tests were parametric and two-tailed, with statistical significance set at $p < 0.05$.

Results

Distribution of Young's modulus

The mean value and standard deviation of E_{calc} for the subject group was 0.25 ± 0.10 MPa and for Eiopg was 0.29 ± 0.06 MPa. The data for both values is normally distributed and the frequency distribution is shown in Fig 1.




Fig 1. Distribution of Young's modulus in 100 young healthy eyes as calculated using the Orssengo and Pye formula and Goldmann tonometry results only (Eiopg) or Goldmann and Pascal DCT results combined (Ecalc).

<https://doi.org/10.1371/journal.pone.0224824.g001>

Intercorrelation of parameters

A summary of the intercorrelations of parameters is shown in Table 2.

CCC was negatively correlated with Ecalc ($r = -0.199$, $p = 0.0467$) and Eiopg ($r = -0.197$, $p = 0.0492$) but not IOPG, PDCT or IOPTcalc.

There was no significant Pearson correlation between IOPG and CCT ($r = 0.010$, $p = 0.921$) as shown in Fig 2A, nor IOPG and central corneal curvature (CCC) ($r = -0.169$, $p = 0.093$) as

Table 2. Significance of Pearson correlations for the variables (* = $p < 0.05$, ** = $p < 0.001$).

	CCC (mm)	IOPG (mmHg)	PDCT (mmHg)	IOPTcalc (mmHg)	CCT (mm)	Ecalc (MPa)	Eiopg (MPa)
CCC (mm)	-----	ns	ns	ns	*	*	*
IOPG (mmHg)	ns	-----	**	**	ns	**	**
PDCT (mmHg)	ns	**	-----	**	ns	ns	**
IOPTcalc (mmHg)	ns	**	**	-----	**	**	**
CCT (mm)	*	ns	ns	**	-----	**	**
Ecalc (MPa)	*	**	ns	**	**	-----	**
Eiopg (MPa)	*	**	**	**	**	**	-----

<https://doi.org/10.1371/journal.pone.0224824.t002>





Fig 2. Correlation between Goldmann IOP and (a) central corneal thickness ($r = 0.010$, $p = 0.921$); (b) central corneal curvature ($r = -0.169$, $p = 0.092$).

<https://doi.org/10.1371/journal.pone.0224824.g002>

shown in Fig 2B. IOPG was significantly correlated with PDCT ($r = 0.5969$, $p < 0.0001$), Ecalc ($r = 0.7468$, $p < 0.0001$), Eiopg ($r = 0.900$, $p < 0.0001$) and IOPTcalc ($r = 0.900$, $p < 0.0001$).

The PDCT results were significantly correlated with Eiopg ($r = 0.526$, $p < 0.0001$) and IOPTcalc ($r = 0.526$, $p < 0.0001$) but not with CCT, CCC or Ecalc results, suggesting that the PDCT IOP is importantly independent of these three variables.

There was a positive correlation between IOPG and PDCT ($r = 0.598$, $p < 0.001$, Fig 3A), PDCT and IOPTcalc ($r = 0.525$, $p < 0.001$, Fig 3B), with the mean PDCT result higher than the IOPTcalc by 1.99mmHg.

IOPTcalc was positively correlated with IOPG ($r = 0.900$, $p < 0.0001$), and negatively correlated with PDCT ($r = 0.526$, $p < 0.0001$) and CCT ($r = -0.413$, $p < 0.001$), but not correlated with CCC.

CCT was positively correlated with CCC ($r = 0.230$, $p < 0.05$), negatively correlated with IOPTcalc ($r = -0.4131$, $p < 0.0001$), Ecalc ($r = -0.502$, $p < 0.0001$) and Eiopg ($r = -0.409$, $p < 0.0001$), but not correlated with IOPG or PDCT.

Young's modulus (Ecalc) was positively correlated with IOPG ($r = 0.749$, $p < 0.001$, Fig 4A). Ecalc was also correlated to CCT ($r = -0.502$, $p < 0.001$, Fig 4B) and weakly to corneal curvature ($r = -0.199$, $p = 0.05$, Fig 4C). The latter correlation should be viewed with suspicion, as this relationship was not found in the contralateral eye ($r = -0.11$, $p = 0.27$). Ecalc was correlated with Eiopg ($r = 0.890$, $p < 0.001$, Fig 4D), but there is proportional bias as shown in the Bland Altman plot[27] in Fig 5. However, Ecalc was not correlated with PDCT ($r = 0.093$, $p = 0.36$).

Eiopg was positively correlated with PDCT ($r = 0.526$, $p < 0.001$), IOPTcalc ($r = 0.998$, $p < 0.001$), Ecalc ($r = 0.890$, $p < 0.0001$) and IOPG ($r = 0.900$, $p < 0.001$) and negatively correlated with CCC ($r = -0.197$, $p = 0.049$) and CCT ($r = -0.409$, $p < 0.0001$).

Discussion

1. CCT vs CCC

The association between CCT and CCC was not strong, but was statistically significant in both eyes of the subjects. The finding that thicker corneas tended to have flatter CCCs has also been reported by Kiely et al[28], Hovding[29], and Tomidokoro[30]. Studies reported by Lowe[31], Tomlinson[32] and Hamilton and Pye[25] found no correlation between CCT and CCC, and their study was conducted with similar instrumentation and similar population to that reported in this study.

2. Goldmann vs Pascal DCT tonometry

The tonometer recognised by the International Standards Organisation[33] as the reference tonometer is the Goldmann tonometer, and yet in 1993 Whitacre and Stein[34] identified as many as 26 potential sources of error with this tonometry technique. The PDCT tonometer claims to be independent of corneal properties due to the design of the footplate and contour matching of the corneal front surface, and the results with this tonometer have been shown to be in good agreement with manometer studies conducted *ex vivo* and *in vivo*[35,36,37].

In 19 of the subjects, the difference in IOP measured by the Goldmann and PDCT was ≥ 4 mmHg and, in most instances, PDCT IOP was greater than IOPG, and this difference contributed to the larger spread of results for Ecalc as shown in Fig 1.





Fig 3. Correlation between Pascal DCT and (a) Goldmann IOP ($r = 0.597$, $p < 0.001$); (b) calculated IOPT ($r = 0.525$, $p < 0.001$).

<https://doi.org/10.1371/journal.pone.0224824.g003>

If the Goldman and PDCT tonometers are accurate measures of a patient's true IOP, it would be expected that their tonometry results would be equivalent for the mean CCT and CCC ie the average eye values for this particular subject group. However, the PDCT values were 1.8mmHg higher than the those of IOPG. This discrepancy has been noted in the past, and previous studies have found PDCT values to be higher than IOPG by 0.8mmHg[38], 1mmHg[39], 1.7mmHg[40], 2.3mmHg[41], 2.34mm Hg[42] and 3.98mmHg[43]. This difference in results between the two instruments has been found to be linked to the corneal resistance factor measured by the Ocular Response Analyzer by Kotecha et al[41], but another possible cause for this IOP measurement difference might be the surface tension developed at the edge of the Goldmann prism when measuring IOP, which should not apply to the PDCT technique. There are published works which suggest that the surface tension of the tear film for the GAT probe might vary from 0.11mmHg[44,45], 0.26mmHg[46] 4.15mmHg[47], 4.67mmHg[48], to 4.7mmHg[49] and hence a difference of 1.9mmHg between the two tonometry techniques could be as a result of surface tension acting during the IOPG measurement.

Hamilton and Pye[38] (2008) felt that the difference in mean values between the Pascal DCT and Goldmann tonometers could possibly be explained by slight calibration differences between the two instruments, interobserver variations or population differences. The difference between the IOP measurements determined by Goldmann and PDCT were not correlated with CCC ($r = -0.07$, $p = 0.55$).

3. Young's modulus of the cornea *in vivo*:

In this paper, the calculation of Young's modulus is based upon the cornea acting as a linear elastic material on the basis of a small applied load (around 1.5gf) and a small corneal indentation by the Goldmann tonometry probe of approximately 150 μ m.

There are two values for Young's modulus presented in this paper. Ecalc is based on the clinical measurements of a patient's CCC, CCT, IOPG and PDCT, and it is assumed that the PDCT result is a true measure of IOP and that IOPG is a measure of the cornea's response to a specific applied load.

Eiopg was calculated from the CCC, CCT and IOPG results, and is based on the assumption that IOPG is equivalent to IOPT when the cornea has average values for CCT and CCC (called the calibration cornea (Orssengo and Pye)[26]. As CCC has little effect on IOPG results, Eiopg can be calculated in clinical practice from the CCT and IOPG results alone.

The results for Eiopg and Ecalc are similar to those reported in a similar cohort of subjects reported in other papers and these results are shown in Table 3 for comparison.

Ecalc and Eiopg are strongly correlated ($r = 0.890$, $p < 0.001$) but Ecalc was, on average, 0.04MPa less than Eiopg. Hence the results are not interchangeable but, in the clinical situation, Eiopg may be a useful indicator of Young's modulus when determined with healthy normal corneas. However, if the IOPG result is likely to be significantly affected by the physiological state of the cornea such as oedema[50,51], refractive surgery[52,53] or other causes, Ecalc should provide a more valid result.

In the young healthy adults reported in this study, it was surprising that there was no apparent significant effect of CCT on IOPG as shown in Fig 2, but Ecalc had a significant effect upon IOPG as shown in Fig 4A, with Ecalc decreasing as CCT increases. By using the data from this study and the Orssengo and Pye algorithms[26], the effects of Ecalc and CCT upon IOPG can be calculated, and are shown in Fig 6. Over the 200 μ m range of corneal thicknesses







Fig 4. Correlation between Ecalc and (a) Goldmann IOP ($r = 0.748$, $p < 0.001$); (b) central corneal thickness ($r = -0.502$, $p < 0.001$); (c) central corneal curvature ($r = -0.199$, $p = 0.047$); (d) Eiopg ($r = 0.890$, $p < 0.001$).

<https://doi.org/10.1371/journal.pone.0224824.g004>

presented in Fig 6, the total effect of central corneal thickness on IOPG was estimated as 8.9mmHg, and of Ecalc was 11.0mmHg. It is apparent that Ecalc has a greater effect on IOPG measurement error than CCT, and this confirms a similar finding published by Hamilton and Pye[20]. In the young healthy normal subject group of this study, the measurement error in IOPG predicted by changes in CCT are negated by changes in Ecalc, supporting the predictions regarding the significant effect of Young's modulus on IOPG made by Liu and Roberts [54].

The above effect of Ecalc on IOPG could be related to the behaviour of the number of layers of corneal collagen or the effect of hydration upon the biomechanical behaviour of the cornea and the subsequent effects on IOPG as discussed by Hatami-Marbini and Etebu[55].

However, these results appear to be related the structural stiffness of the cornea, which has been described by Palko and Liu[56] by the equation $k = t.E$, where k = stiffness, t = the central corneal thickness and E is the Young's modulus of the tissue. If k is calculated using Eiopg and CCT, there is no correlation between CCT and k as shown in Fig 7A, and $k = 0.16 \pm 0.03$ N/mm. If k is calculated using Ecalc, there is significant relationship between the two variables ($p < 0.001$), but there is considerable scatter of the data as shown in Fig 7B.




Fig 5. Bland Altman plot of the relationship between Ecalc and Eiopg demonstrating proportional bias.

<https://doi.org/10.1371/journal.pone.0224824.g005>

Table 3. Published values for the Young’s modulus for human corneas of human subjects of similar age to those in this study.

Author	Technique	In vivo or ex vivo	Number of subjects	Age of subjects (years)	Young’s modulus (MPa) (\pm SD)
Hamilton and Pye 2008 ²⁵	Clinical measurements and Orssengo-Pye algorithm	In vivo	100	22.0 \pm 2.9	Eiopg 0.29 \pm 0.06
Knox Cartwright et al 2011 ¹⁶	Radial shearing speckle pattern interferometer	Ex vivo	1	24	0.28
Lam et al 2015 ²²	Corneal indentation	In vivo	29	23.4 \pm 1.7	Etangent 0.76 \pm 0.16
Shih et al 2015 ²³	Corvis ST based on water-filled spherical diaphragm dynamics	In vivo	10	25.2 \pm 2.15	0.44 \pm 0.37
Sit et al 2017 ²⁴	Ultrasound surface wave elastography	In vivo	20	51.4 \pm 7.2	0.69 \pm 0.11
Pye 2019	Clinical measurements Orssengo-Pye algorithm	In vivo	100	21.1 \pm 2.0	Ecalc 0.25 \pm 0.10 Eiopg 0.29 \pm 0.06

<https://doi.org/10.1371/journal.pone.0224824.t003>




Fig 6. Figure demonstrates the actual Goldmann tonometry error obtained in this subject group (Net Result), the Goldmann tonometry error calculated from E iopg (Effect of E) and from the error due to central corneal thickness as calculated from the Orssengo and Pye (Effect of CCT).

<https://doi.org/10.1371/journal.pone.0224824.g006>





Fig 7. Correlation between corneal stiffness as calculated from Eiopg and (a) central corneal thickness ($r = -0.123$, $p = 0.224$); (b) Ecalc and central corneal thickness ($r = -0.388$, $p < 0.001$).

<https://doi.org/10.1371/journal.pone.0224824.g007>

The corneal stiffness results suggest that applanation tonometry results in young healthy normals are influenced by the corneal stiffness ie the CCT and E combined rather than one variable alone. Whether this relationship applies in a similar fashion for subjects in older age groups is yet to be investigated.

The value for k of 0.16N/mm is greater than the value of 0.079N/mm obtained by Lam et al [22] who used a corneal indentation device. Whilst Lam et al [22] used a similarly aged subject cohort, their sample size was considerably smaller than in this study, and the cornea was indented to depth of 1mm , which is greater than that which occurs during Goldmann tonometry. It might also be possible that the anterior cornea is stiffer than the cornea as a whole, and this could also contribute to the discrepancy in values.

As can be seen from Fig 8A, there is a very strong relationship between corneal stiffness (calculated from Eiopg and CCT) and the Goldmann tonometry values. This strong correlation may be expected as the IOPG results were used in the calculation of these corneal stiffness values, but this result indicates the strength of the relationship between corneal stiffness and IOPG. There is a similarly strong correlation between IOPG and corneal stiffness calculated using Ecalc ($r = 0.801$, $p < 0.001$ see Fig 8B).

Conclusion

This paper presents a linear elastic model to estimate the Young's modulus of the human cornea *in vivo*, although the cornea is thought to be anisotropic, nonlinear, viscoelastic [57] and hyperelastic [58]. However, many of the techniques to obtain these results above are performed *ex vivo* and do not use Goldmann tonometry, which applies a relatively small force (1.5 gf) over an area of 7.35mm^2 , with a small degree of central anterior corneal deformation (approximately $150\mu\text{m}$ when the measurement is taken), and the force is applied over a longer time period than the non-contact tonometers.

This paper presents two methods for estimating the Young's modulus of the cornea *in vivo* which involve the use of currently available clinical instrumentation. The most accurate method is Ecalc which involves the use of the Pascal DCT as well as IOPG, and relies upon the Pascal DCT providing a more accurate measurement of the true IOP in all instances.

There is a strong correlation between Ecalc and Eiopg in young healthy adults, although the results are not interchangeable and, as CCC has no effect on Eiopg, it may be feasible to calculate Eiopg from CCT and IOPG alone in patients with normal corneas. As CCT and IOPG are often measured routinely in clinical practice, this would enable Eiopg to be calculated relatively easily in the clinical setting.

Young's modulus is an important descriptor of the mechanical behaviour of a material and being able to estimate this value for the individual patient in clinical practice may enable further developments in corneal modelling, biomechanics and bioengineering (such as the development of artificial corneas). It may also provide further information regarding improvements to corneal surgery, comparison of tonometers and tonometry methods, methods for altering anterior surface corneal topography (surgery and contact lenses), corneal aging, early diagnosis of corneal conditions (such as keratoconus) and the effects of topical drugs on corneal behaviour and IOP measurement. The behaviour of the cornea may also be a surrogate for what is happening elsewhere in the eye, such as the management and diagnosis of glaucoma, myopia development and the effects of systemic diseases on the eye, and a number of these issues are currently being investigated.





Fig 8. Correlation between corneal stiffness and IOPG calculated from (a) Eiopg ($r = 0.989$, $p < 0.001$); (b) Ecalc ($r = 0.801$, $p < 0.001$).

<https://doi.org/10.1371/journal.pone.0224824.g008>

These results suggest that corneal stiffness affects applanation tonometry results, and that both CCT and E combine to affect the findings. As such, it would be unwise to consider applying an IOPG correction factor based on only one of these two variables in subjects of this age group.

Supporting information

S1 Dataset. Data for all of the subjects in both eyes is included in this dataset.
(XLSX)

Acknowledgments

The author wishes to acknowledge Kirsten Hamilton for her assistance with the ethics proposal and especially Nguyet Nguyen, John Nguyen and Jane Le for their assistance in the collection of the data.

Author Contributions

Conceptualization: David C. Pye.

Formal analysis: David C. Pye.

Investigation: David C. Pye.

Methodology: David C. Pye.

Project administration: David C. Pye.

Resources: David C. Pye.

Software: David C. Pye.

Supervision: David C. Pye.

Writing – original draft: David C. Pye.

Writing – review & editing: David C. Pye.

References

1. Girard MJ, Dupps WJ, Baskaram M, Scarcelli G, Yun SH, Quigley HA et al. Translating ocular biomechanics into clinical practice: current state and future prospects. *Curr Eye Res* 2015; 40:1–18. <https://doi.org/10.3109/02713683.2014.914543> PMID: 24832392
2. Dupps WJ, Roberts CJ. Corneal biomechanics: a decade later. *J Cataract Ref Surg* 2014; 40:857.
3. Kotecha A. What biomechanical properties of the cornea are relevant to the clinician? *Surv Ophthalmol* 2007; 52 Suppl 2:S109–14.
4. Hon Y, Chen G-Z, Lu S-H, Lam DCC, Lam AKC. High myopes have lower normalised corneal tangent moduli (less 'stiff' corneas) than low myopes. *Ophthalmic Physiol Opt* 2016; 37:42–50. <https://doi.org/10.1111/opo.12335> PMID: 27873338
5. Wan K, Cheung SW, Wolffsohn JS, Orr JB, Cho P. Role of corneal biomechanical properties in predicting of speed of myopia progression in children wearing orthokeratology lenses or single-vision spectacles. *BMJ Open Ophth* 2018; 3:e000204. <https://doi.org/10.1136/bmjophth-2018-000204> eCollection 2018. PMID: 30687783
6. Ramm L, Herber R, Spoerl E, Pillunat LE, Terai N. Measurement of corneal biomechanical properties in diabetes mellitus using the Ocular Response Analyzer and the Corvis ST. *Cornea* 2019; 5:595–99.
7. Wu N, Chen Y, Yu X, Li M, Wen W, Sun X. Changes in corneal biomechanical properties after long-term topical prostaglandin therapy. *PLoS One*. 2016 May 17; 11(5):e0155527. <https://doi.org/10.1371/journal.pone.0155527> eCollection 2016. PMID: 27187282

8. Zhao Y, Shen Y, Yan Z, Tian M, Zhao J, Zhou X. Relationship among stiffness, thickness, and biomechanical parameters measured by the Corvis ST, Pentacam and ORA in keratoconus. *Front Physiol*. 2019 Jun 13; 10:740. <https://doi.org/10.3389/fphys.2019.00740> eCollection 2019. PMID: 31263429
9. Mahendradas P, Francis M, Vala R, Poornachandra B, Kawali A, Sheythy R et al. Quantification of ocular biomechanics in ocular manifestations of systemic autoimmune diseases. *Ocul Immunol Inflamm*. 2018 Aug 7:1–11. <https://doi.org/10.1080/09273948.2018.1501491> [Epub ahead of print]. PMID: 30084704
10. Luz A, Faria-Correia F, Salomao MQ, Lopes BT, Ambrosio R Jr. Corneal biomechanics: Where are we? *J Curr Ophthalmol* 2016;97–8. <https://doi.org/10.1016/j.joco.2016.07.004> PMID: 27579450
11. Woo SL, Kobayashi AS, Schlegel WA, Lawrence C. Nonlinear material properties of intact cornea and sclera. *Exp Eye Res* 1972; 14:29–39. [https://doi.org/10.1016/0014-4835\(72\)90139-x](https://doi.org/10.1016/0014-4835(72)90139-x) PMID: 5039845
12. Smolek MK. Holographic interferometry of intact and radially excised human eye-bank corneas. *J Cataract Refract Surg* 1994; 20: 277–86. [https://doi.org/10.1016/s0886-3350\(13\)80578-0](https://doi.org/10.1016/s0886-3350(13)80578-0) PMID: 8064603
13. Hjortdahl JO. Biomechanical studies of the human cornea. *Acta Ophthalmol Scand* 1995; 77:364–5.
14. Elsheikh A, Wang D, Pye D. Determination of the modulus of elasticity of the human cornea. *J Refract Surg* 2007; 23:808–18. PMID: 17985801
15. Tanter M, Touboul D, Gennisson JL, Bercoff J, Fink M. high-resolution quantitative imaging of corneal elasticity using supersonic shear imaging. *IEE Trans Med Imag* 2009; 28:1881–93.
16. Knox Cartwright NE, Tyrer JR, Marshall J. Age-related differences in the elasticity of the human cornea. *Invest Ophthalmol Vis Sci* 2011;4324–9. <https://doi.org/10.1167/iovs.09-4798> PMID: 20847118
17. Last JA, Thomasy SM, Croasdale CR, Russell P, Murphy CJ. Compliance profile of the human cornea as measured by atomic force microscopy. *Micron* 43 2012;1293–98. <https://doi.org/10.1016/j.micron.2012.02.014> PMID: 22421334
18. Ford MR, Sinha Roy A, Rollins AM, Dupps WJ Jr. serial biomechanical comparison of edematous, normal, and collagen crosslinked human donor corneas using optical coherence tomography. *J Cataract Refract Surg* 2014; 40:1041–7. <https://doi.org/10.1016/j.jcrs.2014.03.017> PMID: 24767794
19. Zappone B, Patil NJ, Lombardo M, Lombardo G. Transient viscous response of the human cornea probed with the surface force apparatus. *PLoS One*. 2018 May 25; 13(5):e0197779. <https://doi.org/10.1371/journal.pone.0197779> eCollection 2018. PMID: 29799859
20. Scarcellis G, Yun SH. *In vivo* Brillouin optical microscopy of the human eye. *Opt Express* 2012; 20:9197–202. <https://doi.org/10.1364/OE.20.009197> PMID: 22513631
21. Yun SH, Chernyak D. Brillouin microscopy: assessing ocular tissue biomechanics. *Curr Opin Ophthalmol* 2018; 29:299–305. <https://doi.org/10.1097/ICU.0000000000000489> PMID: 29771749
22. Lam AK, Hon Y, Leung LK, Lam DC. Repeatability of a novel corneal indentation device for corneal biomechanical measurement. *Ophthalmic Physiol Opt* 2015; 35:455–61. <https://doi.org/10.1111/opo.12219> PMID: 26094834
23. Shih P-J, Cao H-J, Hunag C-J, Wang I-J, Yen J-Y. A corneal elastic dynamic model derived from Scheimpflug imaging technology. *Ophthalmic Physiol Opt* 2015; 35:663–72. <https://doi.org/10.1111/opo.12240> PMID: 26353939
24. Sit AJ, Lin S-C, Kazemi A, McLaren JW, Pruet CM. *In vivo* noninvasive measurement of Young's modulus in human eyes: a feasibility study. *J Glaucoma* 2017; 26:967–73. <https://doi.org/10.1097/IJG.0000000000000774> PMID: 28858155
25. Hamilton KE, Pye DC. Young's modulus in normal corneas and the effect on applanation tonometry. *Optom Vis Sci* 2008; 6:445–50.
26. Orssengo G, Pye DC. Determination of the true intraocular pressure and modulus of elasticity of the human cornea *in vivo*. *Bull Math Biol* 1999; 61:551–72. <https://doi.org/10.1006/bulm.1999.0102> PMID: 17883231
27. Bland JM, Altman DG. Statistical methods for assessing agreement between two methods of clinical measurement. *Lancet* 1986; Feb 8; 1(8476):307–10. PMID: 2868172
28. Kiely PM, Carney LG, Smith G. Diurnal variation of corneal topography and thickness. *Am J Optom Physiol Optics* 1982; 59:976–82.
29. Hovding G. A clinical study of the association between thickness and curvature of the central cornea. *Acta Ophthalmol (Copenhagen)* 1983; 61:461–6.
30. Tomidokoro A, Araie M, Iwase A; The Tajimi Study Group. Corneal thickness and relating factors in a population-based study in Japan: The Tajimi Study. *Am J Ophthalmol* 2007; 144:152–4. <https://doi.org/10.1016/j.ajo.2007.02.031> PMID: 17601447
31. Lowe RF. Central corneal thickness. *Brit J Ophthalmol* 1969; 53:824–6.

32. Tomlinson A. A clinical study of the central and peripheral thickness and curvature of the central cornea. *Acta Ophthalmol (Copenhag)* 1972; 50:73–82.
33. International Standards Organisation (2009): Ophthalmic Instruments—Tonometers (ISO 8612:2009). Geneva: International Standard Organization.
34. Whitacre MM, Stein R. Sources of error with use of Goldmann-type tonometers. *Surv Ophthalmol* 1993; 38:1–30. [https://doi.org/10.1016/0039-6257\(93\)90053-a](https://doi.org/10.1016/0039-6257(93)90053-a) PMID: 8235993
35. Kniestedt C, Nee M, Stamper RL. Dynamic contour tonometry. *Arch Ophthalmol* 2004; 122:1287–93. <https://doi.org/10.1001/archophth.122.9.1287> PMID: 15364707
36. Kniestedt, Lin S, Chloe J, Bostrom A, Nee M, Stamper RL. Clinical comparison of contour and applanation tonometry and their relationship to pachymetry. *Arch Ophthalmol* 2005; 123:1532–37. <https://doi.org/10.1001/archophth.123.11.1532> PMID: 16286615
37. Kniestedt C, Nee M, Stamper RL. Accuracy of dynamic contour tonometry compared with applanation tonometry in human cadaver eyes of different hydration states. *Graefes Arch Clin Exp Ophthalmol* 2005; 243:359–66. <https://doi.org/10.1007/s00417-004-1024-6> PMID: 15864625
38. Hamilton KE, Pye DC, Kao L, Pham N, Tran A-Q N. The effect of corneal edema on dynamic contour and Goldmann tonometry. *Optom Vis Sci* 2008; 86:451–6.
39. Pasche M, Wilmsmeyer S, Lautebach S, Funk K. Dynamic contour tonometry versus Goldmann applanation tonometry: a comparative study. *Graefes Arch Clin Exp Ophthalmol* 2005; 24:763–7.
40. Kaufman C, Bachmann LM, Thiel MA. Comparison of dynamic contour tonometry with goldmann applanation tonometry. *Invest Ophthalmol Vis Sci* 2004; 45:3118–21. <https://doi.org/10.1167/iovs.04-0018> PMID: 15326129
41. Kotecha A, White E, Schlottmann PG, Garway-Heath DF. Intraocular pressure measurement precision with the Goldmann applanation, dynamic contour, and ocular response analyzer tonometers. *Ophthalmology* 2010; 117:730–7. <https://doi.org/10.1016/j.ophtha.2009.09.020> PMID: 20122737
42. Schneider E, Grehn F. Intraocular pressure measurement-comparison of dynamic contour tonometry and Goldmann applanation tonometry. *J Glaucoma* 2006; 15:2–6. <https://doi.org/10.1097/01.jgg.0000196655.85460.d6> PMID: 16378009
43. Andreanos K, Koutsandrea C, Papconstantinou D, Kotoulas A, Dimitrakis P, Moschos MM. Comparison of Goldmann applanation and pascal dynamic contour tonometry in relation to central corneal thickness and corneal curvature. *Clin Ophthalmol* 2016; 10:2477–84. <https://doi.org/10.2147/OPHT.S115203> PMID: 28003737
44. Elsheikh A, Wang D. Numerical modelling of corneal biomechanical behaviour. *Comput Methods in Biomech Biomed Engin* 2007; 10:85–95.
45. Elsheikh A, Wang D, Kotecha A, Brown M, Garway-Heath D. Evaluation of Goldman applanation tonometry using a nonlinear finite element ocular model. *Annals Biomed Eng* 2006; 34:1628–40.
46. Osborne SF, Williams R, Batterbury M, Wong D. Does the surface property of a disposable applanation tonometer account for its underestimation of intraocular pressure when compared with the Goldmann tonometer? *Graefes Arch Clin Exp Ophthalmol* 2007; 245:555–9. <https://doi.org/10.1007/s00417-006-0380-9> PMID: 16912886
47. Schwartz NJ, Mackay RS, Sackman JR. A theoretical and experimental study of the mechanical behaviour of the cornea with the application to the measurement of intraocular pressure. *Bull Math Biophys* 1966; 28: 585–631.
48. Chihara E. Assessment of true intraocular pressure.: the gap between theory and practical data. *Surv Ophthalmol* 2008; 53:203–18. <https://doi.org/10.1016/j.survophthal.2008.02.005> PMID: 18501267
49. Kwon TH, Ghaboussi J, Pecknold DA, Hashash YM. Effect of cornea material stiffness on measured intraocular pressure. *J Biomech* 2008; 41:1707–13. <https://doi.org/10.1016/j.jbiomech.2008.03.004> PMID: 18455173
50. Kaufman HE. Pressure measurement: which tonometer? *Invest Ophthalmol* 1972; 11:80–85. PMID: 5009106
51. Simon G, Small RH, Ren Q, Parel J-M, Ing ETS. Effect of corneal hydration on Goldmann applanation tonometry and corneal topography. *Refract Corneal Surg* 1993; 9:110–17. PMID: 8494810
52. Schneider E, Grehn F. Intraocular pressure measurement-comparison of dynamic contour tonometry and Goldmann applanation tonometry. *J Glaucoma* 2006; 15:2–6. <https://doi.org/10.1097/01.jgg.0000196655.85460.d6> PMID: 16378009
53. Sadigh AL, Fouladi RF, Hashemi H, Beheshtnejad AH. A comparison between Goldmann applanation tonometry and dynamic contour tonometry after photorefractive keratectomy. *Graefes Arch Clin Exp Ophthalmol* 2013; 251:603–8. <https://doi.org/10.1007/s00417-012-2142-1> PMID: 22940796

54. Liu J, Roberts CJ. Influence of corneal biomechanical properties on intraocular pressure measurement: quantitative analysis. *J Cataract Ref Surg* 2005; 31:146–55.
55. Hatami-Marbini H, Etebu E. Hydration dependent biomechanical properties of the corneal stroma. *Exp Eye Res* 2013; 116:47–54. <https://doi.org/10.1016/j.exer.2013.07.016> PMID: 23891861
56. Palko JR, Liu J. Definitions and concepts. In: Roberts CJ, Liu J, editors. *Corneal Biomechanics: From Theory to Practice*. 1st ed. Amsterdam. Kugler Publications; 2016 p7–8.
57. Palko JR, Liu J. Definitions and concepts. In: Roberts CJ, Liu J, editors. *Corneal Biomechanics: From Theory to Practice*. 1st ed. Amsterdam: Kugler Publications; 2016. p1–24.
58. Elsheikh A. *Understanding Corneal Biomechanics Through Experimental Assessment and Numerical Simulation*. New York. Nova Science Publications. 2010. p42.