

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

1-[(1*S*,6*R*,7*S*,9*R*)-8,8-Dibromo-5,5,9-trimethyltricyclo[4.4.0.1^{7,9}]decan-1-yl]-ethanoneMohamed Zaki,^{a*} Ahmed Benharref,^a Lahcen El Ammari,^b Mohamed Saadi^b and Moha Berraho^a^aLaboratoire de Chimie Biomoléculaires, Substances Naturelles et Réactivité, URAC16, Faculté des Sciences, Semlalia, BP 2390 Bd My Abdellah, 40000 Marrakech, Morocco, and ^bLaboratoire de Chimie du Solide Appliquée, Faculté des Sciences, Université Mohammed V-Agdal, BP 1014, Avenue Ibn Battouta, Rabat, Morocco

Correspondence e-mail: berraho@uca.ma


Received 6 March 2014; accepted 10 March 2014

Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; R factor = 0.031; wR factor = 0.060; data-to-parameter ratio = 18.6.

The title compound, $\text{C}_{16}\text{H}_{24}\text{Br}_2\text{O}$, was synthesized by three steps from β -himachalene (3,5,5,9-tetramethyl-2,4a,5,6,7,8-hexahydro-1*H*-benzocycloheptene), which was isolated from essential oil of the Atlas cedar *cedrus atlantica*. The asymmetric unit contains two independent molecules with almost identical conformations. Each molecule is built up from two fused six-membered rings, one having a chair conformation and the other a boat conformation, and an additional three-membered ring arising from the reaction of himachalene with dibromocarbene. In the crystal, there are no significant intermolecular interactions present. The absolute structure of the title compound was confirmed by resonance scattering.

Related literature

For background to the reactivity and biological properties of β -himachalene, see: El Haib *et al.* (2011); El Jamili *et al.* (2002); Daoubi *et al.* (2004). For a related structure, see: Benharref *et al.* (2013). For ring conformational analysis, see: Cremer & Pople (1975).



Experimental

Crystal data

$\text{C}_{16}\text{H}_{24}\text{Br}_2\text{O}$
 $M_r = 392.17$
 Orthorhombic, $P2_12_12_1$
 $a = 13.5013$ (19) Å
 $b = 14.042$ (2) Å
 $c = 17.213$ (3) Å
 $V = 3263.3$ (8) Å³
 $Z = 8$
 Mo $K\alpha$ radiation
 $\mu = 4.96$ mm⁻¹
 $T = 298$ K
 $0.5 \times 0.03 \times 0.03$ mm

Data collection

Bruker X8 APEX CCD area-detector diffractometer
 Absorption correction: multi-scan (*SADABS*; Bruker, 2009) $T_{\min} = 0.557$, $T_{\max} = 0.747$
 22468 measured reflections
 6542 independent reflections
 5262 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.040$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.031$
 $wR(F^2) = 0.060$
 $S = 1.02$
 6540 reflections
 351 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.27$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.39$ e Å⁻³
 Absolute structure: Flack & Bernardinelli (2000)
 Absolute structure parameter: -0.005 (8)

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 2012); software used to prepare material for publication: *WinGX* (Farrugia, 2012).

The authors thank the Unit of Support for Technical and Scientific Research (UATRS, CNRST) for the X-ray measurements.

Supporting information for this paper is available from the IUCr electronic archives (Reference: BT6967).

References

- Benharref, A., Ourhriss, N., El Ammari, L., Saadi, M. & Berraho, M. (2013). *Acta Cryst.* **E69**, o933–o934.
 Bruker (2009). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
 Daoubi, M., Duran-Patron, R., Hmamouchi, M., Hernandez-Galan, R., Benharref, A. & Isidro, G. C. (2004). *Pest Manag. Sci.* **60**, 927–932.
 El Haib, A., Benharref, A., Parrès-Maynadié, S., Manoury, E., Urrutigoity, M. & Gouygou, M. (2011). *Tetrahedron Asymmetry*, **22**, 101–108.
 El Jamili, H., Auhmani, A., Dakir, M., Lassaba, E., Benharref, A., Pierrot, M., Chiaroni, A. & Riche, C. (2002). *Tetrahedron Lett.* **43**, 6645–6648.
 Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
 Flack, H. D. & Bernardinelli, G. (2000). *J. Appl. Cryst.* **33**, 1143–1148.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supplementary materials

Acta Cryst. (2014). E70, o444 [doi:10.1107/S1600536814005431]

1-[(1*S*,6*R*,7*S*,9*R*)-8,8-Dibromo-5,5,9-trimethyltricyclo[4.4.0.1^{7,9}]decan-1-yl]ethanone

Mohamed Zaki, Ahmed Benharref, Lahcen El Ammari, Mohamed Saadi and Moha Berraho

1. Comment


The bicyclic sesquiterpene β -himachalene is the main constituent of the essential oil of the Atlas cedar (*Cedrus atlantica*) (El Haib *et al.*, 2011). The reactivity of this sesquiterpene and its derivatives has been studied extensively by our team in order to prepare new products having biological properties (El Jamili *et al.*, 2002; Benharref *et al.*, 2013). Indeed, these compounds were tested, using the food poisoning technique, for their potential antifungal activity against phytopathogen *Botrytis cinerea* (Daoubi *et al.*, 2004). In this paper we present the crystal structure of the title compound, (1*S*, 6*R*,7*S*, 9*R*)-1-acetyl-8,8-dibromo-5,5,9-trimethyltricyclo[4.4.0,1^{7,9}]decane. The asymmetric unit of the title compound contains two independent molecules of similar geometry (Fig. 1). Each molecule contains two fused six-membered rings, which are fused to a three-membered ring as shown in Fig.1. In both molecules, one of the six-membered ring has a chair conformation as indicated by the total puckering amplitude $QT = 0.543$ (4) Å and spherical polar angle $\theta_2 = 177.7$ (4)° with $\varphi_2 = 1(11)^\circ$, whereas the other six-membered ring displays a boat conformation with $QT = 0.725$ (3) Å, $\theta_2 = 92.2$ (2)° (Cremer & Pople, 1975). Owing to the presence of Br atoms, the absolute configuration could be fully confirmed, by refining the Flack parameter as C1(*S*), C6(*R*), C7(*S*) and C9(*R*).

2. Experimental

To obtain the title compound, $\text{BF}_3\text{-Et}_2\text{O}$ (1 ml) was added dropwise to a solution of (1*S*,2*R*,7*R*,8*S*,10*R*)-9,9-dibromo- 1 α , 2 α - epoxy -2, 6, 6, 10-tetramethyltricyclo [5.5.0.0^{8,10}] dodecane (1 g, 2.5 mmol) in 60 ml of dichloromethane at 195 K under nitrogen. The reaction mixture was stirred for two hours at a constant temperature of 195 K and the left at ambient temperature for 24 h. Water (60 ml) was added in order to separate the two phases, and the organic phase was dried and concentrated. The residue obtained was chromatographed on silica-gel eluting with hexane-ethyl acetate (98/2), which allowed the isolation of pure(1*S*, 6*R*,7*S*, 9*R*)-12-acetyl- 8,8-dibromo-5,5,9-trimethyltricyclo[4.4.0,1^{7,9}]decane in a yield of 70% (686 mg, 1.75 mmol). The title compound was recrystallized from its cyclohexane solution.

3. Refinement

All H atoms were fixed geometrically and treated as riding with C—H = 0.96 Å (methyl), 0.97 Å (methylene), 0.98 Å (methine) with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}$ (methylene, methine) or $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}$ (methyl).

**Figure 1**

Perspective view of one of the two molecules in the asymmetric unit of the title compound with the atom-labelling scheme. Displacement ellipsoids are drawn at the 30% probability level. H atoms are represented as small spheres of arbitrary radii.

1-[(1*S*,6*R*,7*S*,9*R*)-8,8-Dibromo-5,5,9-trimethyltricyclo[4.4.0.1^{7,9}]decan-1-yl]ethanone

Crystal data

$C_{16}H_{24}Br_2O$

$M_r = 392.17$

Orthorhombic, $P2_12_12_1$

$a = 13.5013 (19) \text{ \AA}$

$b = 14.042 (2) \text{ \AA}$

$c = 17.213 (3) \text{ \AA}$

$V = 3263.3 (8) \text{ \AA}^3$

$Z = 8$

$F(000) = 1584$

$D_x = 1.596 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 6542 reflections

$\theta = 2.4\text{--}26.4^\circ$

$\mu = 4.96 \text{ mm}^{-1}$

$T = 298$ K $0.5 \times 0.03 \times 0.03$ mm
 Block, colourless

Data collection

Bruker X8 APEX CCD area-detector diffractometer	22468 measured reflections 6542 independent reflections
Radiation source: fine-focus sealed tube	5262 reflections with $I > 2\sigma(I)$
Graphite monochromator	$R_{\text{int}} = 0.040$
φ and ω scans	$\theta_{\text{max}} = 26.4^\circ$, $\theta_{\text{min}} = 2.4^\circ$
Absorption correction: multi-scan (SADABS; Bruker, 2009)	$h = -15 \rightarrow 16$ $k = -12 \rightarrow 17$ $l = -17 \rightarrow 21$
$T_{\text{min}} = 0.557$, $T_{\text{max}} = 0.747$	

Refinement

Refinement on F^2	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H-atom parameters constrained
$R[F^2 > 2\sigma(F^2)] = 0.031$	$w = 1/[\sigma^2(F_o^2) + (0.0104P)^2]$
$wR(F^2) = 0.060$	where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.02$	$(\Delta/\sigma)_{\text{max}} = 0.001$
6540 reflections	$\Delta\rho_{\text{max}} = 0.27 \text{ e } \text{\AA}^{-3}$
351 parameters	$\Delta\rho_{\text{min}} = -0.39 \text{ e } \text{\AA}^{-3}$
0 restraints	Absolute structure: Flack & Bernardinelli (2000)
Primary atom site location: structure-invariant direct methods	Absolute structure parameter: -0.005 (8)
Secondary atom site location: difference Fourier map	

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against all reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on all data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.7516 (2)	1.0732 (2)	-0.03701 (18)	0.0342 (8)
C2	0.6483 (3)	1.1166 (3)	-0.0456 (2)	0.0518 (10)
H2A	0.6508	1.1658	-0.0851	0.062*
H2B	0.6032	1.0676	-0.0636	0.062*
C3	0.6068 (3)	1.1596 (3)	0.0284 (2)	0.0628 (12)
H3A	0.6465	1.2142	0.0431	0.075*
H3B	0.5398	1.1815	0.0189	0.075*
C4	0.6060 (3)	1.0889 (3)	0.0941 (2)	0.0544 (10)
H4A	0.5600	1.0379	0.0819	0.065*
H4B	0.5827	1.1202	0.1409	0.065*
C5	0.7090 (2)	1.0460 (3)	0.10960 (19)	0.0394 (9)
C6	0.7513 (2)	1.0035 (2)	0.03386 (15)	0.0250 (7)
H6	0.8204	0.9864	0.0442	0.030*

C7	0.6976 (2)	0.9125 (2)	0.01156 (17)	0.0298 (8)
H7	0.6264	0.9124	0.0228	0.036*
C8	0.7455 (2)	0.8170 (2)	0.01354 (18)	0.0329 (8)
C9	0.7266 (2)	0.8636 (2)	-0.06406 (17)	0.0320 (8)
C10	0.8094 (2)	0.9159 (2)	-0.1044 (2)	0.0393 (9)
H10A	0.8216	0.8877	-0.1549	0.047*
H10B	0.8696	0.9114	-0.0739	0.047*
C11	0.7797 (3)	1.0195 (3)	-0.11368 (18)	0.0435 (9)
H11A	0.7235	1.0225	-0.1488	0.052*
H11B	0.8340	1.0532	-0.1382	0.052*
C12	0.8323 (3)	1.1502 (3)	-0.0304 (2)	0.0452 (9)
C13	0.9368 (3)	1.1193 (3)	-0.0122 (2)	0.0568 (11)
H13A	0.9824	1.1549	-0.0437	0.085*
H13B	0.9439	1.0526	-0.0231	0.085*
H13C	0.9506	1.1309	0.0417	0.085*
C14	0.7777 (3)	1.1229 (3)	0.1434 (2)	0.0547 (11)
H14A	0.7492	1.1479	0.1902	0.082*
H14B	0.7855	1.1734	0.1062	0.082*
H14C	0.8412	1.0955	0.1547	0.082*
C15	0.6990 (3)	0.9672 (3)	0.1702 (2)	0.0553 (11)
H15A	0.6551	0.9188	0.1512	0.083*
H15B	0.6728	0.9935	0.2175	0.083*
H15C	0.7629	0.9398	0.1802	0.083*
C16	0.6480 (3)	0.8239 (3)	-0.1165 (2)	0.0517 (10)
H16A	0.6738	0.7697	-0.1438	0.077*
H16B	0.6282	0.8717	-0.1532	0.077*
H16C	0.5918	0.8049	-0.0861	0.077*
Br1	0.87872 (2)	0.80148 (3)	0.05270 (2)	0.04524 (10)
Br2	0.66521 (3)	0.71109 (3)	0.04601 (3)	0.05635 (12)
O1	0.8142 (2)	1.23284 (19)	-0.04537 (18)	0.0746 (9)
C17	0.8403 (2)	0.5321 (2)	0.30776 (18)	0.0353 (8)
C18	0.9200 (3)	0.4590 (3)	0.3290 (2)	0.0536 (11)
H18A	0.9638	0.4872	0.3674	0.064*
H18B	0.8881	0.4046	0.3531	0.064*
C19	0.9822 (3)	0.4239 (3)	0.2615 (2)	0.0645 (12)
H19A	1.0214	0.4762	0.2412	0.077*
H19B	1.0274	0.3750	0.2796	0.077*
C20	0.9176 (3)	0.3837 (3)	0.1974 (2)	0.0532 (10)
H20A	0.8836	0.3276	0.2166	0.064*
H20B	0.9593	0.3643	0.1543	0.064*
C21	0.8409 (2)	0.4558 (2)	0.16839 (19)	0.0358 (8)
C22	0.7784 (2)	0.4940 (2)	0.23762 (17)	0.0297 (7)
H22	0.7397	0.5479	0.2178	0.036*
C23	0.7047 (2)	0.4204 (2)	0.26590 (18)	0.0319 (8)
H23	0.7282	0.3544	0.2648	0.038*
C24	0.5956 (2)	0.4328 (2)	0.25512 (19)	0.0365 (8)
C25	0.6391 (2)	0.4460 (2)	0.33457 (19)	0.0331 (8)
C26	0.6622 (2)	0.5442 (3)	0.3653 (2)	0.0411 (9)
H26A	0.6258	0.5554	0.4130	0.049*

H26B	0.6426	0.5920	0.3276	0.049*
C27	0.7733 (2)	0.5508 (3)	0.38072 (18)	0.0437 (9)
H27A	0.7906	0.5052	0.4208	0.052*
H27B	0.7880	0.6139	0.4007	0.052*
C28	0.8851 (3)	0.6296 (3)	0.2910 (2)	0.0501 (10)
C29	0.8180 (3)	0.7092 (3)	0.2652 (2)	0.0614 (11)
H29A	0.8507	0.7691	0.2729	0.092*
H29B	0.7580	0.7077	0.2951	0.092*
H29C	0.8024	0.7014	0.2112	0.092*
C30	0.8932 (3)	0.5378 (3)	0.1257 (2)	0.0517 (10)
H30A	0.9371	0.5699	0.1609	0.078*
H30B	0.8446	0.5820	0.1067	0.078*
H30C	0.9305	0.5128	0.0828	0.078*
C31	0.7714 (3)	0.4076 (3)	0.1084 (2)	0.0541 (11)
H31A	0.8092	0.3880	0.0640	0.081*
H31B	0.7213	0.4521	0.0925	0.081*
H31C	0.7406	0.3530	0.1317	0.081*
C32	0.6193 (3)	0.3715 (3)	0.3973 (2)	0.0520 (10)
H32A	0.5559	0.3833	0.4206	0.078*
H32B	0.6699	0.3751	0.4364	0.078*
H32C	0.6196	0.3091	0.3744	0.078*
Br3	0.54217 (3)	0.53917 (3)	0.19802 (2)	0.04941 (11)
Br4	0.51892 (3)	0.32022 (3)	0.23283 (2)	0.06195 (13)
O2	0.9721 (2)	0.6461 (2)	0.30335 (19)	0.0872 (10)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0425 (19)	0.0318 (19)	0.0284 (19)	-0.0034 (16)	-0.0046 (16)	0.0053 (15)
C2	0.060 (2)	0.041 (2)	0.054 (2)	0.0080 (18)	-0.018 (2)	0.009 (2)
C3	0.048 (2)	0.052 (3)	0.089 (3)	0.025 (2)	-0.011 (2)	-0.007 (2)
C4	0.051 (2)	0.049 (3)	0.063 (3)	0.0153 (19)	0.011 (2)	-0.014 (2)
C5	0.044 (2)	0.040 (2)	0.0341 (19)	0.0017 (17)	0.0072 (16)	-0.0049 (18)
C6	0.0288 (15)	0.0211 (16)	0.0249 (17)	0.0002 (13)	0.0003 (13)	0.0007 (13)
C7	0.0280 (17)	0.030 (2)	0.0310 (18)	0.0021 (14)	0.0019 (14)	-0.0011 (14)
C8	0.0285 (16)	0.0267 (19)	0.0435 (19)	-0.0038 (15)	-0.0005 (15)	0.0008 (15)
C9	0.0354 (18)	0.0330 (19)	0.0277 (18)	-0.0013 (15)	0.0005 (15)	-0.0085 (15)
C10	0.047 (2)	0.041 (2)	0.0297 (19)	-0.0038 (16)	0.0077 (16)	-0.0060 (16)
C11	0.056 (2)	0.043 (2)	0.0315 (19)	-0.0121 (18)	0.0011 (17)	0.0059 (17)
C12	0.069 (3)	0.034 (2)	0.033 (2)	-0.0144 (19)	0.0010 (19)	0.0014 (17)
C13	0.054 (2)	0.052 (3)	0.064 (3)	-0.024 (2)	0.006 (2)	0.001 (2)
C14	0.067 (3)	0.055 (3)	0.042 (2)	-0.004 (2)	0.003 (2)	-0.0179 (19)
C15	0.077 (3)	0.060 (3)	0.0290 (19)	-0.002 (2)	0.0176 (19)	-0.001 (2)
C16	0.054 (2)	0.047 (2)	0.054 (2)	-0.0055 (19)	-0.0146 (19)	-0.0152 (19)
Br1	0.04248 (19)	0.0378 (2)	0.0554 (2)	0.00669 (16)	-0.01088 (17)	0.00135 (19)
Br2	0.0576 (2)	0.0342 (2)	0.0772 (3)	-0.01247 (18)	0.0117 (2)	0.0086 (2)
O1	0.108 (2)	0.0304 (17)	0.086 (2)	-0.0148 (15)	-0.0083 (19)	0.0137 (15)
C17	0.0281 (16)	0.042 (2)	0.0356 (18)	-0.0008 (16)	-0.0081 (15)	0.0005 (17)
C18	0.045 (2)	0.069 (3)	0.047 (2)	0.005 (2)	-0.0138 (18)	0.007 (2)
C19	0.041 (2)	0.077 (3)	0.075 (3)	0.023 (2)	-0.003 (2)	0.012 (2)

C20	0.052 (2)	0.046 (2)	0.061 (2)	0.0181 (19)	0.015 (2)	0.002 (2)
C21	0.0389 (18)	0.031 (2)	0.0372 (18)	0.0057 (16)	0.0062 (16)	-0.0015 (16)
C22	0.0327 (17)	0.0252 (18)	0.0312 (17)	0.0045 (14)	0.0006 (14)	0.0011 (14)
C23	0.0324 (17)	0.0258 (19)	0.0374 (19)	0.0012 (14)	-0.0019 (15)	0.0062 (15)
C24	0.0353 (18)	0.0303 (19)	0.044 (2)	-0.0014 (15)	-0.0059 (16)	0.0011 (16)
C25	0.0297 (17)	0.034 (2)	0.0351 (18)	0.0026 (15)	0.0022 (14)	0.0030 (16)
C26	0.045 (2)	0.045 (2)	0.0334 (19)	0.0048 (18)	0.0052 (16)	-0.0004 (17)
C27	0.052 (2)	0.045 (2)	0.034 (2)	-0.0074 (18)	-0.0054 (17)	0.0005 (17)
C28	0.049 (2)	0.054 (3)	0.047 (2)	-0.018 (2)	-0.0040 (19)	-0.0065 (19)
C29	0.086 (3)	0.036 (2)	0.062 (2)	-0.009 (2)	-0.003 (2)	0.004 (2)
C30	0.051 (2)	0.053 (3)	0.051 (2)	0.000 (2)	0.0137 (19)	0.012 (2)
C31	0.061 (2)	0.062 (3)	0.039 (2)	0.001 (2)	0.006 (2)	-0.015 (2)
C32	0.055 (2)	0.054 (2)	0.047 (2)	0.006 (2)	0.009 (2)	0.018 (2)
Br3	0.03866 (19)	0.0577 (3)	0.0519 (2)	0.00694 (18)	-0.00959 (18)	0.01018 (19)
Br4	0.0534 (2)	0.0531 (3)	0.0794 (3)	-0.0209 (2)	-0.0020 (2)	-0.0105 (2)
O2	0.0579 (18)	0.085 (2)	0.119 (3)	-0.0333 (18)	-0.020 (2)	0.004 (2)

Geometric parameters (Å, °)

C1—C2	1.529 (5)	C17—C28	1.524 (5)
C1—C12	1.540 (5)	C17—C18	1.532 (5)
C1—C6	1.564 (4)	C17—C22	1.563 (4)
C1—C11	1.567 (4)	C17—C27	1.570 (5)
C2—C3	1.516 (5)	C18—C19	1.516 (5)
C2—H2A	0.9700	C18—H18A	0.9700
C2—H2B	0.9700	C18—H18B	0.9700
C3—C4	1.506 (5)	C19—C20	1.515 (5)
C3—H3A	0.9700	C19—H19A	0.9700
C3—H3B	0.9700	C19—H19B	0.9700
C4—C5	1.539 (5)	C20—C21	1.531 (4)
C4—H4A	0.9700	C20—H20A	0.9700
C4—H4B	0.9700	C20—H20B	0.9700
C5—C15	1.527 (5)	C21—C30	1.538 (4)
C5—C14	1.537 (5)	C21—C31	1.550 (5)
C5—C6	1.543 (4)	C21—C22	1.556 (4)
C6—C7	1.519 (4)	C22—C23	1.515 (4)
C6—H6	0.9800	C22—H22	0.9800
C7—C8	1.489 (4)	C23—C24	1.496 (4)
C7—C9	1.523 (4)	C23—C25	1.520 (4)
C7—H7	0.9800	C23—H23	0.9800
C8—C9	1.510 (4)	C24—C25	1.500 (4)
C8—Br2	1.924 (3)	C24—Br4	1.928 (3)
C8—Br1	1.933 (3)	C24—Br3	1.928 (3)
C9—C16	1.501 (4)	C25—C26	1.509 (5)
C9—C10	1.506 (4)	C25—C32	1.528 (4)
C10—C11	1.517 (5)	C26—C27	1.526 (5)
C10—H10A	0.9700	C26—H26A	0.9700
C10—H10B	0.9700	C26—H26B	0.9700
C11—H11A	0.9700	C27—H27A	0.9700
C11—H11B	0.9700	C27—H27B	0.9700

C12—O1	1.213 (4)	C28—O2	1.215 (4)
C12—C13	1.509 (5)	C28—C29	1.506 (5)
C13—H13A	0.9600	C29—H29A	0.9600
C13—H13B	0.9600	C29—H29B	0.9600
C13—H13C	0.9600	C29—H29C	0.9600
C14—H14A	0.9600	C30—H30A	0.9600
C14—H14B	0.9600	C30—H30B	0.9600
C14—H14C	0.9600	C30—H30C	0.9600
C15—H15A	0.9600	C31—H31A	0.9600
C15—H15B	0.9600	C31—H31B	0.9600
C15—H15C	0.9600	C31—H31C	0.9600
C16—H16A	0.9600	C32—H32A	0.9600
C16—H16B	0.9600	C32—H32B	0.9600
C16—H16C	0.9600	C32—H32C	0.9600
C2—C1—C12	111.9 (3)	C28—C17—C18	111.5 (3)
C2—C1—C6	108.8 (3)	C28—C17—C22	112.0 (3)
C12—C1—C6	112.5 (3)	C18—C17—C22	109.3 (3)
C2—C1—C11	109.4 (3)	C28—C17—C27	103.3 (3)
C12—C1—C11	103.3 (3)	C18—C17—C27	109.0 (3)
C6—C1—C11	110.9 (3)	C22—C17—C27	111.6 (2)
C3—C2—C1	114.5 (3)	C19—C18—C17	115.1 (3)
C3—C2—H2A	108.6	C19—C18—H18A	108.5
C1—C2—H2A	108.6	C17—C18—H18A	108.5
C3—C2—H2B	108.6	C19—C18—H18B	108.5
C1—C2—H2B	108.6	C17—C18—H18B	108.5
H2A—C2—H2B	107.6	H18A—C18—H18B	107.5
C4—C3—C2	111.8 (3)	C20—C19—C18	111.1 (3)
C4—C3—H3A	109.3	C20—C19—H19A	109.4
C2—C3—H3A	109.3	C18—C19—H19A	109.4
C4—C3—H3B	109.3	C20—C19—H19B	109.4
C2—C3—H3B	109.3	C18—C19—H19B	109.4
H3A—C3—H3B	107.9	H19A—C19—H19B	108.0
C3—C4—C5	112.4 (3)	C19—C20—C21	112.4 (3)
C3—C4—H4A	109.1	C19—C20—H20A	109.1
C5—C4—H4A	109.1	C21—C20—H20A	109.1
C3—C4—H4B	109.1	C19—C20—H20B	109.1
C5—C4—H4B	109.1	C21—C20—H20B	109.1
H4A—C4—H4B	107.8	H20A—C20—H20B	107.9
C15—C5—C14	107.7 (3)	C20—C21—C30	109.9 (3)
C15—C5—C4	108.8 (3)	C20—C21—C31	109.8 (3)
C14—C5—C4	109.6 (3)	C30—C21—C31	106.6 (3)
C15—C5—C6	109.3 (3)	C20—C21—C22	110.2 (3)
C14—C5—C6	111.6 (3)	C30—C21—C22	110.9 (3)
C4—C5—C6	109.8 (3)	C31—C21—C22	109.4 (3)
C7—C6—C5	111.2 (3)	C23—C22—C21	111.5 (3)
C7—C6—C1	109.3 (2)	C23—C22—C17	109.7 (2)
C5—C6—C1	114.7 (3)	C21—C22—C17	114.8 (2)
C7—C6—H6	107.1	C23—C22—H22	106.8

C5—C6—H6	107.1	C21—C22—H22	106.8
C1—C6—H6	107.1	C17—C22—H22	106.8
C8—C7—C6	123.0 (2)	C24—C23—C22	121.8 (3)
C8—C7—C9	60.1 (2)	C24—C23—C25	59.7 (2)
C6—C7—C9	118.2 (3)	C22—C23—C25	118.1 (3)
C8—C7—H7	114.8	C24—C23—H23	115.3
C6—C7—H7	114.8	C22—C23—H23	115.3
C9—C7—H7	114.8	C25—C23—H23	115.3
C7—C8—C9	61.0 (2)	C23—C24—C25	61.0 (2)
C7—C8—Br2	117.2 (2)	C23—C24—Br4	117.3 (2)
C9—C8—Br2	119.8 (2)	C25—C24—Br4	119.5 (2)
C7—C8—Br1	120.9 (2)	C23—C24—Br3	121.5 (2)
C9—C8—Br1	121.0 (2)	C25—C24—Br3	121.0 (2)
Br2—C8—Br1	109.63 (15)	Br4—C24—Br3	109.45 (15)
C16—C9—C10	115.4 (3)	C24—C25—C26	120.9 (3)
C16—C9—C8	119.4 (3)	C24—C25—C23	59.4 (2)
C10—C9—C8	119.5 (3)	C26—C25—C23	111.6 (3)
C16—C9—C7	120.0 (3)	C24—C25—C32	119.4 (3)
C10—C9—C7	111.4 (3)	C26—C25—C32	114.5 (3)
C8—C9—C7	58.8 (2)	C23—C25—C32	119.3 (3)
C9—C10—C11	108.6 (3)	C25—C26—C27	108.7 (3)
C9—C10—H10A	110.0	C25—C26—H26A	110.0
C11—C10—H10A	110.0	C27—C26—H26A	110.0
C9—C10—H10B	110.0	C25—C26—H26B	110.0
C11—C10—H10B	110.0	C27—C26—H26B	110.0
H10A—C10—H10B	108.3	H26A—C26—H26B	108.3
C10—C11—C1	115.9 (3)	C26—C27—C17	114.6 (3)
C10—C11—H11A	108.3	C26—C27—H27A	108.6
C1—C11—H11A	108.3	C17—C27—H27A	108.6
C10—C11—H11B	108.3	C26—C27—H27B	108.6
C1—C11—H11B	108.3	C17—C27—H27B	108.6
H11A—C11—H11B	107.4	H27A—C27—H27B	107.6
O1—C12—C13	120.5 (4)	O2—C28—C29	119.4 (4)
O1—C12—C1	120.9 (4)	O2—C28—C17	121.5 (4)
C13—C12—C1	118.4 (3)	C29—C28—C17	118.9 (3)
C12—C13—H13A	109.5	C28—C29—H29A	109.5
C12—C13—H13B	109.5	C28—C29—H29B	109.5
H13A—C13—H13B	109.5	H29A—C29—H29B	109.5
C12—C13—H13C	109.5	C28—C29—H29C	109.5
H13A—C13—H13C	109.5	H29A—C29—H29C	109.5
H13B—C13—H13C	109.5	H29B—C29—H29C	109.5
C5—C14—H14A	109.5	C21—C30—H30A	109.5
C5—C14—H14B	109.5	C21—C30—H30B	109.5
H14A—C14—H14B	109.5	H30A—C30—H30B	109.5
C5—C14—H14C	109.5	C21—C30—H30C	109.5
H14A—C14—H14C	109.5	H30A—C30—H30C	109.5
H14B—C14—H14C	109.5	H30B—C30—H30C	109.5
C5—C15—H15A	109.5	C21—C31—H31A	109.5
C5—C15—H15B	109.5	C21—C31—H31B	109.5

H15A—C15—H15B	109.5	H31A—C31—H31B	109.5
C5—C15—H15C	109.5	C21—C31—H31C	109.5
H15A—C15—H15C	109.5	H31A—C31—H31C	109.5
H15B—C15—H15C	109.5	H31B—C31—H31C	109.5
C9—C16—H16A	109.5	C25—C32—H32A	109.5
C9—C16—H16B	109.5	C25—C32—H32B	109.5
H16A—C16—H16B	109.5	H32A—C32—H32B	109.5
C9—C16—H16C	109.5	C25—C32—H32C	109.5
H16A—C16—H16C	109.5	H32A—C32—H32C	109.5
H16B—C16—H16C	109.5	H32B—C32—H32C	109.5
