

SCIENTIFIC REPORTS



OPEN

dFoxO promotes Wingleless signaling in *Drosophila*

Shiping Zhang¹, Xiaowei Guo¹, Changyan Chen¹, Yujun Chen¹, Jikai Li¹, Ying Sun¹, Chenxi Wu^{1,†}, Yang Yang¹, Cizhong Jiang², Wenzhe Li¹ & Lei Xue¹

Received: 17 March 2015

Accepted: 11 February 2016

Published: 03 March 2016

The Wnt/ β -catenin signaling is an evolutionarily conserved pathway that regulates a wide range of physiological functions, including embryogenesis, organ maintenance, cell proliferation and cell fate decision. Dysregulation of Wnt/ β -catenin signaling has been implicated in various cancers, but its role in cell death has not yet been fully elucidated. Here we show that activation of Wg signaling induces cell death in *Drosophila* eyes and wings, which depends on dFoxO, a transcription factor known to be involved in cell death. In addition, dFoxO is required for ectopic and endogenous Wg signaling to regulate wing patterning. Moreover, dFoxO is necessary for activated Wg signaling-induced target genes expression. Furthermore, Arm is reciprocally required for dFoxO-induced cell death. Finally, dFoxO physically interacts with Arm both *in vitro* and *in vivo*. Thus, we have characterized a previously unknown role of dFoxO in promoting Wg signaling, and that a dFoxO-Arm complex is likely involved in their mutual functions, e.g. cell death.

The Wnt/ β -catenin signaling represents one of the most intensively studied pathways, which is tightly associated with cancers, especially colorectal cancer^{1,2}. Upon binding of Wnt ligands to the receptor Frizzled (Fz) and co-receptor LDL-receptor-related protein (LRP), the extracellular signals are transduced via Disheveled (Dsh) to the destruction complex composed of Axin (Axn), Adenomatous polyposis coli (APC) and glycogen synthase kinase 3 (GSK3), thereby preventing the proteasomal degradation of the transcriptional coactivator β -catenin. Stabilized β -catenin thus translocates into the nucleus and associates with the T cell factor (TCF)/Lymphoid-enhancing factor (LEF) family of transcription factors to activate the transcription of target genes^{3–5}. Through this mechanism, Wnt/ β -catenin signaling plays a wide range of functions, including embryogenesis, organ maintenance, cell proliferation and cell fate decisions^{3,6–8}. Wnt signaling is also reported to regulate cell death during *Drosophila* retina and ommatidia development^{9–11}, yet the underlying mechanism has not been fully elucidated, and the downstream regulators remain elusive.

The Forkhead box O (FoxO) transcription factors belong to the large forkhead family proteins, which are characterized by a winged helix DNA binding domain called 'Forkhead box'^{12,13}. The FoxO proteins have been conserved from *C. elegans* to mammals¹⁴. While there is only one FoxO gene in invertebrates, four members have been identified in mammals including FoxO1, FoxO3a, FoxO4 and FoxO6^{13,15}. FoxO1, 3a, and 4 are ubiquitously expressed, whereas FoxO6 expression is confined to the brain¹². FoxO activity is negatively regulated by the Insulin/PI3K/Akt signaling pathway. Activation of Akt (also known as protein kinase B) phosphorylates FoxO and results in nuclear exclusion of FoxO, thereby inhibits its transcriptional activity^{14–16}. However, stress conditions, such as high levels of reactive oxygen species (ROS) or deprivation of growth factors, promote the nuclear localization of FoxO and induce its target genes expression¹⁴. FoxO is critically involved in a variety of physiological processes including cell cycle, cell death and differentiation, DNA repair, oxidative stress response and longevity^{12,17–19}. Dysregulation of FoxO has been associated with many diseases, including immune defects, malignancy, diabetes and Alzheimer's disease (AD)^{13,20}.

β -catenin has been reported to interact with FoxO and enhance its transcriptional activity in mammalian cells and *C. elegans*²¹. On the other hand, FoxO inhibits Wnt signaling by diverting the limited pool of β -catenin from Wnt/TCF to FoxO^{22,23}, leading to embryogenesis and bone formation defects^{24,25}.

¹Institute of Intervention Vessel, Shanghai 10th People's Hospital, Shanghai Key Laboratory of Signaling and Diseases Research, School of Life Science and Technology, Tongji University, 1239 Siping Road, Shanghai 200092, China. ²Shanghai Key Laboratory of Signaling and Diseases Research, School of Life Science and Technology, Tongji University, 1239 Siping Road, Shanghai 200092, China. [†]Present address: Department of Traditional Chinese Medicine, North China University of Science and Technology, 57 South Jianshe Road, Tangshan 063000, China. Correspondence and requests for materials should be addressed to W.L. (email: lwz@tongji.edu.cn) or L.X. (email: lei.xue@tongji.edu.cn)




Figure 1. Activation of Wg signaling induces cell death in *Drosophila*. Light micrographs of *Drosophila* adult eyes and wings, and fluorescent micrographs of 3rd instar eye and wing discs are shown. Compared with the *GMR*-Gal4 control (a,a'), expression of Wg, Dsh or Arm induces cell death in eye discs indicated by AO staining (b'-d') and produces adult eyes with reduced size (b-d). Compared with the *ptc*-Gal4 control (e,e'), expression of Dsh, Arm or Pan induces cell death in wing discs (f'-h') and produces a loss-of-ACV phenotype in adult wings (f-h, the lower panels are high magnification of the boxed areas in upper panels). Scale bars: 100 μ m in (a-d) and (e-h, upper panels); 50 μ m in (a'-d'), (e-h, lower panels) and (e'-h').

Here using *Drosophila melanogaster*, which has reduced genome redundancy and many available genetic tools, we found that activation of Wingless (Wg, *Drosophila* Wnt homolog) signaling induces intensive cell death in *Drosophila* eyes and wings, which depends on dFoxO (*Drosophila* FoxO homolog). In addition, dFoxo is required for Wg signaling to activate target genes expression and execute its endogenous functions in wing patterning. Furthermore, loss of *armadillo* (*arm*, encoding *Drosophila* β -catenin homolog) or *pangolin* (*pan*, encoding *Drosophila* TCF homolog) could also suppress dFoxO-triggered cell death, suggesting a reciprocal effect. Finally, dFoxO physically interacts with Arm, providing a molecular mechanism for the role of dFoxO in promoting Wg signaling. Thus, contradict to the previous studies that FoxO proteins inhibit Wnt signaling, our data point to a positive role of dFoxO in modulating the canonical Wg signaling in *Drosophila*, suggesting FoxO differentially regulates Wnt signaling in a cell context-dependent manner.

Results

Activation of Wg signaling induces cell death in *Drosophila*. To investigate the role of Wg signaling in cell death, we expressed the representative components of this pathway in the *Drosophila* eye, and found that ectopic expression of Wg, Dsh or Arm resulted in massive cell death in 3rd instar eye discs as revealed by acridine orange (AO) staining (Fig. 1a'-d'), and produced eyes with reduced size (Fig. 1a-d). In addition, ectopic expression of Dsh, Arm and Pan under *ptc*-Gal4 produced a loss of anterior cross vein (ACV) phenotype (Fig. 1e-h), accompanied by increased cell death in wing discs (Fig. 1e'-h'). Furthermore, ectopic expression of Dsh or Arm in wing discs by additional drivers including *en*-Gal4, *omb*-Gal4 and *sd*-Gal4 resulted in extensive cell death in




Figure 2. dFoxO is required for Arm-induced cell death. Light micrographs of *Drosophila* adult eyes and fluorescent micrographs of 3rd instar eye discs are shown. Compared with the *GMR*-Gal4 control (a), expression of Arm triggers cell death and produces a small eye phenotype (b), which remains unaffected by expressing LacZ (c), but is partially suppressed by knocking-down of *dfoxo* (d), or in heterozygous *dfoxo*^{21/+} (e) or *dfoxo*^{25/+} (f) background, and fully suppressed in *dfoxo*^{21/dfoxo}²⁵ mutants (g). As positive controls, knocking-down of *arm* (h) or *pan* (i) suppresses the *GMR* > Arm small eye phenotype. *GMR* > Arm-induced AO staining (j) is suppressed partially in heterozygous *dfoxo*^{25/+} mutants (k), and fully in *dfoxo*^{21/dfoxo}²⁵ trans-heterozygous mutants (l). Sample numbers: a, 87; b, 110; c, 100; d, 75; e, 77; f, 56; g, 75; h, 97; i, 78; j, 22; k, 14; l, 17. Scale bars: 100 μ m in (a–i) and 50 μ m in (j–l).

wing pouches, as indicated by AO or TUNEL staining (Fig. S1). Collectively, these results indicate that activation of Wg signaling is able to induce cell death in *Drosophila*.

Wg signaling is necessary to provoke cell death in pupal retinas to sculpture the interommatidial lattice^{9,10}. To investigate whether further activation of the Wg signaling could trigger more cell death in pupal retinas, we performed AO staining at 21 h after pupal formation (AFP). Consistent with previous reports^{9,10}, extensive cell death was observed in *GMR*-Gal4 control retinas, mostly located in the anterior portion (Fig. S2a). Overexpression of Arm did not further increase cell death (Fig. S2b and S2c), suggesting the endogenous Wg signaling is optimally activated, and additional Wg signaling is not able to induce more cell death in this context.

Loss of dFoxO suppresses Arm-induced cell death. It has been reported that FoxO directly binds to β -catenin²¹, and competes with TCF for the limited pool of β -catenin, thereby inhibits Wnt signaling activity^{22,23}. Thus we wonder whether dFoxO also inhibits Wg signaling-induced cell death in *Drosophila*. To our surprise, *GMR* > Arm-induced small eye phenotype (Fig. 2b) was suppressed partially by RNAi-mediated knocking-down

of *dfoxo* (Fig. 2d), or in heterozygous *dfoxo*²¹ or *dfoxo*²⁵ mutants (Fig. 2e,f), and fully in *dfoxo*²¹/*dfoxo*²⁵ trans-heterozygous mutants (Fig. 2g), but remained unaffected by the expression of LacZ (Fig. 2c). As positive controls, loss of *arm* or *pan* fully suppressed *GMR* > Arm-induced small eye phenotype (Fig. 2h,i). Consistently, loss of *dfoxo* also suppressed *GMR* > Wg-induced small eye phenotype (Fig. S3). Moreover, *GMR* > Arm-induced AO staining (Fig. 2j) was suppressed partially in heterozygous *dfoxo*²⁵ mutants (Fig. 2k), and fully in *dfoxo*²¹/*dfoxo*²⁵ trans-heterozygous mutants (Fig. 2l). Thus, dFoxO is positively required for Wg signaling-induced cell death in *Drosophila*.

Our previous study showed Wg mediates JNK signaling induced cell death²⁶, and dFoxO is reported to be a downstream transcription factor of JNK signaling^{27–29}. To test whether JNK is reciprocally required for Wg pathway induced cell death, we overexpressed Wg, Dsh or Arm by *GMR*-Gal4 in a compromised JNK background. We found activated Wg signaling induced small eye phenotype remained unaffected by blocking JNK signaling (Fig. S4a–S4r). As a positive control, *GMR* > Egr induced small eye phenotype was effectively suppressed by abrogating JNK activity (Fig. S4s–S4x). Thus, dFoxO acts independently of the JNK pathway to mediate Wg signaling induced cell death.

Loss of dFoxO suppresses Arm-induced apoptotic gene activation. *reaper* (*rpr*) and *head involution defective* (*hid*) are important pro-apoptotic genes regulating cell death³⁰. To examine whether activated Wg signaling induces cell death through up-regulation of *rpr* and *hid*, we ectopically expressed Arm in the wing pouch driven by *sd*-Gal4. Compared with the controls (Fig. 3a,g), *rpr* and *hid* expressions were significantly up-regulated by activated Wg signaling in the wing pouch (Fig. 3b,h), which were considerably suppressed by knocking-down of *dfoxo* or in heterozygous *dfoxo* mutants (Fig. 3d–f,j–l), but remained unaffected by the expression of GFP (Fig. 3c,i). Taken together, these results suggest that dFoxO is indispensable for Wg signaling induced cell death in *Drosophila*, which contradicts to the previous reports that FoxO impedes Wnt signaling in mammalian cells^{22,23}. Thus, it is possible that FoxO may regulate Wnt signaling differently in a tissue or cell type specific manner.

dFoxO is required for the wing patterning functions of Wg signaling. Wg signaling is one of the profound pathways that regulate wing pattern formation. Elevation of this pathway induces ectopic bristles, whereas inhibition of which results in wing margin notches^{31,32}. Consistent with previous reports, we found that expression of Arm driven by *ptc*-Gal4 produced ectopic bristles between L3 and L4 veins (Fig. 4b, arrow head), and generated a loss-of-ACV phenotype (Fig. 4b). Intriguingly, a similar loss-of-ACV phenotype has been reported as a result of dFoxO overexpression (Fig. S5b)²⁰. We found that both the ectopic bristles and loss-of-ACV phenotypes were considerably suppressed by loss of *dfoxo* (Fig. 4d,e), but remained unchanged by the expression of LacZ (Fig. 4c). Similarly, *sd* > Arm induced ectopic bristles near the wing margin (Fig. 4g, arrow head) were fully suppressed by loss of *dfoxo* (Fig. 4i,j), but remained unaffected by the expression of LacZ (Fig. 4h). Hence, dFoxO is required for ectopic Wg signaling-induced extra bristles and loss-of-ACV phenotypes in the wing.

Interestingly, despite cell death in wing discs, the adult wing sizes were not significantly altered upon Arm overexpression (Fig. S6a and S6b). As Wg signaling also regulates cell proliferation, we speculate that cell death might be compensated by increased cell proliferation. To address this issue, we performed PH3 staining (Fig. S6c–S6f) and BrdU incorporation (Fig. S6g–S6j) in 3rd instar wing discs, but found no significant increase of cell proliferation in regions expressing Arm (Fig. S6k and S6l). Thus, other mechanism may exist to cope with ectopic Wg induced cell death and maintain tissue homeostasis in wing development. Alternatively, activation of cell death program may not necessary lead to cell loss, but instead, cell fate alteration. For instance, activation of apoptosis by *ptc* > Grim and *ptc* > *pelle-IR* produced the same loss-of-ACV phenotype¹⁹ as that of *ptc* > Dsh and *ptc* > Arm (Fig. 1f,g).

On the other hand, compromised Wg signaling impedes proliferation and results in loss of wing tissue, with wing margin notches as the most frequently observed phenotype^{31,33,34}. Consistently, loss of *wg* between L3 and L4 veins by *ptc* > *wg-IR* generated a mild notch phenotype (Fig. 4k), which was suppressed by the expression of dFoxO (Fig. 4l), but enhanced in heterozygous *dfoxo*²¹ or *dfoxo*²⁵ mutants (Fig. 4m,n). In addition, knock down *dfoxo* by two copies of *dfoxo-IR* also produced a weak notch phenotype in the wing margin (Fig. S5c). Hence, dFoxO is physiologically required for the wing patterning functions of endogenous Wg signaling.

dFoxO is required for the activation of Wg pathway target genes. To further confirm that dFoxO is required for Wg signaling activity, we checked the expression of *wingful* (*wf*), a known Wg pathway target gene that mimic *wg* expression pattern, using a *wf*-LacZ reporter (Fig. 5a)³⁵. Activation of Wg signaling by *ptc* > Arm along the A/P compartment boundary of wing discs strongly up-regulated *wf* transcription (Fig. 5b). The elevated *wf* expression was significantly suppressed by the expression of a *dfoxo* RNAi (Fig. 5d) or in heterozygous *dfoxo* mutants (Fig. 5e,f), but not that of GFP (Fig. 5c). Similar results were obtained in salivary glands where loss of *dfoxo* suppressed Arm-induced *wf*-LacZ expression (Fig. 5g–i). To verify the *wf*-LacZ reporter data, we carried out qRT-PCR experiments to analyze the transcription of endogenous *wf* and another Wg signaling target gene *senseless* (*sens*), which is required for bristle formation³⁶. We confirmed that loss of *dfoxo* suppressed *sd* > Arm induced *wf* and *sens* expression in wing discs (Fig. S7). Thus, dFoxO is required for the transcriptional up-regulation of Wg pathway target genes.

Arm binds to dFoxO and is reciprocally required for dFoxO-induced cell death. When Flag-tagged dFoxO and Myc-tagged Arm were co-expressed in *Drosophila* S2 cell, Flag-dFoxO could be immunoprecipitated with Myc-Arm, and *vice versa* (Fig. 6a). Furthermore, when Arm and GFP-dFoxO were co-expressed in the fly eye, Arm could be immunoprecipitated with GFP-dFoxO, and *vice versa* (Fig. 6b), suggesting dFoxO interacts with Arm both *in vitro* and *in vivo*. Furthermore, we co-expressed Flag-dFoxO, Myc-Arm and HA-Pan in S2 cells, and performed co-IP experiments with an anti-Flag antibody. We found that the antibody not only




Figure 3. dFoxO is required for Arm-induced *hid* and *rpr* expression. Compared with the *sd*-Gal4 control (a,g), expression of Arm activates *rpr*-LacZ (b) and *hid*-LacZ (h) expression in the wing pouch, which remain unaffected by the expression of GFP (c,i), but are significantly suppressed by knocking-down of *dfoxo* (d,j), or heterozygous mutation of *dfoxo*²¹ (e,k) or *dfoxo*^{Δ94} (f,l). Scale bars: 100 μm.

precipitated Flag-dFoxO, but also Myc-Arm and HA-Pan (Fig. S8), suggesting all three factors exist in the same complex.

Next we examined whether Arm is reciprocally required for the function of dFoxO. Expression of dFoxO in the eye driven by *GMR*-Gal4 also induced cell death and generated a small eye phenotype²⁸ (Fig. 6d). The *GMR* > dFoxO-triggered small eye phenotype was suppressed by knocking-down of *arm* or *arm* mutation (Fig. 6f-h), but remained unaffected by the expression of LacZ (Fig. 6e). Consistently, loss of *pan* also suppressed dFoxO-induced small eye phenotype (Fig. 6i,j). As a positive control, this phenotype was fully suppressed by knocking-down of *dfoxo* (Fig. 6k). Therefore, dFoxO and Arm are reciprocally required for each other's function, possibly through their physical interaction.

Discussion

In this work, we show that dFoxO promotes Wg signaling-induced cell death, wing pattern formation and target genes expression, while Arm is reciprocally required for dFoxO-induced cell death *in vivo*. Our data are consistent with previous study that β-catenin promotes FoxO activity²¹, but contradict to others that FoxO competes




Figure 4. dFoxO is required for the wing patterning functions of Wg signaling. Light micrographs of *Drosophila* adult wings are shown. Compared with the *ptc*-Gal4 control (a, the right panel shows high magnification of the boxed area containing ACV in the left panel), expression of Arm induces ectopic bristles (indicated by the red arrow head) and loss-of-ACV phenotypes (b), which remain unaffected by the expression of LacZ (c), but are largely suppressed by RNAi-mediated knocking-down of *dfoxo* (d) or heterozygous mutation of *dfoxo*^{Δ94} (e). Compared with the *sd*-Gal4 control (f, the right panel shows high magnification of the boxed area containing anterior-distal wing margin in the left panel), expression of Arm induces ectopic bristles near the wing margin (g, indicated by red arrow heads), which remains unaffected by the expression of LacZ (h), but is suppressed by knocking-down of *dfoxo* (i) or heterozygous mutation of *dfoxo*^{Δ94} (j). knocking-down of *wg* by *ptc*-Gal4 generates a mild wing margin notch phenotype between the L3 and L4 veins (k, indicated by the red box and amplified in the right panel), which is rescued by the expression of dFoxO (l), but is enhanced by heterozygous mutation of *dfoxo*²¹ (m) or *dfoxo*²⁵ (n). Sample numbers: a, 226; b, 201; c, 108; d, 197; e, 160; f, 168; g, 166; h, 146; i, 130; j, 178; k, 160; l, 62; m, 174; n, 226. Scale bars: 100 μm in (a–n) and 50 μm in high magnification figures.

with TCF for the limited pool of β -catenin and thereby inhibits Wnt signaling activity^{22,23}. A possible explanation for this discrepancy is that FoxO may regulate Wg/Wnt signaling in a tissue/cell type specific manner, depending on the presence of other transcriptional activating or repressing factor(s), or the different levels of FoxO and β -catenin/Arm in distinct cellular context. Importantly, while previous results were obtained mainly from the *in vitro* or gain-of-function studies, we have examined both loss-of-function and gain-of-function effects of *dfoxo* on ectopic and endogenous Wg signaling *in vivo*, and thus, the data are more likely to reflect the physiological situation. Moreover, the positive role of dFoxO in Wg signaling-induced cell death is in line with the well-known function of FoxO in promoting cell death.

The present study revealed for the first time that dFoxO acts as a vital regulator of Wg signaling-triggered cell death *in vivo*, which expands our knowledge of Wnt/Wg signaling function and the broad array of FoxOs' regulatory activities. Both FoxO and Wnt/Wg signaling have been implicated in various cancers^{2,3,12,13,37}, and a




Figure 5. dFoxO is required for Wg target gene activation. Light micrographs of *Drosophila* 3rd instar wing discs (a–f) and salivary glands (g–l) with X-Gal staining are shown. Compared with the *ptc*-Gal4 control (a,g), ectopic Arm-induced *wf*-LacZ expression in the wing disc (b, indicated by black arrow heads) or salivary gland (h) remains unaffected by the expression of GFP (c and i), but is significantly suppressed by knocking-down of *dfoxo* (d,j), or heterozygous mutation of *dfoxo*²⁵ (e,k) or *dfoxo*^{Δ94} (f,l). Scale bars: 100 μm in (a–f) and 200 μm in (g–l).

FOXO3a/β-catenin/GSK-3β signaling is essential for cell proliferation and apoptosis in prostate cancer cells³⁸, further study is needed to understand their interaction in tumorigenesis.

Materials and Methods

***Drosophila* strains.** The following lines have been described previously: *UAS-Wg*, *UAS-Dsh*, *UAS-Arm*, *UAS-Pan*, *UAS-wg-IR*, *UAS-arm-IR*^{#1}, *UAS-arm-IR*^{#2}, *arm*^{XM19}, *UAS-pan-IR*^{#1}, *GMR* > *Egr*, *UAS-Bsk*^{DN}, *bsk*^l,




Figure 6. Arm binds to dFoxO and is required for dFoxO-induced cell death. (a) Co-immunoprecipitation between transfected Myc-Arm and Flag-dFoxO in *Drosophila* S2 cells. (b) Co-immunoprecipitation between ectopically expressed Arm and GFP-dFoxO in *Drosophila* eye. (c–k) Light micrographs of *Drosophila* adult eyes are shown. Compared with the *GMR*-Gal4 control (c), expression of dFoxO triggers cell death and produces a small eye phenotype (d), which remains unaffected by the expression of LacZ (e), but is considerably suppressed by the expression of two independent *arm* RNAi (f,g), or heterozygous mutation of *arm* (h), or the expression of two independent *pan* RNAi (i,j). The *GMR* > dFoxO eye phenotype is also suppressed by knocking-down of *dfoxo* (k), which serves as a positive control. Sample numbers from c to k: c, 123; d, 104; e, 74; f, 67; g, 73; h, 64; i, 97; j, 106; k, 53. Scale bars: 100 μ m.

en-Gal4²⁶; *UAS-bsk-IR*²⁹; *UAS-dFoxO-GFP*-3³⁹; *UAS-dfoxo-IR*, *dfoxo* ^{Δ 94}, *ptc*-Gal4, *sd*-Gal4²⁰; *UAS-dFoxO^P*, *UAS-dFoxO^{W2}*, *dfoxo*²¹, *dfoxo*^{25,40}, *UAS-LacZ*⁴¹; *UAS-GFP*⁴², *GMR*-Gal4⁴³, *omb*-Gal4¹⁹, *hid*-LacZ, *rpr*-LacZ⁴⁴, *wf*-LacZ³⁵. The *UAS-Dcr2^{D2}* (24650), *UAS-Arm*⁵² (4783), *UAS-pan-IR*⁵² (26743) and *bsk*² (5283) lines were obtained from the Bloomington stock center.

X-Gal staining. X-Gal staining was done as described⁴⁵, discs and salivary glands were dissected in PBST at the 3rd instar larva stage and fixed in 4% formaldehyde for 15 minutes, rinse the tissue once in PBST buffer containing 3.3 mM K₃Fe(CN)₆ and 3.3 mM K₄Fe(CN)₆, then incubate in the above PBST buffer with 0.2% X-gal at room temperature for 24 hours, photographs were taken under light microscope.

AO staining. AO staining was done as described⁴⁶, discs were dissected at the 3rd instar larva stage, and stained in 1 × 10⁻⁵M AO solution for 5 minutes, photographs were taken under fluorescence microscope.

TUNEL staining. TUNEL staining was done as described²⁶, discs were dissected at 3rd instar larva stage, and stained for TUNEL using the Fluorescein Cell Death Kit (Boster), photographs were taken under Leica confocal microscope.

Immunostaining. Discs were dissected at 3rd instar larva stage in PBS, fixed in 4% formaldehyde for 20 minutes and washed in 0.3% PBST for 3 times, then label with primary antibody overnight, and secondary antibody for 2 hours. Primary antibodies used were 1:100 mouse-anti-BrdU (Sigma) and 1:400 rabbit-anti-PH3 (Cell Signaling Technology, CST), secondary antibodies used were 1:1000 goat anti-mouse CY3 (CST) and 1:1000 goat anti-rabbit CY3 (CST). Photographs were taken under fluorescence microscope.

BrdU incorporation. BrdU incorporation was done as described⁴⁷, discs were dissected at 3rd instar larva stage in Schneider's medium (Sigma), incubated for 40 minutes in Schneider's medium containing 0.2 mg/ml of BrdU, and fixed in 4% formaldehyde, then washed in PBST and hydrolyzed in 2 N HCl. After that the discs were blocked in 10% horse serum and labeled with mouse anti-BrdU and anti-mouse CY3 antibodies, photographs were taken under fluorescence microscope.

Co-immunoprecipitation. S2 cells were transfected with indicated plasmids for 48 h, immunoprecipitation and western analyses were performed as previously described⁴³, pre-cleared cell lysates were incubated with the indicated antibodies followed by precipitation with protein G Sepharose beads (Sigma). Immune complexes were washed with lysis buffer, eluted in 2 × SDS sample buffer, and then subjected to western blot using corresponding antibodies. Fly heads were cut from indicated genotypes and homogenized in lysis buffer, immunoprecipitation and western analyses were performed as above. Antibodies used in this study were as follows: mouse anti-Arm antibody (DSHB), mouse anti-Flag (Sigma), rabbit anti-GFP (CST), mouse anti-Myc (CST), rabbit anti-HA (CST) and goat anti-mouse IgG-HRP (CST).

qRT-PCR. Wing discs were dissected at 3rd instar larva, for each genotype, more than 100 discs were collected and total RNA was isolated using TRIzol (Invitrogen). qRT-PCR was performed as previously described and *rp49* was used as internal control⁴⁸. Primer sequences were
wf fwd: AAGTCGAGCAATGGGCAATGAT,
 rev: TGGAGGAGCGTGTCTTCTG;
sens fwd: CCGAAAAGGAGCATGAACTC,
 rev: CGCTGTTGCTGTGGTGTACT³⁶;
rp49 fwd: CATCCGCCAGCATAACAG,
 rev: CCATTTGTGCGACAGCTTAG⁴⁸.

Statistics. For pupal retina AO staining, wing size measurement and PH3 staining, unpaired t-text analysis was used. For qRT-PCR, one-way analysis of variance test followed by the post Dunnett test was used. A *P* value of less than 0.05 was considered significant.

References

- Clevers, H. & Nusse, R. Wnt/beta-catenin signaling and disease. *Cell* **149**, 1192–1205, doi: 10.1016/j.cell.2012.05.012 (2012).
- Anastas, J. N. & Moon, R. T. WNT signalling pathways as therapeutic targets in cancer. *Nature reviews. Cancer* **13**, 11–26, doi: 10.1038/nrc3419 (2013).
- MacDonald, B. T., Tamai, K. & He, X. Wnt/beta-catenin signaling: components, mechanisms, and diseases. *Developmental cell* **17**, 9–26, doi: 10.1016/j.devcel.2009.06.016 (2009).
- Nusse, R. Wnt signaling in disease and in development. *Cell Res* **15**, 28–32, doi: 10.1038/sj.cr.7290260 (2005).
- Miller, J. R. The Wnts. *Genome biology* **3**, REVIEWS3001 (2002).
- Swarup, S. & Verheyen, E. M. Wnt/Wingless signaling in Drosophila. *Cold Spring Harbor perspectives in biology* **4**, doi: 10.1101/cshperspect.a007930 (2012).
- van Amerongen, R. & Nusse, R. Towards an integrated view of Wnt signaling in development. *Development* **136**, 3205–3214, doi: 10.1242/dev.033910 (2009).
- Cadigan, K. M. & Nusse, R. Wnt signaling: a common theme in animal development. *Genes & development* **11**, 3286–3305 (1997).
- Cordero, J., Jassim, O., Bao, S. & Cagan, R. A role for wingless in an early pupal cell death event that contributes to patterning the Drosophila eye. *Mechanisms of development* **121**, 1523–1530, doi: 10.1016/j.mod.2004.07.004 (2004).
- Lin, H. V., Rogulja, A. & Cadigan, K. M. Wingless eliminates ommatidia from the edge of the developing eye through activation of apoptosis. *Development* **131**, 2409–2418, doi: 10.1242/dev.01104 (2004).
- Cordero, J. B. & Cagan, R. L. Canonical wingless signaling regulates cone cell specification in the Drosophila retina. *Developmental dynamics: an official publication of the American Association of Anatomists* **239**, 875–884, doi: 10.1002/dvdy.22235 (2010).
- Greer, E. L. & Brunet, A. FOXO transcription factors at the interface between longevity and tumor suppression. *Oncogene* **24**, 7410–7425, doi: 10.1038/sj.onc.1209086 (2005).
- Maiese, K., Chong, Z. Z., Shang, Y. C. & Hou, J. A “FOXO” in sight: targeting Foxo proteins from conception to cancer. *Medicinal research reviews* **29**, 395–418, doi: 10.1002/med.20139 (2009).
- Calnan, D. R. & Brunet, A. The FoxO code. *Oncogene* **27**, 2276–2288, doi: 10.1038/onc.2008.21 (2008).
- Zhao, Y., Wang, Y. & Zhu, W. G. Applications of post-translational modifications of FoxO family proteins in biological functions. *Journal of molecular cell biology* **3**, 276–282, doi: 10.1093/jmcb/mjr013 (2011).

16. Puig, O., Marr, M. T., Ruhf, M. L. & Tjian, R. Control of cell number by Drosophila FOXO: downstream and feedback regulation of the insulin receptor pathway. *Genes & development* **17**, 2006–2020, doi: 10.1101/gad.1098703 (2003).
17. Lam, E. W., Francis, R. E. & Petkovic, M. FOXO transcription factors: key regulators of cell fate. *Biochemical Society transactions* **34**, 722–726, doi: 10.1042/BST0340722 (2006).
18. Huang, H. & Tindall, D. J. Dynamic FoxO transcription factors. *Journal of cell science* **120**, 2479–2487, doi: 10.1242/jcs.001222 (2007).
19. Wu, C. *et al.* Pelle Modulates dFoxO-Mediated Cell Death in Drosophila. *PLoS genetics* **11**, e1005589, doi: 10.1371/journal.pgen.1005589 (2015).
20. Wang, X. *et al.* FoxO mediates APP-induced AICD-dependent cell death. *Cell Death and Disease* **5**, e1233, doi: 10.1038/cddis.2014.196 (2014).
21. Essers, M. A. *et al.* Functional interaction between beta-catenin and FOXO in oxidative stress signaling. *Science* **308**, 1181–1184, doi: 10.1126/science.1109083 (2005).
22. Almeida, M., Han, L., Martin-Millan, M., O'Brien, C. A. & Manolagas, S. C. Oxidative stress antagonizes Wnt signaling in osteoblast precursors by diverting beta-catenin from T cell factor- to forkhead box O-mediated transcription. *The Journal of biological chemistry* **282**, 27298–27305, doi: 10.1074/jbc.M702811200 (2007).
23. Hoogeboom, D. *et al.* Interaction of FOXO with beta-catenin inhibits beta-catenin/T cell factor activity. *The Journal of biological chemistry* **283**, 9224–9230, doi: 10.1074/jbc.M706638200 (2008).
24. Xie, X. W., Liu, J. X., Hu, B. & Xiao, W. Zebrafish foxo3b negatively regulates canonical Wnt signaling to affect early embryogenesis. *PLoS one* **6**, e24469, doi: 10.1371/journal.pone.0024469 (2011).
25. Iyer, S. *et al.* FOXOs attenuate bone formation by suppressing Wnt signaling. *The Journal of clinical investigation* **123**, 3409–3419, doi: 10.1172/JCI168049 (2013).
26. Zhang, S. *et al.* The canonical Wg signaling modulates Bsk-mediated cell death in Drosophila. *Cell death & disease* **6**, e1713, doi: 10.1038/cddis.2015.85 (2015).
27. Essers, M. A. *et al.* FOXO transcription factor activation by oxidative stress mediated by the small GTPase Ral and JNK. *The EMBO journal* **23**, 4802–4812, doi: 10.1038/sj.emboj.7600476 (2004).
28. Luo, X., Puig, O., Hyun, J., Bohmann, D. & Jasper, H. Foxo and Fos regulate the decision between cell death and survival in response to UV irradiation. *The EMBO journal* **26**, 380–390, doi: 10.1038/sj.emboj.7601484 (2007).
29. Wu, C. *et al.* Toll pathway modulates TNF-induced JNK-dependent cell death in Drosophila. *Open biology* **5**, 140171, doi: 10.1098/rsob.140171 (2015).
30. Cashio, P., Lee, T. V. & Bergmann, A. Genetic control of programmed cell death in Drosophila melanogaster. *Seminars in cell & developmental biology* **16**, 225–235, doi: 10.1016/j.semcdb.2005.01.002 (2005).
31. Couso, J. P., Bishop, S. A. & Martinez Arias, A. The wingless signalling pathway and the patterning of the wing margin in Drosophila. *Development* **120**, 621–636 (1994).
32. Zhang, J. & Carthew, R. W. Interactions between Wingless and Dfz2 during Drosophila wing development. *Development* **125**, 3075–3085 (1998).
33. Phillips, R. G. & Whittle, J. R. wingless expression mediates determination of peripheral nervous system elements in late stages of Drosophila wing disc development. *Development* **118**, 427–438 (1993).
34. Zeng, Y. A., Rahnema, M., Wang, S., Lee, W. & Verheyen, E. M. Inhibition of Drosophila Wg signaling involves competition between Mad and Armadillo/beta-catenin for dTcf binding. *PLoS one* **3**, e3893, doi: 10.1371/journal.pone.0003893 (2008).
35. Gerlitz, O. & Basler, K. Wingful, an extracellular feedback inhibitor of Wingless. *Genes & development* **16**, 1055–1059, doi: 10.1101/gad.991802 (2002).
36. Eivers, E. *et al.* Mad is required for wingless signaling in wing development and segment patterning in Drosophila. *PLoS one* **4**, e6543, doi: 10.1371/journal.pone.0006543 (2009).
37. Reya, T. & Clevers, H. Wnt signalling in stem cells and cancer. *Nature* **434**, 843–850, doi: 10.1038/nature03319 (2005).
38. Li, Y. *et al.* Regulation of FOXO3a/beta-catenin/GSK-3beta signaling by 3,3'-diindolylmethane contributes to inhibition of cell proliferation and induction of apoptosis in prostate cancer cells. *The Journal of biological chemistry* **282**, 21542–21550, doi: 10.1074/jbc.M701978200 (2007).
39. Wagner, C., Isermann, K. & Roeder, T. Infection induces a survival program and local remodeling in the airway epithelium of the fly. *FASEB journal: official publication of the Federation of American Societies for Experimental Biology* **23**, 2045–2054, doi: 10.1096/fj.08-114223 (2009).
40. Junger, M. A. *et al.* The Drosophila forkhead transcription factor FOXO mediates the reduction in cell number associated with reduced insulin signaling. *Journal of biology* **2**, 20, doi: 10.1186/1475-4924-2-20 (2003).
41. Hu, Y., Han, Y., Wang, X. & Xue, L. Aging-related neurodegeneration eliminates male courtship choice in Drosophila. *Neurobiology of aging*, doi: 10.1016/j.neurobiolaging.2014.02.026 (2014).
42. Ma, X. *et al.* Src42A modulates tumor invasion and cell death via Ben/dUev1a-mediated JNK activation in Drosophila. *Cell death & disease* **4**, e864, doi: 10.1038/cddis.2013.392 (2013).
43. Ma, X. *et al.* Bendless modulates JNK-mediated cell death and migration in Drosophila. *Cell death and differentiation* **21**, 407–415, doi: 10.1038/cdd.2013.154 (2014).
44. Ma, X. *et al.* NOPO modulates Egr-induced JNK-independent cell death in Drosophila. *Cell Res* **22**, 425–431, doi: 10.1038/cr.2011.135 (2012).
45. Xue, L. & Noll, M. Dual role of the Pax gene paired in accessory gland development of Drosophila. *Development* **129**, 339–346 (2002).
46. Ma, X. *et al.* dUev1a modulates TNF-JNK mediated tumor progression and cell death in Drosophila. *Developmental biology* **380**, 211–221, doi: 10.1016/j.ydbio.2013.05.013 (2013).
47. Peng, F. *et al.* Loss of Polo ameliorates APP-induced Alzheimer's disease-like symptoms in Drosophila. *Scientific reports* **5**, 16816, doi: 10.1038/srep16816 (2015).
48. Beck, E. S. *et al.* Regulation of Fasciclin II and synaptic terminal development by the splicing factor beag. *The Journal of neuroscience: the official journal of the Society for Neuroscience* **32**, 7058–7073, doi: 10.1523/JNEUROSCI.3717-11.2012 (2012).

Acknowledgements

We thank Haiyun Song, Ernst Hafen, the Bloomington Drosophila Stock Center and the Vienna Drosophila RNAi Center for fly stocks, members of Xue lab for discussion and critical comments. This work is supported by the National Basic Research Program of China (973 Program) (2011CB943903), National Natural Science Foundation of China (31171413, 31371490, 31571516), the Specialized Research Fund for the Doctoral Program of Higher Education of China (20120072110023 and 20120072120030), the China Postdoctoral Science Foundation (2015M581659), and Shanghai Committee of Science and Technology (09DZ2260100, 14JC1406000).

Author Contributions

S.Z., W.L. and L.X. conceived and designed the experiments. S.Z., X.G., C.C., Y.C., J.L., C.W., Y.Y. and W.L. performed the experiments. S.Z., Y.S., C.J., W.L. and L.X. analyzed the data, S.Z. and L.X. wrote the manuscript.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Zhang, S. *et al.* dFoxO promotes Wingless signaling in *Drosophila*. *Sci. Rep.* **6**, 22348; doi: 10.1038/srep22348 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>