

Rhodococcus equi's Extreme Resistance to Hydrogen Peroxide Is Mainly Conferred by One of Its Four Catalase Genes

Pauline Bidaud¹*, Laurent Hébert¹*, Corinne Barbey¹^{‡a}, Anne-Cécile Appourchaux¹^{‡b}, Riccardo Torelli², Maurizio Sanguinetti², Claire Laugier¹, Sandrine Petry¹*

1 Dozulé Laboratory for Equine Diseases, Unit Bacteriology and Parasitology, ANSES, Goustranville, France, **2** Institute of Microbiology, Catholic University of Sacred Heart, Rome, Italy

Abstract

Rhodococcus equi is one of the most widespread causes of disease in foals aged from 1 to 6 months. *R. equi* possesses antioxidant defense mechanisms to protect it from reactive oxygen metabolites such as hydrogen peroxide (H₂O₂) generated during the respiratory burst of phagocytic cells. These defense mechanisms include enzymes such as catalase, which detoxify hydrogen peroxide. Recently, an analysis of the *R. equi* 103 genome sequence revealed the presence of four potential catalase genes. We first constructed $\Delta katA$ -, $\Delta katB$ -, $\Delta katC$ - and $\Delta katD$ -deficient mutants to study the ability of *R. equi* to survive exposure to H₂O₂ *in vitro* and within mouse peritoneal macrophages. Results showed that $\Delta katA$ and, to a lesser extent $\Delta katC$, were affected by 80 mM H₂O₂. Moreover, *katA* deletion seems to significantly affect the ability of *R. equi* to survive within murine macrophages. We finally investigated the expression of the four catalases in response to H₂O₂ assays with a real time PCR technique. Results showed that *katA* is overexpressed 367.9 times (± 122.6) in response to exposure to 50 mM of H₂O₂ added in the stationary phase, and 3.11 times (± 0.59) when treatment was administered in the exponential phase. In untreated bacteria, *katB*, *katC* and *katD* were overexpressed from 4.3 to 17.5 times in the stationary compared to the exponential phase. Taken together, our results show that *KatA* is the major catalase involved in the extreme H₂O₂ resistance capability of *R. equi*.

Citation: Bidaud P, Hébert L, Barbey C, Appourchaux A-C, Torelli R, et al. (2012) *Rhodococcus equi*'s Extreme Resistance to Hydrogen Peroxide Is Mainly Conferred by One of Its Four Catalase Genes. PLoS ONE 7(8): e42396. doi:10.1371/journal.pone.0042396

Editor: Roy Martin Roop II, East Carolina University School of Medicine, United States of America

Received: May 16, 2012; **Accepted:** July 4, 2012; **Published:** August 6, 2012

Copyright: © 2012 Bidaud et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: P. Bidaud and L. Hébert were funded by a grant awarded by Regional Council of Low-Normandy and by Research and Higher Education Ministry, within the project R25_p3 (CPER 2007–2013). C. Barbey was supported by a thesis grant awarded by Anses and Regional Council of Low-Normandy. Financial support was obtained from Institut Français du Cheval et de l'Équitation. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: sandrine.petry@anses.fr

‡ These authors contributed equally to this work.

‡a Current address: Laboratory of Microbiology Signals and Microenvironment, Normandy University, EA 4312 University of Rouen, IUT Evreux, Evreux, France

‡b Current address: Frank Duncombe Laboratory, Animal Health Department, IFR146 ICORE, Caen, France

Introduction

Rhodococcus equi is a Gram-positive, facultative intracellular pathogen affecting foals up to six months of age. The major manifestation of *R. equi* infection is chronic suppurative broncho-pneumonia, but 50% of contaminated foals also show clinical signs of intestinal disease and occasionally septic arthritis or osteomyelitis [1]. It is now well known that, in addition to being a veterinary concern, *R. equi* is an important opportunistic zoonotic pathogen mainly identified in immunocompromised people, and particularly AIDS patients [2,3]. Interestingly, different cases of infection by *R. equi* have also been reported in immunocompetent patients [4,5].

R. equi pathogenicity is closely linked to the bacterium's ability to survive and replicate in the lung macrophages of infected foals [6], where bacteria are generally killed by the combined action of a low pH (4.0–5.0), hydrolytic enzymes (including proteases, lipases, DNAses and RNAses) and the production of reactive oxygen metabolites such as hydrogen peroxide, H₂O₂ [7]. *R. equi*

has been characterized as being able to survive and multiply within a membrane-bound vacuole inside macrophages. This capability is conferred by the presence of a 80–90 kb virulence plasmid which inhibits i) the maturation of phagosomes, leading to the absence of phagosome acidification and prevents ii) the fusion of phagosomes with the lysosomes containing many hydrolytic enzymes [7]. In addition, *R. equi* must be able to mount an antioxidant defense system to ensure its persistence and multiplication within the highly oxidative environment of host macrophages generated notably by the reactive oxygen species (ROS) producing phagocyte oxidase Nox2 [8]. It has already been shown that *R. equi* is highly resistant to the action of H₂O₂ [9], though plasmid-encoded proteins do not play a role in *R. equi*'s resistance to H₂O₂ [9]. An analysis of the *R. equi* 103 genome revealed the presence of numerous genes involved in oxidative stress resistance. Four genes in particular display homology with catalases [10], which are enzymes responsible for detoxifying H₂O₂ [11].

This was the background for our investigation into the causes of the extreme resistance of *R. equi* to oxidative stress. Our study therefore focused on an analysis of the four potential catalase genes *katA*, *katB*, *katC* and *katD*. We studied the role of these proteins in H₂O₂ resistance and in the ability of *R. equi* to survive within mouse peritoneal macrophages. These results were confirmed with the expression analysis of the catalase genes in response to H₂O₂ exposure.

Results

Identification of Four Genes Potentially Involved in H₂O₂ Resistance

The *in silico* analysis of the genome sequence of the *R. equi* 103 strain [10] revealed the presence of four potential catalase genes, designated *katA*, *katB*, *katC* and *katD*. These four genes are at separate locations on the chromosome and do not seem to share genetic links. Sequence analysis of these proteins shows that KatA, KatB and KatC share the conserved catalytic residues and metal ligand amino acids of members of monofunctional heme catalases [12]. KatD shares the conserved catalytic residues and metal ligand amino acids of members of the nonheme catalases. These nonheme catalases utilize manganese ions instead of ferric heme in their active site and are therefore also known as manganese catalases [13]. No catalase of the bifunctional catalase-peroxidase family was identified. Phylogenetic relationships between the four catalases are represented in Figure 1, which shows that the three heme and manganese catalases constitute a distinct group, and that KatA and KatB are closely related. No signal sequence and/or transmembrane domain were identified for these four catalases by TMHMM [14] or SignalP [15] analysis.

In order to estimate the involvement of each catalase in *R. equi*'s oxidative stress resistance, we constructed corresponding mutants. To define the physiological roles of the four catalases, we studied the growth characteristics of the four deficient mutants. The characteristics were the same as those of the wild type (WT) strain under aerobic growth in BHI medium (data not shown). This similarity demonstrates that the catalases are not essential to cell viability under aerobic conditions.

$\Delta katA$ and $\Delta katC$ are the most Sensitive to H₂O₂ Treatment

To evaluate the effect of exogenous H₂O₂ stress on the survival of each mutant, cells were challenged with 80 mM of H₂O₂ under agitation, and CFUs determined 30 min after treatment (Figure 2). $\Delta katA$ was the most sensitive mutant to H₂O₂ since it did not survive when treated in the stationary phase, and its survival decreased by two orders of magnitude compared to the WT strain when treated in the exponential phase. $\Delta katC$ survived as well as the WT strain in the stationary phase but in contrast, when the cells were in the exponential phase, $\Delta katC$ was the most susceptible to H₂O₂ exposure with a decrease in survival of over three magnitudes compared to the WT strain. Both $\Delta katB$ and $\Delta katD$




Figure 1. Phylogenetic relationships between the catalases of *Rhodococcus equi* 103. Catalases were aligned using Phylogeny.fr (<http://www.phylogeny.fr/>) [37]. doi:10.1371/journal.pone.0042396.g001

had the same resistance capability as the WT strain, whether in exponential or stationary phases. The drop in survival rates for $\Delta katA$ and $\Delta katC$ after exposure to H₂O₂ demonstrate that, in these experimental conditions, only KatA and KatC primarily contribute to the oxidative defense system of *R. equi*.

$\Delta katA$ is the most Susceptible to Elimination by Macrophages

During phagocytosis, phagocytic cells generate superoxide and other ROS involved in antibacterial activity [16]. This study compared the intracellular survival of the WT and $\Delta katA$, $\Delta katB$, $\Delta katC$ and $\Delta katD$ strains inside infected mouse peritoneal macrophages (Figure 3). No significant difference was observed in the levels of *R. equi* strains recovered 8 h post infection, suggesting that the different strains possessed similar abilities to infect macrophages (Figure 3). $\Delta katA$ was shown to be the most susceptible mutant, completely eliminated by macrophages 72 h post infection ($P < 0.05$). $\Delta katB$, $\Delta katC$ and $\Delta katD$ were shown to be more susceptible than the WT strain to macrophage killing at 48- and 72-h time points ($P < 0.05$) but were not completely eliminated from macrophages 72 h post infection. The survival rate of each strain was similar by 24 h post infection but the WT strain's ability to survive intracellularly was greater than that of $\Delta katB$, $\Delta katC$ and $\Delta katD$ at 48- and 72-h time points ($P < 0.05$). The inability of $\Delta katA$ to survive in macrophages beyond 72 h reveals KatA's involvement in the intramacrophage resistance of *R. equi* to H₂O₂.

katA is Overexpressed in Response to H₂O₂ Treatment

We evaluated the effect of a sublethal H₂O₂ concentration (50 mM) defined previously (data not shown) on the expression of *katA*, *katB*, *katC* and *katD* transcripts during exponential (Figure 4A) and stationary phases (Figure 4B). The expression of catalases was measured at several timepoints: 5, 10 and 20 minutes after treatment in the exponential phase and 5, 10, 20, 30 and 60 minutes after treatment in the stationary phase. Results showed that only the *katA* gene was overexpressed in response to H₂O₂ treatment. For example, when H₂O₂ was added in the exponential phase, *katA* was overexpressed 3.11 (± 0.59) times in bacteria treated for 5 min compared to untreated bacteria (Figure 4A). However, when H₂O₂ was added in the stationary phase, *katA* was overexpressed 367.90 (± 122.63) times in bacteria treated for 10 min compared to untreated bacteria (Figure 4B). We did not observe any modifications in the rate of transcripts for *katB*, *katC* or *katD* (Figures 4A and 4B).

katB, *katC* and *katD* are Overexpressed in the Stationary Growth Phase Compared to the Exponential Phase

In untreated bacteria, we observed overexpressions of *katB*, *katC* and *katD* in the stationary phase compared to the exponential phase. Irrespective of exposure to oxidative stress, *katB* was overexpressed 17.55 (± 3.22) times in the stationary phase compared to the exponential phase (Figure 5). *katC* and *katD* were overexpressed 4.51 (± 0.59) and 4.31 (± 0.50) times in the stationary phase compared to the exponential phase, respectively (Figure 5).

Discussion

Resistance to oxidative stress is one of the key processes that allow pathogens to survive within macrophages, which is why pathogenic bacteria have developed different antioxidative defenses to eliminate ROS or repair their damage [17]. *In silico* analysis shows that *R. equi* 103 contains three monofunctional heme catalases (KatA, KatB and KatC) and a monofunctional




Figure 2. Decrease in $\Delta katA$ and $\Delta katC$ survival after exposure to H₂O₂ (80 mM, 30 min). *R. equi* WT, $\Delta katA$, $\Delta katB$, $\Delta katC$ or $\Delta katD$ were grown to exponential (OD₆₀₀ = 0.2) and stationary growth phases (16 h of growth). The number of bacterial cells was standardized by dilution of stationary phase cells before treatment. Cells were challenged with 80 mM of H₂O₂ for 30 min under agitation. Viability was assayed both before (black bars) and after (grey and white bars) H₂O₂ treatment by plating the bacterial cells on BHI agar. The mean values of three independent experiments are represented, and the standard deviations are indicated. doi:10.1371/journal.pone.0042396.g002

manganese catalase (KatD). The present study addressed the issue of the involvement of these potential H₂O₂-degrading catalase enzymes in *R. equi* virulence and demonstrates that survival following exposure to exogenous H₂O₂ and the efficient proliferation of this bacterium in macrophages mainly rely on the activities of KatA, one of its four catalases.

Benoit *et al.* showed that the extreme resistance of *R. equi* to exposure to a high H₂O₂ concentration (100 mM) did not depend on plasmid-encoded proteins [9]. This was confirmed by the analysis of virulence plasmid–chromosome regulatory crosstalk by Letek *et al.* [10] which shows that the presence of the virulence plasmid did not interfere with the expression of catalase genes. These results suggest that chromosomal genes are key to this function, and genomic analysis of the *R. equi* chromosome

identified an unusually high number of catalases: three mono-functional heme catalases (*katA*, *katB* and *katC*) and one manganese catalase (*katD*). Surprisingly, no bifunctional catalase-peroxidase was identified although for numerous pathogens, these enzymes have already been shown to be involved in the infection of macrophages [18–20].

An analysis of the physiological functions of the four catalases following exposure to exogenously-provided H₂O₂ showed that only KatA and KatC are part of the peroxide defense system of *R. equi*. KatA seemed to be more important than KatC under these stress conditions. The fact that *katB* and *katD* mutations did not affect survival following H₂O₂ exposure suggests that either i) these catalases are non functional, ii) they function as a surrogate in the absence of the primary antioxidant, but their role can only be detected in a double or triple catalase mutant [21] or iii) they have specialized roles for protecting cells against oxidative stress perhaps due to different cellular locations. Although no secretion peptide signal was identified for *R. equi* catalases, it has already been shown that a catalase from *B. subtilis* could be secreted independently of the Sec-secretion apparatus [22,23].

Inactivation of catalase-encoding genes had little if any effect on pathogenicity-associated traits of numerous bacteria including *Salmonella typhimurium* [24], *Yersinia pestis* [25], *Staphylococcus aureus* [26], and *Neisseria gonorrhoeae* [27]. This research demonstrates that significantly lower quantities of viable intracellular bacteria were found in the infected murine macrophages of $\Delta katA$ -deficient mutants during the 72-h infection period than in the WT strain. This suggests that the *katA* gene is linked to *R. equi* virulence mechanisms. This observation could be linked to the oxidative microenvironment present within phagosomes, where ROS and reactive nitrogen species are synthesized in huge quantities following macrophage phagocytosis [17,28].

Studies of physiological functions are complex when multiple catalases are present in a single bacterial cell and differences in regulation are a critical factor in understanding the *in vivo* roles of these enzymes. Our results show that only *katA* expression is induced in response to exogenous H₂O₂. This fast and transitory overexpression is induced mainly in the stationary phase (overexpression factor of 367.9 ± 122.6), which is in accordance with the critical role of *katA* in stationary-phase survival. In




Figure 3. $\Delta katA$ is the most susceptible to elimination by macrophages. Mice were infected with 10⁷ to 10⁸ cells of *R. equi* WT, $\Delta katA$, $\Delta katB$, $\Delta katC$ or $\Delta katD$ by intraperitoneal injection. Intramacrophages bacteria viability was evaluated 24-, 48- and 72 h post infection. The data are the mean numbers of viable intracellular bacteria per 10⁵ macrophages ± standard deviations (error bars) for three independent experiments with three wells in each experiment. doi:10.1371/journal.pone.0042396.g003




Figure 4. *katA* is overexpressed after H₂O₂ treatment (50 mM). *R. equi* WT was treated with 50 mM H₂O₂ in (A) the exponential phase (OD₆₀₀=0.4) or (B) the stationary phase. cDNAs derived from total RNA were used for real time PCR. The overexpression factor in treated bacteria vs untreated bacteria, was calculated using the 2^{-ΔΔCt} method [36]. The data are the mean of overexpression factor ± standard deviation (error bars) of triplicate measurements from four reverse transcriptions of two independent experiments. To evaluate overexpressions in the exponential phase (A), we only considered time points 5, 10 and 20 min post exposure to avoid any effects of bacterial transition in the stationary growth phase. doi:10.1371/journal.pone.0042396.g004

contrast, despite its role in the exponential phase, *katC* is not induced in the stationary phase. This phenomenon, already observed for a catalase-peroxidase of *Legionella pneumophila* [11], is




Figure 5. In untreated bacteria, *katB*, *katC* and *katD* are overexpressed in the stationary growth phase. *R. equi* WT was grown in BHI broth without any H₂O₂ treatment and cells were collected at the exponential (OD of 0.4) and stationary (OD of 1) growth phases for total RNA extraction. cDNAs derived from total RNA, were used for real time PCR. The overexpression factor was calculated using the 2^{-ΔΔCt} method [36]. The data are the mean of overexpression factor in stationary phase collected bacteria compared to exponential phase collected bacteria ± standard deviation of triplicate measurements from four reverse transcriptions of two independent experiments. doi:10.1371/journal.pone.0042396.g005

not yet fully understood. In untreated bacteria, *katB*, *katC* and *katD* expressions increase in the stationary growth phase. These overexpressions could prepare the cells for subsequent stresses [29]. In this context, we could hypothesize that the lack of effect of the mutation of these genes in the stationary phase is due to a compensatory catalase mechanism between them or that these catalases play a role in protecting against the oxidative stress generated by growth in a minimal medium as already described [30]. Our results therefore suggest that *katA* is regulated by a specific stress response transcriptional regulator able to sense the presence of H₂O₂, whereas the other three catalases are under the control of a RpoS-like central regulator of gene expression in the stationary phase [31]. Moreover, other unidentified conditions could induce *katB*, *katC* and *katD* expression. We could suggest, for example, that the expression of manganese catalase *katD* is induced by iron starvation as previously described for the manganese catalase of *Pseudomonas aeruginosa* [32].

Conclusions

This study addressed two fundamental questions: how are the four *R. equi* catalases involved in survival i) following external exposure to H₂O₂ and ii) in mouse peritoneal macrophages? Taken together, our results show that *KatA* is the major determinant of resistance to exogenous H₂O₂ exposure in the

stationary phase and, to a lesser extent, in the exponential phase. KatA is also crucial to *R. equi*'s survival in mouse peritoneal macrophages. Of the other three *R. equi* catalases, only KatC plays a role in resistance to external H₂O₂ exposure, but only in the exponential phase.

Materials and Methods

Ethics Statement

The animal experiments (Survival assays in mouse peritoneal macrophages) were performed under a protocol approved by the Institutional Animal Use and Care Committee at Università Cattolica del S. Cuore, Rome, Italy (Permit number: N21, 12/05/2010) and authorized by the Italian Ministry of Health, according to Legislative Decree 116/92, which implemented the European Directive 86/609/EEC on laboratory animal protection in Italy. Animal welfare was routinely checked by veterinarians of the Service for Animal Welfare.

Bacterial Strains, Plasmids and Culture Conditions

Table 1 lists the bacterial strains and plasmids used in this study. The *R. equi* 103 strain used is devoid of its virulence plasmid. Wild type (WT) or mutant *R. equi* strains and the *Escherichia coli* strain used to construct mutants were routinely grown at 37°C in Brain Heart Infusion (BHI) broth and Luria-Bertani media respectively, with vigorous shaking (200 rpm). When required, antibiotics were added to culture media at the following concentrations: apramycin (30 µg/ml); ampicillin (100 µg/ml) [33]. Dilutions were performed in physiological solution (0.9% NaCl).

Construction of Four Catalase Mutants

A double crossover homologous recombination strategy based on suicide vectors derived from pUC19 containing a cassette consisting of the upstream homologous region, the apramycin

resistance gene (*aacC4*) and the downstream homologous region as previously described [33] was used to inactivate the four catalase genes identified in *R. equi*: REQ4750, REQ44520, REQ26870 and REQ26290, respectively named *ΔkatA*, *ΔkatB*, *ΔkatC* and *ΔkatD*. Briefly, for *ΔkatA* construction, a DNA fragment consisting of the *katA* gene and ≈600 bp up-and downstream of the start and stop codons was amplified with primers L39 and L79 containing respectively *EcoRI* and *XbaI* restriction sites (Table S1). Then, the product was inserted into pUC19 using *EcoRI* and *XbaI* restriction sites. To delete the *katA* gene's internal region, an inverse PCR was performed on the resulting pUC19::*katA* plasmid using divergent primers L162 and L163 with *NsiI* restriction site sequence in order to obtain a linear pUC19-*ΔkatA* plasmid containing a 70% deleted version of *katA*. Separately, the *aacC4* gene was amplified using primers L118 and L119 each containing an *NsiI* restriction site. Both pUC19-*ΔkatA* and *aacC4* amplicons were digested by *NsiI* and ligated to obtain a pUC19-*ΔkatA*::*aacC4* plasmid. This plasmid was introduced into *R. equi* by electroporation, and transformants were selected on BHI agar supplemented with 80 µg/ml apramycin. Allelic exchange double recombinants were selected as previously described [33]. The same procedure was applied for the construction of *ΔkatB*, *ΔkatC* and *ΔkatD*. Construction was verified by Southern blot (data not shown). The primers used are listed in Table S1.

Hydrogen Peroxide Challenge Assays

Ten ml of exponentially growing cultures of *R. equi* WT, *ΔkatA*, *ΔkatB*, *ΔkatC* or *ΔkatD* (Optical Density at 600 nm (OD₆₀₀) of 0.2) were harvested by centrifugation and pellets were resuspended in 10 ml of 0.9% (w/v) NaCl. OD₆₀₀ of stationary growing cultures were measured and the volume necessary for obtaining an OD₆₀₀ of 0.2 in a volume of 10 ml was harvested by centrifugation and resuspended in 10 ml of 0.9% (w/v) NaCl. Both cell suspensions were treated with 80 mM H₂O₂ (Riedel de Haën) and bacterium

Table 1. Bacterial strains and plasmids.

| Bacterial species and plasmids | Bacterial strains and plasmid names | Relevant characteristics | Source or reference |
|-------------------------------------|--|---|---------------------|
| <i>R. equi</i> | 103 ⁻ | plasmid-less wild-type strain | [38] |
| | <i>ΔkatA</i> | 103 ⁻ isogenic derivative REQ4750 deletion mutant | This study |
| | <i>ΔkatB</i> | 103 ⁻ isogenic derivative REQ44520 deletion mutant | This study |
| | <i>ΔkatC</i> | 103 ⁻ isogenic derivative REQ26870 deletion mutant | This study |
| | <i>ΔkatD</i> | 103 ⁻ isogenic derivative REQ26290 deletion mutant | This study |
| <i>E. coli</i> | Top10 | <i>F-mcrA Δ(mrr-hsdRMS-mcrBC) T180lacZΔM15 ΔlacX74 recA1 araD139 galU galK Δ(ara-leu)7697 rpsL (Str^r) endA1 nupG</i> | Invitrogen |
| Plasmids | pRHE2 | pUC19 inserted with <i>aacC4</i> | [33] |
| | pUC19:: <i>katA</i> | pUC19 inserted with <i>katA</i> | This study |
| | pUC19:: <i>katB</i> | pUC19 inserted with <i>katB</i> | This study |
| | pUC19:: <i>katC</i> | pUC19 inserted with <i>katC</i> | This study |
| | pUC19:: <i>katD</i> | pUC19 inserted with <i>katD</i> | This study |
| | pUC19- <i>ΔkatA</i> :: <i>aacC4</i> | pUC19 with <i>ΔkatA</i> :: <i>aacC4</i> mutant allele (suicide vector for <i>katA</i> mutagenesis by gene replacement) | This study |
| | pUC19- <i>ΔkatB</i> :: <i>aacC4</i> | pUC19 with <i>ΔkatB</i> :: <i>aacC4</i> mutant allele (suicide vector for <i>katB</i> mutagenesis by gene replacement) | This study |
| | pUC19- <i>ΔkatC</i> :: <i>aacC4</i> | pUC19 with <i>ΔkatC</i> :: <i>aacC4</i> mutant allele (suicide vector for <i>katC</i> mutagenesis by gene replacement) | This study |
| pUC19- <i>ΔkatD</i> :: <i>aacC4</i> | pUC19 with <i>ΔkatD</i> :: <i>aacC4</i> mutant allele (suicide vector for <i>katD</i> mutagenesis by gene replacement) | This study | |

doi:10.1371/journal.pone.0042396.t001

viability was determined immediately before treatment and after 30 min of incubation at 37°C, 200 rpm, by viable counts of bacteria made by serially diluting samples and plating onto BHI agar. Colony-Forming Units (CFU) were enumerated after 48 h of incubation at 37°C by counting two plates at two different dilutions. All experiments were performed in triplicate.

Survival Assays in Mouse Peritoneal Macrophages

The survival of *R. equi* WT, $\Delta katA$, $\Delta katB$, $\Delta katC$ and $\Delta katD$ in mouse peritoneal macrophages was tested by using an *in vivo/in vitro* infection model as described previously [34,35]. Briefly, after 16 h of growth, bacterial cells were pelleted and resuspended in an adequate volume of phosphate-buffered saline (PBS) for injection. Male BALB/c mice (10 weeks old) were infected with 10^7 to 10^8 cells of each strain by intraperitoneal injection. After an 8 h infection period, peritoneal macrophages were collected by a peritoneal wash, centrifuged then suspended in Dulbecco's modified Eagle's medium containing 10 mM HEPES, 2 mM glutamine, 10% bovine fetal serum, and 1X nonessential amino acids supplemented with vancomycin (10 µg/ml) and gentamicin (150 µg/ml). The cell suspension was dispensed into 24-well tissue culture plates and incubated at 37°C under 5% CO₂ for 2 h. Duplicate wells of infected macrophages were lysed with detergent at 24, 48, and 72 h post infection, and lysates were plated on BHI agar to monitor bacterial survival. All experiments were performed in triplicate, and the results were subjected to statistical analysis using one-way analysis of variance with Bonferroni's correction post test using GraphPad Prism version 5.00 for Windows (GraphPad Software, San Diego, CA).

Catalase Overexpression Assays

R. equi WT in mid-exponential (OD₆₀₀ of 0.4) and stationary (OD₆₀₀ of 1) growth phases was treated with 50 mM H₂O₂. Previous results have shown that 50 mM H₂O₂ is a sublethal concentration (data not shown). A control for each growth phase was performed without H₂O₂ treatment. Bacterial cells were incubated (37°C, 200 rpm) for 5, 10, 20, 30 or 60 min after treatment and total bacterial RNA was isolated: cells were harvested by centrifugation, rinsed in PBS 1X and harvested again by centrifugation. Cells were resuspended in 50 mg/ml lysozyme (Sigma) prepared with Tris/HCl-Ethylenediaminetetraacetic acid (EDTA) buffer, and lysed with a Ribolyser (Hybaid). Samples were incubated for 90 min at 37°C. Total RNA was isolated using the RNeasy RNA mini kit (Qiagen) according to the

manufacturer's instructions. A 15-min on-column DNA digestion was also performed. After elution, RNA was treated with Turbo DNA-free (Ambion) according to the manufacturer's instructions. The concentrations and quality of total RNA were evaluated using Experion (Biorad) with the RNA Std Sens kit, according to the manufacturer's instructions. One microgram of total RNA was reverse-transcribed using the Omniscript enzyme (Qiagen), recombinant RNasin (Promega) and random hexamer primers (Invitrogen) according to the manufacturer's instructions.

The resulting cDNAs were used for subsequent PCR amplification with specific primers, designed using Primer3 software. Primer sequences are listed in Table S1 and were used with the QuantiFast SYBR Green PCR mix (Qiagen) according to the manufacturer's instructions, using a Mastercycler® ep realplex (Eppendorf). The quantification of 16S RNA levels was used as an internal control. Two independent bacterial cultures were used to obtain two different RNA extracts. Each RNA was reverse-transcribed twice in order to obtain four cDNAs. Amplification, detection and real-time analysis were performed in triplicate with the four cDNAs. The overexpression factor (OF) in treated bacteria *vs* control, was calculated using the $2^{-\Delta\Delta Ct}$ method [36]. To evaluate the efficiency of the amplification, a standard curve was constructed using the cycle threshold (C_T) versus 10-fold dilution (data not shown). The reaction's specificity is given by detection of the melting temperatures (T_m) of the amplification products *via* the melting curve.

Supporting Information

Table S1 Oligonucleotide primers used in this study. (DOCX)

Acknowledgments

We would like to thank Anthony Madeline, Arnaud Prehu and Sabrina Ribard for their technical assistance. We also wish to thank Delphine Libby-Claybrough, a professional translator and native English speaker, for carefully reading and reviewing the article.

Author Contributions

Conceived and designed the experiments: SP PB LH MS CL. Performed the experiments: PB LH CB RT ACA. Analyzed the data: LH PB SP. Contributed reagents/materials/analysis tools: PB LH CB RT. Wrote the paper: LH PB.

References

- Giguère S, Cohen ND, Keith Chaffin M, Hines SA, Hondalus MK, et al. (2011) *Rhodococcus equi*: clinical manifestations, virulence, and immunity. *Journal of Veterinary Internal Medicine* 25: 1221–1230.
- Hondalus MK (1997) Pathogenesis and virulence of *Rhodococcus equi*. *Vet Microbiol* 56: 257–268.
- Topino S, Galati V, Grilli E, Petrosillo N (2010) *Rhodococcus equi* infection in HIV-infected individuals: case reports and review of the literature. *AIDS Patient Care STDS* 24: 211–222.
- Devi P, Malhotra S, Chadha A (2011) Bacteremia due to *Rhodococcus equi* in an immunocompetent infant. *Indian J Med Microbiol* 29: 65–68.
- Kedlaya I, Ing MB, Wong SS (2001) *Rhodococcus equi* infections in immunocompetent hosts: case report and review. *Clin Infect Dis* 32: E39–46.
- Hondalus MK, Mosser DM (1994) Survival and replication of *Rhodococcus equi* in macrophages. *Infect Immun* 62: 4167–4175.
- von Bargen K, Haas A (2009) Molecular and infection biology of the horse pathogen *Rhodococcus equi*. *FEMS Microbiol Rev* 33: 870–891.
- Ehrt S, Schnappinger D (2009) Mycobacterial survival strategies in the phagosome: defence against host stresses. *Cell Microbiol* 11: 1170–1178.
- Benoit S, Benachour A, Taoji S, Auffray Y, Hartke A (2002) H₂O₂, which causes macrophage-related stress, triggers induction of expression of virulence-associated plasmid determinants in *Rhodococcus equi*. *Infect Immun* 70: 3768–3776.
- Letek M, González P, MacArthur I, Rodríguez H, Freeman TC, et al. (2010) The genome of a pathogenic *Rhodococcus*: cooptive virulence underpinned by key gene acquisitions. *PLoS Genet* 6: e1001145.
- Bandyopadhyay P, Steinman HM (2000) Catalase-peroxidases of *Legionella pneumophila*: cloning of the *katA* gene and studies of KatA function. *J Bacteriol* 182: 6679–6686.
- Loewen PC, Klotz MG, Hassett DJ (2000) Catalase—an “old” enzyme that continues to surprise us. *ASM News* 66: 76–82.
- Amo T, Atomi H, Imanaka T (2002) Unique presence of a manganese catalase in a hyperthermophilic archaeon, *Pyrobaculum calidifontis* VA1. *J Bacteriol* 184: 3305–3312.
- Krogh A, Larsson B, von Heijne G, Sonnhammer EL (2001) Predicting transmembrane protein topology with a hidden Markov model: application to complete genomes. *J Mol Biol* 305: 567–580.
- Petersen TN, Brunak S, von Heijne G, Nielsen H (2011) SignalP 4.0: discriminating signal peptides from transmembrane regions. *Nat Methods* 8: 785–786.
- Hassett DJ, Cohen MS (1989) Bacterial adaptation to oxidative stress: implications for pathogenesis and interaction with phagocytic cells. *FASEB J* 3: 2574–2582.

17. Janssen R, van der Straaten T, van Diepen A, van Dissel JT (2003) Responses to reactive oxygen intermediates and virulence of *Salmonella typhimurium*. *Microbes Infect* 5: 527–534.
18. Bandyopadhyay P, Steinman HM (1998) *Legionella pneumophila* catalase-peroxidases: cloning of the *katB* gene and studies of KatB function. *J Bacteriol* 180: 5369–5374.
19. Manca C, Paul S, Barry CE, 3rd, Freedman VH, Kaplan G (1999) *Mycobacterium tuberculosis* catalase and peroxidase activities and resistance to oxidative killing in human monocytes *in vitro*. *Infect Immun* 67: 74–79.
20. Srinivasa Rao PS, Yamada Y, Leung KY (2003) A major catalase (KatB) that is required for resistance to H₂O₂ and phagocyte-mediated killing in *Edwardsiella tarda*. *Microbiology* 149: 2635–2644.
21. Hebrard M, Viala JPM, Meresse S, Barras F, Aussel L (2009) Redundant Hydrogen Peroxide Scavengers Contribute to *Salmonella* Virulence and Oxidative Stress Resistance. *J Bacteriol* 191: 4605–4614.
22. Hirose I, Sano K, Shioda I, Kumano M, Nakamura K, et al. (2000) Proteome analysis of *Bacillus subtilis* extracellular proteins: a two-dimensional protein electrophoretic study. *Microbiology* 146 (Pt 1): 65–75.
23. Sha Z, Stabel TJ, Mayfield JE (1994) *Brucella abortus* catalase is a periplasmic protein lacking a standard signal sequence. *J Bacteriol* 176: 7375–7377.
24. Buchmeier NA, Libby SJ, Xu Y, Loewen PC, Switala J, et al. (1995) DNA repair is more important than catalase for *Salmonella* virulence in mice. *J Clin Invest* 95: 1047–1053.
25. Han Y, Geng J, Qiu Y, Guo Z, Zhou D, et al. (2008) Physiological and regulatory characterization of KatA and KatY in *Yersinia pestis*. *DNA Cell Biol* 27: 453–462.
26. Cosgrove K, Coutts G, Jonsson IM, Tarkowski A, Kokai-Kun JF, et al. (2007) Catalase (KatA) and alkyl hydroperoxide reductase (AhpC) have compensatory roles in peroxide stress resistance and are required for survival, persistence, and nasal colonization in *Staphylococcus aureus*. *J Bacteriol* 189: 1025–1035.
27. Soler-García AA, Jerse AE (2007) *Neisseria gonorrhoeae* catalase is not required for experimental genital tract infection despite the induction of a localized neutrophil response. *Infect Immun* 75: 2225–2233.
28. Vazquez-Torres A, Jones-Carson J, Mastroeni P, Ischiropoulos H, Fang FC (2000) Antimicrobial actions of the NADPH phagocyte oxidase and inducible nitric oxide synthase in experimental salmonellosis. I. Effects on microbial killing by activated peritoneal macrophages *in vitro*. *J Exp Med* 192: 227–236.
29. Demple B, Halbrook J, Linn S (1983) *Escherichia coli xth* mutants are hypersensitive to hydrogen peroxide. *J Bacteriol* 153: 1079–1082.
30. Steele KH, Baumgartner JE, Valderas MW, Roop RM, 2nd (2010) Comparative study of the roles of AhpC and KatE as respiratory antioxidants in *Brucella abortus* 2308. *J Bacteriol* 192: 4912–4922.
31. Tanaka K, Takayanagi Y, Fujita N, Ishihama A, Takahashi H (1993) Heterogeneity of the principal sigma factor in *Escherichia coli*: the *rhoS* gene product, sigma 38, is a second principal sigma factor of RNA polymerase in stationary-phase *Escherichia coli*. *Proc Natl Acad Sci U S A* 90: 8303.
32. Frederick JR, Elkins JG, Bollinger N, Hassett DJ, McDermott TR (2001) Factors affecting catalase expression in *Pseudomonas aeruginosa* biofilms and planktonic cells. *Appl Environ Microbiol* 67: 1375–1379.
33. Navas J, Gonzalez-Zorn B, Ladron N, Garrido P, Vazquez-Boland JA (2001) Identification and mutagenesis by allelic exchange of *choE*, encoding a cholesterol oxidase from the intracellular pathogen *Rhodococcus equi*. *J Bacteriol* 183: 4796–4805.
34. Gentry-Weeks CR, Karkhoff-Schweizer R, Pikis A, Estay M, Keith JM (1999) Survival of *Enterococcus faecalis* in mouse peritoneal macrophages. *Infect Immun* 67: 2160–2165.
35. Verneuil N, Maze A, Sanguinetti M, Laplace JM, Benachour A, et al. (2006) Implication of (Mn)superoxide dismutase of *Enterococcus faecalis* in oxidative stress responses and survival inside macrophages. *Microbiology* 152: 2579–2589.
36. Livak KJ, Schmittgen TD (2001) Analysis of relative gene expression data using real-time quantitative PCR and the 2^{-Delta Delta C(T)} Method. *Methods* 25: 402–408.
37. Dereeper A, Guignon V, Blanc G, Audic S, Buffet S, et al. (2008) Phylogeny.fr: robust phylogenetic analysis for the non-specialist. *Nucleic Acids Research* 36: W465–W469.
38. De La Pena-Moctezuma A, Prescott JF, Goodfellow M (1996) Attempts to find phenotypic markers of the virulence plasmid of *Rhodococcus equi*. *Can J Vet Res* 60: 29–33.