

Review Article

The Importance of Interferon-Tau in the Diagnosis of Pregnancy

Alicja Kowalczyk ¹, Ewa Czerniawska-Piątkowska ², and Marcjanna Wrzecińska ²

¹Department of Environmental Hygiene and Animal Welfare, Wrocław University of Environmental and Life Sciences, Chelmońskiego 38C, Wrocław, Poland

²Department of Ruminant Science, West Pomeranian University of Technology, Ul. Klemensa Janickiego 29, 71-270 Szczecin, Poland

Correspondence should be addressed to Alicja Kowalczyk; alicja.kowalczyk@upwr.edu.pl

Received 14 March 2021; Accepted 18 August 2021; Published 2 September 2021

Academic Editor: Arturo Ortega

Copyright © 2021 Alicja Kowalczyk et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Several decades of improving dairy cattle towards unilateral utilization of dairy cattle led to enormous progress in the field of milk yield; however, it resulted in a number of unfavorable features, such as reproductive disorders, increased calf mortality, and reduced health. Most cases of embryo loss and/or lost pregnancies occur during the first four to five weeks of gestation; accurate detection for pregnancy during this period is likely to contribute to an improvement in gestation rates. A specific protein, interferon-tau (IFNT), stimulates interferon-stimulated genes (ISGs), and their expression increases during gestation within 21 days after insemination. In bovines, the early conceptus undergoes a phase of rapid growth and elongation before implantation, the latter occurring 2–3 weeks after fertilization. IFNT acts mainly in the endometrium of the luminal epithelium. It is a new type I interferon that regulates several genes encoding uterine-derived factors. They are crucial in the processes of preparing the uterus for placenta attachment, modifying the uterine immune system, and regulating early fetal development. Because IFNT is expressed and induces ISGs in the endometrium during pregnancy recognition, it was reasoned that surrogate markers for pregnancy or IFNT might be present in the blood and provide an indicator of pregnancy status in cattle.

1. Introduction

Enormous success in net milk production has been achieved due to the selection of dairy cattle. This selection was aimed at increasing the milk yield of cows [1]. However, cows characterized by high milk yield show reduced fertility. There is a negative correlation between productive and reproductive rates [2, 3]. This is noticeable in high-yield cows that have a lower calving rate (~34%). In these cows, most miscarriages occur within 16 days of insemination [4]. However, early pregnancy loss is not detectable. It can be detected 21 days after insemination when the cow shows signs of estrus. If pregnancy is not detected at an early stage, it results in an extension of the calving interval and consequently reduced milk production. It causes economic losses [5, 6]. Improving reproductive performance by increasing the pregnancy rate and shortening the calving intervals is a consequence of early pregnancy detection in cows after insemination [7].

In the first four to five weeks after fertilization, most cases of embryo loss and/or pregnancy loss are observed. The precise detection of gestation at this time can improve the pregnancy rate [6]. So far, various studies have been conducted for the development of a rapid and straightforward method for the detection of bovine pregnancy [6, 8–13], but most of them, such as rectal palpation, ultrasonography, and monitoring of circulating progesterone, utilize molecular tests to indicate conception success; however, pregnancy is then confirmed by the detection of the fetus approximately 4 to 5 weeks after insemination [14–16].

1.1. Interferon-Tau. A specific protein such as interferon-tau (IFNT) stimulates interferon-stimulated genes (ISGs), and their expression increases during gestation within 21 days after insemination [17–19]. Cows that did not become pregnant after the insemination were identified by the absence of suboptimal detection of embryo-induced ISGs in the maternal circulation on days 18–20. In order to reduce the calving

interval and improve the profitability of the dairy activity, we can use quick resynchronization of the oestrus cycle of cows that did not become pregnant [20].

IFNT signals use classical Type I interferon receptors 1 and 2 (IFNAR1 and IFNAR2) when activating the JAK-STAT signaling pathway [21]. IFNAR1 and IFNAR2 are expressed in the ovine uterus on days 14-15 in nonpregnant and pregnant ewes [22]. IFNT can also signal via mitogen-activated protein kinase (MAPK) and phosphatidylinositol 3-kinase (PI3K) [23]. IFNT acts to rescue the *corpus luteum* (CL) and maintain the pregnancy. However, IFNT is not stimulated by viral infection [24], and transcriptional factors that have control of IFNA and IFNB do not regulate IFNT expression [25]. The expression of IFNT gene is controlled by transcription factor ETS2 [26]. Additionally, IFNT expression is also increased in the elongation phase by combinatory trophoblastic gene regulators DLX3, CDX2, and GATA2/3 [27, 28].

1.2. Maternal Recognition of Pregnancy. Maternal recognition of pregnancy can be defined as the physiological process whereby the conceptus signals its presence to the maternal system and prolongs the lifespan of the ovarian CL [29]. This process in cows requires that the conceptus be elongated to ensure the correct amount of IFNT. Then, pregnancy is diagnosed, and the luteolytic mechanism of the endometrium is inhibited [24, 30–32]. The effect of IFNT on the endometrium is based on the antiluteolytic activity which maintains CL function and progesterone secretion. It is crucial for the growth and development of the fetus during pregnancy. Mononuclear trophoblast cells of ruminant embryos (embryonic and extraembryonic membranes) secrete interferon tau. This type of interferon is characterized by properties such as antiviral, antiproliferative, and immunomodulatory.

In bovines after conception, there is a phase of rapid embryo growth and elongation. Then, 2-3 weeks after fertilization occurs the implantation [33, 34]. Around the 13th day of pregnancy, the embryo lengthens. Then, the embryo changes from spherical to ovoid, then tubular and filamentous [35, 36]. In order to ensure the secretion of suitable concentrations of IFNT, as well as the recognition by the mother of pregnancy, it is necessary to elongate the embryo. The elongation is also aimed at increasing the maximum area of vascular exchange with the mother's tissues after the implantation [37, 38]. The loss of the embryo may be due to the inability to elongate. It may also affect the reproductive failure of cows [38–40].

Maternal recognition of the pregnancy period where bovine conceptus secretes IFNT to signal its presence within the uterus occurs between days 12 and 26 of pregnancy [41]. However, recent researches have shown that the bovine conceptus signals to the mother previously to this period. Bovine oviduct epithelial cells (BOECs) stimulate in vitro bovine embryos on day 4 to produce IFNT, which then acts on immune cells to promote an anti-inflammatory response in the oviduct. However, in this period, the IFNT was not able to stimulate ISG expression in BOECs [42]. Studies report that the endometrial preparation to embryonic recep-

tivity can occur as early as day 4 due to an increase in steroid hormone concentration from large follicles [43]. Moreover, on day 7 after artificial insemination, embryo-dependent factors are already able to modulate ISGs, prostaglandin biosynthesis, and water channel and solute transport pathways in the endometrium at the uterotubal junction of the uterine horn ipsilateral to the CL [44], demonstrating that IFNT secretion may occur before the classic maternal recognition period of pregnancy.

Figure 1 (Figure 1) shows a diagram of the synthesis and changes in INFT levels during early pregnancy in cattle.

1.3. Interferon-Stimulated Genes. There are three main IFN subclasses, namely, types I, II, and III. Type I IFNs contain at least 9 subfamilies, including IFN-alpha (IFNA), beta (IFNB), delta (IFND), omega (IFNW), epsilon (IFNE), kappa (IFNK), tau (IFNT), zeta (IFNZ), and X (IFNX) [47, 48]. The type I family probably has arisen from the type III family due to the fact that mammalian lines diverged from birds and reptiles about 300 million years ago [49, 50]. In the bovine genome, seven of these nine IFN subfamilies (IFNA, IFNB, IFNE, IFNK, IFNW, IFNT, and IFNX) have been identified [48].

IFNT acts mainly in the endometrium of the luminal epithelium. It is a new type I interferon. IFNT regulates several genes encoding uterine-derived factors. They are crucial in the processes of preparing the uterus for placenta attachment, modifying the uterine immune system, and regulating early fetal development [51–53].

The unique trophoblast-specific expression in ruminants is an essential feature that helps to distinguish IFNT from another type I IFNs. From the start of transcription of the IFNT gene, the first 400 bases and upstream are exceptional for the IFNT gene. There are no virus-induced transcriptional elements in this region [54]. However, in the 5' promoter/enhancer region, there are several elements that control trophoblast transcription, including ETS2 and DLX3 [35, 55]. The transcription of IFNT into trophoblast cells may be restricted by the 5' UTR. These cells are the outermost cells of the developing fetus that will form the outermost layer of the placenta. At the morula and blastocyst stage (6-7 days of gestation of the cow), expression in the bovine embryos begins. Expression continues until trophoblast cells adhere to the uterine epithelium (~day 16 in cattle) [56, 57]. In the morula as well as the blastocyst stage, the production of IFNT begins. However, it increases with the elongation of the embryo (days 14–17 in cattle). Table 1 shows the changes in INFT content in cattle in the early stages of pregnancy.

All IFNT paracrine actions block luteolytic PGF pulses, and they also induce expression of ISGs that may have a biological function during maternal recognition and establishment of pregnancy [58].

1.4. The Role of IFNT and ISGs in Pregnancy Diagnosis. Surrogate markers of pregnancy or IFNT can be used as an indicator of the status of pregnancy in cattle because they may be present in the blood. IFNT may be a good marker as it is




FIGURE 1: Diagram of the synthesis and changes of INFT levels during early pregnancy in a cattle after the expression of INFT transcription factor profiles (source: [45, 46]).

TABLE 1: Changes in INFT content in cattle until embryo implantation.

Stage of pregnancy	INFT content	References
Initial blastomere cleavage (1-6 days)	Forms of trophectoderm	
Blastocyst formation (7-8 days)	Forms of trophectoderm	
Embryo spherical form (9-12 days)	Increase in INFT content, INFT expression	
Extending the embryo of the filamentous form (15-18 days)	Big increase in INFT content	[38, 46]
Implantation (21-24 days)	The highest levels of INFT achieved, maternal diagnosis of pregnancy	
>24 days	A sharp decrease in the INFT content, an exponential increase in the trophectoderm layer of a bovine embryo	

expressed and induces ISG in the endometrium during recognition of gestation.

The IFNT analyses did not reveal a sensitivity exceeding the low ng levels from studies based on ISGs. Detecting ISGs as indicators of pregnancy in the blood is not a new practice [59]. Moreover, there are no ISG-based protein biomarkers available to determine pregnancy status from blood. In pregnant sheep 15 to 30 days after insemination, higher levels of MX1 mRNA in PBMC were found compared to nonpregnant sheep [60]. Also in cattle, a similar relationship was observed between pregnancy and ISG in PBMC [61, 62]. In pregnant cows, mean blood levels of ISG15 mRNA are higher from days 15 to 32 od gestation (maximum level is on the 20th day of gestation) compared to fertilized nonpregnant cows. The identification of ISG15 mRNA in PBMC for several days gave a more accurate prediction of gestation than with a single assay. On one hand, studies in cattle showed an increase in ISG in heifer leukocytes in response to pregnancy on day 18. On the other hand, it was not observed in lactating dairy cows [63]. A beef cow study showed an increase in ISG mRNA between days 15 and 22 od gestation. The maximum concentration was reached on the 20th day of pregnancy [64]. This showed that ISG leukocyte mRNA

levels along with CL ultrasound on day 20 of pregnancy are a more precise tool for predicting pregnancy.

In pregnant Holstein-Friesian cows, researchers defined the shortest time to detection of pregnancy. They defined it by measuring ISGs in blood on days 17 and 18 after AI in nulliparous, primiparous, and multiparous cows as well [63]. Primiparous cows had the highest number of ISG on the 20th day. This correlates with maximal production of IFNT by the elongated embryo. ISG-based detection may be more feasible with nulliparous cows as high false-positive rates were obtained during the lactation period. Higher levels of PBMC ISG15 and RTP4 were shown in Holstein primiparous cows than in multiparous cows on day 19 [65]. The differences in ISG expression between heifers or primiparous and multiparous cows may be explained by the length of the embryos. Because, the length of the embryos is longer in heifers than in cows during pregnancy diagnosis, this influences the amount of IFNT produced [66]. Also, another possibility for better peripheral expression of ISG in heifers may be that they are smaller than multiparous cows. This may have an impact on systemic IFNT levels and consequently on the ISG response in leukocytes. An attempt to improve the accuracy of early pregnancy diagnosis has been made by Green et al. They used the ratio of

ISG expression on day 18 post AI to ISG expression based on a control blood sample collected before AI [63].

2. Conclusion

Interferon-tau (IFNT) is the main signal for the maternal recognition of pregnancy in ruminants and exerts its effects by stimulating the expression of interferon-stimulated genes. The process of embryo lengthening significantly affects the production of an appropriate amount of IFNT, and its detection in peripheral fluids located distal from the uterine vein may soon become a reliable and competitive candidate for a biomarker for early pregnancy detection.

Data Availability

Data availability is not applicable.

Conflicts of Interest

The authors declare no conflict of interest.

Authors' Contributions

We declare that all authors made substantial contributions to this manuscript. A.K. was responsible for the conception of the review, preparation of the manuscript, text translation into English, and editorial preparation. E.C.-P. and M.W. were responsible for the collection of the literature and preparation of the manuscript.

References

- [1] P. A. Oltenacu and B. Algers, "Selection for increased production and the welfare of dairy cows: are new breeding goals needed?," *Ambio*, vol. 34, no. 4, pp. 311–315, 2005.
- [2] M. D. Royal, A. P. F. Flint, and J. A. Woolliams, "Genetic and phenotypic relationships among endocrine and traditional fertility traits and production traits in Holstein-Friesian dairy cows," *Journal of Dairy Science*, vol. 85, no. 4, pp. 958–967, 2020.
- [3] P. Dillon, D. P. Berry, R. D. Evans, F. Buckley, and B. Horan, "Consequences of genetic selection for increased milk production in European seasonal pasture based systems of milk production," *Livestock Science*, vol. 99, no. 2-3, pp. 141–158, 2006.
- [4] M. G. Diskin, S. M. Waters, M. H. Parr, and D. A. Kenny, "Pregnancy losses in cattle: potential for improvement," *Reproduction, Fertility, and Development*, vol. 28, no. 2, pp. 83–93, 2016.
- [5] R. Arbel, Y. Bigun, E. Ezra, H. Sturman, and D. Hojman, "The effect of extended calving intervals in high lactating cows on milk production and profitability," *Journal of Dairy Science*, vol. 84, no. 3, pp. 600–608, 2001.
- [6] P. M. Fricke, "Scanning the future—ultrasonography as a reproductive management tool for dairy cattle," *Journal of Dairy Science*, vol. 85, no. 8, pp. 1918–1926, 2002.
- [7] P. M. Fricke, A. Ricci, J. O. Giordano, and P. D. Carvalho, "Methods for and implementation of pregnancy diagnosis in dairy cows," *Veterinary Clinics of North America: Food Animal Practice*, vol. 32, no. 1, pp. 165–180, 2016.
- [8] R. T. Garmo, A. O. Refsdal, K. Karlberg et al., "Pregnancy incidence in Norwegian red cows using nonreturn to estrus, rectal palpation, pregnancy-associated glycoproteins, and progesterone," *Journal of Dairy Science*, vol. 91, no. 8, pp. 3025–3033, 2008.
- [9] J. E. Romano, K. Bryan, R. S. Ramos, J. Velez, and P. Pinedo, "Effect of early pregnancy diagnosis by per rectum amniotic sac palpation on pregnancy loss, calving rates, and abnormalities in newborn dairy calves," *Theriogenology*, vol. 85, no. 3, pp. 419–427, 2016.
- [10] J. J. Schiavo, R. L. Matuszczak, E. B. Oltenacu, and R. H. Foote, "Milk progesterone in postpartum and pregnant cows as a monitor of reproductive status," *Journal of Dairy Science*, vol. 58, no. 11, pp. 1713–1716, 1975.
- [11] C. L. Schanzenbach, B. Kirchner, S. E. Ulbrich, and M. W. Pfaffl, "Can milk cell or skim milk miRNAs be used as biomarkers for early pregnancy detection in cattle?," *PLoS One*, vol. 12, no. 2, article e0172220, 2017.
- [12] J. Ioannidis and F. X. Donadeu, "Circulating miRNA signatures of early pregnancy in cattle," *BMC Genomics*, vol. 17, no. 1, 2016.
- [13] L. M. Mayo, S. G. Moore, S. E. Pooock, W. J. Silvia, and M. C. Lucy, "Technical note. Validation of a chemical pregnancy test in dairy cows that uses whole blood, shortened incubation times, and visual readout," *Journal of Dairy Science*, vol. 99, no. 9, pp. 7634–7641, 2016.
- [14] P. M. Mercadante, E. S. Ribeiro, C. Risco, and A. D. Ealy, "Associations between pregnancy-associated glycoproteins and pregnancy outcomes, milk yield, parity, and clinical diseases in high-producing dairy cows," *Journal of Dairy Science*, vol. 99, no. 4, pp. 3031–3040, 2016.
- [15] G. T. Fosgate, B. Motimele, A. Ganswindt, and P. C. Irons, "A Bayesian latent class model to estimate the accuracy of pregnancy diagnosis by transrectal ultrasonography and laboratory detection of pregnancy-associated glycoproteins in dairy cows," *Preventive Veterinary Medicine*, vol. 145, pp. 100–109, 2017.
- [16] D. P. Bartel, "MicroRNAs: genomics, biogenesis, mechanism, and function," *Cell*, vol. 116, no. 2, pp. 281–297, 2004.
- [17] T. R. Hansen, L. D. P. Sinedino, and T. E. Spencer, "Paracrine and endocrine actions of interferon tau (IFNT)," *Reproduction*, vol. 154, no. 5, pp. F45–F59, 2017.
- [18] J. F. Oliveira, L. E. Henkes, R. L. Ashley et al., "Expression of interferon (IFN)-stimulated genes in extrauterine tissues during early pregnancy in sheep is the consequence of endocrine IFN-tau release from the uterine vein," *Endocrinology*, vol. 149, no. 3, pp. 1252–1259, 2008.
- [19] T. E. Spencer, N. Forde, P. Dorniak, T. R. Hansen, J. J. Romero, and P. Lonergan, "Conceptus-derived prostaglandins regulate gene expression in the endometrium prior to pregnancy recognition in ruminants," *Reproduction*, vol. 146, no. 4, pp. 377–387, 2013.
- [20] A. De Vries, "Economic value of pregnancy in dairy cattle," *Journal of Dairy Science*, vol. 89, no. 10, pp. 3876–3885, 2006.
- [21] M. Binelli, P. Subramaniam, T. Diaz et al., "Bovine Interferon- τ stimulates the Janus kinase-signal transducer and activator of transcription pathway in bovine endometrial epithelial Cells1," *Biology of Reproduction*, vol. 64, no. 2, pp. 654–665, 2001.
- [22] C. S. Rosenfeld, C. S. Han, A. P. Alexenko, T. E. Spencer, and R. M. Roberts, "Expression of interferon receptor subunits,

- IFNAR1 and IFNAR2, in the ovine uterus," *Biol. Reprod.*, vol. 67, no. 3, pp. 847–853, 2002.
- [23] L. C. Platanius, "Mechanisms of type-I- and type-II-interferon-mediated signalling," *Nature Reviews. Immunology*, vol. 5, no. 5, pp. 375–386, 2005.
- [24] R. M. Roberts, Y. Chen, T. Ezashi, and A. M. Walker, "Interferons and the maternal-conceptus dialog in mammals," *Seminars in Cell & Developmental Biology*, vol. 19, no. 2, pp. 170–177, 2008.
- [25] D. W. Leaman, J. C. Cross, and R. M. Roberts, "Multiple regulatory elements are required to direct trophoblast interferon gene expression in choriocarcinoma cells and trophoblast," *Molecular Endocrinology*, vol. 8, no. 4, pp. 456–468, 1994.
- [26] T. Ezashi, A. D. Ealy, M. C. Ostrowski, and R. M. Roberts, "Control of interferon-tau gene expression by Ets-2," *Proceedings of the National Academy of Sciences*, vol. 95, no. 14, pp. 7882–7887, 1998.
- [27] T. Ezashi and K. Imakawa, "Transcriptional control of IFNT expression," *Reproduction*, vol. 154, no. 5, pp. F21–F31, 2017.
- [28] M. S. Kim, K. S. Min, H. H. Seong et al., "Regulation of conceptus interferon-tau gene subtypes expressed in the uterus during the peri-implantation period of cattle," *Animal Reproduction Science*, vol. 190, pp. 39–46, 2018.
- [29] F. W. Bazer, M. A. Mirando, T. L. Ott et al., "Roles of ovine trophoblast protein-1 and oestradiol/prolactin in the establishment of pregnancy in sheep and pigs," *Reproduction, Fertility, and Development*, vol. 4, no. 3, pp. 335–340, 1992.
- [30] T. E. Spencer and F. W. Bazer, "Ovine interferon tau suppresses transcription of the estrogen receptor and oxytocin receptor genes in the ovine endometrium," *Endocrinology*, vol. 137, no. 3, pp. 1144–1147, 1996.
- [31] R. M. Roberts, A. D. Ealy, A. P. Alexenko, C. S. Han, and T. Ezashi, "Trophoblast interferons," *Placenta*, vol. 20, no. 4, pp. 259–264, 1999.
- [32] T. E. Spencer, G. A. Johnson, F. W. Bazer, R. C. Burghardt, and M. Palmirini, "Pregnancy recognition and conceptus implantation in domestic ruminants: roles of progesterone, interferons and endogenous retroviruses," *Reproduction, Fertility and Development*, vol. 19, no. 1, pp. 65–78, 2007.
- [33] J. Wang, M. Guillomot, and I. Hue, "Organisation cellulaire des cellules trophoblastiques dans l'elongation du conceptus ovin," *Comptes Rendus Biologies*, vol. 332, no. 11, pp. 986–997, 2009.
- [34] O. Sandra, G. Charpigny, L. Galio, and I. Hue, "Preattachment embryos of domestic animals: insights into development and paracrine secretions," *Annual Review of Animal Biosciences*, vol. 5, no. 1, pp. 205–228, 2017.
- [35] K. Brooks, G. Burns, and T. E. Spencer, "Conceptus elongation in ruminants: roles of progesterone, prostaglandin, interferon tau and cortisol," *Journal of Animal Science and Biotechnology*, vol. 5, no. 1, p. 53, 2014.
- [36] C. V. Barnwell, P. W. Farin, C. M. Ashwell, W. T. Farmer, S. P. Galphin Jr., and C. E. Farin, "Differences in mRNA populations of short and long bovine conceptuses on day 15 of gestation," *Molecular Reproduction and Development*, vol. 83, no. 5, pp. 424–441, 2016.
- [37] L. Blomberg, K. Hashizume, and C. Viebahn, "Blastocyst elongation, trophoblastic differentiation, and embryonic pattern formation," *Reproduction*, vol. 135, no. 2, pp. 181–195, 2008.
- [38] N. Forde and P. Lonergan, "Interferon-tau and fertility in ruminants," *Reproduction*, vol. 154, no. 5, pp. F33–F43, 2017.
- [39] T. W. Geary, "Management strategies to reduce embryonic loss," in *Range Beef Cow Symposium*, University of Nebraska, Animal Science Department, Rapid City, South Dakota, 2005.
- [40] M. C. Wiltbank, G. M. Baez, A. Garcia-Guerra et al., "Pivotal periods for pregnancy loss during the first trimester of gestation in lactating dairy cows," *Theriogenology*, vol. 86, no. 1, pp. 239–253, 2016.
- [41] C. E. Farin, K. Imakawa, T. R. Hansen et al., "Expression of trophoblastic interferon genes in sheep and cattle," *Biology of Reproduction*, vol. 43, no. 2, pp. 210–218, 1990.
- [42] A. K. Talukder, M. B. Rashid, M. S. Yousef et al., "Oviduct epithelium induces interferon-tau in bovine day-4 embryos, which generates an antiinflammatory response in immune cells," *Scientific Reports*, vol. 8, no. 1, p. 7850, 2018.
- [43] F. S. Mesquita, R. S. Ramos, G. Pugliesi et al., "The receptive endometrial transcriptomic signature indicates an earlier shift from proliferation to metabolism at early diestrus in the cow," *Biology of Reproduction*, vol. 93, pp. 12–51, 2015.
- [44] M. Sponchiado, N. S. Gomes, P. K. Fontes et al., "Pre-hatching embryo-dependent and -independent programming of endometrial function in cattle," *PLOS ONE*, vol. 12, no. 4, article e0175954, 2017.
- [45] K. Imakawa, R. Bai, K. Nakamura, and K. Kusama, "Thirty years of interferon-tau research; past, present and future perspective," *Animal Science Journal*, vol. 88, no. 7, pp. 927–936, 2017.
- [46] A. D. Ealy and Q. E. Yand, "Control of interferon-tau expression during early pregnancy in ruminants," *American Journal of Reproductive Immunology*, vol. 61, no. 2, pp. 95–106, 2009.
- [47] C. D. Krause and S. Pestka, "Evolution of the class 2 cytokines and receptors, and discovery of new friends and relatives," *Pharmacology and Therapeutics*, vol. 106, no. 3, pp. 299–346, 2005.
- [48] A. M. Walker and R. M. Roberts, "Characterization of the bovine type I IFN locus: rearrangements, expansions, and novel subfamilies," *BMC Genomics*, vol. 10, no. 1, p. 187, 2009.
- [49] A. L. Hughes, "The evolution of the type I interferon gene family in mammals," *Journal of Molecular Evolution*, vol. 41, no. 5, pp. 41539–41548, 1995.
- [50] J. P. Levraud, P. Boudinot, I. Colin et al., "Identification of the zebrafish IFN receptor: implications for the origin of the vertebrate IFN system," *Journal of Immunology*, vol. 178, no. 7, pp. 4385–4394, 2007.
- [51] M. S. Kim, K. S. Min, and K. Imakawa, "Regulation of Interferon-stimulated Gene (<i>ISG</i>), <i>ISG15</i>, and <i>MX1</i> and <i>MX2</i> by Conceptus Interferons (IFNTs) in Bovine Uterine Epithelial Cells," *Asian-Australian Journal of Animal Sciences*, vol. 26, no. 6, pp. 795–803, 2013.
- [52] C. L. Gilfeather and C. O. Lemley, "Effects of interferon-tau and steroids on cytochrome P450 activity in bovine endometrial epithelial cells," *Reproduction in Domestic Animals*, vol. 51, no. 3, pp. 415–420, 2016.
- [53] M. M. Meyerholz, K. Mense, H. Knaack, O. Sandra, and M. Schmicke, "Pregnancy-induced ISG-15 and MX-1 gene expression is detected in the liver of holstein-friesian heifers during late peri-implantation period," *Reproduction in Domestic Animals*, vol. 51, no. 1, pp. 175–177, 2016.

- [54] D. W. Leaman, J. C. Cross, and R. M. Roberts, "Genes for the trophoblast interferons and their distribution among mammals," *Reproduction, Fertility, and Development*, vol. 4, no. 3, pp. 349–353, 1992.
- [55] K. Imakawa, M. S. Kim, F. Matsuda-Minehata et al., "Regulation of the ovine interferon-tau gene by a blastocyst-specific transcription factor, Cdx2," *Molecular Reproduction and Development*, vol. 73, no. 5, pp. 559–567, 2006.
- [56] A. D. Ealy, S. F. Larson, L. Liu et al., "Polymorphic forms of expressed bovine interferon-tau genes: relative transcript abundance during early placental development, promoter sequences of genes and biological activity of protein products," *Endocrinology*, vol. 142, no. 7, pp. 2906–2915, 2001.
- [57] H. M. Kubisch, S. Sirisathien, P. Bosch et al., "Effects of developmental stage, embryonic interferon-tau secretion and recipient synchrony on pregnancy rate after transfer of in vitro produced bovine blastocysts," *Reproduction in Domestic Animals*, vol. 39, no. 2, pp. 120–124, 2004.
- [58] T. R. Hansen, K. J. Austin, D. J. Perry, J. K. Pru, M. G. Teixeira, and G. A. Johnson, "Mechanism of action of interferon-tau in the uterus during early pregnancy," *Journal of Reproduction and Fertility: Supplement*, vol. 54, pp. 329–339, 1999.
- [59] T. R. Hansen and B. Rueda, "Identifying early pregnancy in cattle," *Cow Country January 16–17*, 1991, <http://www.drcouncil.org/>.
- [60] S. J. Yankey, B. A. Hicks, K. G. Carnahan et al., "Expression of the antiviral protein Mx in peripheral blood mononuclear cells of pregnant and bred, non-pregnant ewes," *Journal of Endocrinology*, vol. 170, no. 2, pp. R7–R11, 2001.
- [61] H. Han, K. J. Austin, L. A. Rempel, and T. R. Hansen, "Low blood ISG15 mRNA and progesterone levels are predictive of non-pregnant dairy cows," *Journal of Endocrinology*, vol. 191, no. 2, pp. 505–512, 2006.
- [62] P. L. Monteiro Jr., E. S. Ribeiro, R. P. Maciel et al., "Effects of supplemental progesterone after artificial insemination on expression of interferon-stimulated genes and fertility in dairy cows," *Journal of Dairy Science*, vol. 97, pp. 4907–4921, 2014.
- [63] J. C. Green, C. S. Okamura, S. E. Poock, and M. C. Lucy, "Measurement of interferon-tau (IFN-tau) stimulated gene expression in blood leukocytes for pregnancy diagnosis within 18–20d after insemination in dairy cattle," *Animal Reproduction Science*, vol. 121, no. 1-2, pp. 24–33, 2010.
- [64] G. Pugliesi, B. T. Miagawa, Y. N. Paiva, M. R. Franca, L. A. Silva, and M. Binelli, "Conceptus-induced changes in the gene expression of blood immune cells and the ultrasound-accessed luteal function in beef cattle: how early can we detect pregnancy?," *Biology of Reproduction*, vol. 91, p. 95, 2014.
- [65] L. D. Sinedino, P. M. Honda, L. R. Souza et al., "Effects of supplementation with docosahexaenoic acid on reproduction of dairy cows," *Reproduction*, vol. 153, no. 5, pp. 707–723, 2017.
- [66] D. K. Berg, J. van Leeuwen, S. Beaumont, M. Berg, and P. L. Pfeffer, "Embryo loss in cattle between days 7 and 16 of pregnancy," *Theriogenology*, vol. 73, no. 2, pp. 250–260, 2010.